



Universiteit  
Leiden  
The Netherlands

## **Novel formulations and delivery strategies for inactivated polio vaccines : new routes with benefits**

Kraan, H.B.

### **Citation**

Kraan, H. B. (2018, October 18). *Novel formulations and delivery strategies for inactivated polio vaccines : new routes with benefits*. Retrieved from <https://hdl.handle.net/1887/66318>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/66318>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/66318> holds various files of this Leiden University dissertation.

**Author:** Kraan, H.B.

**Title:** Novel formulations and delivery strategies for inactivated polio vaccines : new routes with benefits

**Issue Date:** 2018-10-18

# Development of thermostable lyophilized inactivated polio vaccine

Heleen Kraan <sup>1</sup>, Paul van Herpen <sup>1</sup>, Gideon Kersten <sup>1,2</sup>,  
Jean-Pierre Amorij <sup>1</sup>

<sup>1</sup> Intravacc (Institute for Translational Vaccinology), Bilthoven, The Netherlands

<sup>2</sup> Division of Drug Delivery Technology, Leiden Academic Center for Drug Research, Leiden University, Leiden, The Netherlands

## ABSTRACT



The aim of the current study was to develop a dried inactivated polio vaccine (IPV) formulation with minimal loss during the drying process and improved stability when compared with the conventional liquid IPV.

Extensive excipient screening was combined with the use of a Design of Experiment (DoE) approach in order to achieve optimal results with high probability. Although it was shown earlier that the lyophilization of a trivalent IPV while conserving its antigenicity is challenging, we were able to develop a formulation that showed minimal loss of potency during drying and subsequent storage at higher temperatures.

This study showed the potential of a highly stable and safe lyophilized polio vaccine, which might be used in developing countries without the need of a cold-chain.

## INTRODUCTION

Poliomyelitis is a highly infectious disease, which mainly affects young children. The disease, caused by any one of three serotypes of poliovirus (type 1, type 2 or type 3) has no specific treatment, but can be prevented through vaccination.

Currently, the live attenuated oral poliomyelitis vaccine (Sabin OPV) is the vaccine of choice to prevent polio outbreaks and stop transmission of wild polioviruses, especially in the remaining endemic countries. However, a major concern is the ability of OPV to revert to a form that can cause paralysis, so-called vaccine-associated paralytic poliomyelitis (VAPP). Permanent use of OPV would continue to generate circulating vaccine-derived polioviruses (cVDPVs) that will inevitably lead to new outbreaks [1]. Therefore, the new endgame strategy of the Global Polio Eradication Initiative (GPEI) includes the introduction of an inactivated polio vaccine (IPV) into all routine immunization programs followed by phased withdrawal of OPV [2]. In most of the high-income countries, IPV based on Salk strains is already the present preferred way to eliminate the risk of VAPP and cVDPVs.

To achieve global polio eradication, an (improved) IPV must be efficacious, inexpensive, safe to manufacture, and easy to administer [3]. The feasibility of current IPV in developing countries is limited, because IPV is more expensive than OPV and is administered through injections only. In order to extend the availability of IPV, the World Health Organization (WHO) and the Institute for Translational Vaccinology (Intravacc) in the Netherlands have developed a non-commercial IPV for technology transfer to developing countries [4]. Because the containment of the wild-type Salk poliovirus during production might be an issue, especially in developing countries, the new vaccine will be based on the traditional Sabin OPV strains (sIPV). For reduction in costs, Intravacc is developing sIPV formulations that show dose sparing by using an adjuvant [5] and/or other immunization routes [6, 7].

Vaccine delivery encompasses both administration of the vaccine formulation to specific sites and delivery of the antigen to and activation of relevant cells of the immune system [8]. Since alternative delivery methods and improved vaccine formulations have the potential to make vaccine delivery easier and safer [9, 10], several alternative vaccine delivery methods are currently being developed. However, the focus in vaccine development has been on optimization of the immunological properties, while stability issues are minimally addressed. Most vaccines, IPV included, are insufficiently stable to allow them to be purified, transported and stored at unrefrigerated conditions [11, 12].

One way to improve the storage stability of (s)IPV might be conversion into the dry state as is known to improve the stability of biopharmaceuticals [13]. An increased shelf life is not only of use for the final product for use within three months to two years, but also for stockpiling (1-10 or more years of storage). Particularly after polio eradication, a stockpile of polio vaccines is required to anticipate the potential risk of new polio outbreaks caused by circulating VDPV (even after OPV cessation) [14, 15] or bioterrorism attacks. In order to achieve an optimal vaccine stockpile, various issues need to be considered. The shelf life is paramount, because a delayed expiration time will reduce the stockpile costs [16]. Moreover, storage of dried materials at ambient temperature, including concomitant costs (e.g., reformulation costs), is cost effective compared with storage options at low temperatures. In addition, the ability to develop solid antigen formulations is crucial for new vaccine delivery routes including dermal delivery by coated or dissolving microneedles, parenteral delivery by powders or dissolvable needles, and pulmonary delivery of powders [8]. Technologies for producing dried biologicals include vacuum drying, air-drying, coating, spray (freeze-)drying and foam-drying [17-19]. One of the oldest and commonly used techniques is freeze-drying, also called lyophilization. However, during the lyophilization process the proteinaceous vaccine is subjected to freezing and drying stress by which its activity can be lost. Therefore, cryoprotective and lyoprotective agents are required. Many compounds, such as sugars, polymers, amino acids and surfactants, have been shown to improve the stability of biopharmaceuticals during lyophilization and subsequent storage [20].

Aim of the current study is to design IPV in the dry state with maintenance of the potency. A potency indicating parameter is D-antigenicity, which can be determined in ELISA using specific antibodies as stated in the European Pharmacopeia. Lyophilization of polio vaccines has been shown to be challenging since earlier studies failed to obtain a stable product with preservation of all three serotypes [21-23]. We describe the development of an IPV formulation by selecting excipients that <sup>i</sup> minimize potency loss upon drying and subsequent reconstitution, and <sup>ii</sup> increase the stability of IPV at elevated temperatures. Extensive excipient screening was combined with the use of a Design of Experiments (DoE) approach in order to achieve optimal results with high probability.

## MATERIALS AND METHODS

### Materials

Trivalent IPV, containing the inactivated Mahoney strain for type 1, MEF for type 2 and Saukett for type 3, was obtained from the process development department of Intravacc (Bilthoven, The Netherlands). The ten times concentrated trivalent bulk used in this study was determined at a nominal concentration of 400-80-320 DU/mL by ELISA as described [24].

The excipients sucrose, D-sorbitol, D-trehalose dihydrate, mannitol, L-glutamic monosodium salt monohydrate (MSG), glycine, myo-inositol, magnesiumchloride hexahydrate, lithium chloride and ovalbumin were all purchased from Sigma (St. Louis, MO). Peptone (vegetable) and dextran (6 kDa, from *Leuconostoc ssp*) were from Fluka (Buchs, Switzerland) and sodium chloride was from Merck (Darmstadt, Germany). To prepare 10 mM McIlvaine buffer, 10 mM citric acid (Sigma-Aldrich, St. Louis, MO) was added to 10 mM disodium hydrogen phosphate dehydrate ( $\text{Na}_2\text{HPO}_4$ ) (Fluka, Buchs, Switzerland) in a ratio of 1:6 resulting in a pH-value of 7.0. All excipients used were of reagent quality or higher grade.

### Methods

#### Dialysis

Unless otherwise indicated, the trivalent IPV bulk material was dialyzed against 10 mM McIlvaine buffer (pH 7.0) using a 10 kDa molecular weight cut-off, low-binding regenerated cellulose membrane dialysis cassette (Slide-A-Lyzer®, Pierce, Thermo Scientific, Rockford, IL) to replace the buffer components of the IPV bulk (M199 medium).

#### Formulations

All excipients were dissolved in McIlvaine buffer at a double concentration of the indicated end concentration. The dialyzed IPV was mixed 1:1 by volume with the formulation to be tested. Subsequently, 2 mL neutral glass injection vials (Müller + Müller, Holzminden, Germany) were filled with 0.2 mL of the IPV-excipient mixtures and half-closed with 13 mm pre-dried (overnight at 90°C) rubber stoppers (type V9250 from Helvoet Pharma, Alken, Belgium).

## Drying processes

The filled (0.2 mL/vial) and half-stopped vials were loaded into a pilot freeze dryer (freeze-drying unit sublimator 2-3-3, Zirbus) at a shelf temperature of -50°C, or at a shelf of 4°C and subsequently frozen to -50°C by reducing the shelf temperature at a rate of 1°C/min. These different processes will be denoted as fast and slow freezing, respectively. The vials were kept at a temperature of -50°C for two hours. For the primary drying phase, the shelf temperature was increased at a rate of 0.2°C/h to -45°C (while decreasing the chamber pressure to 0.045 mbar) followed by drying for 3h. The secondary drying phase was performed by further increasing the shelf temperature at 0.02°C/min to 25°C while decreasing the chamber pressure to 0.01 mbar, followed by 24h drying at 25°C. At the end of the cycle, the vials were closed under vacuum, sealed with alu-caps and kept at 4°C until analysis.

In literature, different vacuum drying processes are described [25-27]. The vacuum drying process used in current study was slightly adapted, due to the characteristics of IPV. Briefly, the vials were loaded at shelves of 15°C and kept at that temperature for 10 minutes. The chamber pressure was reduced to 1 mbar in different ramping steps of 15 minutes and starting at a 25 mbar chamber pressure. The temperature was decreased to -10°C for one hour at 0.05 mbar and for one hour at 0.03 mbar so that freezing of the formulations was prevented since product temperature was kept above the ice-nucleation temperature of the formulations. Subsequently, shelf temperature was increased at 0.05°C/min to 30°C. At the end of the cycle, the vials were closed under vacuum, sealed with alu-caps and kept at 4°C until analysis.

## Design of Experiments (DoE)

The Design of Experiments models for D-antigen recoveries measured by ELISA were evaluated in Modde 9.1 software (Umetrics AB, Umea, Sweden) to establish the stability profiles of the lyophilized IPV. In a first pilot experiment, the effect of common used stabilizers, i.e., sucrose (0-20% w/v), trehalose (0-20% w/v), mannitol (0-10% w/v), dextran (0-10% w/v) and NaCl (0-63 mM), was determined using a D-optimal design containing of 22 different formulations and three replicates of the center point ([supplemental data, table S1](#)). For the screening of some excipients, a full factorial DoE was performed around sorbitol, MgCl<sub>2</sub>, monosodium glutamate (MSG) and mannitol, all within the concentration range from 0 to 10% w/v. In this full factorial design 24 formulations were tested and three replicates of the center point ([Table 2](#)). For the optimization experiment a central composite circumscribed



(CCC) design was used around sorbitol (8 to 12% w/v),  $MgCl_2$  (5 to 12% w/v) and MSG (5 to 12% w/v). The CCC design consisted of eight corner experiments, six axial experiments and three replicated center points (Table 3). The models were fitted using partial least squares (PLS) regression and subsequently optimized by deleting non-significant terms leading to a model with the best model performance parameters, i.e., goodness of fit ( $R^2$ ), goodness of prediction ( $Q^2$ ), model validity and reproducibility.

## D-antigen ELISA

A sandwich enzyme-linked immunosorbent assay (ELISA) was used to quantify D-antigen units (DU) of the lyophilized polio vaccine formulations as described by Ten Have *et al.* (2012) [28]. Briefly, microtiter plates were coated with serotype-specific bovine anti-polio serum (Bilthoven Biologicals, Bilthoven, The Netherlands). After washing dilutions of IPV-formulations were added (in duplicate). After a 30 minutes incubation at 37°C under gentle shaking, plates were washed and a mixture of serotype-specific anti-poliovirus monoclonal antibody (mab 3-4-E4 (type 1), 3-14-4 (type 2), 1-12-9 (type 3); all from Intravacc, Bilthoven, The Netherlands) and HRP-labeled anti-mouse IgG (GE Healthcare, Buckinghamshire, UK) was added and incubated for 30 minutes at 37°C while shaking. Subsequently, plates were washed followed by addition of ELISA HighLight signal reagent (ZomerBloemen BV, Zeist, The Netherlands). Chemiluminescence was measured for 10-15 minutes by using a luminometer (Berthold Centro LB960). The signal at maximum intensity was used to calculate D-antigen content relative to the reference standard. Unless otherwise indicated, D-antigen recovery values were shown as normalized values for liquid formulations prior to lyophilization.

## Dynamic Light Scattering (DLS)

Particle size measurements were performed using a Zetasizer Nano-ZS system (Malvern Instruments, Malvern, UK). DLS measurements were done in triplicate with 0.5 mL liquid (undialyzed) IPV bulk at an operating temperature of 25°C. Homogeneity of the size distribution was reflected in the polydispersity index (Pdl), which ranges between 0.0 (fully homogeneous size distribution) and 1.0 (random size distribution).

## Moisture content analysis

The water content was determined using a Karl Fischer coulometric titrimeter (Model CA-06 Moisture meter, Mitsubishi). The samples were weighted, subsequently reconstituted in

the Karl-Fischer reagent, Hydrana Coulomat A (Fluka, Buchs, Switzerland), and injected into the titration vessel. Each vial was measured in triplicate. The empty vials were weighted and the water content was calculated based on the water content measured by the titrimer, the weight of the lyophilized product in the vial, the reconstitution volume of the reagent, titration volume and the water content of the blank titration.

## Differential Scanning Calorimetry (DSC)

The thermodynamic behavior of the formulations was determined by differential scanning calorimetry (DSC). Aluminum DSC pans were filled with the liquid formulations and subjected to a controlled temperature program in a differential scanning calorimeter (DSC Q100, TA Instruments). The samples were cooled to  $-70^{\circ}\text{C}$  at a rate of  $10^{\circ}\text{C}/\text{min}$ , kept isothermal for 2 min, and subsequent heated from  $0^{\circ}\text{C}$  to  $20^{\circ}\text{C}$  at a heating rate of  $10^{\circ}\text{C}/\text{min}$ . The sample chamber was purged with nitrogen gas (50 mL/min). The glass transition temperatures ( $T_g$ ) were determined as the midpoint of the discontinuities in the heat flow curves using thermal analysis software (Universal Analysis 2000, TA Instruments).

# RESULTS & DISCUSSION


## Common lyoprotectants

The first design of experiment approach was based on the most commonly used lyoprotectants, i.e., sucrose, trehalose and mannitol, in combination with dextran, with and without NaCl (Table S1, supplemental data). These excipients are known to provide physical and biochemical stabilization as well as appropriate structural properties to the cake structure, during and after lyophilization [29-32].

A partial least squares (PLS) regression model was fitted and optimized per serotype, which resulted in valid models to predict the D-antigen recoveries directly after lyophilization according to the model performance parameters. For serotype 1, 2 and 3, the  $Q^2$  values were, respectively, 0.650, 0.592 and 0.671, while the  $R^2$  values were 0.905, 0.873 and 0.929. The effects of the different stabilizers, after optimization (excluding non-significant parameters) on the D-antigen recovery after lyophilization are presented in figure 1A.

Without the addition of stabilizers, D-antigen recoveries of type 1, 2 and 3 after lyophilization were 9, 11 and 2%, respectively (Table S1, supplemental data). Both sucrose and mannitol are able to stabilize all serotypes to a certain extent during the lyophilization process (Figure

1A). Dextran has a negative effect on type 1 and 3 during the lyophilization process, whereas the addition of NaCl has no significant effect on the D-antigen recovery, independent of serotype (Figure 1A). Best results, with recoveries of approximately 55%, 85% and 50% for serotype 1, 2 and 3 respectively, were obtained with formulations containing sucrose and/or trehalose in combination with mannitol (Figure 1B). Among the three serotypes, type 2 was the least affected during lyophilization, resulting in a maximum recovery of 85% after lyophilization. This serotype is known to be the most stable at higher temperatures [33].




**Figure 1** Stabilizing potential of the excipients sucrose (SUC), trehalose (TRE), mannitol (MAN), dextran (DEX) and NaCl on D-antigen recovery directly after lyophilization. Main and interaction effects that contribute (per serotype) to the best model, according to the model performance parameters ( $Q^2=0.650$ ,  $R^2=0.905$  (type 1);  $Q^2=0.592$ ,  $R^2=0.873$  (type 2);  $Q^2=0.671$ ,  $R^2=0.929$  (type 3)), are depicted in coefficient plots (A). Surface response plots of the D-antigen recovery for each serotype based on formulations containing sucrose and trehalose in combination with 10% mannitol (without dextran or NaCl) (B).

This first experiment illustrated the complexity of lyophilizing a trivalent polio vaccine retaining its antigenicity and displayed that each serotype may behave different within the same formulations. It has been reported earlier that lyophilization of polio vaccines is challenging. For example, Nagel *et al.* revealed that with the excipients sorbitol and peptone relative potencies of 70% for type 1 and 50% for type 2 could be obtained after lyophilization of IPV, while all activity of type 3 was destroyed [22]. In addition, Pollard *et al.* exhibited that lyophilization of the wild type poliovirus resulted in almost complete inactivation, although it was not determined whether D-antigenicity was also negatively affected [34].

## Drying methods

Literature indicates that vacuum-drying, a drying process without a freezing step, can be used to stabilize IPV [35]. With the use of disaccharides sucrose and trehalose as stabilizing agents, it is feasible to obtain a dried (or highly viscous) polio vaccine by vacuum-drying without affecting its potency. As a result, we decided to investigate the impact of different drying processes on the integrity of IPV. Trehalose and sucrose based IPV formulations were vacuum dried and compared with the same formulations that underwent different lyophilization processes. One lyophilization process starting with a slow freezing step (cooling of shelves from 4°C to -50°C, at 1°C/min) and one with a fast freezing step (shelves pre-cooled to -50°C as has been used in the pilot experiment). The conventional trivalent IPV formulation, without



**Figure 2** D-antigen recovery of dried IPV using different drying methods, i.e., vacuum drying (V) or lyophilization with low (FD-low) or high freezing rate (FD-high). Common used stabilizing sugars sucrose (10% w/v) and trehalose (10% w/v) were compared with the formulation without additives (negative control). Mean values (n=3) and SD are shown.

**Table 1** Formulations tested in an excipient screening experiment (S). D-antigen recoveries were determined directly after lyophilization.

	Sugars	Polyols	Amino acids	Proteins	Other	D-antigen recovery (%)		
						T1	T2	T3
S0	-	-	-	-	-	12	17	2
S1 *	-	7% sorbitol 7% mannitol	2% MSG 2% glycine	7% ovalbumin	-	66	89	73
S2 *	-	7% sorbitol 7% mannitol	2% MSG 2% glycine	7% ovalbumin	10 mM EDTA	66	81	77
S3 *	-	7% sorbitol 7% mannitol	2% MSG 2% glycine	-	-	73	95	74
S4	3% sucrose 3% dextran 3% myo-inositol	-	-	3% ovalbumin	-	27	58	12
S5	5% sucrose	5% sorbitol 5% mannitol	2% glycine 3% lysine 3% arginine	-	-	15	49	5
S6 *	5% sucrose	5% sorbitol 5% mannitol	3% MSG 2% glycine 3% lysine	-	-	65	86	76
S7	5% sucrose 5% trehalose	-	3% lysine 3% alanine	-	-	34	67	36
S8	5% sucrose 5% trehalose	-	3% lysine 3% alanine	-	0.01% Tween80	38	82	43
S9	5% sucrose	-	3% lysine 3% alanine	-	3% Ca- lactobionate	36	36	24
S10	5% sucrose	-	3% lysine 3% alanine	-	3% rec. gelatin	45	72	46
S11 *	-	5% sorbitol	-	5% peptone	2% MgCl <sub>2</sub>	80	79	75
S12 *	-	5% sorbitol	-	5% peptone	1% LiCl	86	100	87
S13	5% sucrose 5% trehalose	-	-	5% peptone	-	29	62	26

\* Formulation selected for stability testing (as shown in figure 3)

any additives, showed recoveries <15% for all serotypes after vacuum drying or lyophilization (Figure 2). The vacuum drying process yielded highly viscous IPV formulations with a water content of respectively 9% or 12% for the formulations containing sucrose or trehalose, whereas lyophilization yielded formulations with a water content of <1%. While more than 55% of the antigen is intact after vacuum drying of IPV containing 10% trehalose, both lyophilization processes resulted in almost complete loss of D-antigenicity. These results depict both one of the main disadvantages of vacuum drying as well as the opportunity to dry without freezing stresses. The high moisture content is caused by the relatively low specific surface area during vacuum drying, which results in an extremely slow secondary drying when compared to lyophilization. As a result, the risk of sugar crystallization and/or phase

separation in the rubbery state exists [13].


The poor recoveries depicted in [figure 2](#) suggest that neither trehalose nor sucrose alone was able to protect IPV against both the freezing and drying stresses during lyophilization. Although the different freezing rates of the lyophilization processes did not show significant differences in D-antigenicity, particle size measurements after freeze thawing revealed differences in IPV particle size that were dependent on freezing rate. Slowly frozen IPV had a size of  $44.7 \pm 2.1$  nm with a Pdl of  $0.485 \pm 0.062$ , whereas fast frozen IPV remains at a particle size of  $38.7 \pm 1.1$  nm with a Pdl of  $0.166 \pm 0.009$  (similar to IPV bulk prior to freeze thawing). Thus, IPV appeared to be most resistant to freeze thawing when a fast freezing rate was applied, which induced less aggregation than slow freezing. For that reason, a fast freezing step was selected for optimization of the IPV formulation for lyophilization.

## Extensive excipient screening

In order to obtain an IPV formulation that is suitable for lyophilization we performed a more extensive excipient screening. The selection of excipients for the screening ([Table 1](#)) was based on findings from literature [20, 22, 36-40] and earlier unpublished data.

In general, the formulations containing sorbitol showed high recoveries directly after lyophilization. Especially formulations containing sorbitol, mannitol and monosodium glutamate (MSG) stabilized the IPV during the process of freezing and subsequent dehydration with D-antigen recoveries of >65% for all serotypes ([Table 1](#); [S1-S3](#), [S6](#)). Another notable formulation is the combination of sorbitol, peptone and the salts LiCl or MgCl<sub>2</sub> ([Table 1](#); [S11 and S12](#)) indicating that this combination of excipients is able to protect the IPV during lyophilization. The best formulations, which showed D-antigen recoveries of >60% directly after lyophilization for all serotypes, were selected and subsequently tested for stability ([Figure 3](#)). In general, the tested formulations so far showed disastrous recoveries after incubation at higher temperatures ([Figure 3B-C](#)) and even after incubation for a month at ambient temperature, a large drop in antigenicity was observed ([Figure 3D](#)).

The appropriate performance of sorbitol combined with mannitol and MSG indicates that the presence of polyols in combination with MSG stabilizes the IPV during lyophilization in a similar way as the disaccharides sucrose and/or trehalose (combined with mannitol) did as shown in the pilot study. In earlier lyophilization studies, sorbitol was used as excipient in combination with peptone [22], which showed again to be a valuable combination in this experiment, whether or not in the presence of a salt like MgCl<sub>2</sub> or LiCl. Peptone-



**Figure 3** D-antigen recoveries of the best formulations from the screening experiment based on recoveries directly after lyophilization (>60% for all serotypes). Panel A shows the D-antigen recoveries directly after lyophilization. Panel B, C and D show the recoveries after incubation for one week at 45°C, two weeks at 37°C, and 4 weeks at room temperature, respectively. The formulations are described in table 1.

containing lyophilized IPV formulations, however, showed limited storage stability at elevated temperatures (Figure 3B-C). Furthermore, peptone is poorly defined and heterogeneous in composition; therefore, it was chosen to exclude peptone further in the formulation design.  $MgCl_2$ 's stabilizing potential has been described for fluid oral polio vaccine. As a result, a number of manufacturers use  $MgCl_2$  to stabilize their OPV (39). Thus,  $MgCl_2$  could also be a critical additive in a dried IPV formulation.


**Table 2** Excipients sorbitol, magnesium chloride ( $MgCl_2$ ), monosodium glutamate (MSG) and mannitol (all 0-10% w/v) examined in a full factorial design (D). Glass transition temperature ( $T_g'$ ) of the liquid formulation before lyophilization and residual moisture content (RMC) of the dried cake were determined.

	Sorbitol	$MgCl_2$	MSG	Mannitol	$T_g'$ (°C)	RMC (%)
D1	-	-	-	-	n.d.	0.2
D2	10%	-	-	-	-43.4	2.4
D3	-	10%	-	-	n.d.	45.9
D4	10%	10%	-	-	n.d.	20.6
D5	-	-	10%	-	-47.0	7.4
D6	10%	-	10%	-	-41.2	1.8
D7	-	10%	10%	-	-58.6	4.0
D8	10%	10%	10%	-	-49.6	7.6
D9	-	-	-	10%	-34.6	1.1
D10	10%	-	-	10%	-40.5	0.3
D11	-	10%	-	10%	n.d.	16.1
D12	10%	10%	-	10%	-51.6	17.1
D13	-	-	10%	10%	-41.0	2.8
D14	10%	-	10%	10%	-39.7	2.3
D15	-	10%	10%	10%	-51.2	8.2
D16	10%	10%	10%	10%	-48.2	9.7
D17	5%	5%	5%	5%	-48.1	5.8
D18	5%	5%	5%	5%	-47.7	17.7
D19	5%	5%	5%	5%	-47.7	9.7





## Design of experiments – full factorial design

Based on the results of our study so far, we selected the most promising excipients to screen them further using a DoE approach and get more insight in the IPV stabilizing potential of these additives. Therefore, a full factorial design was performed around the excipients sorbitol, MSG,  $MgCl_2$  and mannitol (all in the concentration range of 0-10% w/v) (Table 2). The mixture of sorbitol, mannitol and MSG has already shown its capacity to provide valuable protection during the lyophilization process. In addition, mannitol was selected as a common bulking agent due to its excellent cake-forming property and the option to apply annealing when needed [41, 42]. As mentioned above,  $MgCl_2$  is already a proven stabilizer for OPV and therefore we examined its contribution in a dried IPV formulation as well.



**Figure 4** The stabilizing effect of sorbitol, magnesium chloride ( $MgCl_2$ ), monosodium glutamate (MSG) and mannitol was investigated in a screening experiment using a DOE approach. Mean D-antigen recoveries and standard deviations ( $n=3$ ) directly after lyophilization (A) and after incubation for one week at 45°C (B) are shown.

**Figure 4A** shows that sorbitol seemed to be an important excipient for the stabilization of both type 1 and 2 during lyophilization with recoveries up to 89% (Formulation D2 and D8). D-antigen recoveries after one week accelerated stability testing clearly illustrate the benefit of

**A****B**

**Figure 5** Stabilizing potential of the excipients sorbitol (SOR), MgCl<sub>2</sub>, monosodium glutamate (MSG) and mannitol (MAN) on the D-antigen recovery directly after lyophilization. Main and interaction effects that contribute (per serotype) to the best fitted model, according to their model performance parameters ( $Q^2=0.685$ ,  $R^2=0.923$  (type 1);  $Q^2=0.577$ ,  $R^2=0.877$  (type 2);  $Q^2=0.575$ ,  $R^2=0.824$  (type 3)) are shown in coefficient plots (A). Surface response plots of the D-antigen recovery for each serotype based on formulations containing MSG and MgCl<sub>2</sub> in combination with 10% sorbitol (without mannitol) (B).

sorbitol, as well as MgCl<sub>2</sub> and MSG in the formulation (Figure 4A, formulation D8). Regression models ( $R^2>0.65$  and  $Q^2>0.5$ ) of these data confirm the findings that are described above. The stabilizing effects of the excipients on D-antigen recovery directly after lyophilization are depicted per serotype in coefficient plots after model optimization (Figure 5A). For all serotypes, there is a main effect of sorbitol, whereas MSG showed to be significant beneficial for the D-antigen recovery of type 1 and 3. However, sorbitol showed an interaction with mannitol that negatively affected IPV recovery after lyophilization, implying that the addition of mannitol to the IPV formulation for lyophilization is not desired. Accelerated stability


testing revealed that  $MgCl_2$  and MSG are important stabilizers for all serotypes, indicated by significant main effects and interaction between these excipients. This means that the addition of  $MgCl_2$  or MSG (10% w/v) results in an increase of 7-13% in D-antigen recovery, while the combination of these excipients would boost the D-antigenicity with 6-10%. The inclusion of  $MgCl_2$  to the IPV formulation increases the residual water content (Table 2) after lyophilization significantly. This excipient is also responsible for reducing the glass transition temperatures ( $T_g'$ ) of the formulation before drying. The sorbitol and MSG interaction factor is able to increase the  $T_g'$  significantly (data not shown). As such, formulations might be optimized for  $T_g'$  by increasing the sorbitol-MSG content in order to be capable to lyophilize at higher temperatures resulting in shorter process time.

Again, it was observed that there are some divergences between the serotypes regarding their preference for stabilizers during lyophilization, whereas all serotypes seemed to prefer the

**Table 3** Sorbitol (8-12% w/v), magnesium chloride ( $MgCl_2$ : 5-12% w/v) and monosodium glutamate (MSG: 5-12% w/v) tested in a central composite circumscribed design. D-antigen recoveries were determined per serotype directly after lyophilization and after accelerated stability testing (one week at 45°C).

	Sorbitol	$MgCl_2$	MSG	D-antigen recovery (%) Directly after lyophilization			D-antigen recovery (%) Stability 45°C		
				T1	T2	T3	T1	T2	T3
O1	8%	5%	5%	98.6	96.5	85.7	91.9	97.0	55.2
O2	12%	5%	5%	88.9	88.7	65.2	87.6	91.6	73.7
O3	8%	12%	5%	76.0	82.4	33.6	62.5	49.2	12.2
O4	12%	12%	5%	89.4	91.7	59.1	95.8	90.8	48.2
O5	8%	5%	12%	90.6	81.9	70.7	86.0	81.4	57.5
O6	12%	5%	12%	79.3	82.5	62.1	80.2	77.0	39.4
O7	8%	12%	12%	84.9	86.7	64.6	92.0	84.6	63.5
O8	12%	12%	12%	87.9	82.5	73.0	90.3	86.7	69.8
O9	6.6%	8.5%	8.5%	90.0	99.3	84.8	93.3	90.6	73.0
O10	13.4%	8.5%	8.5%	86.2	85.3	67.4	86.1	85.3	69.3
O11	10%	2.6%	8.5%	77.5	72.4	47.4	74.4	60.7	29.9
O12	10%	14.4%	8.5%	89.1	84.6	63.6	91.7	72.7	40.0
O13	10%	8.5%	2.6%	98.1	85.2	49.2	93.0	83.3	42.2
O14	10%	8.5%	14.4%	84.7	70.3	66.3	74.8	81.2	35.5
O15	10%	8.5%	8.5%	86.6	88.4	68.2	90.1	91.1	67.3
O16	10%	8.5%	8.5%	92.2	90.8	75.1	93.5	89.1	69.2
O17	10%	8.5%	8.5%	96.0	98.3	76.7	91.7	94.0	57.0

presence of MSG and  $\text{MgCl}_2$  during stability testing. The surface response plots demonstrate that for type 1, the highest DU recoveries after lyophilization were found in a formulation containing 10% sorbitol in the presence of the highest amounts of MSG, regardless the  $\text{MgCl}_2$  concentration, whereas both additives have no significant effect on type 2 (Figure 5B). Though, type 3 showed to be most delicate for small differences in excipient concentrations with maximal D-antigen recoveries of more than 80% with a formulation containing 10% sorbitol, 5-10%  $\text{MgCl}_2$  and 4-8% MSG.




**Figure 6** Contour plots show the effect of MSG (5-12% w/v) and  $\text{MgCl}_2$  (5-12% w/v) in combination with 8% (A), 10% (B) or 12% w/v sorbitol (C) on the D-antigen recovery after lyophilization and subsequent accelerated stability testing for serotype 3.

## Optimization

With the purpose to optimize the formulation, a response surface methodology design was implemented. Hence, a central composite circumscribed (CCC) [43] study set-up with the factors sorbitol (8-12% w/v), MSG (8-12% w/v) and  $\text{MgCl}_2$  (5-10% w/v) was designed (Table 3) and D-antigen recovery was determined directly after lyophilization. An extensive stability study was included here as well to test whether the dried IPV formulation has an improved stability when compared with the conventional liquid IPV.

Within the design of the study, all formulations resulted in comparable recoveries after lyophilization and subsequent storage (Table 3). This indicates the formulation robustness of the formulation based on sorbitol (8-12% w/v), MSG (8-12% w/v) and  $\text{MgCl}_2$  (5-10% w/v). Due to these small differences in D-antigen recovery after lyophilization and stability assessment between the tested formulations, it was not possible to obtain a valid model to describe the data. With all formulations, satisfying stabilization was achieved for serotypes 1 and 2, which showed recoveries of between 75% and 100% directly after lyophilization and only a small loss (0% to maximal 15%) during accelerated stability testing (Table 3). As mentioned above, type 3 showed to be the determining aspect in the decision for the final concentrations of the three excipients in our IPV formulation. Therefore, we decided to focus on the stability data of this serotype. PLS regression describes the accelerated stability data (one-week incubation at 45°C) well with high values for model validity and reproducibility (respectively 0.67 and 0.86). However, the predicting power of the model is limited ( $Q^2 = 0.43$  instead of  $Q^2 > 0.5$ ). Despite this limitation, we requested the contour plots (Figure 6) to get an indication of the important parameters to stabilize serotype 3. It seems that the best D-antigen recoveries (>70%) were obtained with the lowest sorbitol (8% w/v) concentration in combination with the highest concentrations of both  $\text{MgCl}_2$  and MSG (>10% w/v) or the highest sorbitol (12% w/v) content combined with relatively low amounts of  $\text{MgCl}_2$  and MSG (<8% w/v) (Figure 6).

This experiment showed the robustness of the formulation since many formulations containing sorbitol,  $\text{MgCl}_2$  and MSG in the tested concentrations showed high recoveries directly after lyophilization and preserved antigenicity during accelerated stability. Based on excellent D-antigen recoveries after lyophilization and subsequent stability testing (Table 3), the formulation containing 10% (w/v) sorbitol, 8.5% (w/v)  $\text{MgCl}_2$  and 8.5% (w/v) MSG was selected for additional extensive stability testing up to 24 weeks at 25°C and 37°C, and up to one month at 45°C. Both the conventional liquid IPV and lyophilized IPV formulation remain



**Figure 7** Stability testing of a lyophilized trivalent IPV formulation containing 10% (w/v) sorbitol, 8.5% (w/v)  $MgCl_2$  and 8.5% (w/v) MSG at 25°C (A), 37°C (B) or up to 4 weeks at 45°C (C). The freeze-dried formulation (FD IPV, pink lines) was compared with the conventional liquid IPV (L IPV, blue lines) and a liquid IPV dialyzed against McIlvaine buffer (L dIPV, grey lines), which is the buffer of choice during lyophilization.

stable during long-term incubation at ambient temperature (Figure 7A). However, at elevated temperatures, the lyophilized IPV formulation revealed its improved stability profile. Despite its relatively high residual moisture content, which was determined at  $4.5 \pm 0.9\%$ , minimal loss was observed for the dried IPV after storage at temperatures above room temperature where the liquid IPV has lost its antigenicity completely (Figure 7B and 7C).

The current study shows the feasibility to convert IPV into the dry state using lyophilization. The focus here was on development and optimization of a dried IPV formulation. However,

the lyophilization process needs optimization as well, since the RMC of our lyophilized IPV formulation exceeds the limit of 3% water content from the European Pharmacopoeia. This high residual water content is probably due to the presence of  $MgCl_2$ , a hexahydrate with strongly bound water, in the IPV formulation. The duration of the secondary drying step dictates the residual moisture level in a lyophilized product [20], so probably prolongation of this drying step and raising the end temperature of the lyophilization process could decrease the RMC and thus could possibly improve the final product.

## CONCLUSION

The aim of the study was to develop a dried IPV formulation with minimal loss during the drying process and improved stability when compared with the conventional liquid IPV, which could allow distribution and storage under unrefrigerated conditions. Extensive screening of a large number of excipients combined with a DoE approach yielded a lyophilized IPV formulation with remaining antigenicity for all serotypes when kept at ambient or even higher temperatures.

Although further improvement and research is still possible, this study showed the potential of a highly stable and safe lyophilized polio vaccine, which could be distributed in developing countries without the need of a cold-chain transport.

## ACKNOWLEDGMENTS

The authors thank Cor van Ingen and Hoang Hirschberg for scientific advice during the screening phase of the study and Antoinette van den Dikkenberg and Kayan Tsoi for technical assistance.

## REFERENCES

- Davis R, Wright PF. Circulating vaccine derived poliovirus and the polio eradication endgame. *The Pan African medical journal*. 2012;12:109. Epub 2012/11/08.
- Polio Eradication & Endgame Strategic Plan 2013-2018. [http://www.polioeradication.org/Portals/0/Document/Resources/StrategyWork/PEESP\\_EN\\_A4.pdf](http://www.polioeradication.org/Portals/0/Document/Resources/StrategyWork/PEESP_EN_A4.pdf): Global Polio Eradication Initiative; 2013 [cited 2013 1 August].
- Ehrenfeld E, Modlin J, Chumakov K. Future of polio vaccines. *Expert Rev Vaccines*. 2009;8(7):899-905.
- Bakker WA, Thomassen YE, van't Oever AG, Westdijk J, van Oijen MG, Sundermann LC, *et al*. Inactivated polio vaccine development for technology transfer using attenuated Sabin poliovirus strains to shift from Salk-IPV to Sabin-IPV. *Vaccine*. 2011;29(41):7188-96. Epub 2011/06/10.
- Westdijk J, Koedam P, Barro M, Steil BP, Collin N, Vedvick TS, *et al*. Antigen sparing with adjuvanted inactivated polio vaccine based on Sabin strains. *Vaccine*. 2013;31(9):1298-304. Epub 2013/01/15.
- Soonawala D, Verdijk P, Wijmenga-Monsuur AJ, Boog CJ, Koedam P, Visser LG, *et al*. Intradermal fractional booster dose of inactivated poliomyelitis vaccine with a jet injector in healthy adults. *Vaccine*. 2013. Epub 2013/06/19.
- Verdijk P, Rots NY, Bakker WA. Clinical development of a novel inactivated poliomyelitis vaccine based on attenuated Sabin poliovirus strains. *Expert Rev Vaccines*. 2011;10(5):635-44. Epub 2011/05/25.
- Amorij JP, Kersten GF, Saluja V, Tonnis WF, Hinrichs WL, Slutter B, *et al*. Towards tailored vaccine delivery: Needs, challenges and perspectives. *J Control Release*. 2012. Epub 2012/01/17.
- Clements CJ, Larsen G, Jodar L. Technologies that make administration of vaccines safer. *Vaccine*. 2004;22(15-16):2054-8.
- Friede M, Aguado MT. Need for new vaccine formulations and potential of particulate antigen and DNA delivery systems. *Adv Drug Deliv Rev*. 2005;57(3):325-31.
- Brandau DT, Jones LS, Wiethoff CM, Rexroad J, Middaugh CR. Thermal stability of vaccines. *J Pharm Sci*. 2003;92(2):218-31. Epub 2003/01/18.
- Chen D, Kristensen D. Opportunities and challenges of developing thermostable vaccines. *Expert Rev Vaccines*. 2009;8(5):547-57. Epub 2009/04/29.
- Amorij JP, Huckriede A, Wilschut J, Frijlink HW, Hinrichs WL. Development of stable influenza vaccine powder formulations: challenges and possibilities. *Pharm Res*. 2008;25(6):1256-73. Epub 2008/03/14.
- Bruce Aylward R, Sutter RW, Cochi SL, Thompson KM, Jafari H, Heymann D. Risk management in a polio-free world. *Risk Anal*. 2006;26(6):1441-8. Epub 2006/12/23.
- Thompson KM, Duintjer Tebbens RJ. The case for cooperation in managing and maintaining the end of poliomyelitis: stockpile needs and coordinated OPV cessation. *Medscape J Med*. 2008;10(8):190. Epub 2008/10/18.
- Tebbens RJ, Pallansch MA, Alexander JP, Thompson KM. Optimal vaccine stockpile design for an eradicated disease: application to polio. *Vaccine*. 2010;28(26):4312-27. Epub 2010/05/01.
- Jangle RD, Pisal SS. Vacuum foam drying: an alternative to lyophilization for biomolecule preservation. *Indian journal of pharmaceutical sciences*. 2012;74(2):91-100. Epub 2013/01/18.
- Kim YC, Quan FS, Compans RW, Kang SM, Prausnitz MR. Stability kinetics of influenza vaccine coated onto microneedles during drying and storage. *Pharmaceutical research*. 2011;28(1):135-44. Epub 2010/04/14.
- Maa YF, Zhao L, Payne LG, Chen D. Stabilization of alum-adjuvanted vaccine dry powder formulations: mechanism and application. *J Pharm Sci*. 2003;92(2):319-32. Epub 2003/01/18.
- Wang W. Lyophilization and development of solid protein pharmaceuticals. *Int J of Pharm*. 2000;203:1-60.
- Kraft LM, Pollard EC. Lyophilization of poliomyelitis virus; heat inactivation of dry MEFI virus. *Proc Soc Exp Biol Med*. 1954;86(2):306-9. Epub 1954/06/01.
- Nagel J, Hekker AC, Hofman B, Cohen H. Some experiments on freeze-drying of inactivated poliomyelitis-vaccines. *Arch Gesamte Virusforsch*. 1963;12:718-20. Epub 1963/01/01.
- Portocala R, Samuel I, Popescu M. Effect of lyophilization on picornaviruses. (Brief report). *Arch Gesamte Virusforsch*. 1969;28(1):97-9. Epub 1969/01/01.
- Westdijk J, Brugmans D, Martin J, van't Oever A, Bakker WA, Levels L, *et al*. Characterization and standardization of Sabin based inactivated polio vaccine: proposal for a new antigen unit for inactivated polio vaccines. *Vaccine*. 2011;29(18):3390-7. Epub 2011/03/15.
- Bieganski RM, Fowler A, Morgan JR, Toner M. Stabilization of active recombinant retroviruses in an amorphous dry state with trehalose. *Biotechnol Prog*. 1998;14(4):615-20. Epub 1998/08/08.
- Mattern M, Winter G, Kohert U, Lee G. Formulation of proteins in vacuum-dried glasses. II. Process and storage stability in sugar-free amino acid systems. *Pharm Dev Technol*. 1999;4(2):199-208. Epub 1999/05/08.
- Rossi S, Buera MP, Moreno S, Chirife J. Stabilization of the restriction enzyme EcoRI dried with trehalose and other selected glass-forming solutes. *Biotechnol Prog*. 1997;13(5):609-16. Epub 1997/10/23.
- Ten Have R, Thomassen YE, Hamzink MR, Bakker WA, Nijst OE, Kersten G, *et al*. Development of a fast ELISA for quantifying polio D-antigen in in-process samples. *Biologicals*. 2012;40(1):84-7. Epub 2011/12/14.
- Johnson RE, Kirchhoff CF, Gaud HT. Mannitol-sucrose mixtures--versatile formulations for protein lyophilization. *J Pharm Sci*. 2002;91(4):914-22. Epub 2002/04/12.
- Maa YF, Ameri M, Shu C, Payne LG, Chen D. Influenza vaccine powder formulation development: spray-freeze-drying and stability evaluation. *J Pharm Sci*. 2004;93(7):1912-23. Epub 2004/06/04.
- Maa YF, Ameri M, Shu C, Zuleger CL, Che J, Osorio JE, *et al*. Hepatitis-B surface antigen



- (HBsAg) powder formulation: process and stability assessment. *Current drug delivery*. 2007;4(1):57-67. Epub 2007/02/03.
32. Zhai S, Hansen RK, Taylor R, Skepper JN, Sanches R, Slater NK. Effect of freezing rates and excipients on the infectivity of a live viral vaccine during lyophilization. *Biotechnol Prog*. 2004;20(4):1113-20. Epub 2004/08/07.
33. Moynihan M, Petersen I. The durability of inactivated poliovirus vaccine: studies on the stability of potency *in vivo* and *in vitro*. *J Biol Stand*. 1982;10(3):261-8. Epub 1982/07/01.
34. Pollard M. The inactivation of poliomyelitis virus by freeze-drying. *Texas reports on biology and medicine*. 1951;9(4):749-54. Epub 1951/01/01.
35. Mayeresse Y, Stephenne J, inventors; GlaxoSmithKline Biologicals, assignee. Immunogenic composition 2004 13 May 2004.
36. Akers MJ, Milton N, Byrn SR, Nail SL. Glycine crystallization during freezing: the effects of salt form, pH, and ionic strength. *Pharmaceutical research*. 1995;12(10):1457-61. Epub 1995/10/01.
37. Berge TO, Jewett RL, Blair WO. Preservation of enteroviruses by freeze-drying. *Applied microbiology*. 1971;22(5):850-3. Epub 1971/11/01.
38. Leal ML, Lopes FJ, Carvalho MH, Moura HA, Soares SA. Study on thermostabilizers for trivalent oral poliomyelitis vaccine. *Mem Inst Oswaldo Cruz*. 1990;85(3):329-38. Epub 1990/07/01.
39. Melnick JL. Thermostability of poliovirus and measles vaccines. *Dev Biol Stand*. 1996;87:155-60. Epub 1996/01/01.
40. Ohtake S, Martin RA, Yee L, Chen D, Kristensen DD, Lechuga-Ballesteros D, *et al*. Heat-stable measles vaccine produced by spray drying. *Vaccine*. 2010;28(5):1275-84. Epub 2009/12/01.
41. Cao W, Xie Y, Krishnan S, Lin H, Ricci M. Influence of process conditions on the crystallization and transition of metastable mannitol forms in protein formulations during lyophilization. *Pharmaceutical research*. 2013;30(1):131-9. Epub 2012/08/22.
42. Mehta M, Bhardwaj SP, Suryanarayanan R. Controlling the physical form of mannitol in freeze-dried systems. *European journal of pharmaceuticals and biopharmaceutics : official journal of Arbeitsgemeinschaft fur Pharmazeutische Verfahrenstechnik eV*. 2013. Epub 2013/05/07.
43. Eriksson L, Johansson E, Kettaneh-Wold N, Wikström C, Wold S. *Design of Experiments; Principles and Applications*: Umetrics AB; 2008.

## SUPPLEMENTAL DATA

**Table S1** Commonly used excipients tested using a design of experiments approach. D-antigen recoveries were determined per serotype by ELISA directly after lyophilization.

	Sucrose	Trehalose	Mannitol	Dextran	NaCl (mM)	D-antigen recovery (%)		
						T1	T2	T3
P1	-	-	-	-	-	8.5	11.3	1.8
P2	20%	20%	-	-	-	30.3	58.0	22.5
P3	20%	-	10%	-	-	45.3	68.3	46.4
P4	-	20%	10%	-	-	44.4	68.5	40.5
P5 *	11.4%	11.4%	5.7%	5.7%	-	30.5	51.1	29.9
P6	-	-	6.7%	10%	-	25.9	54.1	8.9
P7	-	6.7%	10%	10%	-	31.6	62.2	16.3
P8 *	13.3%	13.3%	2.2%	6.7%	-	33.9	57.8	25.9
P9	6.7%	-	10%	10%	-	34.9	58.1	27.5
P10	10%	10%	-	5%	-	22.2	45.0	8.2
P11	10%	10%	5%	-	-	35.5	62.7	34.9
P12	20%	-	-	-	63	32.8	59.3	33.3
P13	-	20%	-	-	63	21.7	47.6	7.1
P14	-	-	10%	-	63	39.8	57.9	44.2
P15	-	-	-	10%	63	8.2	15.1	4.0
P16	10%	-	5%	5%	31.5	31.9	56.7	31.5
P17	10%	10%	5%	5%	63	29.2	49.6	25.2
P18	16%	16%	8%	-	50	42.3	64.6	42.0
P19 *	13.3%	13.3%	-	6.7%	41.7	22.7	43.7	9.9
P20 *	11.4%	-	5.7%	5.7%	35.7	33.1	56.3	32.9
P21	-	16%	8%	8%	50	25.9	49.1	16.2
P22 *	13.3%	6.7%	3.3%	3.3%	41.7	29.2	52.2	27.2
P23	10%	10%	5%	5%	31.5	29.2	49.6	25.2
P24	10%	10%	5%	5%	31.5	30.3	52.8	27.2
P25	10%	10%	5%	5%	31.5	34.0	58.1	28.1

\* Concentrations differ from design due to maximal solubility of the formulation.