

# Recovery of arm-hand function after stroke: developing neuromechanical biomarkers to optimize rehabilitation strategies.

Krogt, J.M. van der

### Citation

Krogt, J. M. van der. (2018, November 6). Recovery of arm-hand function after stroke: developing neuromechanical biomarkers to optimize rehabilitation strategies. Retrieved from https://hdl.handle.net/1887/66816

Version: Not Applicable (or Unknown)

License: License agreement concerning inclusion of doctoral thesis in the

Institutional Repository of the University of Leiden

Downloaded from: <a href="https://hdl.handle.net/1887/66816">https://hdl.handle.net/1887/66816</a>

Note: To cite this publication please use the final published version (if applicable).

## Cover Page



# Universiteit Leiden



The handle <a href="http://hdl.handle.net/1887/66816">http://hdl.handle.net/1887/66816</a> holds various files of this Leiden University dissertation.

Author: Krogt, J.M. van der

Title: Recovery of arm-hand function after stroke: developing neuromechanical

biomarkers to optimize rehabilitation strategies

**Issue Date**: 2018-11-06

Stellingen behorende bij het proefschrift Recovery of arm-hand function after stroke: developing neuromechanical biomarkers to optimize rehabilitation strategies.

#### Hanneke van der Krogt

(Propositions)

- 1. A clear description of pathophysiological concepts supports discussion about and facilitates follow up of movement disorder after stroke. (this thesis)
- 2. Objective and reproducible measurement of body functions in movement disorder after stroke requires biomechanical, electromyographical and system identification techniques. (this thesis)
- 3. To identify the phenotype of a patient in a spectrum of neural and non-neural contributors to movement disorder, multiple measurement and task conditions should be used. (this thesis)
- 4. Invasive, costly and/or labor intensive treatment should not be prescribed without proper information on (loss of) selective muscle activation. (this thesis)
- 5. The connection between pathophysiology and treatment goals in the field of rehabilitation is supported by the framework of the International Classification of Functioning, Disability and Health.
- 6. Goal setting in interdisciplinary rehabilitation should be formulated in a SMART way: specific, measurable, acceptable, realistic and time-based.
- 7. Investment in research and development and realization of rapid implementation will greatly enhance the effectiveness and efficiency of rehabilitation medicine. (position paper VRA, 2015)
- 8. Safe application of medical technology means a safe product, in the hands of a trained user in an environment that can guarantee safe use. (Convenant Veilige Toepassing van Medische Technologie in de medisch specialistische zorg. Utrecht 2016)
- 9. If the wires of the surface EMG electrodes can get entangled, they will get entangled. (Free adaptation from Murphy's law)
- 10. It is a capital mistake to theorize before one has data. Insensibly one begins to twist facts to suit theories, instead of theories to suit facts. (Sherlock Holmes A Scandal in Bohemia. A. Conan Doyle, The adventures of Sherlock Holmes; George Newnes Ltd, London. 1892)
- 11. More harm is done because you do not look / Than from not knowing what is in the book. (Zachary Cope, The Diagnosis of the Acute Abdomen in Rhyme, H. K. Lewis, London. 1947)
- 12. Experience is what you obtain just after you need it. (Dr Baur, intensivist, MC Haaglanden circa 2006)