



Universiteit
Leiden
The Netherlands

Inferno Worlds

Ridden - Harper, A.

Citation

Ridden - Harper, A. (2018, November 21). *Inferno Worlds*. Retrieved from <https://hdl.handle.net/1887/67080>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/67080>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/67080> holds various files of this Leiden University dissertation.

Author: Ridden, - Harper A.

Title: Inferno Worlds

Issue Date: 2018-11-21

Inferno Worlds

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op woensdag 21 november 2018
klokke 10:00 uur

door

Andrew Ridden-Harper
geboren te Christchurch, Nieuw-Zeeland
in 1991

Promotiecommissie

Promotores: Prof. dr. I. A. G. Snellen
Prof. dr. C. U. Keller

Overige leden: Prof. dr. H. J. A. Röttgering
Prof. dr. H. V. J. Linnartz
Prof. dr. M. Fridlund
Dr. D. M. Stam T. U. Delft
Dr. M. Min SRON

Cover design: An artist's impression of a hot rocky exoplanet that has a dust-tail. Cover designed by Andrew Ridden-Harper, using a false colour photograph of the Sun in ultra-violet wavelengths (credit: NASA Solar Dynamics Observatory) to represent the host star. The image of the planet was adapted from an artist's impression of 55 Cancri e (credit: ESA/Hubble, M. Kornmesser).

ISBN: 978-94-6323-396-5

© 2018 Andrew Ridden-Harper

*To my parents for their unwavering
support and encouragement.*

*“Imagination will often carry us to worlds that
never were, but without it we go nowhere.”
– Carl Sagan*

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 1 |
| 1.1 | Summary | 1 |
| 1.2 | Overview of the field | 1 |
| 1.3 | Planet formation | 3 |
| 1.4 | Hot rocky exoplanets | 5 |
| 1.4.1 | Sputtering | 6 |
| 1.4.2 | Possible mineral vapour atmospheres | 6 |
| 1.5 | Characterisation with spectroscopy | 7 |
| 1.6 | Disintegrating rocky exoplanets | 8 |
| 1.6.1 | Mass-loss mechanism | 10 |
| 1.6.2 | Dust particle dynamics | 11 |
| 1.7 | This thesis | 13 |
| 1.7.1 | Chapter 2 | 13 |
| 1.7.2 | Chapter 3 | 14 |
| 1.7.3 | Chapter 4 | 14 |
| 1.7.4 | Chapter 5 | 14 |
| 1.8 | Future outlook | 15 |
| 2 | Search for an exosphere in sodium and calcium in the transmission spectrum of exoplanet 55 Cancri e | 19 |
| 2.1 | Introduction | 21 |
| 2.2 | Observational data | 24 |
| 2.2.1 | UVES data | 25 |
| 2.2.2 | HARPS data | 25 |
| 2.2.3 | HARPS-N data | 26 |
| 2.3 | Data analysis | 26 |
| 2.3.1 | Processing of UVES spectra | 26 |
| 2.3.2 | Processing of HARPS and HARPS-N data | 31 |
| 2.3.3 | Combining the different data sets | 31 |
| 2.3.4 | Injection of artificial planet signals | 32 |

| | | |
|----------|--|-----------|
| 2.4 | Results | 33 |
| 2.4.1 | Sodium | 33 |
| 2.4.2 | Ionized calcium | 35 |
| 2.5 | Discussion and conclusions | 36 |
| 3 | Chromatic transit light curves of disintegrating rocky planets | 47 |
| 3.1 | Introduction | 48 |
| 3.2 | Method: The model | 51 |
| 3.2.1 | Dust dynamics code | 51 |
| 3.2.2 | Particle dynamics simulations | 54 |
| 3.2.3 | Ray tracing with MCMa3D | 55 |
| 3.3 | Results of simulations | 58 |
| 3.3.1 | Modelling the light curve of Kepler-1520 b with a low planet mass | 58 |
| 3.3.2 | Optically thick tail | 62 |
| 3.3.3 | Modelling the light curve of Kepler-1520 b with a planet mass of $0.02 M_{\oplus}$ | 69 |
| 3.3.4 | Modelling the light curve of Kepler-1520 b with a planet mass of $0.02 M_{\oplus}$ and larger maximum height | 70 |
| 3.3.5 | Behaviour of large particles | 75 |
| 3.4 | Wavelength dependence | 79 |
| 3.5 | Constraints on particle ejection velocity | 82 |
| 3.5.1 | Particle trajectories | 84 |
| 3.5.2 | Constraint from the transit depth | 86 |
| 3.5.3 | Polarimetry | 86 |
| 3.6 | Discussion | 88 |
| 3.6.1 | Observational implications | 88 |
| 3.6.2 | Limitations of the model | 88 |
| 3.6.3 | High mass-loss rates | 89 |
| 3.6.4 | Constraints from dynamics | 90 |
| 3.6.5 | Plausibility of volcanic particle ejection mechanism | 90 |
| 3.7 | Summary | 92 |
| 3.8 | Appendix: Derivation of linear relationship between maximum tail height and vertical velocity | 93 |
| 4 | Self-shielding in dust tails of disintegrating rocky exoplanets | 97 |
| 4.1 | Introduction | 98 |
| 4.2 | Method: The model | 99 |
| 4.3 | Results and Discussion | 101 |
| 4.3.1 | General effects of self-shielding | 101 |

| | | |
|----------|--|------------|
| 4.3.2 | Fitting the average transit of Kepler-1520 b with the self-shielding model | 104 |
| 4.3.3 | Reduction of the intrinsic sublimation rate | 104 |
| 4.3.4 | Short-time scale outbursts | 107 |
| 4.3.5 | Highly optically thick regime | 112 |
| 4.4 | Conclusions | 112 |
| 4.5 | Future outlook | 115 |
| 5 | Search for gas from the disintegrating rocky exoplanet K2-22b | 117 |
| 5.1 | Introduction | 118 |
| 5.2 | Observational data | 119 |
| 5.3 | Analyses | 120 |
| 5.4 | Synthetic planet signal injection | 125 |
| 5.5 | Results and discussion | 126 |
| 5.5.1 | Instantaneous gas-mass limits | 126 |
| 5.5.2 | Dust and gas mass-loss comparison | 132 |
| 5.5.3 | Important Caveats: high velocity gas | 133 |
| 5.5.4 | Alternative interpretations | 136 |
| 5.6 | Conclusions and future outlook | 136 |
| 6 | Samenvatting | 143 |
| 6.1 | Gas van hete rotsachtige planeten | 143 |
| 6.2 | Waarnemen van exoplaneet atmosferen | 145 |
| 6.3 | Stofstaarten | 146 |
| 7 | Summary | 149 |
| 7.1 | Gas from hot rocky planets | 149 |
| 7.2 | Observing exoplanet atmospheres | 151 |
| 7.3 | Dust tails | 152 |
| | Curriculum Vitae | 153 |
| | List of publications | 155 |
| | Acknowledgements | 157 |