



Universiteit
Leiden
The Netherlands

Inferno Worlds

Ridden - Harper, A.

Citation

Ridden - Harper, A. (2018, November 21). *Inferno Worlds*. Retrieved from <https://hdl.handle.net/1887/67080>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/67080>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/67080> holds various files of this Leiden University dissertation.

Author: Ridden, - Harper A.

Title: Inferno Worlds

Issue Date: 2018-11-21

Inferno Worlds

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector **Magnificus** prof. mr. C.J.J.M. Stolker,
volgens besluit van het College voor Promoties
te verdedigen op woensdag 21 november 2018
klokke 10:00 uur

door

Andrew Ridden-Harper
geboren te Christchurch, Nieuw-Zeeland
in 1991

Promotiecommissie

Promotores: Prof. dr. I. A. G. Snellen
Prof. dr. C. U. Keller

Overige leden: Prof. dr. H. J. A. Röttgering
Prof. dr. H. V. J. Linmartz
Prof. dr. M. Fridlund
Dr. D. M. Stam T. U. Delft
Dr. M. Min SRON

Cover design: An artist's impression of a hot rocky exoplanet that has a dust-tail. Cover designed by Andrew Ridden-Harper, using a false colour photograph of the Sun in ultra-violet wavelengths (credit: NASA Solar Dynamics Observatory) to represent the host star. The image of the planet was adapted from an artist's impression of 55 Cancri e (credit: ESA/Hubble, M. Kornmesser).

ISBN: 978-94-6323-396-5

© 2018 Andrew Ridden-Harper

*To my parents for their unwavering
support and encouragement.*

“Imagination will often carry us to worlds that never were, but without it we go nowhere.”
– Carl Sagan

Contents

1	Introduction	1
1.1	Summary	1
1.2	Overview of the field	1
1.3	Planet formation	3
1.4	Hot rocky exoplanets	5
1.4.1	Sputtering	6
1.4.2	Possible mineral vapour atmospheres	6
1.5	Characterisation with spectroscopy	7
1.6	Disintegrating rocky exoplanets	8
1.6.1	Mass-loss mechanism	10
1.6.2	Dust particle dynamics	11
1.7	This thesis	13
1.7.1	Chapter 2	13
1.7.2	Chapter 3	14
1.7.3	Chapter 4	14
1.7.4	Chapter 5	14
1.8	Future outlook	15
2	Search for an exosphere in sodium and calcium in the transmission spectrum of exoplanet 55 Cancri e	19
2.1	Introduction	21
2.2	Observational data	24
2.2.1	UVES data	25
2.2.2	HARPS data	25
2.2.3	HARPS-N data	26
2.3	Data analysis	26
2.3.1	Processing of UVES spectra	26
2.3.2	Processing of HARPS and HARPS-N data	31
2.3.3	Combining the different data sets	31
2.3.4	Injection of artificial planet signals	32

2.4	Results	33
2.4.1	Sodium	33
2.4.2	Ionized calcium	35
2.5	Discussion and conclusions	36
3	Chromatic transit light curves of disintegrating rocky planets	47
3.1	Introduction	48
3.2	Method: The model	51
3.2.1	Dust dynamics code	51
3.2.2	Particle dynamics simulations	54
3.2.3	Ray tracing with MCMax3D	55
3.3	Results of simulations	58
3.3.1	Modelling the light curve of Kepler-1520 b with a low planet mass	58
3.3.2	Optically thick tail	62
3.3.3	Modelling the light curve of Kepler-1520 b with a planet mass of $0.02 M_{\oplus}$	69
3.3.4	Modelling the light curve of Kepler-1520 b with a planet mass of $0.02 M_{\oplus}$ and larger maximum height	70
3.3.5	Behaviour of large particles	75
3.4	Wavelength dependence	79
3.5	Constraints on particle ejection velocity	82
3.5.1	Particle trajectories	84
3.5.2	Constraint from the transit depth	86
3.5.3	Polarimetry	86
3.6	Discussion	88
3.6.1	Observational implications	88
3.6.2	Limitations of the model	88
3.6.3	High mass-loss rates	89
3.6.4	Constraints from dynamics	90
3.6.5	Plausibility of volcanic particle ejection mechanism	90
3.7	Summary	92
3.8	Appendix: Derivation of linear relationship between maximum tail height and vertical velocity	93
4	Self-shielding in dust tails of disintegrating rocky exoplanets	97
4.1	Introduction	98
4.2	Method: The model	99
4.3	Results and Discussion	101
4.3.1	General effects of self-shielding	101

4.3.2	Fitting the average transit of Kepler-1520 b with the self-shielding model	104
4.3.3	Reduction of the intrinsic sublimation rate	104
4.3.4	Short-time scale outbursts	107
4.3.5	Highly optically thick regime	112
4.4	Conclusions	112
4.5	Future outlook	115
5	Search for gas from the disintegrating rocky exoplanet K2-22b	117
5.1	Introduction	118
5.2	Observational data	119
5.3	Analyses	120
5.4	Synthetic planet signal injection	125
5.5	Results and discussion	126
5.5.1	Instantaneous gas-mass limits	126
5.5.2	Dust and gas mass-loss comparison	132
5.5.3	Important Caveats: high velocity gas	133
5.5.4	Alternative interpretations	136
5.6	Conclusions and future outlook	136
6	Samenvatting	143
6.1	Gas van hete rotsachtige planeten	143
6.2	Waarnemen van exoplaneet atmosferen	145
6.3	Stofstaarten	146
7	Summary	149
7.1	Gas from hot rocky planets	149
7.2	Observing exoplanet atmospheres	151
7.3	Dust tails	152
Curriculum Vitae		153
List of publications		155
Acknowledgements		157