



Universiteit
Leiden
The Netherlands

Criminal substance abusing adolescents and systemic treatment

Pol, T.M. van der

Citation

Pol, T. M. van der. (2018, November 28). *Criminal substance abusing adolescents and systemic treatment*. Retrieved from <https://hdl.handle.net/1887/67102>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/67102>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/67102> holds various files of this Leiden University dissertation.

Author: Pol, T.M. van der

Title: Criminal substance abusing adolescents and systemic treatment

Issue Date: 2018-11-28

4

MULTIDIMENSIONAL FAMILY THERAPY REDUCES SELF-REPORTED CRIMINALITY AMONG ADOLESCENTS WITH A CANNABIS USE DISORDER

Thimo van der Pol, Craig Henderson, Vincent Hendriks, Michael Schaub,
Henk Rigter

International Journal of Offender Therapy and Comparative Criminology.
2017;62(6):1573-88

ABSTRACT

Introduction

Multidimensional family therapy (MDFT) is an established treatment programme for youth displaying multi-problem behaviour. We examined if MDFT decreased criminal offending among cannabis abusing adolescents, as compared with individual psychotherapy (IP).

Method

In a Western-European randomised controlled trial comparing MDFT with IP, a sample of 169 adolescents with a cannabis disorder completed self-reports on criminal offending. Half indicated they had committed one or more criminal offences in the 90 days before the baseline assessment. Follow-up assessments were at 6 and 12 months after randomisation.

Results

The proportion of adolescents reporting non-delinquency increased during the study period, most so in the MDFT condition. In addition, MDFT lowered the number of violent offences more than IP. This difference was not seen for property crimes.

Conclusions

In cannabis abusing adolescents, MDFT is an effective treatment to prevent and reduce criminal offending. MDFT outperforms individual psychotherapy for violent crimes.

INTRODUCTION

In adolescents, a behavioural problem – such as substance abuse, criminal offending, truancy, or symptoms of (other) mental health disorder – often is part of a broader multi-problem behaviour constellation (Version 7.1; Griffith-Lendering, Huijbregts, Mooijaart, Vollebergh, & Swaab, 2011; Phan et al., 2011; Skeer, McCormick, Normand, Buka, & Gilman, 2009). Common is the combination of substance use disorder and criminal behaviour (delinquency) (Copeland & Swift, 2009; Y. Hser et al., 2001; Husler, Plancherel, & Werlen, 2005). Substance use disorders have been identified as a risk factor for criminal offending. Conversely, criminal offending is a risk factor for the development of substance use disorders (Moffitt et al., 2002).

Comprehensive treatments targeting multiple problems are likely to be more effective in improving the perspective of the youth than treatments targeting a single behavioural problem (Whitmore & Riggs, 2006). Problematic substance use and criminal offending are influenced by similar risk factors (Moffitt et al., 2002; Mulder, Vermunt, Brand, Bullens, & Van Marle, 2012). The two types of problem behaviour respond to the same kinds of treatment (Baldwin et al., 2012; Hogue, Henderson, Ozechowski, & Robbins, 2014).

In forensic settings, treatment often is embedded in broader intervention programmes, which may also include non-therapeutic guidance and counselling, and rehabilitation services targeting school, work, leisure time activities, and housing. A meta-analysis reviewing 28 studies found no evidence that intervention programmes, overall, decreased criminal offending in adolescents (Schwalbe, Gearing, MacKenzie, Brewer, & Ibrahim, 2012). In contrast, another meta-analysis, based on 73 studies, indicated that intervention programmes may be of modest use in preventing recidivism (Wilson & Hoge, 2013). No doubt, this discrepancy in results is partly due to the large heterogeneity of the studies included in these and other meta-analyses. Some of the studies selected for the various analysis samples focused on a disorder a youth might have (such as conduct disorder), others on measures of self-reported or registered (e.g.,

police arrests) offences, and yet others on a specific judicial programme or process (such as cautioning, diversion, probation, detention, post-release rehabilitation). Added to this heterogeneity is the large variety of intervention approaches, ranging from minimal interventions to a score of individual treatments and to family therapy. Comparison of studies was further hampered by differences and weaknesses in the organisation of the intervention programmes considered (Wilson & Hoge, 2013).

Nevertheless, a few conclusions can be drawn. Programmes involving individual treatment of the adolescent may reduce recidivism, though generally the effect is small and transient (C.E. Henderson et al., 2010; Van der Pol, Hoeve, et al., 2017; Van der Stouwe et al., 2014; Von Sydow et al., 2013). On average, cognitive behaviour therapy (CBT) has the best record among individual treatments (Greenberg & Lippold, 2013; Smeets et al., 2015; Von Sydow et al., 2013). Even better treatment results have been obtained with family therapy. In systematic literature reviews (Greenberg & Lippold, 2013; Von Sydow et al., 2013), meta-analyses (Baldwin et al., 2012; Schwalbe et al., 2012; Van der Pol, Hoeve, et al., 2017; Van der Stouwe et al., 2014), and in randomised controlled trials comparing family therapy with CBT (C.E. Henderson et al., 2010; Hendriks et al., 2011; Schaub et al., 2014), family therapy generally outperformed CBT on one or more measures of recidivism or other antisocial behaviour.

An example of a well-established family-based treatment approach is multidimensional family therapy (MDFT) (Galanter, Kleber, & Brady, 2014). MDFT is an outpatient and inpatient treatment programme for adolescents displaying problem behaviour. The term 'multidimensional' means that each major domain in the life of an adolescent is seen as contributing to the incidence and persistence of behavioural problems (through risk factors) and as potentially helpful in resolving such problems (through protective factors). The life domains include the youth him- or herself, parents, family, friends and peers, school and work, and leisure time. MDFT has been found to be more effective than active comparison therapies in various adolescent populations, doses and treatment delivery settings (Greenbaum et al., 2015; Liddle, 2010).

Most findings regarding MDFT are from U.S.-based randomised controlled trials initiated by the developers of this treatment programmes. (Junior-) Ministers of Health from five Western European countries decided to have MDFT independently tested in a European context, in a trial named INCANT (International Cannabis Need of Treatment study) comparing MDFT with individual psychotherapy (IP) (Rigter et al., 2010). INCANT confirmed the pattern of results from American trials. The European therapists delivered MDFT with a high degree of fidelity (Rowe et al., 2013). The therapy improved treatment motivation and lowered cannabis disorder rates in adolescents from outpatient treatment sites in Berlin, Brussels, Geneva, Paris and The Hague (Rigter et al., 2013), and decreased the number of symptoms of externalising disorders (Schaub et al., 2014).

One of the INCANT sites (The Hague) examined the relationship between cannabis use and criminal offending. In delinquent as compared to non-delinquent youth, MDFT outperformed IP in decreasing the number of days on which cannabis was consumed (Hendriks et al., 2012). In U.S.-based studies, MDFT lowered criminal offence rates in adolescents regardless of its effect on substance abuse in Drug Court and diversion settings (Dakof et al., 2015; Liddle, Rowe, Dakof, Henderson, & Greenbaum, 2009). These findings led us to examine MDFT's effect on criminal offending in European adolescents in more detail. In designing INCANT, two of the five INCANT sites – Geneva and The Hague – decided to extend the basic battery of assessments with the Self-Report Delinquency (SRD) Scale. The SRD records the number and types of criminal offences committed by the adolescents over the previous 90 days.

Objectives

The purpose of our study was to examine the effects of MDFT on self-reported criminal offending. We describe here the SRD outcomes for the Geneva and The Hague INCANT sites addressing two hypotheses: (a) across the 12 months follow-up period, both MDFT and IP will decrease the proportion of youth engaged in criminal offences and will reduce the number of offences committed; and (b) MDFT is more effective than IP on both types of outcome measures.

METHOD

Approval

INCANT was approved by medical-ethical committees in all involved countries (Rigter et al., 2010). For Geneva, approval was granted by the Ethical Board for Clinical and Outpatient Research (Medical Association Geneva Canton; Switzerland), and for The Hague by the Medical-Ethical Board for the Mental Health Sector in the Netherlands (METiGG).

Sample and treatment sites

Across the treatment sites in the five countries supporting INCANT, the total number of adolescents recruited for the study was 450. The study flow diagram was published by Rigter et al. (2013). The site (two sub-sites) in The Hague contributed 109 adolescents and the site in Geneva 60, yielding a sample of 169 participants for the current study.

To be included in INCANT, youth (boys and girls) had to be between 13 and 18 years of age and meet criteria for a cannabis use disorder (abuse or dependence) based on the DSM-IV, which was the manual in use when the study was carried out. Dependence and abuse were diagnosed, respectively, if at least 3 of 7 dependence criteria or 1 of 4 abuse criteria had been met. In addition, at least one parent had to indicate that he or she would participate in the treatment if they were randomised to the MDFT condition. Adolescents were excluded if they were requiring inpatient treatment because of psychosis, advanced eating disorder, or severe suicidal ideation (Rigter et al., 2010).

The treatment centres recruited for INCANT were nominated by government officials working together in the INCANT Steering Committee. The sites were visited by MDFT trainers and European project staff and were asked to give presentations and to submit documentation on the mission and funding of the centre, training level and professional background of the therapists, sources of referral of cases, caseload, treatments delivered, and links with research groups. All sites offered outpatient treatment to adolescents with substance use disorders.

The Geneva site was Phénix. In 2004, this foundation created a unit for treating adolescents with substance use disorders. Treatment staff included a psychiatrist, psychologists and social workers. Phénix is a private, non-profit organisation, with treatment costs covered by basic health insurance. There were two treatment sub-sites in The Hague. First, Parnassia Brijder – now called Brijder Addiction Care –, which among other services offers treatment programmes for adolescents with substance abuse problems. The second sub-site was Palmhuis, the forensic unit of De Jutters, which is the child and adolescent mental health institute serving The Hague and the surrounding region. Both sites are private, non-profit organisations, with treatment being paid, at the time of the study, by national, regional and local governments and through insurance funds. The MDFT team was a joint enterprise of the two sub-sites, with therapists (psychologists and social workers) from both organisations being members of the team.

Treatments

At both sites, the therapists were experienced in treating behaviourally troubled adolescents. In Geneva, MDFT was delivered by 3 MDFT certified therapists and IP by another 3 therapists. The corresponding numbers for The Hague were 6 and 12 therapists. The characteristics of these professionals (age, gender, years of experience, background) did not differ between Geneva and The Hague, or between the two treatment conditions (Rowe et al., 2013).

When preparing for INCANT, we assessed the usual treatment provided at each of the recruited sites. Although sites confessed to different theoretical orientations, e.g., mainly psychodynamic in Geneva and cognitive-behavioural in The Hague, in practice treatment as usual was individual psychotherapy (IP), consistently involving enhancement of treatment motivation, sessions with the individual adolescents (not with the parents except to inform them on treatment progress), and relapse prevention (Rigter et al., 2013; Rigter et al., 2010).

MDFT consists of three stages. The first one focuses on intensively enhancing treatment motivation, building multiple therapeutic alliances, and drafting the treatment plan. In stage 2, treatment plan interventions

targeting the youth and his or her family are carried out, including education about adolescence, behavioural development, and risk factors for problem behaviour; relapse prevention; improving family communication and relationships; and strengthening parental educational skills. Stage 3 involves sealing off the treatment, agreeing on a relapse prevention plan, and providing booster sessions if needed.

In INCANT, both MDFT and IP were scheduled to last for 6 months. MDFT was delivered in approximately two sessions per week – in roughly equal proportion to be held with the adolescent, parent, and family (adolescent and parent together). In IP, the number of sessions with the adolescent was matched to be similar to MDFT. However, the total number of IP sessions was lower than for MDFT, as there were no sessions with parents and family. Rowe et al. (2013) present details on the actual treatment dose received; this paper also documents the efforts to evaluate and safeguard treatment integrity and fidelity.

Design

INCANT was a multi-centre phase randomised controlled effectiveness trial with an open-label, parallel group design, running from 2006 to 2010. Assessments were scheduled at baseline – immediately before randomisation and start of treatment – and at 3, 6, 9 and 12 months thereafter (Rigter et al., 2010). The SRD was administered at baseline and 6 and 12 months follow-up. Randomisation occurred immediately after the eligibility of the case had been confirmed at baseline. The INCANT database, at the Department of Public Health of Erasmus Medical Centre in Rotterdam, assigned a code to each new case entered by a site's research assistant and automatically informed her about the allocated treatment. In order to conceal the randomisation process, trial staff was not involved in any step of the procedure (Rigter et al., 2010).

Outcome measures

The measurements were delivered at baseline, at 6 months (if planned), and at 12 months follow-up.

Cannabis use

Frequency of cannabis use by the adolescents was recorded with the Timeline Follow-Back method (TLFB), as adapted and validated for adolescents (Sobell & Sobell, 1992; Holly Barrett Waldron & Charles W Turner, 2008). The TLFB obtains reports of daily cannabis use for the 90 days preceding the assessment, using a calendar and other memory prompts.

Cannabis use disorders (abuse and dependence)

Cannabis use disorders were identified with the Adolescent Diagnostic Interview-Light (ADI-Light). This structured multi-axial interview generating DSM-IV diagnoses has good psychometric properties, as assessed in reliability and validity tests (Winters & Henly, 1993). The ADI-Light was administered at baseline and at 12-month follow-up.

Criminal offences

To trace the number and type of offences committed by the adolescents, we administered the SRD, the Self-Report Delinquency Scale (Elliott, Huizinga, & Ageton, 1985), asking the youth how many and which type of criminal offences they had committed in the past 90 days. We analysed SRD scores for the classes of property crimes and of violent crimes (aggression, violent sexual offence, violent property offence), respectively, and for these types of crimes together (total scores). The SRD scored well in tests of reliability and validity (Elliott et al., 1985).

Statistical analyses*Baseline characteristics*

Between-treatment equivalence was tested with analyses of variance for continuous variables, and chi-square tests for categorical variables.

Changes over time across treatment conditions

Latent growth curve (LGC) modelling with robust maximum likelihood estimation was used to analyse change for each adolescent. The missing at random (MAR) assumption could not be directly evaluated. We explored the reasonableness of the MAR assumption holding with these data by checking if there were significant correlations between key

study variables and a binary variable indicating whether the data were missing or not (1 = missing at follow-up assessment, 0 = not missing). As correlations were negligible ($r < 0.10$), we treated incomplete data as MAR and accounted for it in subsequent models using Full Information Maximum Likelihood estimation (Little & Rubin, 2002). We included a dummy coded variable representing treatment condition (IP = 0; MDFT = 1) in the model to test the equivalence of groups at baseline and the impact of intervention type on change over time (i.e., the intercept and slope growth parameters). Intervention effects were demonstrated by a statistically significant slope parameter, as tested by the pseudo-z test associated with treatment condition. LGC modelling was carried out with Mplus (Version 7.1; Muthén, 2016).

Density plots revealed a high proportion of participants reporting no criminal offences at each follow-up assessment. Therefore, we used a two-part growth modelling approach (E. C. Brown, Catalano, Fleming, Haggerty, & Abbott, 2005; Liddle et al., 2009) to estimate separate but correlated continuous and categorical LGC models. This approach was developed to address non-normality caused by a preponderance of zeros (Olsen & Schafer, 2001). As implemented in Mplus, two-part growth modelling applies a natural log transformation to the continuous outcomes. The modelling approach was successful in bringing skewness and kurtosis below acceptable levels (below 1.5). Further, we used the robust maximum likelihood estimator for all analyses to minimise the impact of non-normality on the results.

Effect sizes

The effect size parameter d for treatment comparisons was computed using Feingold's method for calculating effect sizes with growth curve models (Feingold, 2009). A d in the range of 0.30 - 0.70 indicates that the effect was of moderate size; higher d values reflect strong effects.

RESULTS

Missing values

There were no missing data at baseline. At 12 months follow-up, 28% of the adolescents did not complete the SRD. There were no differences between treatments in this respect ($\chi^2 [1] = 0.40$). However, the rate of missing SRD forms was higher in Geneva than in The Hague, $\chi^2 [1] = 50.62, p < 0.01$.

Baseline

Table 1 presents baseline data for the two sites and two treatment conditions. Both across sites (the columns 'Total') and per site, the adolescents from the two treatment conditions were similar in age and gender, and in characteristics of their parents (divorce rate; prevalence of mental health and substance use problems). However, when comparing the two sites with each other across treatment conditions, the two populations of adolescents differed in foreign descent (rate higher in Geneva; $\chi^2 [4] = 10.88, p = 0.03$), and proportion of youth living with their family (rate higher in The Hague; $\chi^2 [3] = 13.48, p = 0.004$). Also, the two populations were distinct from each other on cannabis use measures (days of use: $F [1, 167] = 9.56, p = 0.002$; proportion of adolescents being dependent on cannabis: $\chi^2 [1] = 17.13, p < 0.001$). Cannabis dependence was more common among the adolescents in Geneva than in The Hague, although the self-reported number of cannabis use days was lower in Geneva. Within sites, the variables mentioned did not statistically differ between the two treatment conditions.

Table 1. Baseline data for the The Hague and Geneva sites.

Variable	The Hague MDFT	The Hague IP	Geneva MDFT	Geneva IP	Total MDFT	Total IP
Mean age \pm SD	16.2 \pm 1.3	16.3 \pm 1.2	16.1 \pm 1.2	16.0 \pm 1.2	16.2 \pm 1.3	16.2 \pm 1.2
Male	80%	80%	90%	93%	83%	85%
From foreign descent	46%	48%	73%	60%	58%	55%
Attending school	77%	74%	67%	70%	74%	73%
Living with family	98%	98%	82%	83%	90%	93%
Parents separated	60%	56%	63%	53%	61%	55%
Parents with mental health or substance use problems	29%	26%	33%	33%	31%	29%
Behaviour						
Total [#] nr. of self-reported criminal offences \pm SD*	5.8 \pm 13.8	6.4 \pm 18.5	8.5 \pm 18.6	8.9 \pm 20.3	6.8 \pm 15.6	7.3 \pm 19.1
Cannabis use days \pm SD*	64 \pm 23	61 \pm 24	47 \pm 25	52 \pm 29	58 \pm 25	58 \pm 26
Cannabis dependence	73%	78%	90%	97%	77%	77%

* In the 90 days before the baseline assessment. SD = standard deviation.

Criminal offences = property and violent crimes taken together.

Change in criminal offence rate

Preliminary analyses

At baseline, 43% of the adolescents said that they had not committed any criminal offence in the preceding 90 days. Forty-eight percent of participants reported they had committed a property crime, and 33% a violent crime. At this point in time, there were no differences between treatment groups in proportion of adolescents reporting any ($X^2(1) = 3.06$, *ns*), property ($X^2(1) = 1.21$, *ns*), or violent ($X^2(1) = 2.83$, *ns*) crimes.

Offences of any type: changes over time across treatment conditions

Over the 12 months follow-up period and across sites, the number of self-reported criminal offences dropped in both the MDFT and IP groups, with 70% and 46%, respectively, with no significant difference noted. Looking per site, the decline in number of offences was similar in the MDFT and IP groups in The Hague (77% versus 72%), but dissimilar in Geneva (35% decrease in the MDFT condition versus 35% increase for IP). Across treatments, the *proportion* of youth reporting no criminal offences (represented by the categorical part of the frequency model) slightly increased from baseline to follow-up assessments (Mean Slope = -0.28, standard error [SE] = 0.16, *pseudo-z* = -1.79, $p = 0.07$). Among those reporting they had engaged in criminal offending during the study period (the continuous part of the model), the *number* of criminal offences decreased over time (Mean Slope = -0.21, SE = 0.08, *pseudo z* = -2.54, $p = 0.01$).

Abstaining from criminal offending: treatment comparisons

Comparing the treatments in the categorical model, more youth receiving MDFT reported abstaining from any type of criminal offence over time than corresponding IP youth (Treatment Slope = -0.70, SE = 0.33, *pseudo z* = -2.14, $p = 0.03$, $d = 0.51$); see Figure 1, Panel 1. This pattern of results held for both property crimes (Slope = 0.34, SE = 0.05, *pseudo z* = 6.92, $p < 0.01$, $d = 4.95$) and violent crimes (Slope = 0.39, SE = 0.05, *pseudo z* = 8.53, $p < 0.01$, $d = 7.53$).

As for the *number* of criminal offences of any type, the two treatments did not differ in the continuous part of the model (Treatment Slope = -0.01, $SE = 0.17$, $pseudo\ z = -0.07$, ns , $d = 0.01$; see Figure 1, Panel 2). The number of *total* criminal offences declined in both treatment groups, with no advantage of MDFT over IP. The decrease in the number of criminal offences was marginally larger in The Hague than in Geneva (Slope = 0.29, $SE = 0.17$, $pseudo\ z = 1.68$, $p = 0.09$, $d = 0.79$).




Figure 1. Change in proportion of youth reporting abstaining from delinquency (Panel 1) and number of delinquent acts among those persisting in delinquent behaviour (Panel 2).

Committing property crimes versus violent crimes: treatment comparisons

We divided the total criminal offences category into property and violent crimes. Among youth engaging in property crimes over the 12-month follow-up period, the number of these offences among those reporting delinquent behaviour dropped over time (Slope = -0.37, $SE = 0.17$, $pseudo\ z = -2.17$, $p = 0.03$); Figure 2. MDFT and IP did not differ in this respect in either the continuous (Slope = -0.11, $SE = 0.22$, $pseudo\ z = -0.50$, ns , $d = 0.10$) or categorical part of the model (Slope = 0.07, $SE = 0.06$, $pseudo\ z = 1.18$, ns , $d = 0.28$).




Figure 2. Change in proportion of youth reporting abstaining from property crimes (Panel 1) and number of property crimes among those committing property crimes (Panel 2).

For violent crimes, the overall *proportion* of adolescents engaging in violent crimes did not change over time (Slope = 0.02, *SE* = 0.03, *pseudo z* = 0.46, *ns*). However, when treatment condition was entered into the calculations, more youth receiving MDFT rather than IP reported to have abstained from violent offences (Slope = 0.10, *SE* = 0.05, *pseudo z* = 2.07, $p = 0.04$, $d = 0.43$).




Figure 3. Change in proportion of youth reporting abstaining from violent crimes (Panel 1) and number of violent crimes among those committing violent crimes (Panel 2).

The *number* of violent crimes among self-reported violent offenders did not change over time (Slope = 0.11, $SE = 0.15$, $pseudo\ z = 0.74$, *ns*). Comparing the two treatments, the adolescents receiving MDFT tended to commit fewer violent offences over time than their IP counterparts, but although the effect size was moderately large, the difference between the treatment groups was not statistically significant (Slope = 0.23, $SE = 0.20$, $pseudo\ z = 1.13$, *ns*, $d = 0.63$); Figure 3.

DISCUSSION

All adolescents in the present study had a cannabis use disorder at baseline, mostly cannabis dependence. Half of them reported having committed one or more criminal offences in the 90 days before the baseline assessment, i.e., the moment of their recruitment in the INCANT randomised trial. Across all youth, the number of self-reported criminal offences per period of 6 months dropped going from baseline to the 12-months follow-up assessment. In other words, both MDFT and IP appeared to be effective in decreasing criminal behaviour, in accordance with our first study hypothesis. MDFT was as effective in this respect as IP, which would appear to run counter to our second study hypothesis, which stated that MDFT would outperform IP. Yet, the second study hypothesis was confirmed in part. Dividing criminal offences into property and violent crimes revealed a treatment difference. The drop in property crimes was similar in the MDFT and IP conditions, but the decrease in violent crimes was larger for MDFT than for IP.

We do not know of any publications clearly showing differential treatment effects on committing property crimes versus violent crimes in adolescents. The offence measures used by Dakof et al. (2015), who found MDFT to be superior to Drug Court group therapy, included 'serious crimes', but without clear distinction between property and violent offences. However, there are epidemiological data suggesting that treatment of delinquent adolescents should be tuned to certain characteristics of these youth. The literature contains many attempts to draft a typology of delinquent youth.

Most often mentioned (disregarding sexual offenders) is the distinction between violent offenders, non-violent (property) offenders, and versatile offenders who commit both violent and property crimes (Lai, Zeng, & Chu, 2016). For these three classes of adolescent offenders, different profiles of risk factors apply (Colins, Vermeiren, Schuyten, & Broekaert, 2009; Lai et al., 2016; Mulder et al., 2012). Most impaired in risk factor exposure and mental and behavioural health are the versatile offenders (Lai et al., 2016), who in our study were labelled as violent offenders, because violent property crimes were classified as violent offences. The excess of risk factors facing violent/versatile offenders appears to be concentrated on the 'mental comorbidity' (Colins et al., 2009), 'family' (e.g., poor parental supervision) and 'peers' (wrong friends) dimensions (Lai et al., 2016; Mulder et al., 2012).

Why would family therapy work out better in reducing delinquency in adolescents than individual therapy? Both types of treatment are effective in decreasing criminal offending. The surplus value of family therapy may be explained by the ambition to have this type of treatment address risk factors not only at the individual level (the adolescent with his or her personality traits and response patterns), but also at the family, peers, school/work and leisure time levels. The latter factors strongly influence the behaviour of an adolescent (Lai et al., 2016; Leve, Chamberlain, & Kim, 2015; Wilson & Hoge, 2013), strengthening the case for family therapy.

MDFT is not the only family therapy with credits in treating criminal youth. From U.S. research, five major programmes have emerged (Leve et al., 2015), with Multisystemic Therapy (MST) and MDFT probably having the best research record in Europe, in addition to good performance in the USA (Van der Pol, Hoeve, et al., 2017; Van der Stouwe et al., 2014). The evidence suggests that MDFT has effect in 'light' cases, but certainly is to be preferred in 'severe' cases. In substance abuse research (Henderson et al., 2010; Rigter et al., 2013), MDFT was as effective as individual psychotherapy in reducing problem behaviour for all cases together. However, MDFT did better than IP in 'severe' cases, however defined. Our present results suggest that the same may be true for the effect of treatment on criminal offending in adolescents. MDFT and IP are both effective in reducing self-reported criminal offences, but MDFT

outperforms IP in violent/versatile offenders, who might have been more severely impaired than the offenders committing property crimes (Colins et al., 2009). Clearly, more research is needed here.

A special finding of our study was that MDFT may not only lower recidivism rates, but also may help to prevent first-time offences. In our trial, the proportion of non-delinquent youth grew somewhat during the study period, most clearly so for adolescents receiving MDFT. A recent meta-analysis confirmed that intervention programmes may prevent (the first incident of) criminal offending. Effective programmes are family-oriented and 'multimodal' (multidimensional, in MDFT's terminology) (Vries, Hoeve, Assink, Stams, & Asscher, 2015). Our findings are in keeping with this conclusion.

A strength of the INCANT trial was that it excluded few adolescents from taking part in the study. The trial's aim was to achieve a high external validity level. The sites differed in many respects, such as in referral practices, i.e., the route of bringing an adolescent and his or her family into contact with a treatment centre. Many Swiss adolescents recruited for INCANT had been referred to the trial by a juvenile judge. So, the high rate of criminal offending in the Geneva youth is not surprising. The adolescents from The Hague, who were regularly referred from non-Justice sources, were probably less impaired than the Geneva youth (Phan et al., 2011). Yet, despite these differences in referral pathways, MDFT appeared to be effective at all sites in all countries (Phan et al., 2011; Rigter et al., 2013). This is confirmed in the present paper. A possible weakness of the study was that the criminal offence data were based on self-report. Self-report data may be biased. However, the jury is still out on the question if supposedly more objective database records (on arrests, convictions) are a better source of information (Kirk, 2006). Database records only contain data on registered criminal offences; self-report invites respondents to also report criminal offences that went unnoticed to police and justice authorities. In a separate paper, we will describe results for a database measure of criminal offending, i.e., police arrests of The Hague INCANT youth in the 3 years following randomisation, which confirmed that MDFT lowers criminal offence rates in adolescents.

From a policy perspective, we would recommend that in juvenile forensic settings treatment programmes are to be implemented that do not focus on just one behavioural problem, but on the common multiplicity of behavioural problems. Also, it is advisable to opt for an evidence-based family therapy rather than an individual treatment targeting the adolescent him- or herself. The broader approach of family therapy is likely to more strongly reduce recidivism rates of serious (violent/versatile) crimes than individual treatment, in addition to having a preventive effect on criminal offending in general.