



Universiteit
Leiden
The Netherlands

Monogenic models of migraine : from clinical phenotypes to pathophysiological mechanisms

Pelzer, N.

Citation

Pelzer, N. (2018, November 28). *Monogenic models of migraine : from clinical phenotypes to pathophysiological mechanisms*. Retrieved from <https://hdl.handle.net/1887/67103>

Version: Not Applicable (or Unknown)

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/67103>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden




The handle <http://hdl.handle.net/1887/67103> holds various files of this Leiden University dissertation.

Author: Pelzer, N.

Title: Monogenic models of migraine : from clinical phenotypes to pathophysiological mechanisms

Issue Date: 2018-11-28



Chapter 10.

Heterozygous *TREX1* mutations in early-onset cerebrovascular disease

N. Pelzer¹, B. de Vries², E.M.J. Boon³, M.C. Kruit⁴, J. Haan^{1,5}, M.D. Ferrari¹, A.M.J.M. van den Maagdenberg^{1,2}, G.M. Terwindt¹

¹Department of Neurology, Leiden University Medical Center, Leiden, the Netherlands

²Department of Human Genetics, Leiden University Medical Center, Leiden, the Netherlands

³Department of Clinical Genetics, Leiden University Medical Center, Leiden, the Netherlands

⁴Department of Radiology, Leiden University Medical Center, Leiden, the Netherlands

⁵Department of Neurology, Alrijne Hospital, Leiderdorp, the Netherlands

Dear Sirs,

We report on 100 patients suspected of cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL)¹ who were analyzed in our international CADASIL referral center, but in whom no *NOTCH3* mutations were found. *TREX1* was considered an excellent next disease gene candidate because of its link with Aicardi-Goutières syndrome (AGS),³ the small vessel disease retinal vasculopathy and cerebral leukodystrophy (RVCL),⁸ and (neuropsychiatric) systemic lupus erythematosus (SLE).⁴⁻⁶ Screening of our patient cohort yielded heterozygous *TREX1* mutations in two patients with early-onset cerebrovascular disease. This expands the clinical spectrum of diseases associated with *TREX1* mutations, and offers a differential diagnosis for CADASIL-like phenotypes.

Patient A, a 53-year-old woman with a history of hypertension, hyperlipidemia and alcohol abuse presented with otherwise unexplained presenile dementia. MRI demonstrated basal ganglia and pontine lacunar infarcts and bilateral confluent white matter lesions (Fig. 1a–c). An extensive analysis for causes of dementia and stroke, including serum autoantibodies (ANCA, ANA) was negative. Her severe retinopathy was labeled as hypertensive. Severely demented, she died at the age of 55 of an aspiration pneumonia. Direct sequencing of the *TREX1* gene detected a heterozygous mutation c.1079A>G; p.Tyr360-Cys (p.Tyr305Cys on *TREX1* isoform B).

Patient B, a 42-year-old heavily smoking man with a past medical history including a splenic artery aneurysm at age 29, hypertension, hyperlipidemia and vascular claudication presented with progressive cognitive dysfunction. His family history was positive for (cardio)vascular disease. MRI showed a cortico-subcortical infarct in the right frontal lobe, ischemic lesions in the basal ganglia, brainstem and corpus callosum, and focal T2 white matter hyperintensities (Fig. 1d–f). Extensive diagnostic work-up did not reveal other causes for his cognitive decline. CSF showed a mononuclear pleocytosis with normal protein levels, but angiography was not suggestive of a cerebral vasculitis. CSF culture was negative and serum autoantibodies (ANCA, ANA, cardiolipines) were absent. He had a deep venous thrombosis in the leg at age 44. At age 46 and 47 he suffered from new brainstem infarcts. Direct sequencing detected a heterozygous *TREX1* mutation c.506G>A; p.Arg169His (p.Arg114His on *TREX1* isoform B).

The two patients described were referred for CADASIL screening because of extensive white matter hyperintensities and infarcts and progressive early-onset (cerebro)vascular disease. No *NOTCH3*




Fig. 1 **a–c** Axial FLAIR MRIs and sagittal T1-weighted MRI of patient A at age 53 show signs of lacunar infarcts in the basal ganglia, bilateral periventricular white matter lesions, and a pontine infarct; **d–f** Axial FLAIR MRIs and sagittal T2-weighted MRI of patient B at age 47 show right frontal, basal ganglia, brainstem and corpus callosum infarcts.

mutations were found. Screening of the *TREX1* gene revealed heterozygous mutations p.Tyr305Cys and p.Arg114His, which were both previously reported in SLE patients.^{5,6} The p.Tyr305Cys mutation affects a highly conserved residue and was not found in large cohorts of healthy controls.^{5,6} The heterozygous p.Arg114His mutation, however, has also been identified in six controls.^{5,6} Nonetheless, when homozygous, this mutation is quite common in AGS.^{2,5} Mutation p.Arg114His affects a highly conserved residue in the catalytic domain of *TREX1* and severely decreases enzymatic activity, suggesting pathogenicity.^{5,7} Enzymatic activity was also (mildly) decreased when co-expressed with wildtype *TREX1* protein, mimicking the situation in heterozygotes.⁷ Mutated *TREX1* protein seems to cause accumulation of cytosolic nucleic acids and subsequent abnormal innate immune responses, which may have damaging effects on the circulatory system.²

Clinical features and MRI abnormalities in our patients were not suggestive of RVCL or AGS. Absent

autoantibodies made SLE unlikely. Nevertheless, we hypothesise that *TREX1* mutations may have caused these phenotypes with adult onset. Even the phenomena referred to as ‘vascular risk factors’ may be part of the phenotypic spectrum of *TREX1* mutations.

We suggest that early-onset cerebrovascular disease can be caused by heterozygous *TREX1* mutations. Further elucidation of pathogenetic mechanisms of *TREX1* mutations may reveal new cerebrovascular disease mechanisms.

Acknowledgments: This work was supported by the European Union Seventh Framework Programme grant agreement number 241779 (NIMBL: <http://www.NIMBL.eu/>) (A.M.J.M.v.d.M). They had no role in the design and conduct of the study, the collection, management, analysis, and interpretation of the data, nor in the preparation, review, or approval of the manuscript.

Conflicts of interest: On behalf of all authors, the corresponding author states that there is no conflict of interest.

Ethical standard: This study has been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

References

1. Chabriat H, Joutel A, Dichgans M, et al. Cadasil. *Lancet Neurol* 2009;8:643–653.
2. Chahwan C, Chahwan R. Aicardi-Goutieres syndrome: from patients to genes and beyond. *Clin Genet* 2012;81:413–420.
3. Crow YJ, Hayward BE, Parmar R, et al. Mutations in the gene encoding the 3′-5′ DNA exonuclease *TREX1* cause Aicardi-Goutieres syndrome at the *AGS1* locus. *Nat Genet* 2006;38:917–920.
4. de Vries B, Steup-Beekman GM, Haan J, et al. *TREX1* gene variant in neuropsychiatric systemic lupus erythematosus. *Ann Rheum Dis* 2010;69:1886–1887.
5. Lee-Kirsch MA, Gong M, Chowdhury D, et al. Mutations in the gene encoding the 3′-5′ DNA exonuclease *TREX1* are associated with systemic lupus erythematosus. *Nat Genet* 2007;39:1065–1067.
6. Namjou B, Kothari PH, Kelly JA, et al. Evaluation of the *TREX1* gene in a large multi-ancestral lupus cohort. *Genes Immun* 2011;12:270–279.
7. Orebaugh CD, Fye JM, Harvey S, et al. The *TREX1* exonuclease R114H mutation in Aicardi-Goutieres syndrome and lupus reveals dimeric structure requirements for DNA degradation activity. *J Biol Chem* 2011;286:40246–40254.
8. Richards A, van den Maagdenberg AM, Jen JC, et al. C-terminal truncations in human 3′-5′ DNA exonuclease *TREX1* cause autosomal dominant retinal vasculopathy with cerebral leukodystrophy. *Nat Genet* 2007;39:1068–1070.