



Universiteit
Leiden
The Netherlands

Impact of plant domestication on spermosphere and rhizosphere microbiome composition

Perez Jaramillo, J.E.

Citation

Perez Jaramillo, J. E. (2019, March 28). *Impact of plant domestication on spermosphere and rhizosphere microbiome composition*. Retrieved from <https://hdl.handle.net/1887/70478>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/70478>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/70478> holds various files of this Leiden University dissertation.

Author: Perez Jaramillo, J.E.

Title: Impact of plant domestication on spermosphere and rhizosphere microbiome composition

Issue Date: 2019-03-28

References

- Abbo S, Pinhasi van-Oss R, Gopher A, Saranga Y, Ofner I, Peleg Z. 2014. Plant domestication versus crop evolution: a conceptual framework for cereals and grain legumes. *Trends Plant Sci* 19(6):351-60.
- Adam E, Bernhart M, Müller H, Winkler J, Berg G. 2018. The *Cucurbita pepo* seed microbiome: genotype-specific composition and implications for breeding. *Plant Soil* 422:35-49.
- Aira M, Gómez-Brandón M, Lazcano C, Bååth E, Domínguez J. 2010. Plant genotype strongly modifies the structure and growth of maize rhizosphere microbial communities. *Soil Biol Biochem* 42:2276-2281.
- Aguilar OM, Riva O, Peltzer E. 2004. Analysis of *Rhizobium etli* and of its symbiosis with wild *Phaseolus vulgaris* supports coevolution in centres of host diversification. *Proc Natl Acad Sci U S A* 101(37):13548-13553.
- Akibode S, Maredia M. 2011. Global and regional trends in production, trade and consumption of food legume crops. (Report submitted to the CGIAR Standing Panel on Impact Assessment). <http://www.cgiar.org/our-research/crop-factsheets/beans/>
- Aleklett K, Hart M. 2013. The root microbiota—a fingerprint in the soil? *Plant Soil* 370:671-686.
- Aleklett K, Leff JW, Fierer N, Hart M. 2015. Wild plant species growing closely connected in a subalpine meadow host distinct root-associated bacterial communities. *PeerJ* 3:e804.
- Amarger N. 2001. Rhizobia in the field. *Adv Agron* 73:109-168.
- Arrieta MC, Stiemsma LT, Amenyogbe N, Brown EM, Finlay B. 2014. The intestinal microbiome in early life: health and disease. *Front Immunol* 5:427.
- Badri DV, Chaparro JM, Zhang R, Shen Q, Vivanco JM. 2013. Application of natural blends of phytochemicals derived from the root exudates of Arabidopsis to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. *J Biol Chem* 288(7):4502-4512.
- Badri DV, Vivanco JM. 2009. Regulation and function of root exudates. *Plant Cell Environ* 32(6):666-81.
- Badri DV, Weir TL, van der Lelie D, Vivanco JM. 2009. Rhizosphere chemical dialogues: plant-microbe interactions. *Curr Opin Biotechnol* 20(6):642-50.
- Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. 2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol* 57:233-66.
- Bakker PAHM, Berendsen RL, Doombos RF, Wintermans PCA, Pieterse CML. 2013. The rhizosphere revisited: root microbiomics. *Front Plant Sci* 4:165.
- Bakker MG, Manter DK, Sheflin AM, Weir TL, Vivanco JM. 2012. Harnessing the rhizosphere microbiome through plant breeding and agricultural management. *Plant Soil* 360(1-2):1-13.
- Barberán A, Bates ST, Casamayor EO, Fierer N. 2012. Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J* 6(2):343-51.
- Barret M, Briand M, Bonneau S, Préveaux A, Valière S, Bouchez O, *et al.* 2015. Emergence shapes the structure of the seed microbiota. *Appl Environ Microbiol* 81(4):1257-66.

- Bastian M, Heymann S, Jacomy M. 2009. Gephi: an open source software for exploring and manipulating networks. International AAAI Conference on Weblogs and Social Media: San Jose, California.
- Beckers B, Op De Beeck M, Weyens N, Van Acker R, Van Montagu M, *et al.* 2016. Lignin engineering in field-grown poplar trees affects the endosphere bacterial microbiome. *Proc Natl Acad Sci U S A* 113:2312-7.
- Bednarek P, Kwon C, Schulze-Lefert P. 2010. Not a peripheral issue: secretion in plant-microbe interactions. *Curr Opin Plant Biol* 13(4):378-87.
- Beebe S, Rengifo J, Gaitan E, Duque MC, Tohme J. 2001. Diversity and origin of Andean landraces of common bean. *Crop Sci* 41(3):854-862.
- Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health. *Trends Plant Sci* 17(8):478-486.
- Berendsen RL, Vismans G, Yu K, Song Y, de Jonge R, Burgman WP, *et al.* 2018. Disease-induced assemblage of a plant-beneficial bacterial consortium. *ISME J* 12(6):1496-1507.
- Berg G, Grube M, Schloter M, Smalla K. 2014. Unraveling the plant microbiome: looking back and future perspectives. *Front Microbiol* 5:148.
- Berg G, Raaijmakers JM. 2018. Saving seed microbiomes. *ISME J* 12(5):1167-1170.
- Berg G, Smalla, K. 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiol Ecol* 68: 1-13.
- Berlemont R, Martiny AC. 2015. Genomic potential for polysaccharide deconstruction in bacteria. *Appl Environ Microbiol* 81(4):1513-19.
- Berry D, Widder S. 2014. Deciphering microbial interactions and detecting keystone species with co-occurrence networks. *Front Microbiol* 5:219.
- Bingol K. 2018. Recent advances in targeted and untargeted metabolomics by NMR and MS/NMR methods. *High Throughput* 7(2):E9.
- Bitocchi E, Bellucci E, Giardini A, Rau D, Rodriguez M, Biagetti E, *et al.* 2013. Molecular analysis of the parallel domestication of the common bean. *Phaseolus vulgaris*. in Mesoamerica and the Andes. *New Phytol* 197:300–313.
- Bitocchi E, Nanni L, Bellucci E, Rossi M, Giardini A, P Spagnoletti Zeuli, *et al.* 2012. Mesoamerican origin of the common bean (*Phaseolus vulgaris* L.) is revealed by sequence data. *Proc Nat Acad Sci U S A* 109(14): E788-E796.
- Blair MW, Soler A, Cortés AJ. 2012. Diversification and population structure in common beans (*Phaseolus vulgaris* L.). *PLoS One* 7(11):e49488.
- Blaser MJ. 2018. The past and future biology of the human microbiome in an age of extinctions. *Cell* 172:1173-1177.
- Boon E, Meehan CJ, Whidden C, Wong DH, Langille MG, Beiko RG. 2013. Interactions in the microbiome: communities of organisms and communities of genes. *FEMS Microbiol Rev* 38(1):90-118.

- Bouffaud M-L, Poirier M-A, Mulle D and Moëgne-Loccoz, Y. 2014. Root microbiome relates to plant host evolution in maize and other Poaceae. *Environ Microbiol* 16:2804-2814.
- Bouwmeester HJ, Roux C, Lopez-Raez JA, Bécard G. 2007. Rhizosphere communication of plants, parasitic plants and AM fungi. *Trends Plant Sci* 12(5):224-30.
- Brewer A, Williamson M. 1994. A new relationship for rarefaction. *Biodivers Conserv* 3:373-379.
- Broughton WJ, Hernández G, Blair M, Beebe S, Gepts P, Vanderleyden J. 2003. Beans (*Phaseolus* spp.) - model food legumes. *Plant Soil* 252:55-128.
- Brown K, DeCoffe D, Molcan E, Gibson DL. 2012. Diet-induced dysbiosis of the intestinal microbiota and the effects on immunity and disease. *Nutrients* 4(8):1095-1119.
- Broz AK, Manter DK, Vivanco JM. 2007. Soil fungal abundance and diversity: another victim of the invasive plant *Centaurea maculosa*. *ISME J* 1(8):763-5.
- Budak H, Kantar M, Kurtoglu KY. 2013. Drought tolerance in modern and wild wheat. *Sci World J* ID:548246.
- Buée M, de Boer W, Martin F, van Overbeek LS, Jurkevitch E. 2009. The rhizosphere zoo: An overview of plant-associated communities of microorganisms, including phages, bacteria, archaea, and fungi, and some of their structuring factors. *Plant Soil* 321:189-212.
- Bulgarelli D, Garrido-Oter R, Münch PC, Weiman A, Dröge J, Pan Y *et al.* 2015. Structure and function of the bacterial root microbiota in wild and domesticated barley. *Cell Host Microbe* 17: 392-403.
- Bulgarelli D, Rott M, Schlaeppi K, van Themaat EVL, Ahmadinejad N, Assenza F *et al.* 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. *Nature* 488:91-95.
- Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P. 2013. Structure and functions of the bacterial microbiota of plants. *Annu Rev Plant Biol* 64: 807-38.
- Burdman S, Volpin H, Kigel J, Kapulnik Y, Okon Y. 1996. Promotion of nod gene inducers and nodulation in common bean. *Phaseolus vulgaris*. roots inoculated with *Azospirillum brasilense* Cd. *Appl Environ Microbiol* 62(8):3030-3.
- Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, *et al.* 2017. Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol* 28:15(3):e2001793.
- Busse HJ. 2016. Review of the taxonomy of the genus *Arthrobacter*, emendation of the genus *Arthrobacter sensu lato*, proposal to reclassify selected species of the genus *Arthrobacter* in the novel genera *Glutamicibacter* gen. nov., *Paeniglutamicibacter* gen. nov., *Pseudoglutamicibacter* gen. nov., *Paenarthrobacter* gen. nov. and *Pseudarthrobacter* gen. nov., and emended description of *Arthrobacter roseus*. *Int J Syst Evol Microbiol* 66(1):9-37
- Buyer JS, Roberts DP, Russek-Cohen E. 1999. Microbial community structure and function in the spermosphere as affected by soil and seed type. *Can J Microbiol* 45(2): 138-144.
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, *et al.* 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7:335-336.

- Cardinale M, Grube M, Erlacher A, Quehenberger J, Berg G. 2015. Bacterial networks and co-occurrence relationships in the lettuce root microbiota. *Environ Microbiol* 17:239-52.
- Carrión VJ, Cordovez V, Tyc O, Etalo DW, de Bruijn I, de Jager VCL, *et al.* 2018. Involvement of Burkholderiaceae and sulfurous volatiles in disease-suppressive soils. *ISME J* 12(9):2307-2321
- Carvalhais LC, Dennis PG, Fedoseyenko D, Hajirezaei MR, Borriss R *et al.* 2011. Root exudation of sugars, amino acids, and organic acids by maize as affected by nitrogen, phosphorus, potassium, and iron deficiency. *J Plant Nutr Soil Sci* 74:3-11.
- Carvalhais LC, Dennis PG, Badri DV, Tyson GW, Vivanco JM, Schenk PM. 2013. Activation of the jasmonic acid plant defence pathway alters the composition of rhizosphere bacterial communities. *PLoS One* 8: e56457.
- Cesco S, Neumann G, Tomasi N, Pinton R, Weisskopf L. 2010. Release of plant-borne flavonoids into the rhizosphere and their role in plant nutrition. *Plant Soil* 329:1-25.
- CGIAR. 2013. Annual progress report 2013: CGIAR Research Program on Grain Legumes, 66 pp.
- Chacón MI, Pickersgill SB, Debouck DG. 2005. Domestication patterns in common bean (*Phaseolus vulgaris* L.) and the origin of the Mesoamerican and Andean cultivated races. *Theor Appl Genet* 110:432-444.
- Chacón-Fuentes M, Parra L, Lizama M, Seguel I, Urzúa A, Quiroz A. 2017. Plant flavonoid content modified by domestication. *Environ Entomol* 46:1080-1089.
- Chagas FO, Pessotti RC, Caraballo-Rodríguez AM, Pupo MT. 2018. Chemical signaling involved in plant-microbe interactions. *Chem Soc Rev* 47(5):1652-1704.
- Chaluvadi S, Bennetzen JL. 2018. Species-associated differences in the below-ground microbiomes of wild and domesticated *Setaria*. *Front Plant Sci* 9:1183.
- Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience* 4:7.
- Chaparro JM, Badri DV, Vivanco JM. 2014. Rhizosphere microbiome assemblage is affected by plant development. *ISME J* 8:790-803.
- Chapelle E, Mendes R, Bakker PA, Raaijmakers JM. 2016. Fungal invasion of the rhizosphere microbiome. *ISME J* 10(1):265-8.
- Chen WM, Laevens S, Lee TM, Coenye T, De Vos P, Mergeay M, *et al.* 2001. *Ralstonia taiwanensis* sp. nov., isolated from root nodules of *Mimosa* species and sputum of a cystic fibrosis patient. *Int J Syst Evol Microbiol* 51:1729-35.
- Chen WM, James EK, Prescott AR, Kierans M, Sprent JI. 2003. Nodulation of *Mimosa* spp. by the beta-proteobacterium *Ralstonia taiwanensis*. *Mol Plant Microbe Interact* 16(12):1051-61.
- Chen YH, Gols R, Benrey B. 2015. Crop domestication and its impact on naturally selected trophic interactions. *Annu Rev Entomol* 60:35-58.
- Chen YH, Shapiro LR, Benrey B, Cibrián-Jaramillo A. 2017. Back to the origin: in situ studies are needed to understand selection during crop diversification. *Front Ecol Evol* 5:125.

- Chen H, Wu H, Yan B, Zhao H, Liu F, Zhang H, *et al.* 2018. Core microbiome of medicinal plant *Salvia miltiorrhiza* seed: A rich reservoir of beneficial microbes for secondary metabolism? *Int J Mol Sci* 19(3):pii: E672
- CIAT-PRG. 2018. <http://genebank.ciat.cgiar.org>
- Clayton JB, Vangay P, Huang H, Ward T, Hillmann BM, Al-Ghalith GA, *et al.* 2016. Captivity humanizes the primate microbiome. *Proc Natl Acad Sci U S A.* 113:10376-81.
- Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y *et al.* 2014. Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res* 42: D633-D642.
- Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, *et al.* 2016. Plant compartment and biogeography affect microbiome composition in cultivated and native *Agave* species. *New Phytol* 209(2):798-811.
- Comas LH, Becker SR, Cruz VMV, Byrne PF, Dierig DA. 2013. Root traits contributing to plant productivity under drought. *Front Plant Sci* 4:442.
- Cook RJ, Thomashow LS, Weller DM, Fujimoto D, Mazzola M, *et al.* 1995. Molecular mechanisms of defense by rhizobacteria against root disease. *Proc Natl Acad Sci U S A* 92(10):4197-4201.
- Cortés AJ, Chavarro MC, Madriñán S, This D, Blair MW. 2012. Molecular ecology and selection in the drought-related *Asr* gene polymorphisms in wild and cultivated common bean (*Phaseolus vulgaris* L.). *BMC Genet* 13:58.
- Coskun D, Britto DT, Shi W, Kronzucker HJ. 2017. How plant root exudates shape the nitrogen cycle. *Trends Plant Sci* 22(8):661-673.
- Crittenden AN, Schnorr SL. 2017. Current views on hunter-gatherer nutrition and the evolution of the human diet. *Am J Phys Anthropol* 63:84-109.
- Cui YS, Im WT, Yin CR, Lee JS, Lee KC, Lee ST. 2007. *Aeromicrobium panaciterrae* sp. nov., isolated from soil of a ginseng field in South Korea. *Int J Syst Evol Microbiol* 57, 687-91.
- Dall'Agnol RF, Plotegher F, Souza RC, Mendes IC, Dos Reis Junior FB, Béna G, *et al.* 2016. *Paraburkholderia nodosa* is the main N₂-fixing species trapped by promiscuous common bean (*Phaseolus vulgaris* L.) in the Brazilian 'Cerradão'. *FEMS Microbiol Ecol* 92(8):fiw108.
- de Bruijn I, Cheng X, de Jager V, Expósito RG, Watrous J, Patel N, *et al.* 2015. Comparative genomics and metabolic profiling of the genus *Lysobacter*. *BMC Genomics* 16:991.
- De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S. *et al.* 2010. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci USA* 107: 14691-96.
- de Weert S, Vermeiren H, Mulders IH, Kuiper I, Hendrickx N, Bloemberg GV, *et al.* 2002. Flagella-driven chemotaxis towards exudate components is an important trait for tomato root colonization by *Pseudomonas fluorescens*. *Mol Plant Microbe Interact* 15(11):1173-80.
- de Werra P, Huser A, Tabacchi R, Keel C, Maurhofer M. 2011. Plant- and microbe-derived compounds affect the expression of genes encoding antifungal compounds in a pseudomonad with biocontrol activity. *Appl Environ Microbiol* 77(8):2807-12.

- Debouck DG, Toro O, Paredes OM, Johnson WC, Gepts P. 1993. Genetic diversity and ecological distribution of *Phaseolus vulgaris* (Fabaceae) in northwestern South America. *Econ Bot* 47:408-423.
- Delgado-Baquerizo M, Maestre FT, Reich PB, Trivedi P, Osanai Y, Liu YR, *et al.* 2016. Carbon content and climate variability drive global soil bacterial diversity patterns. *Ecol Monogr* 86(3):373–390.
- Desiderio F, Bitocchi E, Bellucci E, Rau D, Rodriguez M, Attene G, *et al.* 2013. Chloroplast microsatellite diversity in *Phaseolus vulgaris*. *Front Plant Sci* 3:1-15.
- Dessureault-Rompré J, Nowack B, Schulin R, Luster Jörg. 2006. Modified micro suction cup/rhizobox approach for the in-situ detection of organic acids in rhizosphere soil solution. *Plant Soil* 286:99-107.
- Dicke M. 2009. Behavioural and community ecology of plants that cry for help. *Plant Cell Environ* 32(6):654-65.
- Dotr M, Roehr JT, Ahmed R, Dieterich C. 2012. FLEXBAR—Flexible Barcode and Adapter Processing for Next-Generation Sequencing Platforms. *Biology* 1:895–905.
- Doebley JF, Gaut BS and Smith BD. 2006. The molecular genetics of crop domestication. *Cell* 127(7):1309-1321.
- Dumbrell AJ, Nelson M, Helgason T, Dytham C, Fitter AH. 2010. Relative roles of niche and neutral process in structuring a soil microbial community. *ISME J* 4:337–345.
- Earl DA. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour* 4(2):359-361.
- Eberl L, Vandamme P. 2016. Members of the genus *Burkholderia*: good and bad guys. *F1000Res* 5(F1000 Faculty Rev):1007
- Edgar RC. 2010. Search and clustering hundreds of times faster than BLAST. *Bioinformatics* 26(19):2460-2461.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27(16): 2194-2200.
- Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, *et al.* 2015. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc Natl Acad Sci U S A* 112:E911-20.
- Eisenhauer N, Lanoue A, Strecker T, Scheu S, Steinauer K, Thakur MP, *et al.* 2017. Root biomass and exudates link plant diversity with soil bacterial and fungal biomass. *Sci Rep* 7:44641.
- Elliott GN, Chou JH, Chen WM, Bloembergen GV, Bontemps C, Martínez-Romero E. 2009. *Burkholderia* spp. are the most competitive symbionts of *Mimosa*, particularly under N-limited conditions. *Environ Microbiol* 11(4):762-78.
- Ellouze W, Hamel C, Singh AK, Mishra V, DePauw RM, Knox RE. 2018. Abundance of the arbuscular mycorrhizal fungal taxa associated with the roots and rhizosphere soil of different durum wheat cultivars in the Canadian prairies. *Can J Microbiol* 64(8):527-536.

- Engelhard M, Hurek T, Reinhold-Hurek B. 2000. Preferential occurrence of diazotrophic endophytes, *Azoarcus* spp., in wild rice species and land races of *Oryza sativa* in comparison with modern races. *Environ Microbiol* 2(2):131-41.
- Etienne RS. 2005. A new sampling formula for neutral biodiversity. *Ecol Lett* 8:253-260.
- Evanno G, Regnaut S, Goudet, J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* 14(8): 2611-2620.
- Falagán C, Foesel B, Johnson B. 2017. *Acidicapsa ferrireducens* sp. nov., *Acidicapsa acidiphila* sp. nov., and *Granulicella acidiphila* sp. nov.: novel acidobacteria isolated from metal-rich acidic waters. *Extremophiles*. 21(3):459-469.
- Faust K, Raes J. 2012. Microbial interactions: from networks to models. *Nat Rev Microbiol* 10(8):538-50.
- Fierer N. 2017. Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat Rev Microbiol* 15(10):579-590.
- Fierer N, Bradford MA, Jackson RB. 2007. Toward an ecological classification of soil bacteria. *Ecology* 88(6):1354-64.
- Fierer N, Jackson RB. 2006. The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci U S A* 103(3):626-31.
- Fierer N, Ladau J, Clemente JC, Leff JW, Owens SM, Pollard KS, *et al.* 2013. Reconstructing the microbial diversity and function of pre-agricultural tallgrass prairie soils in the United States. *Science* 342(6158):621-4.
- Ferguson, B.J, Lin MH, Gresshoff PM. 2013. Regulation of legume nodulation by acidic growth conditions. *Plant Signal Behav* 8(3):e23426.
- Flint-Garcia SA. 2013. Genetics and consequences of crop domestication. *J Agric Food Chem* 61(35):8267-76
- Friedman, J. & Alm, E. J. 2012. Inferring correlation networks from genomic survey data. *PLoS Computational Biology*. 8:e1002687.
- Garbeva P, van Veen JA, van Elsas JD. 2004. Microbial diversity in soil: Selection of microbial populations by plant and soil type and implications for disease suppressiveness. *Annu Rev Phytopathol* 42:243-70.
- García-Palacios P, Milla R, Delgado-Baquerizo M, Martín-Robles N, Alvaro-Sánchez M, Wall DH. 2013. Side-effects of plant domestication: ecosystem impacts of changes in litter quality. *New Phytol* 198(2):504-13.
- Gepts P. 2004. Crop domestication as a long-term selection experiment. *Plant Breed Rev* 24(2):1-44.
- Gepts P. 1998. Origin and evolution of common bean: past events and recent trends. *HortScience* 33(7):1124-1130.
- Gepts P, Bliss FA. 1985. F1 hybrid weakness in the common bean. *J Hered* 76:447-450.
- Gepts P, Bliss FA. 1986. Phaseolin variability among wild and cultivated common beans (*Phaseolus vulgaris*) from Colombia. *Econ Bot* 40(4):469-478.

- Gepts P, Debouck D. 1991. Origin, domestication and evolution of the common bean (*Phaseolus vulgaris* L.). In: Shoonhoven A van and Voysest O. eds. Common Beans Research for Crop Improvement. CIAT and CAB International, Wallingford, UK. pp. 27-40.
- Germida JJ, Siciliano SD. 2001. Taxonomic diversity of bacteria associated with the roots of modern, recent and ancient wheat cultivars. *Biol Fertil Soils* 33: 410–415.
- Gomes NCM, Heuer H, Schonfeld J, Costa R, Mendonca-Hagler L, Smalla K. 2001. Bacterial diversity of the rhizosphere of maize (*Zea mays*) grown in tropical soil studied by temperature gradient gel electrophoresis. *Plant Soil* 232:167–80.
- Gomez A, Petrzekova KJ, Burns MB, Yeoman CJ, Amato KR, Vickova K, *et al.* 2016. Gut microbiome of coexisting BaAka Pygmies and Bantu reflects gradients of traditional subsistence patterns. *Cell Rep* 14:2142-53.
- Gómez-Expósito R, Postma J, Raaijmakers JM, de Bruijn I. 2015. Diversity and activity of *Lysobacter* species from disease suppressive soils. *Front Microbiol* 6:1243.
- González I, Déjean S. 2012. CCA: Canonical correlation analysis. R package version 1.2. <http://CRAN.R-project.org/package=CCA>.
- Gotelli NJ, Colwell RK. 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecol Lett* 4:379-391.
- Gross BL, Olsen KM. 2010. Genetic perspectives on crop domestication. *Trends Plant Sci* 15(9):529-537.
- Haichar FZ, Marol C, Berge O, Rangel-Castro JI, Prosser JI, Balesdent J, *et al.* 2008. Plant host habitat and root exudates shape soil bacterial community structure. *ISME J* 2(12):1221-1230.
- Haichar FZ, Santaella C, Heulin T and Achouak W. 2014. Root exudates mediated interactions belowground. *Soil Bio and Biochem* 77:69-80.
- Hajjar R, Hodgkin T. 2007. The use of wild relatives in crop improvement: A survey of developments over the last 20 years. *Euphytica* 156:1-13.
- Hamonts K, Trivedi P, Garg A, Janitz C, Grinyer J, Holford P, *et al.* 2018. Field study reveals core plant microbiota and relative importance of their drivers. *Environ Microbiol* 20(1):124-140.
- Hartman K, van der Heijden MG, Roussely-Provent V, Walser JC, Schlaeppli K. 2017. Deciphering composition and function of the root microbiome of a legume plant. *Microbiome* 5:2.
- Hassan S, Mathesius U. 2012. The role of flavonoids in root-rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. *J Exp Bot* 63(9):3429-44.
- Hassani MA, Durán P, Hacquard S. 2018. Microbial interactions within the plant holobiont. *Microbiome* 6:58.
- Haudry A, Cenci A, Ravel C, Bataillon T, Brunel D, Poncet C, *et al.* 2007. Grinding up wheat: a massive loss of nucleotide diversity since domestication. *Mol Biol Evol* 24(7):1506-1517.
- Hayward AC, Fegan N, Fegan M, Stirling GR. 2010. *Stenotrophomonas* and *Lysobacter*: ubiquitous plant-associated gamma-proteobacteria of developing significance in applied microbiology. *J Appl Microbiol* 108(3):756-70.

- Henry S, Texier S, Hallet S, Bru D, Dambreville C, Chèneby D, *et al.* 2008. Disentangling the rhizosphere effect on nitrate reducers and denitrifiers: insight into the role of root exudates. *Environ Microbiol* 10(11):3082-92.
- Herrera-Flores TS, Acosta-Gallegos JA. 2008. Seed yield of three types of crosses between wild by domesticated genotypes in common bean (*Phaseolus vulgaris* L). *Agricultura Técnica en México* 34(2):167-176.
- Hetrick BAD, Wilson GWT, Cox TS. 1992. Mycorrhizal dependence of modern wheat varieties, landraces, and ancestors. *Can J Bot* 70(10):2032-2040.
- Hetrick BAD, Wilson GWT, Cox TS. 1993. Mycorrhizal dependence of modern wheat cultivars and ancestors: a synthesis. *Can J Bot* 71(3):512-518.
- Hetrick BAD, Wilson GWT, Gill BS, Cox TS. 1995. Chromosome location of mycorrhizal responsive genes in wheat. *Can J Bot* 73:891-897.
- Ho A, Di Lonardo DP, Bodelier PLE. 2017. Revisiting life strategy concepts in environmental microbial ecology. *FEMS Microbiol Ecol* 93(3): fix006.
- Horton MW, Bodenhausen N, Beilsmith K, Meng D, Muegge BD, Subramanian S, *et al.* 2014. Genome-wide association study of Arabidopsis thaliana leaf microbial community. *Nat Commun* 5:5320.
- Iannucci A, Fragasso M, Beleggia R, Nigro F, Papa R. 2017. Evolution of the crop rhizosphere: impact of domestication on root exudates in tetraploid wheat (*Triticum turgidum* L.). *Front Plant Sci* 8:2124.
- Illumina. 2013. 16S Metagenomic sequencing library preparation. http://support.illumina.com/downloads/16s_metagenomic_sequencing_library_preparation.html
- Inceoglu Ö, Abu Al-Soud W, Salles JF, Semenov AV and van Elsas JD. 2011. Comparative analysis of bacterial communities in a potato field as determined by pyrosequencing. *PLoS One* 6(8):e23321.
- Inceoglu Ö, Falcão Salles J, van Overbeek L and van Elsas JD. 2010. Effects of plant genotype and growth stage on the betaproteobacterial communities associated with different potato cultivars in two fields. *Appl Environ Microbiol* 76(11):3675-3684.
- Jackson, LE. 1995. Root architecture in cultivated and wild lettuce (*Lactuca* spp). *Plant Cell Environ* 18:885–894.
- Jabot F, Etienne RF, Chave J. 2008. Reconciling neutral community models and environmental filtering: theory and an empirical test. *Oikos* 117:1308–1320.
- Jaccoud D, Peng K, Feinstein D, Kilian A. 2001. Diversity arrays: a solid state technology for sequence information independent genotyping. *Nucleic Acids Res* 29(4):e25.
- Judy C, Adrian M, Baussard C, Bernard C, Bernaud E, Bourion V, *et al.* 2016. RhizoTubes as a new tool for high throughput imaging of plant root development and architecture: test, comparison with pot grown plants and validation. *Plant Methods* 12:31.
- Jin J, Wang GH, Liu XB, Liu JD, Chen XL and Herbert SJ. 2009. Temporal and spatial dynamics of bacterial community in the rhizosphere of soybean genotypes grown in a black soil. *Pedosphere* 19(6):808-816.

- Johnson WC, Jackson LE, Ochoa O, van Wijk R, Peleman J, St.Clair DA, Michelmore RW. 2000. Lettuce, a shallow-rooted crop, and *Lactuca serriola*, its wild progenitor, differ at QTL determining root architecture and deep soil water exploitation. *Theor Appl Genet* 101:1066–1073.
- Jones DL, Hodge A and Kuzyakov Y. 2004. Plant and mycorrhizal regulation of rhizodeposition. *New Phytol* 163:459–480.
- Jones DL, Nguyen C, Finlay RD. 2009. Carbon flow in the rhizosphere: carbon trading at the soil–root interface. *Plant Soil* 321:5–33.
- Jones RT, Robeson MS, Lauber CL, Hamady M, Knight R, Fierer N. 2009. A comprehensive survey of soil acidobacterial diversity using pyrosequencing and clone library analyses. *ISME J* 3(4):442–53.
- Joshi NA, Fass JN. 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files. Version 1.33. Available at <https://github.com/najoshi/sickle>
- Jousset A, Rochat L, Lanoue A, Bonkowski M, Keel C, Scheu S. 2011. Plants respond to pathogen infection by enhancing the antifungal gene expression of root-associated bacteria. *Mol Plant Microbe Interact* 24(3):352–8.
- Kageyama K, Nelson EB. 2003. Differential inactivation of seed exudate stimulation of *Pythium ultimum* sporangium germination by *Enterobacter cloacae* influences biological control efficacy on different plant species. *Appl Environ Microbiol* 69(2):1114–20
- Kapulnik Y, Kushnir U. 1991. Growth dependency of wild, primitive and modern cultivated wheat lines on vesicular-arbuscular mycorrhiza fungi. *Euphytica* 56:27–36.
- Kawasaki A, Donn S, Ryan PR, Mathesius U, Devilla R, Jones A, *et al.* 2016. Microbiome and exudates of the root and rhizosphere of *Brachypodium distachyon*, a model for wheat. *PLoS One* 11(10):e0164533.
- Kielak AM, Barreto CC, Kowalchuk GA, van Veen JA, Kuramae EE. 2016. The ecology of acidobacteria: Moving beyond genes and genomes. *Front Microbiol* 7:744.
- Kiers ET, Denison RF. 2008. Sanctions, cooperation, and the stability of plant-rhizosphere mutualisms. *Annu Rev Ecol Evol Syst* 39:215–236.
- Kiers ET, Hutton MG, Denison RF. 2007. Human selection and the relaxation of legume defences against ineffective rhizobia. *Proc Biol Sci* 274(1629):3119–26.
- Kim DH, Kaashyap M, Rathore A, Das RR, Parupalli S, Upadhyaya HD, *et al.* 2014. Phylogenetic diversity of *Mesorhizobium* in chickpea. *J Biosci* 39(3):513–7.
- Klaedtke S, Jacques MA, Raggi L, Préveaux A, Bonneau S, Negri V, *et al.* 2016. Terroir is a key driver of seed-associated microbial assemblages. *Environ Microbiol* 18(6):1792–804.
- Koide R, Li M, Lewis J, Irby C. 1988. Role of mycorrhizal infection in the growth and reproduction of wild vs. cultivated plants. I. Wild vs. cultivated oats. *Oecologia* 77:537–543.
- Köllner TG, Held M, Lenk C, Hiltbold I, Turlings TC, Gershenzon J, Degenhardt J. 2008. A maize (E)-beta-caryophyllene synthase implicated in indirect defense responses against herbivores is not expressed in most American maize varieties. *Plant Cell* 20(2):482–94.
- Kolton M, Green SJ, Harel YM, Sela N, Elad Y, Cytryn E. 2012. Draft genome sequence of *Flavobacterium* sp. strain F52, isolated from the rhizosphere of bell pepper (*Capsicum annuum* L. cv. Maccabi). *J Bacteriol* 194:5462–63.

- Koomnok C, Teamroong N, Rerkasem B, Lumyong S. 2007. Diazotroph endophytic bacteria in cultivated and wild rice in Thailand. *ScienceAsia* 33:429-435.
- Köster J, Rahmann S. 2012. Snakemake-a scalable bioinformatics workflow engine. *Bioinformatics* 28(19): 2520–2522.
- Kraft P, Zeggini E, Ioannidis JPA. 2009. Replication in genome-wide association studies. *Stat Sci* 24: 561–573.
- Kramer-Walter KR, Bellingham PJ, Millar TR, Smissen RD, Richardson SJ, Laughlin DC. 2016. Root traits are multidimensional: specific root length is independent from root tissue density and the plant economic spectrum. *J Ecol* 104:1299-1310.
- Kroll S, Agler MT, Kemen E. 2017. Genomic dissection of host-microbe and microbe-microbe interactions for advanced plant breeding. *Curr Opin Plant Biol* 36:71-78.
- Kuczynski J, Stombaugh J, Walters WA, González A, Caporaso JG, Knight R. 2012. Using QIIME to analyze 16S rRNA gene sequences from microbial communities. *Curr Protoc Chem Biol* 1E:5.1–20.
- Kulichevskaya IS, Kostina LA, Valásková V, Rijpstra WI, Damsté JS, de Boer W, *et al.* 2012. *Acidicapsa borealis* gen. nov., sp. nov. and *Acidicapsa ligni* sp. nov., subdivision 1 *Acidobacteria* from *Sphagnum* peat and decaying wood. *Int J Syst Evol Microbiol* 62:1512-20.
- Kuramae EE, Yergeau E, Wong LC, Pijl AS, van Veen JA, Kowalchuk GA. 2012. Soil characteristics more strongly influence soil bacterial communities than land-use type. *FEMS Microbiol Ecol* 79(1):12-24.
- Kwak YS, Bonsall RF, Okubara PA, Paulitz TC, Thomashow LS, *et al.* 2012. Factors impacting the activity of 2, 4-diacetylphloroglucinol-producing *Pseudomonas fluorescens* against take-all of wheat. *Soil Biol Biochem* 54:48–56.
- Lakshmanan V, Kitto SL, Caplan JL, Hsueh YH, Kearns DB, Wu YS *et al.* 2012. Microbe-associated molecular patterns-triggered root responses mediate beneficial rhizobacterial recruitment in *Arabidopsis*. *Plant Physiol* 160:1642-1661.
- Lakshmanan V, Selvaraj G, Bais HP. 2014. Functional soil microbiome: belowground solutions to an aboveground problem. *Plant Physiol* 166(2):689-700.
- Janfranco L, Fiorilli V, Venice F, Bonfante P. 2018. Strigolactones cross the kingdoms: plants, fungi, and bacteria in the arbuscular mycorrhizal symbiosis. *J Exp Bot* 69(9):2175-2188.
- Lareen A, Burton F, Schäfer P. 2016. Plant root-microbe communication in shaping root microbiomes. *Plant Mol Bio* 90(6):575-87.
- Larsson J. 2017. eulerr: Area-Proportional Euler and Venn Diagrams with Circles or Ellipses R package version 3.0.1. <https://cran.r-project.org/package=eulerr>.
- Lauber CL, Hamady M, Knight R, Fierer N. 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microbiol* 75(15):5111-20.
- Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, *et al.* 2015. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* 349:860-864.

- Leff JW, Lynch RC, Kane NC, Fierer N. 2017. Plant domestication and the assembly of bacterial and fungal communities associated with strains of the common sunflower, *Helianthus annuus*. *New Phytol* 214(1):412-423.
- Lehmann A, Barto EK, Powell JR, Rillig MC. 2012. Mycorrhizal responsiveness trends in annual crop plants and their wild relatives-a meta-analysis on studies from 1981 to 2010. *Plant Soil* 355:231–250.
- Lemaire B, Chimphango SBM, Stirton C, Rafudeen S, Honnay O, Smets E. *et al.* 2016. Biogeographical patterns of legume-nodulating *Burkholderia* spp.: from African fynbos to continental scales. *Appl Environ Microbiol* 82(17), 5099-115.
- Lemanceau P, Barret M, Mazurier S, Mondy S, Pivato B, Fort T, *et al.* 2017. Plant communication with associated microbiota in the spermosphere, rhizosphere and phyllosphere. *Adv Bot Res* 82:101-133.
- Lemanceau, P., Blouin, M., Muller, D. & Moëne-Loccoz, Y. 2017. Let the core microbiota be functional. *Trends in Plant Science*. 22(7), 583-595.
- Letunic I, Bork P. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res* 44(Web Server issue):W242–W245.
- Levy A, Salas-Gonzalez I, Mittelviehhaus M, Clingenpeel S, Herrera Paredes S, Miao J, *et al.* Genomic features of bacterial adaptation to plants. *Nat Genet* 50(1):138-150.
- Ley RE, Peterson DA, Gordon JI. 2006. Ecological and evolutionary forces shaping microbial diversity in the human intestine. *Cell* 124(4):837-48.
- Ley RE, Turnbaugh PJ, Klein S, Gordon JI. 2006. Microbial ecology: human gut microbes associated with obesity. *Nature* 444:1022-23.
- Li M, Li D, Tang Y, Wu F, Wang J. 2017. CytoCluster: A Cytoscape Plugin for Cluster Analysis and Visualization of Biological Networks. *Int J Mol Sci* 18(9):1880.
- Ling N, Raza W, Ma J, Huang Q, Shen Q. 2011. Identification and role of organic acids in watermelon root exudates for recruiting *Paenibacillus polymyxa* SQR-21 in the rhizosphere. *Eur J Soil Biol* 47:374–379.
- Liu R, Hong J, Xu X, Feng Q, Zhang D, Gu YR, *et al.* 2017. Gut microbiome and serum metabolome alterations in obesity and after weight-loss intervention *Nat Med* 23:859-68.
- Liu CW, Murray JD. 2016. The role of flavonoids in nodulation host-range specificity: An update. *Plants* 5(3):33.
- López-López A, Rogel MA, Ormeño-Orrillo E, Martínez-Romero J, Martínez-Romero E. 2010. *Phaseolus vulgaris* seed-borne endophytic community with novel bacterial species such as *Rhizobium endophyticum* sp. nov. *Syst Appl Microbiol* 33(6):322-7.
- López-Velasco G, Carder PA, Welbaum GE, Ponder MA. 2013. Diversity of the spinach (*Spinacia oleracea*) spermosphere and phyllosphere bacterial communities. *FEMS Microbiol Lett* 346(2):146-54
- Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S *et al.* 2012. Defining the core *Arabidopsis thaliana* root microbiome. *Nature* 488: 86-90.

- Manteca A, Sanchez J. 2009. *Streptomyces* development in colonies and soils. *Appl Environ Microbiol* 75(9):2920-4.
- Marques JM, da Silva TF, Vollu RE, Blank AF, Ding GC, Seldin L, *et al.* 2014. Plant age and genotype affect the bacterial community composition in the tuber rhizosphere of field-grown sweet potato plants. *FEMS Microbiol Ecol* 88(2):424-35.
- Marschner P, Neumann G, Kania A, Weiskopf L, Lieberei R. 2002. Spatial and temporal dynamics of the microbial community structure in the rhizosphere of cluster roots of white lupin (*Lupinus albus* L.). *Plant Soil* 246: 167-164.
- Martin FM, Uroz S, Barker DG. 2017. Ancestral alliances: Plant mutualistic symbioses with fungi and bacteria. *Science* 356(6340):eaad4501.
- Martin-Robles N, Morente-López J, Milla R. 2015. Digging up the roots of crop evolution. Libro de resúmenes de comunicaciones del 4º Congreso Ibérico de Ecología, Coimbra, Portugal. doi: 10.7818/4IberianEcologicalCongress.
- Martínez-Romero E, Segovia L, Mercante FM, Franco AA, Graham P, Pardo MA. 1991. *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. *Int J Syst Bacteriol* 41(3):417-26.
- Matson PA, Parton WJ, Power AG, Swift MJ. 1997. Agricultural intensification and ecosystem properties. *Science* 277(5325):504-509.
- Masella AP, Bartram AK, Truszkowski JM, Brown DG, Neufeld JD. 2012. PANDAseq: paired-end assembler for illumina sequences. *BMC Bioinformatics* 13: 31.
- Mazzola M, Funnell DL, Raaijmakers JM. 2004. Wheat cultivar-specific selection of 2,4-diacetylphloroglucinol-producing fluorescent pseudomonas species from resident soil populations. *Microbiol Ecol* 48: 338–348.
- McDonald D, Clemente JC, Kuczynski J, Rideout JR, Stombaugh J, Wendel D *et al.* 2012. The Biological Observation Matrix. BIOM. format or: how I learned to stop worrying and love the ome-ome. *Gigascience* 1: 7.
- McKenzie VJ, Song SJ, Delsuc F, Prest TL, Oliverio AM, Korpita TM, *et al.* 2017. The effects of captivity on the mammalian gut microbiome. *Integr Comp Biol.* 57:690-704.
- McMurdie PJ, Holmes S. 2013. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 8(4):e61217.
- McNear Jr DH. 2013. The Rhizosphere - Roots, soil and everything in between. *Nature Education Knowledge* 4(3):1.
- Mendes LW, Kuramae EE, Navarrete AA, van Veen JA, Tsai SM. 2014. Taxonomical and functional microbial community selection in soybean rhizosphere. *ISME J* 8:1577–1587.
- Mendes LW, Raaijmakers JM, de Hollander M, Mendes R, Tsai SM. 2018. Influence of resistance breeding in common bean on rhizosphere microbiome composition and function. *ISME J* 12(1):212-224.
- Mendes R, Garbeva P, Raaijmakers JM. 2013. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic and human pathogenic microorganisms. *FEMS Microbiol Rev* 37:634–663.

- Mendes R, Kruijt M, de Bruijn I, Dekkers E, van der Voort M, Schneider JHM *et al.* 2011. Deciphering the Rhizosphere Microbiome for Disease-Suppressive Bacteria. *Science* 332 (6033):1097-1100.
- Meyer RS, DuVal AE, Jensen HR. 2012. Patterns and processes in crop domestication: an historical review and quantitative analysis of 203 global food crops. *New Phytol* 196:29-48.
- Micallef SA, Channer S, Shiaris MP, Colón-Carmona A. 2009a. Plant age and genotype impact the progression of bacterial community succession in the *Arabidopsis* rhizosphere. *Plant Signal Behav* 4(8):777-80.
- Micallef SA, Shiaris MP and Colón-Carmona A. 2009b. Influence of *Arabidopsis thaliana* accessions on rhizobacterial communities and natural variation in root exudates. *J Exp Bot* 60(6):1729-1742.
- Miller ES, Woese CR, Brenner S. 1991. Description of the erythromycin-producing bacterium *Arthrobacter* sp. strain NRRL B-3381 as *Aeromicrobium erythreum* gen. nov., sp. nov. *Int J Syst Bacteriol* 41(3):363-8.
- Moe LA. 2013. Amino acids in the rhizosphere: From plants to microbes. *Am J Bot* 100(9):1692–1705.
- Mönchgesang S, Strehmel N, Schmidt S, Westphal L, Taruttis F, Müller E, *et al.* 2016. Natural variation of root exudates in *Arabidopsis thaliana*-linking metabolomic and genomic data. *Sci Rep* 6:29033.
- Moulin L, Munive A, Dreyfus B, Boivin-Masson C. 2001. Nodulation of legumes by members of the beta-subclass of Proteobacteria. *Nature* 411(6840):948-50.
- Mounier E, Hallet S, Chêneby D, Benizri E, Gruet Y, Nguyen C, *et al.* 2004. Influence of maize mucilage on the diversity and activity of the denitrifying community. *Environ Microbiol* 6(3):301-12.
- Mueller UG, Sachs JL. 2015. Engineering microbiomes to improve plant and animal health. *Trends Microbiol* 23:606-17.
- Mummey DL, Rillig MC. 2006. The invasive plant species *Centaurea maculosa* alters arbuscular mycorrhizal fungal communities in the field. *Plant Soil* 288:81–90.
- Mutch LA, Young JP. 2004. Diversity and specificity of *Rhizobium leguminosarum* biovar *viciae* on wild and cultivated legumes. *Mol Ecol* 13(8):2435-44.
- Naz AA, Arifuzzaman M, Muzammil S, Pillen K, Léon J. 2014. Wild barley introgression lines revealed novel QTL alleles for root and related shoot traits in the cultivated barley (*Hordeum vulgare* L.) *BMC Genetics* 15:107.
- Naz AA, Ehl A, Pillen K, Léon J. 2012. Validation for root-related quantitative trait locus effects of wild origin in the cultivated background of barley (*Hordeum vulgare* L.). *Plant Breed* 3:392-398.
- Neal AL, Ahmad S, Gordon-Weeks R, Ton J. 2012. Benzoxazinoids in root exudates of maize attract *Pseudomonas putida* to the rhizosphere. *PLoS One* 7(4):e35498.
- Nelson EB. 1990. Exudate molecules initiating fungal responses to seeds and roots. *Plant Soil* 129:61-73.

- Nelson EB. 2004. Microbial dynamics and interactions in the spermosphere. *Annu Rev Phytopathol* 42:271-309.
- Nelson EB. 2018. The seed microbiome: Origins, interactions, and impacts. *Plant Soil* 422:7-24.
- Nevo E, Chen G. 2010. Drought and salt tolerances in wild relatives for wheat and barley improvement. *Plant Cell Environ* 33:670–685.
- Ofek M, Hadar Y, Minz D. 2012. Ecology of root colonizing *Massilia* (Oxalobacteraceae). *PLoS One* 7(7):e40117
- Ofek M, Voronov-Goldman M, Hadar Y, Minz D.. 2014. Host signature effect on plant root-associated microbiomes revealed through analyses of resident vs. active communities. *Environ Microbiol* 16:2157-2167.
- Ofek-Lazar M, Sela N, Goldman-Voronov M, Green SJ, Hadar Y, *et al.* 2014. Niche and host-associated functional signatures of the root surface microbiome. *Nat commun* 5:4950.
- Oksanen J, Kindt R, Legendre P, O’Hara B, Simpson GL, Solymos P *et al.* 2016, 2017. vegan: Community ecology package. R package versions 2.4-0 and 2.4-4. <http://CRAN.R-project.org/package=vegan>
- Olsen KM, Caicedo AL, Polato N, McClung A, McCouch S, Purugganan MD. 2006. Selection under domestication: evidence for a sweep in the rice waxy genomic region. *Genetics* 173(2):975-83.
- Palaisa K, Morgante M, Tingey S, Rafalski A. 2004. Long-range patterns of diversity and linkage disequilibrium surrounding the maize Y1 gene are indicative of an asymmetric selective sweep. *Proc Natl Acad Sci USA* 101(26):9885-90.
- Parks DH, Tyson GW, Hugenholtz P, Beiko RG. 2014. STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics* 30(21):3123-4.
- Paulson JN, Stine OC, Bravo HC, Pop M. 2013. Differential abundance analysis for microbial marker-gene surveys. *Nat Methods* 10:1200–1202.
- Paulson JN, Talukder H, Pop M, Bravo HC. 2016. metagenomeSeq: Statistical analysis for sparse high-throughput sequencing. Bioconductor package: 1.12.0. <http://cceb.umd.edu/software/metagenomeSeq>
- Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL *et al.* 2013. Diversity and heritability of the maize rhizosphere microbiome under field conditions. *Proc Natl Acad Sci U S A* 110(16): 6548–6553.
- Pérez-Jaramillo JE, Carrión VJ, Bosse M, Ferrão LFV, de Hollander M, Garcia AAF, *et al.* 2017. Linking rhizosphere microbiome composition of wild and domesticated *Phaseolus vulgaris* to genotypic and root phenotypic traits. *ISME J* 11:2244–57.
- Pérez-Jaramillo JE, Carrión VJ, de Hollander M, Raaijmakers JM. 2018. The wild side of plant microbiomes. *Microbiome* 6:143.
- Pérez-Jaramillo JE, Mendes R, Raaijmakers JM. 2016. Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant Mol Biol* 90: 635–644.
- Petersen DJ, Srinivasan M, Chanway CP. 1996. *Bacillus polymyxa* stimulates increased *Rhizobium etli* populations and nodulation when co-resident in the rhizosphere of *Phaseolus vulgaris*. *FEMS Microbiol Lett* 142(2-3):271-6.

- Pfeiffer S, Mitter B, Oswald A, Schloter-Hai B, Schloter M, Declerck S. 2017. Rhizosphere microbiomes of potato cultivated in the High Andes show stable and dynamic core microbiomes with different responses to plant development. *FEMS Microbiol Ecol* 93(2): fiw242.
- Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH. 2013. Going back to the roots: the microbial ecology of the rhizosphere. *Nature Rev Microbiol* 11: 789-799.
- Placido DF, Campbell MT, Folsom JJ, Cui X, Kruger GR, Baenziger PS, Harkamal WH. 2013. Introgression of novel traits from a wild wheat relative improves drought adaptation in wheat. *Plant Physiol* 161(4):1806–1819.
- Pongsilp N, Nimnoi P and Lumyong S. 2012. Genotypic diversity among rhizospheric bacteria of three legumes assessed by cultivation-dependent and cultivation-independent techniques. *World J Microbiol Biotechnol* 28(2):615-26.
- Porch TG, Beaver JS, Debouck DG, Jackson SA, Kelly JD. 2013. Use of wild relatives and closely related species to adapt common bean to climate change. *Agronomy* 3:433-461.
- Postma J, Schilder MT, van Hoof RA. 2011. Indigenous populations of three closely related *Lysobacter* spp. in agricultural soils using real-time PCR. *Microb Ecol* 62(4): 948-58.
- Price MN, Dehal PS, Arkin AP. 2009. FastTree: Computing large minimum-evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* 26:1641-1650.
- Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155(2):945-959.
- Puopolo G, Tomada S, Pertot I. 2018. The impact of the omics era on the knowledge and use of *Lysobacter* species to control phytopathogenic micro-organisms. *J Appl Microbiol* 124(1):15-27.
- Purugganan MD, Fuller DQ. 2009. The nature of selection during plant domestication. *Nature* 457:843-848.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, *et al.* 2013. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* 41:D590–D596.
- R Core Team. 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Raaijmakers JM, Paulitz TC, Steinberg C, Alabouvette C, Moënne-Loccoz Y. 2009. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant Soil* 321:341-361.
- Raboin, L-M, Razafimahafaly AHD, Rabenjarisoa MB, Rabary B, Deuserre J, Becquer T. 2016. Improving the fertility of tropical acid soils: liming versus biochar application? A long term comparison in the highlands of Madagascar. *Field Crops Res* 199:99–108.
- Ram SG, Thiruvengadam V, Vinod KK. 2007. Genetic diversity among cultivars, landraces and wild relatives of rice as revealed by microsatellite markers. *J Appl Genet* 48(4):337-45.
- Ramirez KS, Craine JM, Fierer N. 2012. Consistent effects of nitrogen amendments on soil microbial communities and processes across biomes. *Glob Chang Biol* 18:1918-1927.

- Ramirez KS, Knight CG, de Hollander M, Brearley FQ, Constantinides B, *et al.* 2018. Detecting macroecological patterns in bacterial communities across independent studies of global soils. *Nat Microbiol* 3(2):189-196.
- Rasmann S, Köllner TG, Degenhardt J, Hiltbold I, Toepfer S, Kuhlmann U, *et al.* 2005. Recruitment of entomopathogenic nematodes by insect-damaged maize roots. *Nature* 434(7034):732-737.
- Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W *et al.* 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 43(7):e47.
- Rodrigues JL, Pellizari VH, Mueller R, Baek K, Jesus Eda C, Paula FS, *et al.* 2013. Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proc Natl Acad Sci U S A* 110(3):988-993.
- Rodrigues JL, Pellizari VH, Mueller R, Baek K, Jesus Eda C, Paula FS, *et al.* 2013. Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proc Natl Acad Sci U S A* 110(3):988-993.
- Rognes T, Mahé F, Flouris T, Quince C, Nichols B. 2015. VSEARCH version 1.0.16. <https://github.com/torognes/vsearch>
- Rohrbacher F, St-Arnaud M. 2016. Root exudation: The ecological driver of hydrocarbon rhizoremediation. *Agronomy* 6(1):19.
- Rudrappa T, Czymmek KJ, Paré PW, Bais HP. 2008. Root-secreted malic acid recruits beneficial soil bacteria. *Plant Physiol* 148(3):1547-56.
- Ruiz-Pérez CA, Restrepo S, Zambrano MM. 2016. Microbial and functional diversity within the phyllosphere of Espeletia species in an Andean high-mountain ecosystem. *Appl Environ Microbiol* 82:1807–1817.
- Rybakova D, Mancinelli R, Wikström M, Birch-Jensen AS, Postma J, Ehlers RU, *et al.* 2017. The structure of the *Brassica napus* seed microbiome is cultivar-dependent and affects the interactions of symbionts and pathogens. *Microbiome* 5(1):104.
- Saleem M, Law AD, Sahib MR, Pervaiz ZH, Zhang Q. 2018. Impact of root system architecture on rhizosphere and root microbiome. *Rhizosphere* 6:47-51.
- Sánchez PA, Logan JT. 1992. Myths and science about the chemistry and fertility of soils in the tropics. In: Lal R & Sanchez PA. Eds. Myths and Science of Soils in the Tropics, SSSA Special Publication 29. ASA and SSSA, Madison, WI, USA.
- Sangabriel-Conde W, Maldonado-Mendoza IE, Mancera-López ME, Cordero-Ramírez JD, Trejo-Aguilar D, Negrete-Yankelevich S. 2015. Glomeromycota associated with Mexican native maize landraces in Los Tuxtlas, Mexico. *Appl Soil Ecol* 87:63–71.
- Sangabriel-Conde W, Negrete-Yankelevich S, Maldonado-Mendoza IE, Trejo-Aguilar D. 2014. Native maize landraces from Los Tuxtlas, Mexico show varying mycorrhizal dependency for P uptake. *Biol Fertil Soils* 50:405–414.
- Sasse J, Martinoia E, Northen T. 2018. Feed your friends: Do plant exudates shape the root microbiome? *Trends Plant Sci* 23:25-41.
- Schiltz S, Gaillard I, Pawlicki-Jullian N, Thiombiano B, Mesnard F, Gontier E. 2015. A review: what is the spermosphere and how can it be studied? *J Appl Microbiol* 119(6):1467-81.

- Schlaeppli K, Dombrowski N, Oter RG, van Themaat EVL, Schulze-Lefert P. 2014. Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana* relatives. *Proc Natl Acad Sci U S A* 111:585–592.
- Schlub RL, Schmitthenner AF. 1978. Effects of soybean seed coat cracks on seed exudation and seedling quality in soil infested with *Pythium ultimum*. *Phytopathology* 68:1186-1191.
- Schmalenbach I, Körber N, Pillen K. 2008. Selecting a set of wild barley introgression lines and verification of QTL effects for resistance to powdery mildew and leaf rust. *Theor Appl Genet* 117(7):1093–1106.
- Schnorr SL, Candela M, Rampelli S, Centanni M, Consolandi C, Basaglia G, *et al.* 2014. Gut microbiome of the Hadza hunter-gatherers. *Nature Commun* 5:3654.
- Schnorr SL, Sankaranarayanan K, Lewis CM Jr, Warinner C. 2016. Insights into human evolution from ancient and contemporary microbiome studies. *Curr Opin Genet Dev* 41:14-26.
- Schultze M, Kondorosi A. 1998. Regulation of symbiotic root nodule development. *Annu Rev Genet* 32:33-57.
- Schulz-Bohm K, Gerards S, Hundscheid M, Melenhorst J, de Boer W, Garbeva P. 2018. Calling from distance: attraction of soil bacteria by plant root volatiles. *ISME J* 12(5):1252-1262.
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, *et al.* 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Research* 13(11):2498-504.
- Shaposhnikov AI, Morgounov AI, Akin B, Makarova NM, Belimov AA, Tikhonovich IA. Comparative characteristics of root systems and root exudation of synthetic, landrace and modern wheat varieties. *Agricultural Biology*. 2016;51:68-78.
- Shenton M, Iwamoto C, Kurata N, Ikeo K. 2016. Effect of wild and cultivated rice genotypes on rhizosphere bacterial community composition. *Rice* 9(1):42.
- Shi S, Nuccio E, Herman DJ, Rijkers R, Estera K, Li J, *et al.* 2015. Successional trajectories of rhizosphere bacterial communities over consecutive seasons. *mBio* 6:e00746.
- Shi S, Richardson AE, O'Callaghan M, DeAngelis KM, Jones EE, Stewart A, *et al.* 2011. Effects of selected root exudate components on soil bacterial communities. *FEMS Microbiol Ecol* 77(3):600-10.
- Short GE, Lacy ML. 1976. Carbohydrate exudation from pea seeds: Effect of cultivar, seed age, seed color, and temperature. *Phytopathology* 66:182-187.
- Simko I, Atallah AJ, Ochoa OE, Antonise R, Galeano CH, Truco MJ, Michelmore MW. 2013. Identification of QTLs conferring resistance to downy mildew in legacy cultivars of lettuce. *Sci Rep* 3:2875.
- Simon HM, Smith KP, Dodsworth JA, Guenther B, Handelsman J, Goodman RM. 2001. Influence of tomato genotype on growth of inoculated and indigenous bacteria in the spermosphere. *Appl Environ Microbiol* 67(2):514-20.
- Singh SP. 2001. Broadening the genetic base of common bean cultivars: a review. *Crop Sci* 41:1659–1675.

- Singh P, Mehrotra RS. 1980. The influence of cultivar and temperature on carbohydrate and amino acid exudation from gram seeds and on pre-emergence damping-off by *Rhizoctonia bataticola*. *Plant Soil* 55:261-268.
- Smalla K, Wieland G, Buchner A, Zock A, Parzy J, Kaiser S, Roskot N, Heuer H and Berg G. 2001. Bulk and rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis: plant-dependent enrichment and seasonal shifts revealed. *Appl Environ Microbiol* 67(10):4742-4751.
- Smith KP, Handelsman J, Goodman RM. 1997. Modeling dose-response relationships in biological control: Partitioning host responses to the pathogen and biocontrol agent. *Phytopathology* 87(7):720-729.
- Smith KP, Goodman RM. 1999. Host variation for interactions with beneficial plant-associated microbes. *Annu Rev Phytopathol* 37:473-491.
- Smykal P, Nelson MN, Berger JD, von Wettberg EJ. 2018. The impact of genetic changes during crop domestication. *Agronomy* 8:119.
- Sonnante G, Stockton T, Nodari RO, Becerra-Velásquez VL, Gepts P. 1994. Evolution of genetic diversity during the domestication of common bean (*Phaseolus vulgaris* L.). *Theor Appl Genet* 89:629-635.
- Stopnisek N, Bodenhausen N, Frey B, Fierer N, Eberl L, Weisskopf L. 2014. Genus-wide acid tolerance accounts for the biogeographical distribution of soil *Burkholderia* populations. *Environ Microbiol* 16(6):1503-12.
- Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, *et al.* 2018. MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. *Proc Natl Acad Sci U S A* 115(22):E5213-E5222.
- Sugiyama A, Ueda Y, Takase H and Yazaki K. 2014a. Pyrosequencing assessment of rhizosphere fungal communities from a soybean field. *Can J Microbiol* 60(10):687-690.
- Sugiyama A, Ueda Y, Zushi T, Takase H and Yazaki K. 2014b. Changes in the bacterial community of soybean rhizospheres during growth in the field. *PLoS One* 9(6): e100709.
- Szoboszlay M, Lambers J, Chappell J, Kupper JV, Moe LA, McNear Jr DH. 2015. Comparison of root system architecture and rhizosphere microbial communities of Balsas teosinte and domesticated corn cultivars. *Soil Bio and Biochem* 80:34-44.
- Talbi C, Delgado MJ, Girard L, Ramírez-Trujillo A, Caballero-Mellado J, Bedmar EJ. 2010. *Burkholderia phymatum* strains capable of nodulating *Phaseolus vulgaris* are present in Moroccan soils. *Appl Environ Microbiol* 76(13):4587-91.
- Tardy V, Chabbi A, Charrier X, de Berranger C, Reignier T, Dequiedt S, *et al.* 2015. Land use history shifts in situ fungal and bacterial successions following wheat straw input into the soil. *PLoS One* 10(6):e0130672.
- Thomas F, Hehemann JH, Rebuffet E, Czejzek M, Michel G. 2011. Environmental and gut bacteroidetes: The food connection. *Front Microbiol* 2:93.
- Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, *et al.* 2017. A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* 551(7681):457-463.

- Tian L, Zhou X, Ma L, Xu S, Nasir F, Tian C. 2017. Root-associated bacterial diversities of *Oryza rufipogon* and *Oryza sativa* and their influencing environmental factors. *Arch Microbiol* 199(4):563-571.
- Toju H, Peay KG, Yamamichi M, Narisawa K, Hiruma K, Naito K, *et al.* 2018. Core microbiomes for sustainable agroecosystems. *Nat Plants* 4(5):247-257.
- Toro O, Thome J, Debouck DG. 1990. Wild bean (*Phaseolus vulgaris* L.): Description and distribution. Centro Internacional de Agricultura Tropical (CIAT); International Board for Plant Genetic Resources (IBPGR), Cali, Colombia. 106 pp. ISBN: 958-9183-22-0.
- Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. 2006. An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 444(7122):1027-1031.
- Turner TR, James EK, Poole PS. 2013a. The plant microbiome. *Genome Biol* 14:209.
- Turner TR, Ramakrishnan K, Walshaw J, Heavens D, Alston M, Swarbreck D, *et al.* 2013b. Comparative metatranscriptomics reveals kingdom level changes in the rhizosphere microbiome of plants. *ISME J* 7(12):2248-2258.
- Uksa M, Schloter M, Endesfelder D, Kublik S, Engel M, Kautz T, *et al.* 2015. Prokaryotes in subsoil-evidence for a strong spatial separation of different phyla by analysing co-occurrence networks. *Front Microbiol* 6:1269.
- Vandenkoornhuysen P, Quaiser A, Duhamel M, Le Van A, Dufresne A. 2015. The importance of the microbiome of the plant holobiont. *New Phytol* 206(4):1196-206.
- Viaene T, Langendries S, Beirinckx S, Maes M, Goormachtig S. 2016. Streptomyces as a plant's best friend? *FEMS Microbiol Ecol* 92(8):pii: fiw119
- Wallenstein MD. 2017. Managing and manipulating the rhizosphere microbiome for plant health: A systems approach. *Rhizosphere* 3:230-232.
- Walters WA, Jin Z, Youngblut N, Wallace JG, Sutter J, Zhang W, *et al.* 2018. Large-scale replicated field study of maize rhizosphere identifies heritable microbes. *Proc Natl Acad Sci U S A* 115(28):7368-7373.
- Wang E, Schornack S, Marsh JF, Gobbato E, Schwessinger B, Eastmond P, *et al.* 2012. A common signaling process that promotes mycorrhizal and oomycete colonization of plants. *Curr Biol* 22(23):2242-6.
- Wang L, Cao Y, Wang ET, Qiao YJ, Jiao S, Liu ZS, *et al.* 2016. Biodiversity and biogeography of rhizobia associated with common bean (*Phaseolus vulgaris* L.) in Shaanxi Province. *Syst Appl Microbiol* 39(3):211-219.
- Wang ZH, Fang H, Chen M. 2017. Effects of root exudates of woody species on the soil anti-erodibility in the rhizosphere in a karst region, China. *PeerJ* 5:e3029.
- Ward NL, Challacombe JF, Janssen PH, Henrissat B, Coutinho PM, Wu M *et al.* 2009. Three genomes from the phylum Acidobacteria provide insight into the lifestyles of these microorganisms in soils. *Appl Environ Microbiol* 75(7):2046-56.
- Waters MT, Gutjahr C, Bennett T, Nelson DC. 2017. Strigolactone signaling and evolution. *Annu Rev Plant Biol* 68:291-322.
- Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

- Weinert N, Piceno Y, Ding GC, Meincke R, Heuer H, Berg G, *et al.* 2011. PhyloChip hybridization uncovered an enormous bacterial diversity in the rhizosphere of different potato cultivars: many common and few cultivar-dependent taxa. *FEMS Microbiol Ecol* 75(3):497-506.
- Weese DJ, Heath KD, Dentinger BT, Lau JA. 2015. Long-term nitrogen addition causes the evolution of less-cooperative mutualists. *Evolution* 69(3):631-42.
- Wemheuer, F, Kaiser K, Karlovsky P, Daniel R, Vidal S, Wemheuer B. 2017. Bacterial endophyte communities of three agricultural important grass species differ in their response towards management regimes. *Sci Rep* 7:40914.
- Weston LA, Mathesius U. 2013. Flavonoids: their structure, biosynthesis and role in the rhizosphere, including allelopathy. *J Chem Ecol* 39(2):283-97.
- Wickham, H. 2009. ggplot2: elegant graphics for data analysis. New York: Springer-Verlag. 260 pp.
- Wieland G, Neumann R, Backhaus H. 2001. Variation of microbial communities in soil, rhizosphere, and rhizoplane in response to crop species, soil type, and crop development. *Appl Environ Microbiol* 67(12):5849-54.
- Wilhelm RC, Cardenas E, Leung H, Szeitz A, Jensen LD, Mohn WW. 2017. Long-term enrichment of stress-tolerant cellulolytic soil populations following timber harvesting evidenced by multi-omic stable isotope probing. *Front Microbiol* 8:537.
- Wintermans PC, Bakker PA, Pieterse CM. 2016. Natural genetic variation in Arabidopsis for responsiveness to plant growth-promoting rhizobacteria. *Plant Mol Biol* 90(6):623-34.
- Wissuwa M, Mazzola M, Picard C. 2009. Novel approaches in plant breeding for rhizosphere related traits. *Plant Soil* 321:409–430.
- Wolińska A, Kuźniar A, Zielenkiewicz U, Izak D, Szafranek-Nakonieczna A, Banach A, *et al.* 2017. Bacteroidetes as a sensitive biological indicator of agricultural soil usage revealed by culture-independent approach. *Appl Soil Ecol* 119:128–137.
- Xing X, Koch AM, Jones AM, Ragone D, Murch S, Hart MM. 2012. Mutualism breakdown in breadfruit domestication. *Proc Biol Sci* 22:279(1731):1122-30.
- Xu Y, Wang G, Jin J, Liu J, Zhang Q, Liu Xiaobing. 2009. Bacterial communities in soybean rhizosphere in response to soil type, soybean genotype, and their growth stage. *Soil Biol Biochem* 41:919–925.
- Yeoh YK, Dennis PG, Paungfoo-Lonhienne C, Weber L, Brackin R, Ragan MA, *et al.* 2017. Evolutionary conservation of a core root microbiome across plant phyla along a tropical soil chronosequence. *Nat Commun* 8(1):215.
- Yeoh YK, Paungfoo-Lonhienne C, Dennis PG, Robinson N, Ragan MA, Schmidt S, *et al.* 2016. The core root microbiome of sugarcane cultivated under varying nitrogen fertilizer application. *Environ Microbiol* 18(5):1338-51.
- Yin C, Hulbert SH, Schroeder KL, Mavrodi O, Mavrodi D, Dhingra A, *et al.* 2013. Role of bacterial communities in the natural suppression of *Rhizoctonia solani* bare patch disease of wheat (*Triticum aestivum* L.). *Appl Environ Microbiol* 79:7428-7438.
- Yoon JH, Lee CH, Oh TK. 2005. *Aeromicrobium alkaliterrae* sp. nov., isolated from an alkaline soil, and emended description of the genus *Aeromicrobium*. *Int J Syst Evol Microbiol* 55:2171-5.

- Zachow C, Müller H, Tilcher R, Berg G. 2014. Differences between the rhizosphere microbiome of *Beta vulgaris* ssp. *maritima* - ancestor of all beet crops - and modern sugar beets. *Front Microbiol* 5:415.
- Zgadzaj R, Garrido-Oter R, Jensen DB, Koprivova A, Schulze-Lefert P, Radutoiu S. 2016. Root nodule symbiosis in *Lotus japonicus* drives the establishment of distinctive rhizosphere, root, and nodule bacterial communities. *Proc Natl Acad Sci U S A* 113:E7996-E8005.
- Zhalnina K, Louie KB, Hao Z, Mansoori N, da Rocha UN, Shi S, *et al.* Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. *Nat Microbiol* 3(4):470-480.
- Zhang N, Wang D, Liu Y, Li S, Shen Q, Zhang R. 2014. Effects of different plant root exudates and their organic acid components on chemotaxis, biofilm formation and colonization by beneficial rhizosphere-associated bacterial strains. *Plant Soil* 374:689-700.
- Zhao J, Bodner G, Rewald B, Leitner D, Nagel KA, Nakhforoosh A. 2017. Root architecture simulation improves the inference from seedling root phenotyping towards mature root systems. *J Exp Bot* 68(5):965-982.
- Zhu Y, McBride MJ. 2017. The unusual cellulose utilization system of the aerobic soil bacterium *Cytophaga hutchinsonii*. *Appl Microbiol Biotechnol* 101(19):7113-7127.
- Zhu Y-G, Smith SE, Barritt AR, Smith FA. 2001. Phosphorus (P) efficiencies and mycorrhizal responsiveness of old and modern wheat cultivars. *Plant Soil* 237:249-255.
- Zwanenburg B, Pospíšil T, Čavar Zeljković S. 2016. Strigolactones: new plant hormones in action. *Planta* 243(6):1311-26.