



Universiteit  
Leiden  
The Netherlands

## **Inverse Jacobian and related topics for certain superelliptic curves**

Somoza Henares, A.

### **Citation**

Somoza Henares, A. (2019, March 28). *Inverse Jacobian and related topics for certain superelliptic curves*. Retrieved from <https://hdl.handle.net/1887/70564>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/70564>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/70564> holds various files of this Leiden University dissertation.

**Author:** Somoza Henares, A.

**Title:** Inverse Jacobian and related topics for certain superelliptic curves

**Issue Date:** 2019-03-28

# INTRODUCTION

Given an *elliptic curve*  $E$  over  $\mathbb{C}$ , there exists a *lattice*  $\Lambda \subseteq \mathbb{C}$  such that the group  $E(\mathbb{C})$  of complex points on  $E$  is isomorphic to the complex analytic group  $\mathbb{C}/\Lambda$ . This link between elliptic curves and one-dimensional complex tori is called the Uniformization Theorem, and one can explicitly find the curve corresponding to a given lattice with the *Weierstrass  $\wp$ -function*, its derivative, and the *Eisenstein series*.

Similarly, given an algebraic curve  $C$  of genus  $g$ , one associates to it a *principally polarized abelian variety*  $J(C)$ , the *Jacobian of  $C$* . Over  $\mathbb{C}$ , the Jacobian  $J(C)$  is isomorphic to a  $g$ -dimensional complex torus  $\mathbb{C}^g/\Lambda$  for a lattice  $\Lambda$  of full rank in  $\mathbb{C}^g$ .

This determines a map  $J$  from the set  $\mathbf{M}_g$  of isomorphism classes of algebraic curves of genus  $g$  to the set  $\mathbf{A}_g$  of principally polarized abelian varieties of dimension  $g$ , and one may wonder if there exists an explicit inverse to this map, as in the case of elliptic curves. We call this the *inverse Jacobian problem*.

This problem has been solved for curves of genus 2 [37, 50] and genus 3 [1, 9, 16, 21, 48, 52, 53]. However, for genus  $g \geq 4$  there is the additional obstruction that not all principally polarized abelian varieties are Jacobians of curves, hence in order to solve the inverse Jacobian problem one needs to study the image by  $J$  of  $\mathbf{M}_g$  in  $\mathbf{A}_g$ . The problem of describing  $J(\mathbf{M}_g)$  is known as the *Riemann-Schottky problem*.

In this thesis we treat these two problems for two families of *superelliptic curves*, that is, curves of the form  $y^k = \prod_{i=1}^l (x - \alpha_i)$ . We focus on the family of *Picard curves*, with  $(k, l) = (3, 4)$  and genus 3, where we solve the inverse Jacobian problem, and the family of *cyclic plane quintic curves* (CPQ curves), with  $(k, l) = (5, 5)$  and genus 6, where we solve both problems.

In Chapter 1 we first introduce some background on abelian varieties, Jacobians of curves, and Riemann theta constants, and then we present an inverse Jacobian algorithm for Picard curves. Note that Picard curves have genus 3, hence there is no obstruction to the inverse Jacobian problem.

Since Picard curves are plane quartic curves, the inverse Jacobian problem for Picard curves could be solved using the formulas for plane quartics given

in [52], but focusing on a smaller family of curves allows us to present a more efficient solution for the family of interest.

This was originally done by Koike and Weng in [16], but their exposition presents some mistakes that we address and correct here. This chapter is based on joint work with Joan-Carles Lario, see also [21].

In Chapter 2 we present an inverse Jacobian algorithm for CPQ curves. We follow a strategy analogous to the one in Chapter 1 for the case of Picard curves.

In Chapter 3 we address the Riemann-Schottky problem for CPQ curves, that is, we characterize the principally polarized abelian varieties that are Jacobians of CPQ curves. First we use a generalization of the classical theory of *complex multiplication* due to Shimura [39] to study how the existence of the automorphism of CPQ curves  $(x, y) \mapsto (x, \exp(2\pi i/5)y)$  affects the structure of the Jacobians. Then we solve a class number one problem for higher-dimensional Hermitian lattices over  $\mathbb{Z}[\zeta_5]$ , which is key to solving the Riemann-Schottky problem for CPQ curves.

Finally, in Chapter 4 we present one application for the above algorithms: constructing curves such that their Jacobians have complex multiplication. This has previously been done for genus 2 [51, 47] and genus 3 [1, 13, 16, 21, 53]. Here we extend the methods of Kılıçer [12] to determine a complete list of CM-fields whose ring of integers occurs as the endomorphism ring over  $\mathbb{C}$  of the Jacobian of a CPQ curve defined over  $\mathbb{Q}$ .

In particular, this allows us to list conjectural models for all CPQ curves over  $\mathbb{Q}$  whose Jacobians have the maximal order of a degree-12 CM-field as endomorphism ring over  $\mathbb{C}$ . Our list contains the correct number of curves, which are defined over  $\mathbb{Q}$  and numerically correct up to high precision.