



<https://openaccess.leidenuniv.nl>

### **License: Article 25fa pilot End User Agreement**

This publication is distributed under the terms of Article 25fa of the Dutch Copyright Act (Auteurswet) with explicit consent by the author. Dutch law entitles the maker of a short scientific work funded either wholly or partially by Dutch public funds to make that work publicly available for no consideration following a reasonable period of time after the work was first published, provided that clear reference is made to the source of the first publication of the work.

This publication is distributed under The Association of Universities in the Netherlands (VSNU) 'Article 25fa implementation' pilot project. In this pilot research outputs of researchers employed by Dutch Universities that comply with the legal requirements of Article 25fa of the Dutch Copyright Act are distributed online and free of cost or other barriers in institutional repositories. Research outputs are distributed six months after their first online publication in the original published version and with proper attribution to the source of the original publication.

You are permitted to download and use the publication for personal purposes. All rights remain with the author(s) and/or copyrights owner(s) of this work. Any use of the publication other than authorised under this licence or copyright law is prohibited.

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the Library through email: [OpenAccess@library.leidenuniv.nl](mailto:OpenAccess@library.leidenuniv.nl)

### **Article details**

Greene, T., Gelkopf, M., Epskamp, S., & Fried, E. I. (2018), Dynamic networks of PTSD symptoms during conflict, *Psychological Medicine* 48 (14): 2409-2417  
Doi: 10.1017/S0033291718000351

# Dynamic networks of PTSD symptoms during conflict

Talya Greene<sup>1</sup>, Marc Gelkopf<sup>1,2</sup>, Sacha Epskamp<sup>3</sup> and Eiko Fried<sup>3</sup>

<sup>1</sup>Department of Community Mental Health, University of Haifa, Haifa, Israel; <sup>2</sup>NATAL, Israel Trauma Center for Victims of Terror and War, Tel Aviv, Israel and <sup>3</sup>Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands

## Original Article

**Cite this article:** Greene T, Gelkopf M, Epskamp S, Fried E (2018). Dynamic networks of PTSD symptoms during conflict. *Psychological Medicine* **48**, 2409–2417. <https://doi.org/10.1017/S0033291718000351>

Received: 24 July 2017

Revised: 24 January 2018

Accepted: 24 January 2018

First published online: 28 February 2018

### Key words:

ESM; intensive longitudinal assessment; multilevel VAR; network analysis; posttraumatic stress; trauma; war

### Author for correspondence:

Talya Greene, E-mail: [tgreene@univ.haifa.ac.il](mailto:tgreene@univ.haifa.ac.il)

### Abstract

**Background.** Conceptualizing posttraumatic stress disorder (PTSD) symptoms as a dynamic system of causal elements could provide valuable insights into the way that PTSD develops and is maintained in traumatized individuals. We present the first study to apply a multilevel network model to produce an exploratory empirical conceptualization of dynamic networks of PTSD symptoms, using data collected during a period of conflict.

**Methods.** Intensive longitudinal assessment data were collected during the Israel–Gaza War in July–August 2014. The final sample ( $n = 96$ ) comprised a general population sample of Israeli adult civilians exposed to rocket fire. Participants completed twice-daily reports of PTSD symptoms via smartphone for 30 days. We used a multilevel vector auto-regression model to produce contemporaneous and temporal networks, and a partial correlation network model to obtain a between-subjects network.

**Results.** Multilevel network analysis found strong positive contemporaneous associations between hypervigilance and startle response, avoidance of thoughts and avoidance of reminders, and between flashbacks and emotional reactivity. The temporal network indicated the central role of startle response as a predictor of future PTSD symptomatology, together with restricted affect, blame, negative emotions, and avoidance of thoughts. There were some notable differences between the temporal and contemporaneous networks, including the presence of a number of negative associations, particularly from blame. The between-person network indicated flashbacks and emotional reactivity to be the most central symptoms.

**Conclusions.** This study suggests various symptoms that could potentially be driving the development of PTSD. We discuss clinical implications such as identifying particular symptoms as targets for interventions.

## Introduction

The network perspective offers a novel way of understanding the dynamics of psychopathology (McNally, 2016; Borsboom, 2017). According to this framework, symptoms do not primarily result as passive consequences of underlying mental disorders. Rather, causally connected symptoms may interact with each other over time, potentially producing mental disorders as emergent phenomena. The new field of network psychometrics has been used in recent years to investigate the complex structure of various psychiatric disorders (Fried *et al.* 2017b), including depression (Fried *et al.* 2016), psychosis (Isvoranu *et al.* 2016), schizophrenia (Levine & Leucht, 2016), and anxiety (Beard *et al.* 2016), among others.

Several studies have recently used network analysis to examine the symptom-level structure of posttraumatic stress disorder (PTSD) in cross-sectional data (McNally *et al.* 2015; Bryant *et al.* 2017; Knefel *et al.* 2016; Afzali *et al.* 2017a, 2017b; Armour *et al.* 2017; Birkeland & Heir, 2017; Fried *et al.* 2017a; Mitchell *et al.* 2017; Spiller *et al.* 2017; Sullivan *et al.* 2018). These studies indicate the central roles that various symptoms may play in driving PTSD networks by being highly connected to other PTSD symptoms. For example, these studies suggest the central role of re-experiencing symptoms in the acute phase following traumatic injury (Bryant *et al.* 2017); of hypervigilance, concentration problems and foreshortened future in Chinese earthquake survivors 5 years after the disaster (McNally *et al.* 2015); and of intrusive thoughts, anger, and detachment in survivors of mass violence (Sullivan *et al.* 2018). Two studies in military veterans found that negative trauma-related emotions were among the most central symptoms (Armour *et al.* 2017; Mitchell *et al.* 2017).

Only three studies have previously assessed DSM-5 PTSD network structure (Afzali *et al.* 2017a; Armour *et al.* 2017; Mitchell *et al.* 2017), and, crucially, as all of these studies were cross-sectional, none of them explored the temporal dynamics of PTSD, which many researchers have described as one of the major shortcomings of the prior literature (Hamaker, 2012; Fisher, 2015; Fried & Cramer, 2017; Bos *et al.* 2017; Hamaker & Wichers, 2017; Nelson *et al.* 2017). Conceptualizing PTSD symptoms as a dynamic interacting network, rather than limiting the

focus to cross-sectional associations between symptoms, allows for deeper exploration of potential causal pathways. This could provide valuable insights into the way that PTSD builds or is maintained in individuals who have traumatic experiences.

Intensive longitudinal assessments such as Experience Sampling Methodology (ESM) collect data about the daily lives of participants repeatedly over time, often using handheld computers or smartphones (Bolger *et al.* 2003; Ebner-Priemer & Kubiak, 2010; Bolger & Laurenceau, 2013). ESM has been used to investigate various psychiatric disorders, including PTSD (Chun, 2016), anxiety (Walz *et al.* 2014), and depression (Armeij *et al.* 2015), among others. ESM entails multiple methodological advantages (Santangelo *et al.* 2013; Trull & Ebner-Priemer, 2013), as data are collected in real-life settings, with high ecological validity, increased accuracy, and minimized recall bias. Repeated short-term assessments can detect variations in the presence and severity of variables and reveal dynamic processes between them (Ebner-Priemer & Trull, 2009; Myin-Germeys *et al.* 2009; Bolger & Laurenceau, 2013).

Network models can be used to investigate such dynamic processes in ESM data from one participant (vector autoregression models; VAR) or data from multiple participants (multilevel VAR; Epskamp *et al.* 2016). These models produce temporal networks depicting a directed network of the lagged associations of symptoms from one time point to the next, for which Granger-causal connections between symptoms are inferred (Schuurman *et al.* 2016). Temporal networks can identify symptoms with a high 'out-strength', that is symptoms which are most predictive of other symptoms at the next time point, and those with a high 'in-strength', that is symptoms which are most predicted by other symptoms at the previous time point. Recent extensions include contemporaneous networks, for which the within-measurement associations are represented, and a between-persons network, modeling relations between the stable means of participants, for both of which 'cliques' or sub-networks of closely connected symptoms can be identified.

Using ESM data for network analysis is relatively new; ESM data have previously been used for network studies of psychiatric disorders, but almost exclusively for individual-level ( $n = 1$ ) studies (Bak *et al.* 2016; David *et al.* 2017; Epskamp *et al.* 2017), or to investigate idiographic symptom dynamics (Fisher *et al.* 2017), although multilevel VAR using ESM data has been applied to investigate various psychological phenomena, such as positive and negative affect (e.g. Epskamp *et al.* 2016).

The current study uses ESM data from multiple individuals in a multilevel framework to investigate PTSD symptoms. As multilevel models focus on the dynamics of the relationships between symptoms over time, they could further help us to elucidate the processes by which PTSD symptoms build up and interact with each other, and potentially reinforce themselves. This could have important clinical implications (McNally *et al.* 2015; McNally, 2016; Borsboom, 2017); identifying symptoms with high out-strength, and to a lesser extent high in-strength, as well as detecting closely connected groups of symptoms, could eventually help clinicians focus on symptoms that might 'drive' PTSD in its early stages.

We present the first study to apply a multilevel network model to investigate the dynamics of symptoms that comprise the DSM-5 PTSD construct (APA, 2013), using data collected during a period of trauma exposure among Israeli civilians exposed to rocket fire. It is important to note that collecting ESM data regarding traumatic stress symptoms during a conflict situation

is unparalleled in the literature, and we believe that this study, therefore, has the potential to make unique and important contributions to the understanding of mental health during ongoing trauma exposure. The study aims to produce an exploratory empirical conceptualization of dynamic networks of PTSD symptoms, with the goal to investigate the network structure of PTSD symptoms during an extended period of trauma exposure and identify symptoms that closely cluster together, as well as those with high out-strength and high in-strength.

## Method

### Participants and study design

The current study is part of a larger study conducted during and after the 50-day July–August 2014 Israel–Gaza armed conflict (Gelkopf *et al.* 2017; Greene *et al.* 2017; Lapid Pickman *et al.* 2017), during which more than 4500 rockets and mortar shells were fired from Gaza toward Israeli communities, and Israel Defence Forces conducted ground and air attacks on Gaza.

The sample for the current study comprised a general population sample of Israeli adult civilians living in communities exposed to rocket fire ( $n = 114$ ). Participants were recruited via advertisements in local colleges/universities, local organizations and communities, and social networks, and via snowballing methods. Potential participants contacted the research team directly, and study explanation and oral informed consent were conducted over the telephone.

Participants entered the study gradually, between days 8 and 24 of the conflict. Participants were asked to complete ESM daily diary assessments twice a day, morning and evening at predetermined times, for 30 days, and could reply within 2 h. The questionnaires were administered via a personalized email link to participants' smartphones and referred to experiences, reactions, and symptoms occurring since the last assessment.

Participants were remunerated for their participation. The design and procedure were approved by the University of Haifa, Faculty of Social Welfare and Health Sciences Ethics Committee.

### Measures

*Traumatic stress symptoms* were measured using the PTSD Checklist for DSM-5 (PCL-5; Weathers *et al.* 2013). This 20-item self-report tool asks participants to indicate the extent to which they experienced each PTSD symptom, on a four-point Likert scale ranging from 0 = not at all to 3 = very much. The original version was adapted for the current study so that the timeframe for experiencing each symptom was changed from 'in the past month' to 'since the last time you replied' (a minimum of 8 h gap). This measure has good psychometric properties in its original form (Blevins *et al.* 2015).

### Data analysis

We included data from participants providing over 20 valid responses (one-third of potential responses). Of the sample, 84.2% ( $n = 96$ ) met this inclusion criterion.

Demographic characteristics of the sample are presented in Table 1.

The mlVAR package (version 0.3.3) in R was used to estimate three networks (Epskamp *et al.* 2016): a contemporaneous network, which is a Gaussian graphical model (GGM) that depicts

**Table 1.** Demographic characteristics of the sample

|                     | Categories   | <i>n</i> | %    | Mean (s.d.)   |
|---------------------|--|----------|------|---|
| Gender              | Female   | 68       | 70.8 |   |
|                     | Male   | 28       | 29.2 |   |
| Age                 |  |          |      | 30.1 (9.0)  |
| Country of Birth    | Israel   | 80       | 83.3 |   |
|                     | Outside of Israel  | 16       | 16.7 |   |
| Relationship status | Single   | 27       | 28.1 |   |
|                     | Has a partner  | 28       | 29.2 |   |
|                     | Married  | 39       | 40.6 |   |
|                     | Divorced/separated   | 2        | 2.1  |   |
| Have children       | Yes  | 40       | 41.7 |   |
|                     | No   | 56       | 58.3 |   |
| Religion            | Jewish   | 94       | 97.9 |   |
|                     | Christian and other  | 2        | 2.1  |   |
| Religiosity status  | Secular  | 58       | 60.4 |   |
|                     | Traditional  | 21       | 21.9 |   |
|                     | Religious or Orthodox  | 17       | 17.7 |   |
| Education           | High school (no diploma)   | 4        | 4.2  |   |
|                     | High school (with diploma)                                       | 34       | 35.4 |   |
|                     | Academic (including student)                                     | 54       | 56.3 |   |
|                     | Other professional training                                      | 4        | 4.2  |   |
| Employment          | Full-time  | 31       | 32.3 |   |
|                     | Part-time  | 13       | 13.5 |   |
|                     | Temporary jobs   | 10       | 10.4 |   |
|                     | Student  | 24       | 25.0 |   |
|                     | Unemployed   | 5        | 5.2  |   |
|                     | Other  | 13       | 13.5 |   |
| PTSD symptoms       | Endorsed PTSD DSM-5 criteria (other than duration) at least once | 40       | 41.7 |   |
|                     | Per person mean PCL sum score                                    |          |      | 10.18 (7.87) Range of mean PCL sum score (0.20–32.57) |


within-time-window edges (associations) between nodes (DSM-5 PTSD symptoms) that remain after controlling for temporal associations, akin to a multilevel partial correlation network; a temporal network, which is a directed network of regression coefficients that depicts the lagged associations between symptoms from one measurement point to the next and a between-persons network which is a GGM that depicts the variance-covariance structure of participants' means (Epskamp *et al.* 2016, 2017).

Decomposing the variance in these three networks allows for different insight in the covariation and potential dynamics: the temporal network shows if a deviation from a person's mean predicts a deviation from a person's mean in another variable at the next measurement occasion. As a result, temporal networks give insight into Granger causality (Granger, 1969). Contemporaneous networks show if deviations from a person's mean in two variables predict one-another in the same measurement occasion (Epskamp *et al.* 2017). Finally, the between-subjects network models the covariation between means of participants and is a more stable way of estimating networks that can be obtained from cross-

sectional data, while providing a network that can be used for comparison with previous cross-sectional studies (Epskamp *et al.* 2016).

In our analysis, to make the model estimable we estimated forced random effects of incoming edges to a single node (temporal) or edges connected to the same node (contemporaneous) to be orthogonal. In the contemporaneous and between-subjects networks, we used an 'and' rule in retaining significant edges: an edge was retained if both regressions on which the edge was based were significant ( $\alpha = 0.05$ ).

In the current analysis, the between-persons network included numerous negative edges, which are likely spurious connections given that the contemporaneous and temporal network had far fewer negative connections (see online Supplementary Material). We, therefore, conducted an alternative between-persons network analysis in the R Bootnet package using graphical LASSO and extended Bayesian information criterion to select the optimal regularization parameter, for which rather than using the mean edge scores from each individual to depict the network, we used the mean symptom scores for each participant.



**Fig. 1.** Dynamic PTSD networks during the conflict. *Note:* Blue edges denote positive associations between nodes, red edges denote negative associations between nodes. Edge thickness represents the degree of association.

## Results

Figure 1 depicts the contemporaneous, temporal and between-person networks.

### Contemporaneous network

Several important things stand out in the contemporaneous network (Fig. 1 and Table 2). First, contemporaneous connections were nearly all positive in nature, and we identified especially strong associations between hypervigilance and exaggerated startle response (partial correlation edge weight=0.36), avoidance of thoughts and amnesia (0.30), and loss of interest and detachment (0.27).

Second, the DSM-5 symptom clusters were fairly well recovered, with strong within-cluster associations, and weaker between cluster associations.

Third, in a few cases, the contemporaneous network did not fully correspond to the DSM-5 cluster conceptualization. For example, amnesia had stronger associations to the avoidance cluster than it did to its own cluster of negative mood and cognitions,

and anger had stronger associations to the negative mood and cognitions cluster than to the arousal cluster to which it belongs.

Finally, the negative mood and cognitions appear to include two ‘cliques’ – clusters of closely connected symptoms; the first clique comprised negative beliefs, blame, and negative emotions which were all significantly connected to each other, and the second clique was made up of loss of interest, detachment, and restricted affect.

### Temporal network

There were several notable aspects of the temporal network (see Fig. 1). First, there were a substantial number of negative edges, specifically from blame, flashbacks, avoidance of thoughts, nightmares, and self-destructive behavior, which differed from the contemporaneous network. All of the lagged associations from blame to other symptoms were negative. Avoidance of thoughts had a negative lagged association with sleep disturbances.

Exaggerated startle response had by far the strongest out-strength (0.83), which is the sum of absolute significant temporal edge weights extending out from each symptom, excluding

**Table 2.** Contemporaneous network: significant edge weights

|                            | Intr. thoughts | Nightmares | Flashbacks | Emo. React. | Phys. React. | Avoid thoughts | Avoid remind. | Amn.  | Neg. beliefs | Blame | Neg. emo. | Loss interest | Detachment | Res. Affect | Anger | Self-destruct. | Hyper-vig. | Start. | Conc. diff. | Sleep dist. |
|----------------------------|----------------|------------|------------|-------------|--------------|----------------|---------------|-------|--------------|-------|-----------|---------------|------------|-------------|-------|----------------|------------|--------|-------------|-------------|
| Intrusive thoughts         | .              | 0.19       | ns         | 0.19        | 0.07         | ns             | ns            | ns    | ns           | ns    | ns        | ns            | ns         | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Nightmares                 | 0.19           | .          | 0.22       | ns          | 0.13         | ns             | ns            | ns    | ns           | ns    | ns        | ns            | ns         | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Flashbacks                 | ns             | 0.22       | .          | 0.25        | 0.09         | 0.07           | ns            | ns    | ns           | ns    | ns        | ns            | ns         | ns          | ns    | 0.04           | ns         | ns     | ns          | ns          |
| Emotional reactivity       | 0.19           | ns         | 0.25       | .           | 0.13         | ns             | ns            | 0.13  | ns           | ns    | ns        | ns            | ns         | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Physiological reactivity   | 0.07           | 0.13       | 0.09       | 0.13        | .            | ns             | ns            | ns    | ns           | ns    | 0.12      | ns            | ns         | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Avoidance of thoughts      | ns             | ns         | 0.07       | ns          | ns           | .              | 0.24          | 0.30  | ns           | 0.06  | ns        | ns            | ns         | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Avoidance of reminders     | ns             | ns         | ns         | ns          | ns           | 0.24           | .             | 0.18  | ns           | ns    | ns        | ns            | 0.06       | ns          | ns    | ns             | ns         | ns     | -0.03       | ns          |
| Amnesia                    | ns             | ns         | ns         | 0.13        | ns           | 0.30           | 0.18          | .     | 0.08         | ns    | ns        | ns            | -0.05      | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Negative beliefs           | ns             | ns         | ns         | ns          | ns           | ns             | ns            | 0.08  | .            | 0.19  | 0.22      | ns            | 0.05       | ns          | ns    | 0.06           | ns         | ns     | ns          | ns          |
| Blame                      | ns             | ns         | ns         | ns          | ns           | 0.06           | ns            | ns    | 0.19         | .     | 0.14      | ns            | 0.03       | ns          | ns    | ns             | ns         | ns     | ns          | ns          |
| Negative emotions          | ns             | ns         | ns         | ns          | 0.12         | ns             | ns            | ns    | 0.22         | 0.14  | .         | ns            | ns         | 0.10        | 0.08  | ns             | 0.07       | 0.05   | ns          | 0.05        |
| Loss of interest           | ns             | ns         | ns         | ns          | ns           | ns             | ns            | ns    | ns           | ns    | ns        | .             | 0.27       | 0.19        | 0.09  | ns             | ns         | ns     | 0.11        | 0.05        |
| Detachment                 | ns             | ns         | ns         | ns          | ns           | ns             | 0.06          | -0.05 | 0.05         | 0.03  | ns        | 0.27          | .          | 0.21        | 0.13  | ns             | ns         | ns     | 0.06        | ns          |
| Restricted affect          | ns             | ns         | ns         | ns          | ns           | ns             | ns            | ns    | ns           | ns    | 0.10      | 0.19          | 0.21       | .           | 0.18  | ns             | ns         | ns     | 0.08        | ns          |
| Anger                      | ns             | ns         | ns         | ns          | ns           | ns             | ns            | ns    | ns           | ns    | 0.08      | 0.09          | 0.13       | 0.18        | .     | 0.06           | 0.06       | ns     | 0.07        | ns          |
| Self-destructive behavior  | ns             | ns         | 0.04       | ns          | ns           | ns             | ns            | ns    | 0.06         | ns    | ns        | ns            | ns         | ns          | 0.06  | .              | 0.09       | ns     | ns          | ns          |
| Hypervigilance             | ns             | ns         | ns         | ns          | ns           | ns             | ns            | ns    | ns           | ns    | 0.07      | ns            | ns         | ns          | 0.06  | 0.09           | .          | 0.36   | 0.06        | 0.11        |
| Startle                    | ns             | ns         | ns         | ns          | ns           | ns             | ns            | ns    | ns           | ns    | ns        | ns            | ns         | ns          | ns    | ns             | 0.36       | .      | 0.15        | ns          |
| Concentration difficulties | ns             | ns         | ns         | ns          | ns           | ns             | -0.03         | ns    | ns           | ns    | ns        | 0.11          | 0.06       | 0.08        | 0.07  | ns             | 0.06       | 0.15   | .           | 0.17        |
| Sleep disturbance          | ns             | ns         | ns         | ns          | ns           | ns             | ns            | ns    | ns           | ns    | 0.05      | 0.05          | ns         | ns          | ns    | ns             | 0.11       | ns     | 0.17        | .           |

Note: ns denotes non-significant edge

autocorrelations (see Table 3). In other words, the more startle response an individual had at one measurement, the more likely the individual was to report other symptoms at the next. Other variables with high out-strength included restricted affect (0.54), blame (0.51), negative emotions (0.34), and avoidance of thoughts (0.32).

The variables with the highest in-strength centrality – the extent to which variables are influenced by other variables in the network at the previous measurement – were sleep disturbance (0.37), and loss of interest (0.37).

Consistent with prior temporal studies of psychopathological dynamics, there were a considerable number of autocorrelations, which were stronger than the cross-lagged effects. Exaggerated startle response had the strongest autoregressive coefficient (0.27), meaning that knowing the startle response at a given timepoint has strong predictive power about startle response at the next timepoint. Hypervigilance also had a high autoregressive coefficient (0.22).

Finally, a number of interesting feedback loops emerged. For example, sleep disturbance predicted concentration difficulties, which in turn predicted restricted affect, which predicts sleep disturbance. Negative emotions predicted negative beliefs, which went to avoidance of thoughts, and back again to negative emotions.

### Between-persons network

The between-persons network closely resembled the contemporaneous network, albeit more sparsely connected. As with the contemporaneous network, associations between symptoms were mostly positive, with particularly strong associations between the two avoidance symptoms, between negative beliefs and negative emotions, and between hypervigilance and startle. Standardized node strength centrality was calculated (see Fig. 2), representing the sum of all edge weights of a given node. Flashbacks followed by emotional cue reactivity were the two most central symptoms in this network.

### Discussion

This study assessed dynamic PTSD symptom networks in Israeli civilians during the 2014 Israel–Gaza War. There were strong positive contemporaneous associations between hypervigilance and exaggerated startle response, avoidance of thoughts and avoidance of reminders, and between flashbacks and emotional reactivity. There were some notable differences between the temporal and contemporaneous networks, which underlines the importance of analyzing both separately. These differences included the presence of a number of negative associations, particularly from blame. The temporal network also indicated the central role of startle response as a predictor of PTSD symptomatology at the next timepoint. The between-persons network showed similar patterns of associations to the contemporaneous network.

Contemporaneous associations between symptoms largely reflected the DSM-5 clusters, other than the alterations in negative mood and cognitions cluster. Previous PTSD network studies found that amnesia was only weakly associated with other PTSD symptoms (McNally *et al.* 2015; Armour *et al.* 2017; Birkeland & Heir, 2017; Fried *et al.* 2017a; Spiller *et al.* 2017). The current findings indicated, however, that this was not the case, but rather that amnesia was more closely related to the avoidance cluster, than to the cognitive symptoms from its own

**Table 3.** In-strength and out-strength of symptoms in the temporal network

| Symptom                    | In-strength (sum of absolute values of significant edges to symptom) | Out-strength (sum of absolute values of significant edges from symptom) |
|----------------------------|--|---|
| Intrusive thoughts         | 0  | 0.05  |
| Nightmares                 | 0  | 0.18  |
| Flashbacks                 | 0.08   | 0.15  |
| Emotional reactivity       | 0.29   | 0.08  |
| Physiological reactivity   | 0.12   | 0   |
| Avoidance of thoughts      | 0.28   | 0.32  |
| Avoidance of reminders     | 0.13   | 0.1   |
| Amnesia                    | 0.16   | 0   |
| Negative beliefs           | 0.11   | 0.25  |
| Blame                      | 0.12   | 0.51  |
| Negative emotions          | 0.30   | 0.34  |
| Loss of interest           | 0.37   | 0.08  |
| Detachment                 | 0  | 0.04  |
| Restricted affect          | 0.31   | 0.54  |
| Anger/irritability         | 0.19   | 0.05  |
| Self-destructive behavior  | 0.17   | 0.04  |
| Hypervigilance             | 0.31   | 0.19  |
| Startle response           | 0.18   | 0.83  |
| Concentration difficulties | 0.29   | 0.23  |
| Sleep disturbance          | 0.37   | 0.12  |

cluster, and had no significant associations with any of the negative mood items. Perhaps amnesia could be better conceptualized as an additional and perhaps involuntary form of avoidance, rather than as a negative cognition and mood symptom, at least in the context of the ongoing threat.

The pattern of connections from the remaining symptoms from the negative cognition and mood cluster supported a model in which numbing of responsiveness items (loss of interest, detachment, and restricted affect) are split from the increased negative mood and cognition items (negative mood, blame, negative cognitions), something that has also been found in previous investigations of the dimensional structure of DSM-5 PTSD (Armour *et al.* 2015; Pietrzak *et al.* 2015).

It is important to interpret the contemporaneous network in a nuanced manner (Epskamp *et al.* 2017). While ‘contemporaneous’ would seem to imply concurrent associations, in the current study there were relatively large time windows between assessments (8–16 h), so contemporaneous networks are not really ‘contemporaneous’. Rather, they depict associations between symptoms, including potentially causal associations that took place over a number of hours, and may therefore also reflect temporal processes. To

## Between-persons network standardized node strength centrality




Fig. 2. Between-persons network standardized node strength centrality.

assess true contemporaneous networks, far more frequent assessments are needed, and participants should be asked how they are feeling ‘right now’, rather than ‘since the last assessment’.

In the temporal network, the blame of self or others surprisingly predicted lower symptoms at the next measurement, in contrast to the positive associations they showed in the contemporaneous network. The blame item, which was added to the PTSD criteria in DSM-5, actually captures two very different phenomena (Greene, *under review*); blame of self, in which the individual gives an internal attribution to the adversity, and blame of others, which gives an external attribution to the adversity. Attributions theory posits that making external attributions for negative events is protective (Kelley & Michela, 1980). While we cannot ascertain in this study whether the participants were reporting external or internal blame, it could be speculated that in the context of war and conflict, the blame would be more likely to be externally attributed. This may then explain the negative lagged association from blame to avoidance of thoughts, avoidance of reminders, negative emotions, loss of interest, detachment, restricted affect, startle, concentration difficulties and sleep disturbance, suggesting that blaming an external source may be an effective coping mechanism.

Symptoms with strong out-strengths can be seen as those with the highest potential to impact other symptoms within the network, and are therefore likely to be particularly important in the dynamics of PTSD development. Symptoms with strong in-strength, on the other hand, are those that are most affected by other symptoms in the network. For interventions, it has

been speculated that targeting symptoms with strong out-strength might provide an effective way of preventing or reducing future PTSD symptomatology (McNally, 2016; Armour *et al.* 2017; Bos *et al.* 2017).

In the current study, startle response had a greater out-strength than all other symptoms, which may suggest that targeting startle response could help reduce other symptoms. While arousal symptoms have been generally indicated as key drivers of future psychopathology in longitudinal studies of PTSD clusters (Schell *et al.* 2004; Marshall *et al.* 2006; Doron-LaMarca *et al.* 2015), startle has not specifically been found to be a central symptom in previous PTSD network studies with the exception of Spiller *et al.* (2017). It is likely that startle in the context of an ongoing threat is a different phenomenon compared with a startle when the stressor has ceased. Future studies could investigate whether startle has strong out-strength in other lower-stress environments. Additionally, restricted affect, blame, negative emotions, and avoidance of thoughts, also had high out-strength, all of which should be explored as the potential for interventions. It might also be argued that targeting symptoms that have both high in-strength and out-strength, such as sleep disturbance, could especially disrupt the network, facilitating an overall reduction in symptomatology.

Many symptoms in the temporal network had large autoregressive coefficients, i.e. predicting themselves at the next measurement, such as startle response and hypervigilance. This may indicate that these ‘symptoms’ may be more trait-like than others.



It could also be that these strong autocorrelations represent the first signs of a 'critical slowing down', whereby pathological responses gradually crystallize (Wichers *et al.* 2016).

The between-persons network largely resembled the contemporaneous network, although with fewer and weaker connections between the symptoms. This network is the one that can be most directly compared to previous cross-sectional network studies, and there are a number of similarities between them and the current between-persons network. Hypervigilance and startle have been found to be highly connected in a number of studies (e.g., McNally *et al.* 2015; Armour *et al.* 2017; Birkeland & Heir, 2017; Spiller *et al.* 2017). Flashbacks were identified as central symptoms by (Armour *et al.* 2017), while emotional cue reactivity was found to be the most central symptom by Spiller *et al.* (2017). Bryant *et al.* (2017) focused on early symptomatology and also noted the centrality of the re-experiencing symptoms (in a DSM-IV acute traumatic stress symptoms network). These findings are in line with our current finding of the centrality of both flashbacks and emotional cue reactivity. Interestingly, we found strong associations between the two DSM-5 avoidance symptoms, however other DSM-5 PTSD network studies do not find this strong association, relative to the associations between others symptoms. Furthermore, amnesia, while not being one of the most central symptoms, did not have especially weak connections, as found previously (Fried *et al.* 2017a). It is important to investigate whether these differences are because symptoms or reactions that arise during a period of trauma exposure are essentially different from those that emerge at a later stage when exposure has ceased.

There are some limitations of this study. We did not specifically include a sample with a diagnosis of PTSD, thus limiting our ability to generalize as to the associations between symptoms in those meeting DSM-5 PTSD criteria. Data were collected peritraumatically, during a conflict period, and were based on assessments conducted during both the first and second months since the traumatic stressor began, therefore the findings may not necessarily be generalizable to PTSD networks after exposure has ceased. Additionally, there are some methodological limitations with the multilevel network analysis model (Epskamp *et al.* 2016): there are no adequate fit statistics to assess the model; it only models linear processes yet there may be non-linear trends to consider; and it assumes stationarity, treating the properties of the network as constant over time, but this may not be the case, given what we know about PTSD resilience and recovery patterns (Bonanno & Mancini, 2012; Bryant *et al.* 2015), including in this particular sample (Gelkopf *et al.* 2017; Greene *et al.* 2017). Further development of time-varying multilevel network models may better clarify disorder development over time. Finally, there are other key variables that we would have liked to include in the model such as gender, trauma history, and severity of exposure to the conflict; however, power is a key limitation with these models (Fried & Cramer, 2017), and we did not have a large enough sample size to include these covariates.

This study provides a unique contribution to the PTSD research field through the novel use of ESM methodology to assess DSM-5 PTSD symptoms in individuals during a period of conflict, and through its employment of multilevel network analyses to explore contemporaneous and temporal network connections. While this study does not depict 'classic' PTSD, it is certainly relevant for many people around the world who live in situations of ongoing stress and trauma exposure. This study indicates that multilevel network analyses can provide novel and

useful insights into the ways that different symptoms interact with each other, as well as reinforcing themselves. More studies in this vein can elucidate the complex processes by which PTSD symptoms crystallize into disorder and indicate possible causal mechanisms that drive the development of PTSD. This, in turn, could have clinical implications for identifying the relative importance of symptoms as targets for interventions.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291718000351>.

**Acknowledgements.** The authors would like to thank Liron Lapid Pickman and Shulamit Grinapol for their contribution to establishing the study. This work was supported by The Brain and Behavior Research Foundation (Narsad Young Investigator Award, T.G., grant number 23524); The Moshe Hess Foundation (T.G. and M.G.); NATAL: Israel Trauma Center for Victims of Terror and War (M.G.); and the Israel Science Foundation (T.G. and M.G., grant number 1244/16).

**Declaration of interest.** None.

**Ethical standards.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

## References

- Afzali MH, Sunderland M, Batterham PJ, Carragher N, Callear A and Slade T (2017a) Network approach to the symptom-level association between alcohol use disorder and posttraumatic stress disorder. *Social Psychiatry and Psychiatric Epidemiology* 52(3), 329–339.
- Afzali MH, Sunderland M, Teesson M, Carragher N, Mills K and Slade T (2017b) A network approach to the comorbidity between posttraumatic stress disorder and major depressive disorder: the role of overlapping symptoms. *Journal of Affective Disorders* 208, 490–496.
- APA (2013). *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)*. Washington, DC: American Psychiatric Association.
- Armey MF, Schatten HT, Haradhvala N and Miller IW (2015) Ecological momentary assessment (EMA) of depression-related phenomena. *Current Opinion in Psychology* 4, 21–25.
- Armour C, Tsai J, Durham TA, Charak R, Biehn TL, Elhai JD *et al.* (2015) Dimensional structure of DSM-5 posttraumatic stress symptoms: support for a hybrid anhedonia and externalizing behaviors model. *Journal of Psychiatric Research* 61, 106–113.
- Armour C, Fried EI, Deserno MK, Tsai J and Pietrzak RH (2017) A network analysis of DSM-5 posttraumatic stress disorder symptoms and correlates in US military veterans. *Journal of Anxiety Disorders* 45, 49–59.
- Bak M, Drukker M, Hasmi L and van Os J (2016) An  $n = 1$  clinical network analysis of symptoms and treatment in psychosis. *PLoS ONE* 11, e0162811.
- Beard C, Millner A, Forgeard M, Fried E, Hsu K, Treadway M *et al.* (2016) Network analysis of depression and anxiety symptom relationships in a psychiatric sample. *Psychological Medicine* 46, 3359.
- Birkeland MS and Heir T (2017) Making connections: exploring the centrality of posttraumatic stress symptoms and covariates after a terrorist attack. *European Journal of Psychotraumatology* 8, 1333387.
- Blevins CA, Weathers FW, Davis MT, Witte TK and Domino JL (2015) The posttraumatic stress disorder checklist for DSM-5 (PCL-5): development and initial psychometric evaluation. *Journal of Traumatic Stress* 28, 489–498.
- Bolger N and Laurenceau J (2013) *Intensive Longitudinal Methods. An Introduction to Diary and Experience Sampling Research*. New York: Guilford.
- Bolger N, Davis A and Rafaeli E (2003) Diary methods: capturing life as it is lived. *Annual Review of Psychology* 54, 579–616.
- Bonanno GA and Mancini AD (2012) Beyond resilience and PTSD: mapping the heterogeneity of responses to potential trauma. *Psychological Trauma: Theory, Research, Practice, and Policy* 4, 74.

- Borsboom D** (2017) A network theory of mental disorders. *World Psychiatry* **16**, 5–13.
- Bos F, Snippe E, de Vos S, Hartmann J, Simons C, van der Krieke L et al.** (2017) Can We jump from cross-sectional to dynamic interpretations of networks? Implications for the network perspective in psychiatry. *Psychotherapy and Psychosomatics* **86**, 175.
- Bryant RA, Nickerson A, Creamer M, O'Donnell M, Forbes D, Galatzer-Levy I et al.** (2015) Trajectory of post-traumatic stress following traumatic injury: 6-year follow-up. *The British Journal of Psychiatry* **206**, 417–423.
- Bryant RA, Creamer M, O'Donnell M, Forbes D, McFarlane AC, Silove D et al.** (2017). Acute and chronic posttraumatic stress symptoms in the emergence of posttraumatic stress disorder: a network analysis. *JAMA Psychiatry* **74**(2), 135–142.
- Chun CA** (2016) The expression of posttraumatic stress symptoms in daily life: a review of experience sampling methodology and daily diary studies. *Journal of Psychopathology and Behavioral Assessment* **38**, 406–420.
- David SJ, Marshall AJ, Evanovich EK and Mumma GH** (2017) Intraindividual dynamic network analysis—implications for clinical assessment. *Journal of Psychopathology and Behavioral Assessment*, 1–14.
- Doron-LaMarca S, Niles BL, King DW, King LA, Pless Kaiser A and Lyons MJ** (2015) Temporal associations among chronic PTSD symptoms in US combat veterans. *Journal of Traumatic Stress* **28**, 410–417.
- Ebner-Priemer UW and Kubiak T** (2010) The decade of behavior revisited: future prospects for ambulatory assessment. *European Journal of Psychological Assessment* **26**, 151–153.
- Ebner-Priemer UW and Trull TJ** (2009) Ecological momentary assessment of mood disorders and mood dysregulation. *Psychological Assessment* **21**, 463.
- Epskamp S, Waldorp LJ, Möttus R and Borsboom D** (2016) Discovering psychological dynamics: the Gaussian graphical model in cross-sectional and time-series data. arXiv:1609.04156.
- Epskamp S, van Borkulo CD, van der Veen DC, Servaas MN, Isvoranu A-M, Riese H et al.** (2017) Personalized network modeling in psychopathology: the importance of contemporaneous and temporal connections. *Clinical Psychological Science*. <https://doi.org/10.1177/2167702617744325>
- Fisher AJ** (2015) Toward a dynamic model of psychological assessment: implications for personalized care. *Journal of Consulting and Clinical Psychology* **83**, 825.
- Fisher AJ, Reeves JW, Lawyer G, Medaglia JD and Rubel JA** (2017) Exploring the idiographic dynamics of mood and anxiety via network analysis. *Journal of Abnormal Psychology* **126**, 1044.
- Fried EI and Cramer AO** (2017) Moving forward: challenges and directions for psychopathological network theory and methodology. *Perspectives on Psychological Science* **12**(6), 999–1020.
- Fried EI, Epskamp S, Nesse RM, Tuerlinckx F and Borsboom D** (2016) What are 'good' depression symptoms? Comparing the centrality of DSM and non-DSM symptoms of depression in a network analysis. *Journal of Affective Disorders* **189**, 314–320.
- Fried EI, Eidhof M, Palic S, Costantini G, Huisman-van Dijk H, Bockting CL et al.** (2017a) Replicability and generalizability of PTSD networks: a cross-cultural multisite study of PTSD symptoms in four trauma patient samples. *Clinical Psychological Science* [accepted].
- Fried EI, van Borkulo CD, Cramer AO, Boschloo L, Schoevers RA and Borsboom D** (2017b) Mental disorders as networks of problems: a review of recent insights. *Social Psychiatry and Psychiatric Epidemiology* **52**, 1–10.
- Gelkopf M, Lapid L, Grinapol S, Werbeloff N, Carlson E and Greene T** (2017) Peritraumatic reaction courses during War: gender, serious mental illness, and exposure. *Psychiatry: Interpersonal & Biological Processes*.
- Granger CW** (1969) Investigating causal relations by econometric models and cross-spectral methods. *Econometrica: Journal of the Econometric Society* **424–438**.
- Greene T** (under review) Blame, PTSD and DSM-5: An urgent need for clarification. *European Journal of Psychotraumatology*.
- Greene T, Gelkopf M, Grinapol S, Werbeloff N, Carlson E and Lapid L** (2017) Trajectories of traumatic stress symptoms during conflict: a latent class growth analysis. *Journal of Affective Disorders*.
- Hamaker E** (2012) Why researchers should think “within-person”: a paradigmatic rationale. *Handbook of research methods for studying daily life*, 43–61.
- Hamaker EL and Wichers M** (2017) No time like the present: discovering the hidden dynamics in intensive longitudinal data. *Current Directions in Psychological Science* **26**, 10–15.
- Isvoranu A-M, Borsboom D, van Os J and Guloksuz S** (2016) A network approach to environmental impact in psychotic disorder: brief theoretical framework. *Schizophrenia Bulletin* **42**, 870–873.
- Kelley HH and Michela JL** (1980) Attribution theory and research. *Annual Review of Psychology* **31**, 457–501.
- Knefel M, Tran US and Lueger-Schuster B** (2016) The association of post-traumatic stress disorder, complex posttraumatic stress disorder, and borderline personality disorder from a network analytical perspective. *Journal of Anxiety Disorders* **43**, 70–78.
- Lapid Pickman L, Greene T and Gelkopf M** (2017) Sense of threat as a mediator of peritraumatic stress symptom development during wartime: an experience sampling study. *Journal of Traumatic Stress* **30**, 372–380.
- Levine SZ and Leucht S** (2016) Identifying a system of predominant negative symptoms: network analysis of three randomized clinical trials. *Schizophrenia Research* **178**, 17–22.
- Marshall GN, Schell TL, Glynn SM and Shetty V** (2006) The role of hyperarousal in the manifestation of posttraumatic psychological distress following injury. *Journal of Abnormal Psychology* **115**, 624.
- McNally RJ** (2016) Can network analysis transform psychopathology? *Behaviour Research and Therapy* **86**, 95–104.
- McNally RJ, Robinaugh DJ, Wu GW, Wang L, Deserno MK and Borsboom D** (2015) Mental disorders as causal systems a network approach to posttraumatic stress disorder. *Clinical Psychological Science* **3**, 836–849.
- Mitchell K, Wolf E, Bovin M, Lee L, Green J, Rosen R, et al.** (2017) Network models of DSM-5 posttraumatic stress disorder: implications for ICD-11. *Journal of Abnormal Psychology*.
- Myin-Germeys I, Oorschot M, Collip D, Lataster J, Delespaul P and van Os J** (2009) Experience sampling research in psychopathology: opening the black box of daily life. *Psychological Medicine* **39**, 1533–1547.
- Nelson B, McGorry PD, Wichers M, Wigman JT and Hartmann JA** (2017) Moving from static to dynamic models of the onset of mental disorder: a review. *JAMA Psychiatry* **74**, 528–534.
- Pietrzak RH, Tsai J, Armour C, Mota N, Harpaz-Rotem I and Southwick SM** (2015) Functional significance of a novel 7-factor model of DSM-5 PTSD symptoms: results from the national health and resilience in veterans study. *Journal of Affective Disorders* **174**, 522–526.
- Santangelo PS, Ebner-Priemer UW and Trull TJ** (2013) Experience sampling methods in clinical psychology. *The Oxford Handbook of Research Strategies for Clinical Psychology*, 188–210.
- Schell TL, Marshall GN and Jaycox LH** (2004) All symptoms are not created equal: the prominent role of hyperarousal in the natural course of post-traumatic psychological distress. *Journal of Abnormal Psychology* **113**, 189.
- Schuurman NK, Ferrer E, de Boer-Sonnenschein M and Hamaker EL** (2016) How to compare cross-lagged associations in a multilevel autoregressive model. *Psychological Methods* **21**, 206.
- Spiller TR, Schick M, Schnyder U, Bryant RA, Nickerson A and Morina N** (2017) Symptoms of posttraumatic stress disorder in a clinical sample of refugees: a network analysis. *European Journal of Psychotraumatology* **8**, 1318032.
- Sullivan CP, Smith AJ, Lewis M and Jones RT** (2018) Network analysis of ptsd symptoms following mass violence. *Psychological Trauma: Theory, Research, Practice, and Policy* **10**(1), 58.
- Trull TJ and Ebner-Priemer U** (2013) Ambulatory assessment. *Annual Review of Clinical Psychology* **9**, 151.
- Walz LC, Nauta MH and aan het Rot M** (2014) Experience sampling and ecological momentary assessment for studying the daily lives of patients with anxiety disorders: a systematic review. *Journal of Anxiety Disorders* **28**, 925–937.
- Weathers F, Litz B, Keane T, Palmieri P, Marx B and Schnurr P** (2013) The PTSD checklist for DSM-5 (PCL-5). *Scale available from the National Center for PTSD*. [www.ptsd.va.gov](http://www.ptsd.va.gov).
- Wichers M, Groot PC, Psychosystems E and Group E** (2016) Critical slowing down as a personalized early warning signal for depression. *Psychotherapy and Psychosomatics* **85**, 114–116.