



Universiteit
Leiden
The Netherlands

Into the darkness : forging a stable path through the gravitational landscape

Papadomanolakis, G.

Citation

Papadomanolakis, G. (2019, September 19). *Into the darkness : forging a stable path through the gravitational landscape*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/78471>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/78471>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/78471> holds various files of this Leiden University dissertation.

Author: Papadomanolakis, G.

Title: Into the darkness : forging a stable path through the gravitational landscape

Issue Date: 2019-09-19

3 On the stability conditions for theories of modified gravity in the presence of matter fields

3.1 Introduction

In the previous Chapter we started to study the set of conditions an extension of gravity must satisfy in order to guarantee a viable cosmological scenario. This was done by considering the model in a vacuum and as such does not represent our universe as it contains matter fields which are gravitationally coupled to the new d.o.f.. In fact, the stability conditions might be altered by the presence of the additional matter fields, thus changing the viability space of the theory [37, 67, 112–115]. Identifying the correct viability requirements is important when testing MG theories with cosmological data by using statistical tools [22, 23, 116, 117], as they can reduce the viability space one needs to explore. Additionally they can even dominate over the constraining power of observational data as recently shown in the case of designer $f(R)$ -theory on w CDM background [23]. Therefore, in this Chapter we proceed to fix this deficiency and to quantify the modifications induced by the new fields.

With the aim to obtain general results, we will employ once more the Effective Field Theory of Dark Energy and Modified Gravity [14, 15]. This EFToDE/MG approach has been implemented into the Einstein Boltzmann solver, CAMB/CosmoMC [20, 21, 118], creating *EFT-CAMB/EFTCosmoMC* [22, 23, 39, 80, 82, 119] (<http://www.eftcamb.org/>), providing a perfect tool to test gravity models through comparison with observational data. *EFTCAMB* comes with a built-in module to explore the viability space of the underlying theory of gravity, which then can be used as *priors*. The results of the present work have a direct application as they can be employed to improve the current *EFTCAMB* viability requirements but not limited to it as they can be easily mapped to other parametrizations [80].

3 Stability conditions in the presence of matter

For the matter sector we chose to employ the Sorkin-Schutz action, which allows one to treat general matter fluids [120, 121]. Among many models used to describe matter Lagrangians and which have been extensively used and investigated in the past years [37, 67, 112–115], we choose to follow the recent arguments in ref. [122]. Indeed, it has been shown that such an action, along with an appropriate choice for the matter field, describes the dynamics of all matter fluids avoiding some problems which might arise when including pressure-less matter fluids, like dust or cold dark matter (CDM), which are relevant in the evolution of the Universe.

Previously, a stability analysis of the EFToDE/MG in the presence of matter had been done in [37]. However, in our work we present also the conditions which allow to avoid tachyonic instabilities and we analyse all possible sub-cases concerning the stability conditions. Furthermore, due to the different choice of matter action, modifications can be seen when one includes pressureless fluids.

With this machinery, we proceed to derive the viability constraints one needs to impose on the free parameters of the theory by focusing on three sources of possible instabilities, ghost, gradient and tachyonic instabilities. We will proceed while retaining the full generality of the EFToDE/MG approach, i.e. without limiting to specific models. However, where relevant, we will make connections to specific theories, such as low-energy Hořava gravity [35, 36, 55] and beyond Horndeski models [66] and we will analyse the results within the context of these models.

This Chapter is based on the work in [32]: *On the stability conditions for theories of modified gravity in the presence of matter fields* with A. De Felice and N. Frusciante. In section 3.2, we briefly recap the EFToDE/MG formalism we use to parametrize the DE/MG models with one extra scalar d.o.f.. In section 3.3, we introduce the Sorkin-Schutz action to describe the dynamics of matter fluids and we discuss the advantage of using this action with respect to previous approaches. We also work out the corresponding continuity equation and second order perturbed action. In section 3.4, we work out the action for both gravity and matter fields up to second order in perturbations. Then, we calculate and discuss the stability requirements to avoid ghost instabilities (section 3.4.1), to guarantee positive speeds of propagation (section 3.4.2) and to prevent tachyonic instabilities (section 3.4.3). Finally, we conclude in section 4.6.

3.2 The Effective Field Theory of Dark Energy and Modified Gravity

The EFToDE/MG has been proposed as a unifying framework to study the dynamic and evolution of linear order perturbations of a broad

3.2 The Effective Field Theory of Dark Energy and Modified Gravity

class of DE/MG theories [14, 15]. This approach encompasses all the theories of gravity exhibiting one extra scalar and dynamical d.o.f. and admits a well-defined Jordan frame. As discussed in the Introduction, the EFToDE/MG is constructed in the unitary gauge by expanding around the Friedmann-Lemaitre-Robertson-Walker (FLRW) metric. Each operator is accompanied by a time dependent function dubbed *EFT function*. The explicit form of the perturbed EFToDE/MG action is the following:

$$\begin{aligned} \mathcal{S}_{EFT} = \int d^4x \sqrt{-g} \left[\frac{m_0^2}{2} (1 + \Omega(t)) R^{(4)} + \Lambda(t) - c(t) \delta g^{00} + \frac{M_2^4(t)}{2} (\delta g^{00})^2 \right. \\ \left. - \frac{\bar{M}_1^3(t)}{2} \delta g^{00} \delta K - \frac{\bar{M}_2^2(t)}{2} (\delta K)^2 - \frac{\bar{M}_3^2(t)}{2} \delta K_{\mu\nu} \delta K_{\mu\nu} \right. \\ \left. + \frac{\hat{M}^2(t)}{2} \delta g^{00} \delta R^{(3)} + m_2^2(t) (g^{\mu\nu} + n^\mu n^\nu) \partial_\mu g^{00} \partial_\nu g^{00} \right], \quad (3.1) \end{aligned}$$

where m_0^2 is the Planck mass, $g_{\mu\nu}$ is the four dimensional metric and g is its determinant, δg^{00} is the perturbation of the upper time-time component of the metric, n_μ is the normal vector to the constant-time hypersurfaces, $R^{(4)}$ and $R^{(3)}$ are respectively the trace of the four dimensional and three dimensional Ricci scalar, $K_{\mu\nu}$ is the extrinsic curvature and K is its trace. Finally, with $\delta A = A - A^{(0)}$ we indicate the linear perturbation of the quantity A and $A^{(0)}$ is the corresponding background value.

Moreover, it has been shown that appropriate combinations of the EFT functions in action (5.1) allows one to describe specific classes of DE/MG models. We group such combinations as follows:

- $M_2^2 = -\bar{M}_3^2 = 2\hat{M}^2$ and $m_2^2 = 0$: Horndeski [78] or Generalized Galileon class of models [79] (and all the models belonging to them);
- $M_2^2 + \bar{M}_3^2 = 0$ and $m_2^2 = 0$: Beyond Horndeski class of models [66];
- $m_2^2 \neq 0$: Lorentz violating theories (e.g. low-energy Hořava gravity [35, 36, 55]).

For a detailed guide to map a specific theory into the EFToDE/MG language we refer the reader to the previous Chapter as well as refs. [14, 15, 37, 65]. Finally, an extended version of the above EFToDE/MG action has been presented in the previous Chapter which includes operators with higher than second order spatial derivatives.

In the following we will briefly recap the construction of the EFToDE/MG action up to second order in terms of the scalar metric perturbations as it will be the starting point for the stability analysis.

3 Stability conditions in the presence of matter

Because of the unitary gauge in action (3.1), it is natural to choose the Arnowitt-Deser-Misner (ADM) formalism [19] to write the line element, which reads:

$$ds^2 = -N^2 dt^2 + h_{ij}(dx^i + N^i dt)(dx^j + N^j dt), \quad (3.2)$$

where $N(t, x^i)$ is the lapse function, $N^i(t, x^i)$ the shift and $h_{ij}(t, x^i)$ is the metric tensor of the three dimensional spatial slices. Proceeding with the expansion around a flat FLRW background, the metric can be written as:

$$ds^2 = -(1 + 2\delta N)dt^2 + 2\partial_i\psi dt dx^i + a^2(1 + 2\zeta)\delta_{ij}dx^i dx^j, \quad (3.3)$$

where as usual $\delta N(t, x^i)$ is the perturbation of the lapse function, $\partial_i\psi(t, x^i)$ and $\zeta(t, x^i)$ are the scalar perturbations respectively of N_i and h_{ij} and a is the scale factor. Then, the scalar perturbations of the quantities involved in the action (3.1) are:

$$\begin{aligned} \delta g^{00} &= 2\delta N, \\ \delta K &= -3\dot{\zeta} + 3H\delta N + \frac{1}{a^2}\partial^2\psi, \\ \delta K_{ij} &= a^2\delta_{ij}(H\delta N - 2H\zeta - \dot{\zeta}) + \partial_i\partial_j\psi, \\ \delta K_j^i &= (H\delta N - \dot{\zeta})\delta_j^i + \frac{1}{a^2}\partial^i\partial_j\psi, \\ \delta R^{(3)} &= -\frac{4}{a^2}\partial^2\zeta, \end{aligned} \quad (3.4)$$

where we have made use of the following definitions of the normal vector and extrinsic curvature:

$$n_\mu = N\delta_\mu^0, \quad K_{\mu\nu} = h_\mu^\lambda\nabla_\lambda n_\nu, \quad (3.5)$$

with $h^{\mu\nu} = g^{\mu\nu} + n^\mu n^\nu$, $H \equiv \frac{1}{a}\frac{da}{dt}$ is the Hubble function and dots are the derivatives with respect to time. Then, the action (3.1) can be explicitly expanded in terms of metric scalar perturbations up to second order and after some manipulations, we obtain the following final form:

$$\begin{aligned} S_{EFT}^{(2)} &= \int dt d^3x a^3 \left\{ -\frac{F_4(\partial^2\psi)^2}{2a^4} - \frac{3}{2}F_1\dot{\zeta}^2 + m_0^2(\Omega + 1)\frac{(\partial\zeta)^2}{a^2} \right. \\ &\quad - \frac{\partial^2\psi}{a^2}(F_2\delta N - F_1\dot{\zeta}) + 4m_2^2\frac{[\partial(\delta N)]^2}{a^2} + \frac{F_3}{2}\delta N^2 \\ &\quad \left. + \left[3F_2\dot{\zeta} - 2(m_0^2(\Omega + 1) + 2\hat{M}^2)\frac{\partial^2\zeta}{a^2} \right] \delta N \right\}, \end{aligned} \quad (3.6)$$

where we have defined

$$\begin{aligned}
 F_1 &= 2m_0^2(\Omega + 1) + 3\bar{M}_2^2 + \bar{M}_3^2, \\
 F_2 &= HF_1 + m_0^2\dot{\Omega} + \bar{M}_1^3, \\
 F_3 &= 4M_2^4 + 2c - 3H^2F_1 - 6m_0^2H\dot{\Omega} - 6H\bar{M}_1^3, \\
 F_4 &= \bar{M}_2^2 + \bar{M}_3^2,
 \end{aligned} \tag{3.7}$$

and other terms vanish because of the background equations of motion. This result will be considered along with the matter sector which will be presented in the next section in order to facilitate the complete study of conditions that guarantee that a gravity theory, in presence of matter fields, is free from instabilities.

3.3 The Matter Sector

The goal of the present work is to investigate the emergence of instabilities in modified theories of gravity under the influence of matter fluids and subsequently set appropriate stability conditions. Therefore, a crucial step is to make the appropriate choice for the matter action, S_m . Moreover, the generality of the EFTtoDE/MG approach in describing the gravity sector makes that even for the matter action there is an equally general treatment. It is common in literature to choose for the matter Lagrangian a k-essence like form, $P(\mathcal{X})$ [37, 111–114, 123, 124], to model the matter d.o.f. where $\mathcal{X} \equiv \chi_{;\mu}\chi^{;\mu}$ is the kinetic term of the field χ . However, this choice displays problematic behaviours which motivates us to decide for a different action. The easiest way of identifying those issues is to consider the corresponding action for $P(\mathcal{X})$ when it has been specialized to a dust fluid. In that case it can be easily shown that the action diverges. Subsequently, in ref. [122], it has been shown that the real problem arising in the K-essence like matter Lagrangian lies in the choice of the canonical field one uses to describe the d.o.f. of the fluid. The usual choice for the fluid variable, the velocity v_m , satisfies a closed first order equation of motion, which requires only one independent initial condition. Then, the dust fluid would have only one d.o.f. (rather than two) and for that field the action tends to blow up as the speed of propagation goes to zero ($c_{s,d}^2 \rightarrow 0$). Instead, the appropriate variable for the fluid is the matter density perturbation, δ_m .

In order to avoid the issues described above we choose the Sorkin-Schutz action, see refs. [120, 121] which is well defined for a dust component and can describe in full generality perfect fluids. As observed above the appropriate fluid variable is the density perturbation which is exactly the one employed by this action and thus satisfies a second order equation

3 Stability conditions in the presence of matter

of motion as will be evident in the following. The Sorkin-Schutz matter action reads:

$$S_m = - \int d^4x [\sqrt{-g} \rho(n) + J^\nu \partial_\nu \ell], \quad (3.8)$$

where ρ is the energy density, which depends on the number density n , ℓ is a scalar field, whereas J^ν is a vector with weight one. Additionally, we define n as

$$n = \sqrt{\frac{J^\alpha J^\beta g_{\alpha\beta}}{g}}. \quad (3.9)$$

Then, the four velocity vector u^α is defined as

$$u^\alpha = \frac{J^\alpha}{n\sqrt{-g}}, \quad (3.10)$$

and satisfies the usual relation $u^\alpha u_\alpha = -1$. Variation of the matter Lagrangian with respect to J^α leads to

$$u_\alpha = \frac{1}{\partial\rho/\partial n} \partial_\alpha \ell, \quad (3.11)$$

while taking its variation with respect to the metric we find that the stress energy tensor can be defined as

$$T_{\alpha\beta} \equiv \frac{2}{\sqrt{-g}} \frac{\delta S_m}{\delta g^{\alpha\beta}} = n \frac{\partial\rho}{\partial n} u_\alpha u_\beta + \left(n \frac{\partial\rho}{\partial n} - \rho \right) g_{\alpha\beta}, \quad (3.12)$$

which is a barotropic perfect fluid with pressure given by

$$p \equiv n \frac{\partial\rho}{\partial n} - \rho. \quad (3.13)$$

Let us notice that a particular choice for the density, i.e. $\rho \propto n^{1+w}$, allows to have the usual relation $p = w\rho$, where w is the barotropic coefficient. Finally, by varying the matter action with respect to ℓ , one gets the conservation constraint

$$\partial_\alpha J^\alpha = 0. \quad (3.14)$$

On a flat FLRW background the above relation gives $J^0 = \mathcal{N}_0$, where \mathcal{N}_0 is the total particle number and from Eq. (3.9) we have $n = \mathcal{N}_0/a^3$.

Let us now proceed to write the matter action (3.8) up to second order in the scalar fields by using the metric scalar perturbations in Eq. (3.3). For the fluid variables we proceed to expand them as follows

$$\begin{aligned} J^0 &= \mathcal{N}_0 + \delta J, \\ J^i &= \frac{1}{a^2} \partial^i \delta j, \\ \ell &= - \int^t \frac{\partial\rho}{\partial n} dt' - \frac{\partial\rho}{\partial n} v_m, \end{aligned} \quad (3.15)$$

3.4 Study of Stability conditions

where v_m is the velocity of the matter species. Furthermore, we note that since

$$\rho = \bar{\rho} + \frac{\partial \rho}{\partial n} \left(-\frac{3\mathcal{N}_0}{a^3} \zeta + \frac{\delta J}{a^3} \right) \equiv \bar{\rho} + \delta \rho, \quad (3.16)$$

where $\bar{\rho}$ is the density at the background, one can obtain

$$\delta J = \frac{a^3 \bar{\rho} \delta_m}{\partial \rho / \partial n} + 3\mathcal{N}_0 \zeta, \quad (3.17)$$

where, as usual, $\delta_m = \delta \rho / \bar{\rho}$. We can thus rewrite δJ in terms of δ_m in the perturbed matter action. Finally, we can use the equation of motion for δj

$$\delta j = -\mathcal{N}_0 (\psi + v_m) \quad (3.18)$$

in order to eliminate it in favour of v_m and ψ .

Combining the above results and after some integrations by parts, we obtain the action for the scalar perturbations up to second order:

$$\begin{aligned} S_m^{(2)} = & \int dt d^3x a^3 \left[-\frac{n\rho_{,n}(\partial v)^2}{2a^2} + \left(\frac{3H(n\rho_{,n}^2 - n\bar{\rho}\rho_{,nn} - \bar{\rho}\rho_{,n})\delta_m}{\rho_{,n}} \right. \right. \\ & \left. \left. + \frac{n\rho_{,n}\partial^2\psi}{a^2} - 3n\rho_{,n}\dot{\zeta} - \bar{\rho}\dot{\delta}_m \right) v_m - \frac{\rho_{,nn}\bar{\rho}^2\delta_m^2}{2\rho_{,n}^2} - \bar{\rho}\delta N\delta_m \right]. \end{aligned} \quad (3.19)$$

Notice that the velocity v_m can always be integrated out, as $n\rho_{,n} = \bar{\rho} + p \neq 0$.

3.4 Study of Stability conditions

In this section we present the main bulk of our work, i.e. the study of the general conditions that a gravity theory has to satisfy in order to be free from instabilities when additional matter fields are considered. These set of requirements include: no ghost instabilities, positive speeds of propagation (squared) and no tachyonic instabilities as presented in the Introduction. Recently, it has been shown that physical stability plays an important role when testing specific gravity models with cosmological data [23, 125]. In particular, the *EFTCAMB* patch [22, 23] includes a specific module with the task to identify the viable parameter space of a selected theory. The results of the present work can be used to improve such modules and improve on the efficiency of the selection process.

To achieve this goal we consider the general EFToDE/MG parametrization presented in section 3.2 in the presence of two different matter fluids, described by the action (3.19), for which we made the following, realistic,

3 Stability conditions in the presence of matter

choices: a pressure-less fluid, i.e. cold dark matter/dust (d) and radiation (r). A treatment which includes two general fluids complicates the process substantially and we do not expect to learn much more in such a case. So the relevant action required in order to proceed is of the following form:

$$\begin{aligned}
S^{(2)} &= \frac{1}{(2\pi)^3} \int dt d^3ka^3 \left\{ \bar{\rho}_d \left(-\frac{k^2\psi}{a^2} - 3\dot{\zeta} - \dot{\delta}_d \right) v_d + \bar{\rho}_r \left(-\frac{4}{3} \frac{k^2\psi}{a^2} \right. \right. \\
&- \left. \left. 4\dot{\zeta} - \dot{\delta}_r \right) v_r - \bar{\rho}_d \frac{(kv_d)^2}{2a^2} - \frac{2}{3} \bar{\rho}_r \frac{(kv_r)^2}{a^2} - \frac{F_4}{2a^4} (k^2\psi)^2 \right. \\
&+ \left. \left(\frac{2k^2\zeta \left(2\hat{M}^2 + m_0^2(\Omega + 1) \right)}{a^2} + 3F_2\dot{\zeta} \right) \delta N + \left(\delta N F_2 - F_1\dot{\zeta} \right) \frac{k^2\psi}{a^2} + \right. \\
&+ \frac{4m_2^2(k\delta N)^2}{a^2} + \frac{m_0^2(\Omega + 1)(k\zeta)^2}{a^2} - \frac{4}{3} \bar{\rho}_r \frac{(kv_r)^2}{2a^2} - \bar{\rho}_d \delta N \delta_d \\
&\left. + \frac{1}{2} F_3 \delta N^2 - \frac{3}{2} F_1 \dot{\zeta}^2 - \frac{\bar{\rho}_r}{8} \delta_r^2 - \bar{\rho}_r \delta N \delta_r \right\}, \tag{3.20}
\end{aligned}$$

where we have Fourier transformed the spatial coordinates and we have considered the following relations for the number densities:

$$n_d = \bar{\rho}_d, \quad n_r = (\bar{\rho}_r)^{\frac{3}{4}}, \tag{3.21}$$

being $\bar{\rho}_d, \bar{\rho}_r$ respectively the density of dust and radiation at background.

An action constructed in such a way admits only three d.o.f. described by $\{\zeta, \delta_d, \delta_r\}$. Therefore in the above action we notice the presence of four Lagrange multipliers $\delta N, \psi, v_d$ and v_r . Consequently, we proceed with the removal of the latter by using the constraint equations obtained after the variations of the action with respect to the Lagrange multipliers. The resulting set of constraint equations is:

$$\begin{aligned}
&\bar{\rho}_r \left(-\frac{4}{3} \frac{k^2\psi}{a^2} - 4\dot{\zeta} - \dot{\delta}_r \right) - \frac{4}{3} \bar{\rho}_r \frac{k^2 v_r}{a^2} = 0, \\
&\bar{\rho}_d \left(-\frac{k^2\psi}{a^2} - 3\dot{\zeta} - \dot{\delta}_d \right) - \bar{\rho}_d \frac{k^2 v_d}{a^2} = 0, \\
&\frac{2k^2\zeta \left(2\hat{M}^2 + m_0^2(\Omega + 1) \right)}{a^2} + 3F_2\dot{\zeta} + \frac{8m_2^2 k^2 \delta N}{a^2} + F_2 \frac{k^2\psi}{a^2} - \bar{\rho}_d \delta_d \\
&+ F_3 \delta N - \bar{\rho}_r \delta_r = 0, \\
&-\bar{\rho}_d v_d - \frac{4}{3} \bar{\rho}_r v_r + \delta N F_2 - F_1 \dot{\zeta} - \frac{F_4}{a^2} k^2 \psi = 0. \tag{3.22}
\end{aligned}$$

After solving for the auxiliary fields and substituting the results back into action (3.20), we get an action containing only the three dynamical

d.o.f. $\{\zeta, \delta_d, \delta_r\}$:

$$S^{(2)} = \frac{1}{(2\pi)^3} \int d^3k dt a^3 \left(\dot{\vec{\chi}}^t \mathbf{A} \dot{\vec{\chi}} - k^2 \vec{\chi}^t \mathbf{G} \vec{\chi} - \dot{\vec{\chi}}^t \mathbf{B} \vec{\chi} - \vec{\chi}^t \mathbf{M} \dot{\vec{\chi}} \right), \quad (3.23)$$

where we have defined the dimensionless vector:

$$\vec{\chi}^t = (\zeta, \delta_d, \delta_r), \quad (3.24)$$

and the matrix components are listed in Appendix 3.6. In the next sections we will derive the stability conditions one needs to impose on the above action in order to guarantee the viability of the underlying theory of gravity.

Before proceeding with this in-depth analysis of the final action we present the background equations corresponding to our set-up:

$$\begin{aligned} E_1 &\equiv 3m_0^2 \left[1 + \Omega + a \frac{d\Omega}{da} \right] H^2 + \Lambda - 2c - \sum \bar{\rho}_i = 0, \\ E_2 &\equiv m_0^2 (1 + \Omega) (3H^2 + 2\dot{H}) + 2m_0^2 H \dot{\Omega} + m_0^2 \ddot{\Omega} + \sum_i p_i + \Lambda = 0, \\ E_i &\equiv \dot{\bar{\rho}}_i + 3H(\bar{\rho}_i + p_i) = 0. \end{aligned} \quad (3.25)$$

where the Friedmann equations have been supplemented by the continuity equations for the fluids. Finally, in order to close the system of equations, one needs to use the well-known equations of state for dust and radiation. As a side comment, from the background equations it is not possible to define in general a modified gravitational constant because c and Λ can be functions of H^2 . The latter statement is clear when looking at the mapping of specific theories in the EFToDE/MG language as done in Chapter 2.

3.4.1 The presence of ghosts

A negative kinetic term of a field is usually considered as a pathology of the theory, since the high energy vacuum is unstable to the spontaneous production of particles [29]. Such a pathology must be constrained demanding for a positive kinetic term.

Recently in ref. [126], it has been shown that such a constraint has to be imposed only in the high energy regime, in other words, an infrared ghost does not lead to a catastrophic vacuum collapse. On the contrary it was shown that it corresponds to a well known physical phenomenon, the Jeans instability.

In fact, expanding the ghost conditions in high- k one can show that, when using appropriate field re-definitions, the sub-leading terms can

3 Stability conditions in the presence of matter

be recast into the form of a Jeans mass instability, and viceversa. For example, the Hamiltonian $\mathcal{H} = -P^2 + Q^2$ (where a ghost is present), can be recast into $\mathcal{H} = p^2 - q^2$ (with negative squared speed of propagation and/or tachyonic mass), upon using the trivial canonical transformation $Q = p$, $P = -q$. Therefore, we will consider only the constraints coming from the high- k behaviour for the ghost conditions as only in this regime they correspond to a true theoretical instability and not to a hidden physical phenomenon. As for the tachyonic squared mass (i.e. negative mass), it is problematic only when the time of evolution of the instability is much larger than H^2 . We will elaborate on the latter in section 3.4.3.

Although the EFTtoDE/MG approach has been discussed in the context of energies smaller than the cut-off of the theory, $\Lambda_{\text{cut-off}}$, here and in the following we will assume that we can still perform a high- k expansion, namely we assume that in this regime we have $H \ll k/a \ll \Lambda_{\text{cut-off}}$. This assumption is assumed to be valid at least for medium-low redshifts, those for which we can apply all the known cosmological-data constraints, namely BBN, CMB, BAO, etc.

In action (3.23) we have a non-diagonal kinetic matrix for the three fields, i.e. $\mathcal{L} \ni A_{ij} \dot{\chi}_i \dot{\chi}_j$. As previously mentioned, in order to guarantee the absence of ghosts, one needs to demand the high- k limit of the kinetic matrix to be positive definite. It is clear that one case encompassing all viable theories does not exist as a result of the wide range of operators which depend differently on the momentum. In particular, one has to pay attention to the operators accompanying \bar{M}_2^3 , \bar{M}_2^2 and m_2^2 , which exhibit a higher order dependence on k . Therefore, we will present a number of clear sub-cases which we consider relevant

We can identify a few cases:

1. In this case all the functions in the Lagrangian are present, in particular $m_2^2 \neq 0$ and $F_4 \neq 0$. As a reference we note that the low-energy Hořava gravity belongs to this general case. Expanding at high- k , we find

$$\mathcal{G}_1 = \frac{(F_1 - 3F_4) a^3 F_1}{2F_4} > 0, \quad (3.26)$$

$$\mathcal{G}_l = \frac{a^5 \bar{\rho}_l^2}{2k^2(\bar{\rho}_l + p_l)} > 0, \quad (3.27)$$

where the index l indicates the matter components, i.e. dust and radiation, $\mathcal{G}_1 \equiv \text{Det}(\mathbf{A})/(A_{22}A_{33} - A_{23}^2)$, $\mathcal{G}_r = A_{33}$ and $\mathcal{G}_d = A_{22} - A_{23}^2/A_{33}$. The \mathcal{G}_l conditions represent the standard matter no-ghost conditions, which are trivially satisfied.

2. $F_4 = 0 = m_2^2$. This case corresponds to the well known class of

3.4 Study of Stability conditions

beyond Horndeski theories. We find:

$$\mathcal{G}_1 = \frac{(F_1 F_3 + 3 F_2^2) F_1 a^3}{2 F_2^2} > 0, \quad (3.28)$$

$$\mathcal{G}_l = \frac{a^5 \bar{\rho}_l^2}{2 k^2 (\bar{\rho}_l + p_l)} > 0. \quad (3.29)$$

3. $F_4 = 0$ and $m_2^2 \neq 0$. The ghost conditions change into

$$\mathcal{G}_1 = \frac{4 F_1^2 m_2^2 k^2 a}{F_2^2} > 0, \quad (3.30)$$

$$\mathcal{G}_d = \frac{F_2^2 a^5 \bar{\rho}_d}{2 k^2 (F_2^2 - 8 m_2^2 \bar{\rho}_d)} > 0, \quad (3.31)$$

$$\mathcal{G}_r = \frac{9 a^5 (F_2^2 - 8 m_2^2 \bar{\rho}_d) \bar{\rho}_r}{8 k^2 [3 F_2^2 - 8 m_2^2 (3 \bar{\rho}_d + 4 \bar{\rho}_r)]} > 0, \quad (3.32)$$

and in this case the matter no-ghost conditions get non-trivially modified. In particular, we find $0 < m_2^2 < F_2^2 / [8(\bar{\rho}_d + 4\bar{\rho}_r/3)]$. Such condition prevents m_2^2 to be arbitrarily large ensuring the stability of the theory. One might wonder about the role of spatial gradients of the lapse in the stability of matter, since in the action (3.20) there is no direct coupling between matter and gravity. However, the spatial gradient of the lapse turns out to be proportional to δ_d^2 , δ_r^2 and ζ^2 through eqs. (3.22), then it is directly involved in the above ghost conditions. In this sense there is a ‘‘coupling’’ between gravity and matter fields.

4. $m_2^2 = 0$ and $F_4 \neq 0$. In this case we have

$$\mathcal{G}_1 = \frac{a^3 (F_1 - 3 F_4) (F_1 F_3 + 3 F_2^2)}{2 (F_2^2 + F_4 F_3)} > 0, \quad (3.33)$$

$$\mathcal{G}_l = \frac{a^5 \bar{\rho}_l^2}{2 k^2 (\bar{\rho}_l + p_l)} > 0. \quad (3.34)$$

5. $F_1 = 0$. In this case the no-ghost conditions can be written as:

$$\mathcal{G}_1 = -\frac{9 F_2^2 a^5}{16 m_2^2 k^2} > 0, \quad (3.35)$$

$$\mathcal{G}_l = \frac{a^5 \bar{\rho}_l^2}{2 k^2 (\bar{\rho}_l + p_l)} > 0, \quad (3.36)$$

so that $m_2^2 < 0$.

3 Stability conditions in the presence of matter

6. Cases: $F_1 = 3F_4$, or $F_1 = 0 = F_2$, or $m_2^2 = 0 = F_1F_3 + 3F_2^2$. In this cases the determinant of the kinetic matrix identically vanishes. This behaviour, in general, leads to strong coupling, so that this class of theories cannot be considered as a valid EFT.

A final remark on the first two cases, which are the most noticeable since they are strictly related to well known models: the presence of matter fluids does not affect the form of the ghost conditions, indeed, we recover the same results as in the previous Chapter where no matter fluids were included, once the high- k limit has been taken. However, let us note that the parameters space identified by these conditions can change because of the evolution of the scale factor, which in turns is the solution of different Friedmann equations. Moreover, no-ghost conditions have been previously obtained in presence of matter fields described by a $P(\mathcal{X})$ action as in refs. [37, 111, 123, 124] (and references therein). Such results are obtained for the variable v_m and they can be safely applied for all matter fluids but not for dust. Indeed, in the specific case of pressureless fluids ($w \rightarrow 0$) the ghost condition turns out to be ill defined. This can be explained by the fact that the no-ghost conditions need to be derived at the level of the action, which diverges in this limit. From a physical point of view this is related to a "bad" choice of physical variable which has to describe the matter d.o.f. as we discussed in section 3.3. However, in some cases they can be extended to non relativistic matter species as for eg. in ref. [111], where the authors use for the barotropic coefficient of these species the case $w = 0^+$ which implies a small yet non-negligible pressure and speed of propagation. In conclusion, by using appropriate precautions in some cases present in literature one can find some of the above results, mostly related to case 2. In this sense our results are more general and robust.

3.4.2 The speeds of propagation

We will now proceed with the study of the speeds of propagation associated to the scalar d.o.f. in action (3.23). As usual, their positivity guarantees the avoidance of any potential gradient instabilities at high- k . Hereafter, we will consider the action purely in the high- k limit. This is a necessary step in order to obtain the physical speeds of propagation. Indeed, if one does not assume the high- k limit the resulting "speeds of propagation" would be complicated and non-local expressions due to the complex dependence on the momentum of the action (3.23) and the interaction between the three fields. Of course, in order to study the gradient instability in full generality one needs to work out such expressions. However, let us say that in such case the fields do not decouple from each

3.4 Study of Stability conditions

other and it turns out to be very difficult to obtain analytical expressions for the speeds of propagation. Moreover, the regime in which the gradient instability manifests itself faster and thus becomes potentially dangerous within the lifetime of the universe is in the high- k limit, thus justifying our restriction to such a regime.

In order to achieve this, it is necessary to diagonalise the kinetic matrix, therefore we will proceed with the following field redefinition:

$$\begin{aligned}\zeta &= \Psi_1, & \delta_d &= \Psi_2 k - \frac{A_{12}A_{33} - A_{13}A_{23}}{A_{22}A_{33} - A_{23}^2} \Psi_1, \\ \delta_r &= k\Psi_3 + \frac{A_{12}A_{23} - A_{13}A_{22}}{A_{22}A_{33} - A_{23}^2} \Psi_1 - \frac{A_{23}}{A_{33}} \Psi_2 k.\end{aligned}\quad (3.37)$$

The k dependence of the transformation is a convenient choice in order to obtain, in the high- k limit, a scale invariant kinetic matrix and the new kinetic matrix, $\mathcal{L} \ni a^3 K_{ij} \dot{\Psi}_i \dot{\Psi}_j$, is now diagonal without approximations. Finally, we get a Lagrangian of the form:

$$\begin{aligned}\mathcal{L}^{(2)} &= K_{11} \dot{\Psi}_1^2 + K_{22} \dot{\Psi}_2^2 + K_{33} \dot{\Psi}_3^2 + Q_{12} (\dot{\Psi}_1 \Psi_2 - \dot{\Psi}_2 \Psi_1) \\ &+ Q_{13} (\dot{\Psi}_1 \Psi_3 - \dot{\Psi}_3 \Psi_1) + Q_{23} (\dot{\Psi}_2 \Psi_3 - \dot{\Psi}_3 \Psi_2) - \mathcal{M}_{ij} \Psi_i \Psi_j,\end{aligned}\quad (3.38)$$

where the kinetic matrix coefficients are:

$$\begin{aligned}K_{11} &= \frac{A_{33}A_{12}^2 - 2A_{13}A_{23}A_{12} + A_{13}^2A_{22} + A_{11}(A_{23}^2 - A_{22}A_{33})}{A_{23}^2 - A_{22}A_{33}}, \\ K_{22} &= k^2 \left(A_{22} - \frac{A_{23}^2}{A_{33}} \right), \\ K_{33} &= k^2 A_{33} \quad K_{ij} = 0 \quad \text{with } i \neq j,\end{aligned}\quad (3.39)$$

and the Q_{ij} and \mathcal{M}_{ij} matrix coefficients will be specified in the following case by case.

Due to the different scaling with k of the operators in action (3.23), it is necessary to analyse the sub-cases identified before separately. As it will become clear every sub-case exhibits a different behaviour, as expected.

1. General case ($m_2^2 \neq 0$ and $F_4 \neq 0$). The kinetic matrix elements at high- k read

$$\begin{aligned}K_{11} &= \frac{F_1(F_1 - 3F_4)}{2F_4} + \mathcal{O}(k^{-2}), & K_{22} &= \frac{1}{2}a^2\bar{\rho}_d + \mathcal{O}(k^{-2}), \\ K_{33} &= \frac{3}{8}a^2\bar{\rho}_r + \mathcal{O}(k^{-2}),\end{aligned}\quad (3.40)$$

which are scale invariant. In its full generality, the action reduces

3 Stability conditions in the presence of matter

in such a limit to a system of three decoupled fields:

$$\begin{aligned}
S^{(2)} &= \frac{1}{(2\pi)^3} \int dk^3 dt \frac{a^3}{8} \left\{ 4a^2 \bar{\rho}_d \dot{\Psi}_2^2 + 3a^2 \bar{\rho}_r \dot{\Psi}_3^2 + \frac{4F_1 (F_1 - 3F_4)}{F_4} \dot{\Psi}_1^2 \right. \\
&- \frac{k^2}{a^2} \left[\frac{2 \left(-4m_0^2 (\Omega + 1) (m_2^2 - \hat{M}^2) + 4\hat{M}^4 + m_0^4 (\Omega + 1)^2 \right)}{m_2^2} \Psi_1^2 \right. \\
&+ \left. \left. a^2 \bar{\rho}_r \Psi_3^2 \right] + \mathcal{O}(k^{-1}) \right\}, \tag{3.41}
\end{aligned}$$

from which it is easy to read off the Q_{ij} and \mathcal{M}_{ij} coefficients. Then, for high- k , the elements Q_{ij} are corrections and the matrix \mathcal{M}_{ij} becomes diagonal. This decoupling is very helpful when obtaining the speeds of propagation from the Euler-Lagrange equations:

$$\begin{aligned}
&\frac{F_1 (F_1 - 3F_4)}{F_4} \ddot{\Psi}_1 + \frac{k^2}{a^2} \left(\frac{-4m_0^2 (\Omega + 1) (m_2^2 - \hat{M}^2) + 4\hat{M}^4 + m_0^4 (\Omega + 1)^2}{2m_2^2} \right) \Psi_1 \\
&+ \left(\frac{F_1^2 (3F_4 H - \dot{F}_4) + F_1 F_4 (2\dot{F}_1 - 9F_4 H)}{F_4^2} - 3\dot{F}_1 \right) \dot{\Psi}_1 \approx 0, \\
&\ddot{\Psi}_2 + 2H \dot{\Psi}_2 \approx 0, \\
&3\ddot{\Psi}_3 + 3H \dot{\Psi}_3 + \frac{k^2}{a^2} \Psi_3 \approx 0. \tag{3.42}
\end{aligned}$$

It is now straightforward to isolate the three speeds of propagation and look at their functional dependence:

$$\begin{aligned}
c_{s,g}^2 &= \frac{F_4 \left(-4m_0^2 (\Omega + 1) (m_2^2 - \hat{M}^2) + 4\hat{M}^4 + m_0^4 (\Omega + 1)^2 \right)}{2F_1 m_2^2 (F_1 - 3F_4)}, \\
c_{s,d}^2 &= 0, \quad c_{s,r}^2 = \frac{1}{3}, \tag{3.43}
\end{aligned}$$

where we have used the suffix "g" to indicate the speed of propagation associated to the d.o.f. of the gravity sector. It is clear that when we consider all the operators active, including the higher order in spatial derivative operators, one gets a completely decoupled system where the fields do not influence each other and evolve separately.

2. Case $F_4 = 0 = m_2^2$. After applying the fields re-definitions (3.37),

3.4 Study of Stability conditions

we get in the large k -limit the following action:

$$\begin{aligned}
S^{(2)} &= \frac{1}{(2\pi)^3} \int dk^3 dt a^3 \left\{ \frac{1}{2} F_1 \left(\frac{F_1 F_3}{F_2^2} + 3 \right) \dot{\Psi}_1^2 + \frac{1}{2} a^2 \bar{\rho}_d \dot{\Psi}_2^2 \right. \\
&+ \frac{3}{8} a^2 \bar{\rho}_r \dot{\Psi}_3^2 - k^2 \frac{\bar{\rho}_r}{8} \Psi_3^2 + k \frac{1}{2 F_2} \left(4 \hat{M}^2 + 2 m_0^2 (\Omega + 1) - F_1 \right) \\
&\times \left[\bar{\rho}_d \left(\Psi_2 \dot{\Psi}_1 - \Psi_1 \dot{\Psi}_2 \right) + \bar{\rho}_r \left(\Psi_3 \dot{\Psi}_1 - \Psi_1 \dot{\Psi}_3 \right) \right] \\
&+ \frac{k^2}{3 a^2 F_2^2} \left[-3 F_1 F_2 H \left(2 \hat{M}^2 + m_0^2 (\Omega + 1) \right) + (6 \bar{\rho}_d + 8 \bar{\rho}_r) \right. \\
&\times \left. \left(2 \hat{M}^2 + m_0^2 (\Omega + 1) \right)^2 + 3 m_0^2 \left[F_1 (\Omega + 1) \dot{F}_2 \right. \right. \\
&- \left. \left. F_2 \left(F_1 \dot{\Omega} + (\Omega + 1) \dot{F}_1 \right) + F_2^2 (\Omega + 1) \right] - 6 \left(F_2 \left(\dot{F}_1 \hat{M}^2 \right. \right. \right. \\
&+ \left. \left. \left. 2 F_1 \hat{M} \dot{M} \right) - F_1 \dot{F}_2 \hat{M}^2 \right) \right] \Psi_1^2 \left. \right\} + \mathcal{O}(k^{-2}).
\end{aligned} \tag{3.44}$$

As it is clear, the resulting action in the high- k limit exhibits some substantial deviations from the previous case. The complication arises due to the fact that now the fields are coupled in antisymmetric configurations. This will force us to change approach when obtaining the speeds of propagation. Namely, we will choose firstly to Fourier transform the time component in the Lagrangian by using $(\partial_t \rightarrow -i\omega)$ and then proceed to obtain the dispersion relations. This will yield the following:

$$\mathcal{L}^{(2)} \sim \vec{\Psi}^T \begin{pmatrix} \frac{1}{2} F_1 \left(\frac{F_3 F_1}{F_2^2} + 3 \right) \omega^2 - \frac{k^2}{a^2} \mathcal{G}_{11} & -i\omega k \mathcal{B}_{12} & -i\omega k \mathcal{B}_{13} \\ i\omega k \mathcal{B}_{12} & \frac{1}{2} a^2 \bar{\rho}_d \omega^2 & 0 \\ i\omega k \mathcal{B}_{13} & 0 & \frac{3}{8} a^2 \bar{\rho}_r \omega^2 - k^2 \frac{\bar{\rho}_r}{8} \end{pmatrix} \cdot \vec{\Psi}, \tag{3.45}$$

where \mathcal{G}_{11} and \mathcal{B}_{ij} can be read off from the action and we have defined the field vector $\vec{\Psi}$. Now, setting the determinant of the above matrix to zero and considering that $\omega^2 = \frac{k^2}{a^2} c_s^2$ in the high- k limit, we obtain the following results:

$$\begin{aligned}
c_{s,d}^2 &= 0, \\
(3c_s^2 - 1) \bar{\rho}_r \left[\bar{\rho}_d (c_s^2 (F_3 F_1^2 + 3 F_2^2 F_1) - 2 a^2 F_2^2 \mathcal{G}_{11}) - 4 \mathcal{B}_{12}^2 F_2^2 \right] \\
&- 16 c_s^2 \mathcal{B}_{13}^2 F_2^2 \bar{\rho}_d = 0
\end{aligned} \tag{3.46}$$

with $F_2 \neq 0$ and where c_s^2 is the double solution of the dispersion relation obtained after observing that the dust speed of propagation,

3 Stability conditions in the presence of matter

$c_{s,d}^2$ is zero. It is clear that, while the speed of propagation of the dust component remains unaffected by the presence of radiation and gravity, the last dispersion relation manifests the clear interaction between radiation and gravity, which modifies their speeds of propagation. Hence, this result shows us clearly that the interaction with matter can affect the gravity sector in a very deep way.

The only case in which the gravity sector and the radiation one completely decouple is when the following condition applies:

$$4\hat{M}^2 + 2m_0^2(\Omega + 1) - F_1 = 0. \quad (3.47)$$

In this case from (3.46) the standard speed of propagation for the radiation is recovered and the speed of gravity is

$$c_{s,g}^2 = \frac{2F_2^2 \mathcal{G}_{11}}{F_3 F_1^2 + 3F_2^2 F_1}. \quad (3.48)$$

Let us notice that the condition (3.47) is trivially satisfied for the Horndeski class of models. In Eq. (3.48), \mathcal{G}_{11} depends on the background densities of dust and radiation, then one can use the background equations (3.25) to eliminate the dependence from the densities of the matter fluids, thus obtaining

$$\begin{aligned} c_{s,g}^2 &= \frac{2}{F_1 (3F_2^2 + F_1 F_3)} \times \left(2cF_1^2 + 2m_0^2 F_1^2 \dot{H}(\Omega + 1) \right. \\ &+ F_1^2 H \left(F_2 - m_0^2 \dot{\Omega} \right) + m_0^2 F_1^2 \ddot{\Omega} - 2m_0^2 F_2^2 (\Omega + 1) - F_1^2 \dot{F}_2 \\ &\left. + 2F_2 F_1 \dot{F}_1 \right). \end{aligned} \quad (3.49)$$

Even though the radiation and the dust sector appear unaltered there is some interplay between gravity and the matter sector. Although the above expression for the speed of propagation of the gravity mode holds both in the vacuum and matter case, the parameters space defined through Eq. (3.49) changes drastically in the two cases. Indeed, firstly one has to consider a different evolution for the scale factor, $a(t)$ accordingly to the corresponding Friedmann equations, secondly in the vacuum case Eq. (3.49) simplifies because a combination of terms turns to be zero due to the Friedmann equations. Instead, such combination of terms when matter is included gives a non zero contribution.

The same result for this sub-case has been obtained in ref. [37, 124], starting from a $P(\mathcal{X})$ action for the matter sector and the v_m variable. It is important to note that, in contrast to the no-ghost

3.4 Study of Stability conditions

conditions, the results also agree for the case of dust. This can be explained by the fact that the speed of propagation can be obtained at the level of the equations of motion, hence avoiding the issues plaguing the action, described in the previous sections.

3. Case $F_4 = 0$ and $m_2^2 \neq 0$. The action at high- k reads

$$\begin{aligned}
S^{(2)} &= \frac{1}{(2\pi)^3} \int dk^3 dt a^3 \left\{ \frac{4k^2 F_1^2 m_2^2}{a^2 F_2^2} \dot{\Psi}_1^2 + \frac{a^2 \bar{\rho}_d F_2^2}{2F_2^2 - 16\bar{\rho}_d m_2^2} \dot{\Psi}_2^2 \right. \\
&+ \frac{9a^2 \bar{\rho}_r (F_2^2 - 8\bar{\rho}_d m_2^2)}{8(-24\bar{\rho}_d m_2^2 + 3F_2^2 - 32m_2^2 \bar{\rho}_r)} \dot{\Psi}_3^2 - \frac{k^2 \bar{\rho}_r}{8} \Psi_3^2 \\
&- \frac{128k^2 \bar{\rho}_d^2 m_2^4 \bar{\rho}_r}{9(F_2^2 - 8\bar{\rho}_d m_2^2)^2} \Psi_2^2 - \frac{128k^4 F_1^2 m_2^4 \bar{\rho}_r}{9a^4 F_2^4} \Psi_1^2 - \frac{8k^3 F_1 m_2^2 \bar{\rho}_r}{3a^2 F_2^2} \Psi_1 \Psi_3 \\
&+ \frac{256k^3 a \bar{\rho}_d F_1 m_2^4 \bar{\rho}_r}{9a^3 F_2^4 - 72\bar{\rho}_d F_2^2 m_2^2} \Psi_1 \Psi_2 + k(16F_1 F_2 m_2^2 H \\
&+ (4F_2 (F_2 \hat{M}^2 - 2m_2^2 \dot{F}_1) + 2m_0^2 F_2^2 (\Omega + 1) \\
&- F_1 (16F_2 m_2 \dot{m}_2 - 16m_2^2 \dot{F}_2 + F_2^2)) \\
&\times \left[\frac{\bar{\rho}_d}{2(F_2^2 - 8\bar{\rho}_d F_2 m_2^2)} (\Psi_2 \dot{\Psi}_1 - \Psi_1 \dot{\Psi}_2) \right. \\
&+ \frac{3\bar{\rho}_r}{2F_2 (-24\bar{\rho}_d m_2^2 + 3F_2^2 - 32m_2^2 \bar{\rho}_r)} (\Psi_3 \dot{\Psi}_1 - \Psi_1 \dot{\Psi}_3) \left. \right] \\
&+ \left. \frac{8k^2 \bar{\rho}_d m_2^2 \bar{\rho}_r}{3F_2^2 - 24\bar{\rho}_d m_2^2} \Psi_2 \Psi_3 \right\} + O(k^{-2}). \tag{3.50}
\end{aligned}$$

We find that the solutions of the discriminant equation

$$\det \left(\frac{c_s^2 k^2}{a^2} K_{ij} - \mathcal{M}_{ij} \right) = \frac{a^5 c_s^4 \bar{\rho}_m (c_s^2 \rho_{r,n} - n_r \rho_{r,nn}) \bar{\rho}_r^2 m_2^2 F_1^2 k^6}{(-8m_2^2 n_r \rho_{r,n} - 8m_2^2 \bar{\rho}_m + F_2^2) n_r \rho_{r,n}^2}, \tag{3.51}$$

reduce to

$$c_{s,g}^2 = 0, \quad c_{s,d}^2 = 0, \quad c_{s,r}^2 = \frac{1}{3}. \tag{3.52}$$

The results for this case can be found in the limit $F_4 \rightarrow 0$ for the general case discussed above.

4. Case $F_4 \neq 0$ and $m_2^2 = 0$. The action for this sub-case at high- k

3 Stability conditions in the presence of matter

reads

$$\begin{aligned}
S^{(2)} &= \frac{1}{(2\pi)^3} \int dk^3 dt a^3 \left\{ \frac{(3F_2^2 + F_1 F_3)(F_1 - 3F_4)}{2(F_2^2 + F_3 F_4)} \dot{\Psi}_1^2 \right. \\
&+ \frac{1}{2} a^2 \bar{\rho}_d \dot{\Psi}_2^2 + \frac{3}{8} a^2 \bar{\rho}_r \dot{\Psi}_3^2 - k^2 \frac{\bar{\rho}_r (F_2^2 + F_3 F_4 + 4F_4 \bar{\rho}_r)}{8(F_2^2 + F_3 F_4)} \Psi_3^2 \\
&- \frac{k^2 \bar{\rho}_d^2 F_4}{2(F_2^2 + F_3 F_4)} \Psi_2^2 - k^4 \frac{2F_4 (2\hat{M}^2 + m_0^2(\Omega + 1))^2}{a^4 (F_2^2 + F_3 F_4)} \Psi_1^2 \\
&+ k^3 \frac{2F_4 (2\hat{M}^2 + m_0^2(\Omega + 1))}{a^2 (F_2^2 + F_3 F_4)} (\bar{\rho}_r \Psi_3 \Psi_1 + \bar{\rho}_d \Psi_1 \Psi_2) \\
&+ k \frac{F_2 (-F_1 + 3F_4 + 4\hat{M}^2 + 2m_0^2(\Omega + 1))}{2(F_2^2 + F_3 F_4)} \left[\bar{\rho}_d (\Psi_2 \dot{\Psi}_1 - \Psi_1 \dot{\Psi}_2) \right. \\
&+ \left. \bar{\rho}_r (\Psi_3 \dot{\Psi}_1 - \Psi_1 \dot{\Psi}_3) \right] - \frac{k^2 \bar{\rho}_d F_4 \bar{\rho}_r}{(F_2^2 + F_3 F_4)} \Psi_2 \Psi_3 \left. \right\} + \mathcal{O}(k^{-2}), \tag{3.53}
\end{aligned}$$

where the kinetic terms K_{11}, K_{22}, K_{33} are of order $\mathcal{O}(k^0)$ for high values of k and the elements Q_{12} and Q_{13} are of order k and cannot be neglected. Furthermore, the leading component of \mathcal{M}_{11} is of order k^4 . Therefore now we need to consider the discriminant equation as

$$\text{Det}(\omega^2 K_{ij} - i\omega Q_{ij} - \mathcal{M}_{ij}) = 0, \tag{3.54}$$

this equation can be recast as

$$\omega^6 + \left(\mathcal{A} \frac{k^4}{a^4} + \mathcal{O}(k^2) \right) \omega^4 + \left(\mathcal{B} \frac{k^6}{a^6} + \mathcal{O}(k^4) \right) \omega^2 = 0, \tag{3.55}$$

with

$$\mathcal{A} = 4 \frac{((\Omega + 1) m_0^2 + 2\hat{M}^2)^2 F_4}{(3F_4 - F_1)(F_1 F_3 + 3F_2^2)}, \quad \mathcal{B} = -\frac{1}{3} \mathcal{A}. \tag{3.56}$$

For high- k , we find the following solutions:

- One solution can be found by assuming $\omega^2 = W k^4 / a^4$. In this case we find

$$(W + \mathcal{A}) W^2 \frac{k^{12}}{a^{12}} + \mathcal{O}(k^{10}) = 0, \tag{3.57}$$

3.4 Study of Stability conditions

which is verified by $W = -\mathcal{A}$, so that

$$\omega^2 = -\mathcal{A} \frac{k^4}{a^4}, \quad c_{s,g}^2 = -4\mathcal{A} \frac{k^2}{a^2}, \quad (3.58)$$

or

$$c_{s,g}^2 = \frac{16F_4 \left((\Omega + 1) m_0^2 + 2\hat{M}^2 \right)^2}{(F_1 F_3 + 3F_2^2)(F_1 - 3F_4)} \frac{k^2}{a^2}, \quad (3.59)$$

which tends to large values.

- The other two solutions of Eq. (3.55) can be found by assuming $\omega^2 = Wk^2/a^2$, so that

$$(\mathcal{A}W^2 + \mathcal{B}W) \frac{k^8}{a^8} + \mathcal{O}(k^6) = 0, \quad (3.60)$$

which implies the following standard results

$$c_{s,d}^2 = 0, \quad c_{s,r}^2 = \frac{1}{3}. \quad (3.61)$$

5. Case $F_1 = 0$. The action reads:

$$\begin{aligned} S^{(2)} &= \frac{1}{(2\pi)^3} \int dk^3 dt a^3 \left[-\frac{9a^2 F_2^2}{16m_2^2} \dot{\Psi}_1^2 + \frac{\bar{\rho}_d}{2} a^2 \dot{\Psi}_2^2 + \frac{3}{8} \bar{\rho}_r a^2 \dot{\Psi}_3^2 \right. \\ &\quad - \frac{k^2 \bar{\rho}_r}{8} \psi_3^2 + k^2 \frac{\bar{\rho}_d \left(2\hat{M}^2 + m_0^2 (\Omega + 1) \right)}{4m_2^2} \psi_1 \Psi_2 \\ &\quad + k^2 \frac{\bar{\rho}_r \left(4\hat{M}^2 + 2m_0^2 (\Omega + 1) + 8m_2^2 \right)}{8m_2^2} \Psi_1 \Psi_3 \\ &\quad \left. - k^4 \frac{4\hat{M}^4 - 4m_0^2 (\Omega + 1) \left(m_2^2 - \hat{M}^2 \right) + m_0^4 (\Omega + 1)^2}{4a^2 m_2^2} \Psi_1^2 \right] \\ &\quad + \mathcal{O}(k^{-2}). \end{aligned} \quad (3.62)$$

In this case, the matrix Q_{ij} can be neglected, but the \mathcal{M}_{11} coefficient has a term in k^4 , therefore we have the discriminant equation

$$\text{Det}(\omega^2 K_{ij} - \mathcal{M}_{ij}) = 0, \quad (3.63)$$

which leads to

$$\omega^6 + \left(\mathcal{A} \frac{k^4}{a^4} + \mathcal{O}(k^2) \right) \omega^4 + \left(-\frac{1}{3} \mathcal{A} \frac{k^6}{a^6} + \mathcal{O}(k^4) \right) \omega^2 + \mathcal{O}(k^6) = 0. \quad (3.64)$$

3 Stability conditions in the presence of matter

with

$$\mathcal{A} = \frac{4 \left(-4m_0^2(\Omega + 1) \left(m_2^2 - \hat{M}^2 \right) + 4\hat{M}^4 + m_0^4(\Omega + 1)^2 \right)}{9F_2^2}. \quad (3.65)$$

Once more we have a solution

$$\omega^2 = -\mathcal{A} \frac{k^4}{a^4}, \quad (3.66)$$

which leads to

$$c_{s,g}^2 = \frac{16}{9} \frac{4 \left(m_2^2 - \hat{M}^2 \right) \left(1 + \Omega \right) m_0^2 - \left(1 + \Omega \right)^2 m_0^4 - 4 \hat{M}^4}{F_2^2} \frac{k^2}{a^2}, \quad (3.67)$$

whereas the other two solutions are found to be

$$c_{s,d}^2 = 0, \quad c_{s,r}^2 = \frac{1}{3}. \quad (3.68)$$

In summary, in this section we have derived the speeds of propagation for the three dynamical fields describing our system. In general for all the sub-cases analysed we have found that in the high- k regime the three d.o.f. decouple and the resulting speeds of propagation are unaltered with respect to the vacuum case. This can be easily verified by considering the high- k limit of the results in Chapter 2. Only one case stands aside, the beyond Horndeski case. In this case the dust field completely decouples from the other fields, while the radiation and gravity fields are coupled and their speeds result to be modified. We also recall that even in the cases the expressions for the speeds of propagation do not differ from the respective cases in vacuum, the parameters space may change accordingly to a different evolution of the scale factor, which in turns is the solution of different background equations. In conclusion, for all the cases analysed we demand a positive speed of propagation in order to guarantee the viability of the underlying theory of gravity.

3.4.3 Tachyonic and Jeans instabilities

The final aspect of our work which tends to be the one least studied in the literature in the context of MG theories and especially in the EFToDE/MG framework, is the study of the canonical mass of the fields and consequently the boundedness of the Hamiltonian at low momenta. These results are related to the usual tachyonic and Jeans instabilities, the latter being characteristic of the fluids components.

3.4 Study of Stability conditions

In this section we will restrict the analysis to the EFToDE/MG action in the presence of only one matter fluid. We choose the dust over radiation because we know that the dust component clusters and hence for our purpose it might show an interesting behaviour related to the Jeans instability. Thus, the results presented here will be applicable during the dust-dominated era and onwards when the dynamics of the two d.o.f. starts to play a role. A second fluid can be straightforwardly added, but it makes the procedure substantially more difficult.

One can obtain the action for the EFToDE/MG with a dust component by setting $\delta_r = 0$ in the action (3.20) and $\bar{\rho}_r = 0$ in the remaining functions. Now, let us assume the no-ghost conditions hold, and proceed to rewrite the action in its canonical form. The first step is to diagonalise the (2x2) kinetic matrix as in the previous section by making the following field redefinitions

$$\begin{aligned}\zeta &= \Psi_1, \\ \delta_d &= k\Psi_2 - \frac{A_{12}\Psi_1}{A_{22}},\end{aligned}\tag{3.69}$$

with the following diagonal terms:

$$\bar{K}_{11} = A_{11} - \frac{A_{12}^2}{A_{22}}, \quad \bar{K}_{22} = k^2 A_{22},\tag{3.70}$$

where A_{ij} are the ones defined in Appendix 3.6 after setting $\bar{\rho}_r = 0$. Next, the canonical form is obtained by normalising the fields accordingly to:

$$\begin{aligned}\Psi_1 &= \frac{1}{\sqrt{2\bar{K}_{11}}} \bar{\Psi}_1, \\ \Psi_2 &= \frac{1}{\sqrt{2\bar{K}_{22}}} \bar{\Psi}_2.\end{aligned}\tag{3.71}$$

After grouping the different terms and performing a number of integrations by parts, we obtain the Lagrangian as:

$$\mathcal{L}^{(2)} = \frac{a^3}{2} \left[\dot{\bar{\Psi}}_1^2 + \dot{\bar{\Psi}}_2^2 + \bar{B}(t, k) (\dot{\bar{\Psi}}_1 \bar{\Psi}_2 - \dot{\bar{\Psi}}_2 \bar{\Psi}_1) - \bar{C}_{ij}(t, k) \bar{\Psi}_i \bar{\Psi}_j \right],\tag{3.72}$$

where we refer the reader to the Appendix 3.6 for the functional forms of the \bar{B}, \bar{C}_{ij} coefficients.

In order to obtain the mass eigenvalues we need to proceed with the diagonalization of the mass matrix C_{ij} while keeping the canonical form of the action. For this purpose we consider a field rotation via an orthogonal matrix, in the following way:

$$\begin{aligned}\bar{\Psi}_1 &= \cos \alpha \Phi_1 + \sin \alpha \Phi_2, \\ \bar{\Psi}_2 &= -\sin \alpha \Phi_1 + \cos \alpha \Phi_2.\end{aligned}\tag{3.73}$$

3 Stability conditions in the presence of matter

Now, it is possible to choose α in a very specific way in order to diagonalize the mass matrix. This leads to the following relation:

$$\tan(2\alpha) = \beta \equiv -\frac{2\bar{C}_{12}}{\bar{C}_{11} - \bar{C}_{22}}, \quad (3.74)$$

accompanied by:

$$\frac{d[\tan(2\alpha)]}{dt} = 2[1 + \tan^2(2\alpha)]\dot{\alpha} \quad \Longleftrightarrow \quad \dot{\alpha} = \frac{\dot{\beta}}{2(1 + \beta^2)}. \quad (3.75)$$

Then, the Lagrangian becomes

$$\mathcal{L}^{(2)} = \frac{a^3}{2} \left[\dot{\Phi}_1^2 + \dot{\Phi}_2^2 + B(t, k) (\dot{\Phi}_1 \Phi_2 - \dot{\Phi}_2 \Phi_1) - \mu_1(t, k) \Phi_1^2 - \mu_2(t, k) \Phi_2^2 \right], \quad (3.76)$$

with the following definitions:

$$\begin{aligned} B &= \bar{B} + 2\dot{\alpha}, \\ \mu_1 &= -\dot{\alpha}^2 - \bar{B}\dot{\alpha} + \frac{(\bar{C}_{11} - \bar{C}_{22})^2 + 4C_{12}^2}{\bar{C}_{11} - \bar{C}_{22}} \cos^2 \alpha + \frac{\bar{C}_{11}\bar{C}_{22} - 2\bar{C}_{12}^2 - \bar{C}_{22}^2}{\bar{C}_{11} - \bar{C}_{22}}, \\ \mu_2 &= -\dot{\alpha}^2 - \bar{B}\dot{\alpha} - \frac{(\bar{C}_{11} - \bar{C}_{22})^2 + 4C_{12}^2}{\bar{C}_{11} - \bar{C}_{22}} \cos^2 \alpha + \frac{\bar{C}_{11}^2 - \bar{C}_{11}\bar{C}_{22} + 2\bar{C}_{12}^2}{\bar{C}_{11} - \bar{C}_{22}}. \end{aligned} \quad (3.77)$$

It is straightforward to obtain the energy function (which is equal in value to the Hamiltonian, see e.g. [127] for details) which reduces to a formally simple form (see Appendix 3.7), namely

$$H(\Phi_i, \dot{\Phi}_i) = \frac{a^3}{2} \left[\dot{\Phi}_1^2 + \dot{\Phi}_2^2 + \mu_1(t, k) \Phi_1^2 + \mu_2(t, k) \Phi_2^2 \right], \quad (3.78)$$

so that the Hamiltonian will be unbounded from below if the eigenvalues satisfy $\mu_1 < 0$ or $\mu_2 < 0$, for example on the line ($\dot{\Phi}_i = 0$). Naively one could then proceed and constrain the mass eigenvalues to be non-negative. From a physical point of view this condition must be considered too stringent as there is a very well known phenomenon related to a negative mass, the Jeans Instability. As this corresponds to a negative mass but with a slow enough evolution rate in order to be countered by gravity we can reformulate the condition avoiding a catastrophic tachyon instability. For cosmological purposes, in order for a theory to be viable we will proceed to demand the eigenvalues, if negative, to satisfy the condition $|\mu_i(t, 0)| \lesssim H^2$, so that the evolution rate of the instability will not affect the whole stability of the system for time-intervals much shorter than the Hubble time (see also ref. [126]). Finally, one would

expect that the μ_2 eigenvalue can become negative as the dust sector must exhibit a Jeans instability in order for structure to form in our universe.

- **Minimally coupled quintessence model in presence of a dust fluid**

We will now proceed to exemplify the previous, rather abstract, approach by studying a specific model in the presence of dust: minimally coupled quintessence, which has the following action [71]

$$S_\phi = \int d^4x \sqrt{-g} \left[\frac{m_0^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] + S_m, \quad (3.79)$$

where ϕ is the scalar field and V the corresponding potential. The above can be mapped in the EFTtoDE/MG formalism by making the following correspondence as was done in Chapter 2 and the following refs. [14, 15, 75]

$$c = \frac{1}{2} \dot{\phi}_0^2, \quad \Lambda = \frac{1}{2} \dot{\phi}_0^2 - V(\phi_0), \quad \left\{ \Omega, \hat{M}^2, \bar{M}_2^2, \bar{M}_3^2, M_1^3, M_2^4 \right\} = 0, \quad (3.80)$$

where $\phi_0(t)$ is the background value of the scalar field.

Let us now consider that the minimally coupled quintessence model can be also parametrized by assuming that the modification to the gravity sector can be recast as a DE perfect fluid by introducing the following:

$$w_{\text{DE}}(a) \equiv \frac{P_{\text{DE}}}{\bar{\rho}_{\text{DE}}} = \frac{\dot{\phi}^2/2 - V(\phi)}{\dot{\phi}^2/2 + V(\phi)}, \quad (3.81)$$

and assuming that the DE density has the standard perfect fluid form

$$\bar{\rho}_{\text{DE}} = 3m_0^2 H_0^2 \Omega_{\text{DE}}^0 \exp \left[-3 \int_1^a \frac{(1 + w_{\text{DE}}(a))}{a} da \right], \quad (3.82)$$

with $H_0, \Omega_{\text{DE}}^0$ be the present day values of the Hubble and density parameter respectively. Then, the Friedmann equation simply reads:

$$3m_0^2 H^2 = \bar{\rho}_d + \bar{\rho}_{\text{DE}}. \quad (3.83)$$

and the EFT functions can be written accordingly as

$$c = \frac{1}{2} \bar{\rho}_{\text{DE}} (1 + w_{\text{DE}}), \quad \Lambda = w_{\text{DE}} \bar{\rho}_{\text{DE}}. \quad (3.84)$$

This choice for the parametrization makes the whole treatment of the minimally coupled quintessence case more handy. Indeed, this will allow

3 Stability conditions in the presence of matter




Figure 3.1: The figures show the behaviours of the mass eigenvalues μ_1/H^2 (blue dashed line), μ_2/H^2 (orange dot dashed line) for minimally coupled quintessence on a CPL background. *Left panel:* stable tachyonic configuration with $w_0 = -0.9$ and $w_a = 0.009$. *Right panel:* unstable tachyonic configuration with $w_0 = -2.9$ and $w_a = -2$. For this figure the cosmological parameters are chosen to be: $\Omega_{\text{DE}}^0 = 0.69$, $\Omega_d^0 = 0.31$, $H_0 = 67.74$ [128]. See section 3.4.3 for the whole discussion.

us to rewrite the mass eigenvalues (3.95), presented in appendix ??, purely in terms of the fluid parameter, i.e. $\mu_i(w_{\text{DE}})$. As a general remark, from the expressions (3.95) it becomes clear that in general the mass eigenvalues tend to be quite complicated. More complicated theories, especially the non minimally coupled ones, will be substantially harder to treat, yet not impossible. Having the explicit results of the mass eigenvalues for the minimally coupled quintessence model, we want now to proceed and gain some intuition regarding their behaviour compared to H^2 .

We will make a specific choice of the DE equation of state, out of the many options, which will help in illustrating different behaviours:

- The CPL parametrization [129, 130]: $w_{\text{DE}}(a) = w_0 + w_a(1 - a)$, where w_0 and w_a are constant and indicate, respectively, the value and the derivative of w_{DE} today;

as illustrative examples, for the values of $\{w_0, w_a\}$ in the DE equation of state we choose two sets, one for which the system is free from tachyonic instability and one where the gravity sector shows an unstable configuration since a tachyonic instability is manifest. The results are illustrated in figure 3.1. In the left panel, for the choice of the parameters $w_0 = -0.9$ and $w_a = 0.009$, we notice that the eigenvalue associated to the gravity sector (i.e. μ_1) is always positive and approximately of the same order as H^2 . On the contrary, the eigenvalue associated with the dust sector (μ_2), here plotted in its absolute value, is negative and $|\mu_2| \ll H^2$. This, of course, has to be expected as it is a manifestation of the well known Jeans instability which allows structure to form. In the right panel, we chose a rather unrealistic set of the parameters in order to show a tachyonic instability, namely $w_0 = -2.9$ and $w_a = -2$. Both the

eigenvalues oscillate and switch the sign very fast at early time, then μ_2 becomes positive around $\log(a) > -2.9$ and μ_1 turns to be firstly negative and finally positive at very late time. In this case since the eigenvalue associated to gravity is negative (for most of the time) and additionally $|\mu_1| \gg H^2$, this implies that the tachyonic instability evolves very fast, resulting in an unstable system. The dust eigenvalue on the other hand is positive during the matter dominated era which means that matter does not cluster.

The above discussion concerns only the tachyonic and Jeans instabilities, thus can not be considered exhaustive. In order to complete the set of stability conditions for minimally coupled quintessence one needs to study the ghost conditions and the speeds of propagation, as presented in the previous sections, which in the case of minimally coupled quintessence simply reduce to $w_{\text{DE}}(a) > -1$.

3.5 Conclusion

In this Chapter we presented a thorough analysis of the viability conditions which guarantee the stability of the scalar d.o.f. in the presence of matter fields. These conditions guarantee the avoidance of ghosts and tachyonic instabilities, supplemented with a positive speed of propagation. The study of the viability of specific gravity theories in vacuum or in the presence of matter fields has already yielded an extensive literature. However, our results are more general and directly applicable to most of the well known models which are of cosmological interest.

For the gravity sector, we employed the general EFT approach for DE/MG, which has the advantage of being a model independent parametrization of gravity theories with one extra scalar d.o.f. while at the same time preserving a direct link with a wide class of theoretical models which can be explicitly mapped into this formalism. In order to describe the standard perfect fluids we chose the Sorkin-Schutz action which has been shown to be well behaved in contrast to other choices made in the past when considering pressureless fluids. In detail, we specialised to the case where the matter fluids are dust (or CDM) and radiation. From these starting blocks we proceeded to derive the action up to second order in scalar perturbations accompanied by the background equations. Finally, we moved to the study of the viability requirements which we will summarise and discuss in the following.

After constructing the Lagrangian for the perturbations one can straightforwardly guarantee the absence of ghosts by imposing the positivity of the kinetic term, or matrix in case more than one field is considered as in the present chapter. In deriving such conditions we have considered

3 Stability conditions in the presence of matter

only the Lagrangian in the high- k regime, following the recent results in ref. [126] where it was shown that only the high energy terms can turn out in catastrophic instabilities. Sub-leading terms can be recast in mass-like terms through appropriate field redefinitions. Because the EFToDE/MG approach encompasses a variety of DE/MG models, which in some cases show different and non trivial k -dependence, it is not possible to obtain one general result applicable to all possible theories. Therefore, we have identified five relevant sub-cases for which we have worked out the corresponding no-ghost conditions. In particular, two of the aforementioned sub-cases correspond to well known theoretical models, i.e. low-energy Hořava gravity and beyond Horndeski, while the remaining three do not correspond to any specific class of theories but can be useful in a model independent study. In general, we found three no-ghost conditions for each of the sub-cases. Out of those, the conditions corresponding to matter fields were trivially satisfied. Finally, we have also identified conditions which lead to strong coupling regimes, thus excluding these theories from an effective description.

The next step was to identify the speeds of propagation of the three d.o.f., in the high- k limit, for the sub-cases mentioned before. In order to avoid gradient instabilities it was then required that they have a positive sign. Depending on the sub-case the results change drastically because the momentum dependence of various terms differs in each sub-case. In general, the gravity speed of propagation does not depend on fluid variables once one consider theories with higher (than second) order spatial derivatives. In particular, for the sub-case to which low-energy Hořava gravity belongs, we find that at high- k the d.o.f. are completely decoupled and the speeds of propagation of dust and radiation components are unaffected by the coupling to gravity, leading to the standard results. On the contrary, the sub-case corresponding to beyond Horndeski shows non trivial modifications as only the dust speed of propagation stays unaltered. Furthermore, when specifying to Horndeski, the dust and radiation reduce to the standard results while the speed of propagation associated to the gravity mode still is modified with respect to the vacuum case. The three remaining sub-cases exhibit unaltered matter speeds and the gravity one is not influenced by the matter sector. A surprising result is the sub-case corresponding to $F_4 = 0, m_2^2 \neq 0$, for which the gravitational sector has a vanishing speed of sound.

In the last part of the Chapter we considered the tachyonic instability. This instability is tightly related with the Hamiltonian being unbounded as was shown in the extensive calculation in Section 3.4.3 . Only for this case we have simplified the approach by assuming only the dominant matter fluid, Cold Dark Matter. From this starting point, we have identified the two eigenvalues (μ_i) of the system which need to be constrained

3.6 Appendix A: Matrix coefficients

in the limit $k \rightarrow 0$ in order to guarantee the Hamiltonian of the system is bounded from below. These conditions then correspond to the ones one needs to apply in order to avoid tachyonic instabilities. A stringent condition is then to demand both the eigenvalues to be positive definite. On the other hand, it is well known that the dust fluid exhibits a Jeans instability, which is necessary in order to allow structures formation. Therefore, it is more realistic to allow them to become negative in sign but with a evolution rate which sufficiently slow in order to avoid the system to become unviable. In other words this translates to demanding that, in case $\mu_i < 0$, we have $|\mu_i| \ll H^2$. Due to the complexity of these results, we have chosen to exemplify our findings by studying the minimally coupled quintessence case. We have parametrized the gravity modification in terms of the equation of state for DE, i.e. $w_{\text{DE}}(a)$ and then through the appropriate mapping, we were able to write $\mu_i(w_{\text{DE}})$. By choosing the CPL parametrization for the DE fluid and two sets of values for the DE parameters we then presented, figure 3.1, we two typical situations, i.e. the case in which the tachyonic instability shows up and the theory becomes pathological and a stable case exhibiting a Jeans instability for the dust sector.

Concluding we would like to stress that, in the case of ghost and gradient instabilities, we presented the results in a model independent way. This implies that it covers the results existing in the literature, typically derived in the context of a specific theory, and generalises them to cover cases left unexplored. We proved this by comparing our results with the ones in the literature. Additionally we considered in depth the tachyonic instability, in the presence of matter, a result completely novel and of severe importance as the usual two conditions are not enough to produce a stable theory. This will be discussed more in depth in Chapter 5 where we will proceed to implement these results in *EFTCAMB* and test them. That will be the final step of this line of work as then they can be employed in depth in studying models of DE/MG.

3.6 Appendix A: Matrix coefficients

For completeness in this Appendix we will explicitly list the matrix coefficients used in sections 3.4, 3.4.2 and 3.4.3. Let us start by considering the action (3.23) introduced in section 3.4:

$$S^{(2)} = \frac{1}{(2\pi)^3} \int d^3k dt a^3 \left(\dot{\bar{\chi}}^t \mathbf{A} \dot{\bar{\chi}} - k^2 \bar{\chi}^t \mathbf{G} \bar{\chi} - \dot{\bar{\chi}}^t \mathbf{B} \bar{\chi} - \bar{\chi}^t \mathbf{M} \dot{\bar{\chi}} \right), \quad (3.85)$$

where the coefficients are

$$A_{11} = \frac{3(F_1 - 3F_4)(k^2 a^2 (-8m_2^2(3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_2^2 + F_1 F_3) - a^4 F_3(3\bar{\rho}_d + 4\bar{\rho}_r) + 8k^4 F_1 m_2^2)}{2k^2 a^2 (-8m_2^2(3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - 2a^4 F_3(3\bar{\rho}_d + 4\bar{\rho}_r) + 48k^4 m_2^2 F_4},$$

3 Stability conditions in the presence of matter

$$\begin{aligned}
A_{12} = A_{21} &= -\frac{3a^2\bar{\rho}_d(F_1-3F_4)(a^2F_3+8k^2m_0^2)}{2k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-2a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+48k^4m_0^2F_4}, \\
A_{13} = A_{31} &= -\frac{3a^2(F_1-3F_4)\bar{\rho}_r(a^2F_3+8k^2m_0^2)}{2k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-2a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+48k^4m_0^2F_4}, \\
A_{22} &= \frac{a^2\bar{\rho}_d(k^2a^2(3F_3F_4+3F_2^2-32m_0^2\bar{\rho}_r)-4a^4F_3\bar{\rho}_r+24k^4m_0^2F_4)}{2k^2(k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4)}, \\
A_{23} = A_{32} &= \frac{3a^4\bar{\rho}_d\bar{\rho}_r(a^2F_3+8k^2m_0^2)}{2k^2(k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4)}, \\
A_{33} &= \frac{9a^2\bar{\rho}_r(k^2a^2(-8m_0^2\bar{\rho}_d+F_3F_4+F_2^2)-a^4F_3\bar{\rho}_d+8k^4m_0^2F_4)}{8k^2(k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4)}, \\
B_{11} &= -\frac{6k^4F_2(F_1-3F_4)(2\hat{M}^2+m_0^2(\Omega+1))}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4}, \\
B_{12} &= \frac{3k^2a^2F_2\bar{\rho}_d(F_1-3F_4)}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4}, \\
B_{13} &= \frac{3k^2a^2F_2(F_1-3F_4)\bar{\rho}_r}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4}, \\
B_{22} &= \frac{3a^4F_2\bar{\rho}_d^2}{k^2a^2(8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)-3F_3F_4-3F_2^2)+a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)-24k^4m_0^2F_4}, \\
B_{21} &= \frac{6k^2a^2F_2\bar{\rho}_d(2\hat{M}^2+m_0^2(\Omega+1))}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4}, \\
B_{23} = B_{32} &= \frac{3a^4F_2\bar{\rho}_d\bar{\rho}_r}{k^2a^2(8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)-3F_3F_4-3F_2^2)+a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)-24k^4m_0^2F_4}, \\
B_{33} &= \frac{3a^4F_2\bar{\rho}_r^2}{k^2a^2(8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)-3F_3F_4-3F_2^2)+a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)-24k^4m_0^2F_4}, \\
B_{31} &= \frac{6k^2a^2F_2\bar{\rho}_r(2\hat{M}^2+m_0^2(\Omega+1))}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4},
\end{aligned} \tag{3.86}$$

$$\begin{aligned}
G_{11} &= \left\{ k^2a^2 \left(m_0^2(\Omega+1) \left(-8(3\bar{\rho}_d+4\bar{\rho}_r) \left(m_0^2 - \hat{M}^2 \right) + 3F_3F_4 \right. \right. \right. \\
&+ \left. \left. 3F_2^2 \right) + 8\hat{M}^4(3\bar{\rho}_d+4\bar{\rho}_r) + 2m_0^4(\Omega+1)^2(3\bar{\rho}_d+4\bar{\rho}_r) \right) \\
&- m_0^2a^4F_3(\Omega+1)(3\bar{\rho}_d+4\bar{\rho}_r) - 6k^4F_4 \left(-4m_0^2(\Omega+1) \left(m_0^2 - \hat{M}^2 \right) \right. \\
&+ \left. \left. 4(\hat{M}^2)^2 + m_0^4(\Omega+1)^2 \right) \right\} / \left\{ a^2 \left(k^2a^2(8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r) \right. \right. \right. \\
&- \left. \left. 3F_3F_4 - 3F_2^2) + a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r) - 24k^4m_0^2F_4 \right) \right\},
\end{aligned} \tag{3.87}$$

$$\begin{aligned}
G_{12} = G_{21} &= -\frac{\bar{\rho}_d(2\hat{M}^2+m_0^2(\Omega+1))(-a^2(3\bar{\rho}_d+4\bar{\rho}_r)+3k^2\bar{M}_2^2+3k^2\bar{M}_3^2)}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4}, \\
G_{13} = G_{31} &= -\frac{\bar{\rho}_r(2\hat{M}^2+m_0^2(\Omega+1))(-a^2(3\bar{\rho}_d+4\bar{\rho}_r)+3k^2\bar{M}_2^2+3k^2\bar{M}_3^2)}{k^2a^2(-8m_0^2(3\bar{\rho}_d+4\bar{\rho}_r)+3F_3F_4+3F_2^2)-a^4F_3(3\bar{\rho}_d+4\bar{\rho}_r)+24k^4m_0^2F_4},
\end{aligned}$$

3.6 Appendix A: Matrix coefficients

$$\begin{aligned}
G_{22} &= \frac{3a^2 F_4 \bar{\rho}_d^2}{2k^2 a^2 (-8m_2^2 (3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - 2a^4 F_3 (3\bar{\rho}_d + 4\bar{\rho}_r) + 48k^4 m_2^2 F_4}, \\
G_{23} = G_{32} &= \frac{3a^2 F_4 \bar{\rho}_d \bar{\rho}_r}{2k^2 a^2 (-8m_2^2 (3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - 2a^4 F_3 (3\bar{\rho}_d + 4\bar{\rho}_r) + 48k^4 m_2^2 F_4}, \\
G_{33} &= \frac{\bar{\rho}_r (a^2 (-4(m_2^2 (6\bar{\rho}_d + 8\bar{\rho}_r) - 3F_4 \bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) + 24k^2 m_2^2 F_4)}{8(k^2 a^2 (-8m_2^2 (3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - a^4 F_3 (3\bar{\rho}_d + 4\bar{\rho}_r) + 24k^4 m_2^2 F_4)}, \\
M_{11} = M_{12} = M_{21} = M_{13} = M_{31} &= 0, \\
M_{22} &= -\frac{a^4 \bar{\rho}_d^2 (3\bar{\rho}_d + 4\bar{\rho}_r)}{2k^2 a^2 (-8m_2^2 (3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - 2a^4 F_3 (3\bar{\rho}_d + 4\bar{\rho}_r) + 48k^4 m_2^2 F_4}, \\
M_{23} = M_{32} &= -\frac{a^4 \bar{\rho}_d \bar{\rho}_r (3\bar{\rho}_d + 4\bar{\rho}_r)}{2k^2 a^2 (-8m_2^2 (3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - 2a^4 F_3 (3\bar{\rho}_d + 4\bar{\rho}_r) + 48k^4 m_2^2 F_4}, \\
M_{33} &= -\frac{a^4 \bar{\rho}_r (3\bar{\rho}_d + 4\bar{\rho}_r) (F_3 + 4\bar{\rho}_r)}{8(k^2 a^2 (-8m_2^2 (3\bar{\rho}_d + 4\bar{\rho}_r) + 3F_3 F_4 + 3F_2^2) - a^4 F_3 (3\bar{\rho}_d + 4\bar{\rho}_r) + 24k^4 m_2^2 F_4)}.
\end{aligned} \tag{3.88}$$

Now, we write down the matrix coefficients of eq. (3.72) in section 3.4.3:

$$\mathcal{L}^{(2)} = \frac{a^3}{2} \left[\dot{\Psi}_1^2 + \dot{\Psi}_2^2 + \bar{B}(t, k) (\dot{\Psi}_1 \bar{\Psi}_2 - \dot{\Psi}_2 \bar{\Psi}_1) - \bar{C}_{ij}(t, k) \bar{\Psi}_i \bar{\Psi}_j \right], \tag{3.89}$$

where

$$\begin{aligned}
\bar{B} &= -\frac{k \left(A_{22} \left(-2\dot{A}_{12} + B_{12} - B_{21} \right) + 2A_{12} \dot{A}_{22} \right)}{4A_{22} \sqrt{\bar{K}_{11}} \sqrt{\bar{K}_{22}}}, \\
\bar{C}_{12} = \bar{C}_{21} &= \{k \left(A_{22} \left(A_{12} \left(\bar{K}_{11} \left(4\bar{K}_{22} \left(\ddot{A}_{22} + \dot{B}_{22} - 2M_{22} \right) + 2\dot{A}_{22} \dot{\bar{K}}_{22} \right) \right. \right. \right. \\
&\quad \left. \left. \left. - 2\bar{K}_{22} \dot{A}_{22} \dot{\bar{K}}_{11} - 8k^2 G_{22} \bar{K}_{11} \bar{K}_{22} \right) 4\bar{K}_{11} \bar{K}_{22} \dot{A}_{12} \dot{A}_{22} \right) \right. \\
&\quad \left. + A_{22}^2 \left(-\bar{K}_{11} \left(2\bar{K}_{22} \left(2\ddot{A}_{12} + \dot{B}_{12} + \dot{B}_{21} \right) + \dot{\bar{K}}_{22} \left(2\dot{A}_{12} - B_{12} + B_{21} \right) \right) \right) \right. \\
&\quad \left. + \bar{K}_{22} \dot{\bar{K}}_{11} \left(2\dot{A}_{12} - B_{12} + B_{21} \right) + 8k^2 G_{12} \bar{K}_{11} \bar{K}_{22} \right) - 4A_{12} \bar{K}_{11} \bar{K}_{22} \dot{A}_{22}^2 \\
&\quad \left. - 6A_{22} \bar{K}_{11} \bar{K}_{22} aH \left(A_{22} \left(2\dot{A}_{12} + B_{12} + B_{21} \right) \right. \right. \\
&\quad \left. \left. - 2A_{12} \left(\dot{A}_{22} + B_{22} \right) \right) \right) \} / \{16aA_{22}^2 \bar{K}_{11}^{3/2} \bar{K}_{22}^{3/2}\}, \\
\bar{C}_{11} &= \left[6A_{22} \bar{K}_{11} H \left(A_{22} A_{12} \left(A_{12} \dot{\bar{K}}_{11} + B_{12} \bar{K}_{11} + B_{21} \bar{K}_{11} \right) \right. \right. \\
&\quad \left. \left. - A_{22}^2 \left(A_{11} \dot{\bar{K}}_{11} + B_{11} \bar{K}_{11} \right) - A_{12}^2 B_{22} \bar{K}_{11} \right) \right. \\
&\quad \left. + 2A_{12} A_{22} \bar{K}_{11} \left(A_{12} \left(-\dot{A}_{22} \dot{\bar{K}}_{11} + \bar{K}_{11} \left(2M_{22} - \dot{B}_{22} \right) + 2k^2 G_{22} \bar{K}_{11} \right) \right. \right. \\
&\quad \left. \left. + \bar{K}_{11} \dot{A}_{22} \left(4\dot{A}_{12} - B_{12} + B_{21} \right) \right) + A_{22}^3 \left(-2\bar{K}_{11} \left(\dot{A}_{11} \dot{\bar{K}}_{11} + A_{11} \ddot{\bar{K}}_{11} \right) \right. \right. \\
&\quad \left. \left. + 3A_{11} \dot{\bar{K}}_{11}^2 - 2\bar{K}_{11}^2 B'_{11} + 4k^2 G_{11} \bar{K}_{11}^2 \right) + A_{22}^2 \left(2A_{12} \bar{K}_{11} \left(2\dot{A}_{12} \dot{\bar{K}}_{11} \right. \right. \right.
\end{aligned}$$

3 Stability conditions in the presence of matter

$$\begin{aligned}
& + \text{bar}K_{11} \left(\dot{B}_{12} + \dot{B}_{21} \right) - 4k^2 G_{12} \bar{K}_{11} \left) + 2\bar{K}_{11}^2 \dot{A}_{12} \left(-2\dot{A}_{12} + B_{12} - B_{21} \right) \\
& + A_{12}^2 \left(2\bar{K}_{11} \ddot{K}_{11} - 3\dot{K}_{11}^2 \right) - 4A_{12}^2 \bar{K}_{11}^2 \dot{A}_{22}^2 \Big] / \{ 8A_{22}^3 \bar{K}_{11}^3 \} , \\
\bar{C}_{22} & = \frac{k^2}{8\bar{K}_{22}^3} \left[-2\bar{K}_{22} \left(\dot{A}_{22} \dot{K}_{11} + A_{22} \ddot{K}_{11} \right) + 3A_{22} \dot{K}_{11}^2 + 4k^2 G_{22} \bar{K}_{22}^2 \right. \\
& \left. - 6\bar{K}_{22} H \left(A_{22} \dot{K}_{11} + B_{22} \bar{K}_{22} \right) + \bar{K}_{22}^2 \left(4M_{22} - 2\dot{B}_{22} \right) \right] . \tag{3.90}
\end{aligned}$$

Note that the $A_{ij}, B_{ij}, G_{ij}, M_{ij}$ matrix components that appear in the last four coefficients have been obtained from the full expressions defined above by setting $\bar{\rho}_r = 0$.

3.7 Appendix B: Obtaining the Hamiltonian

In this appendix, we will present the derivation of the Hamiltonian used in section 3.4.3 and explain why the antisymmetric matrix B does not affect the unboundedness of the Hamiltonian. For this purpose we start from the Lagrangian (3.76):

$$\mathcal{L}^{(2)} = \frac{a^3}{2} \left[\dot{\Phi}_1^2 + \dot{\Phi}_2^2 + B(t, k) (\dot{\Phi}_1 \Phi_2 - \dot{\Phi}_2 \Phi_1) - \mu_1(t, k) \Phi_1^2 - \mu_2(t, k) \Phi_2^2 \right] . \tag{3.91}$$

Defining, the canonical momenta as:

$$\begin{aligned}
p_1 & = a^3 \left(\dot{\Phi}_1 + B(t, k) \Phi_2 \right) , \\
p_2 & = a^3 \left(\dot{\Phi}_2 - B(t, k) \Phi_1 \right) , \tag{3.92}
\end{aligned}$$

the Hamiltonian can be written as follows

$$\begin{aligned}
H & = \left\{ p_1 \left(\frac{p_1}{a^3} - B\Phi_2 \right) + p_2 \left(\frac{p_2}{a^3} + B\Phi_1 \right) - \frac{a^3}{2} \left[\left(\frac{p_1}{a^3} - B\Phi_2 \right)^2 + \left(\frac{p_2}{a^3} + B\Phi_1 \right)^2 \right. \right. \\
& \left. \left. + B \left[\left(\frac{p_1}{a^3} - B\Phi_2 \right) \Phi_2 - \left(\frac{p_2}{a^3} + B\Phi_1 \right) \Phi_1 \right] - \mu_1 \Phi_1^2 - \mu_2 \Phi_2^2 \right] \right\} \\
& = \frac{a^3}{2} \left[\left(\frac{p_1}{a^3} - B\Phi_2 \right)^2 + \left(\frac{p_2}{a^3} + B\Phi_1 \right)^2 + \mu_1 \Phi_1^2 + \mu_2 \Phi_2^2 \right] . \tag{3.93}
\end{aligned}$$

Now, in terms of $\{\dot{\Phi}_i, \Phi_i\}$ the above Hamiltonian becomes (3.78):

$$H(\Phi_i, \dot{\Phi}_i) = \frac{a^3}{2} \left[\dot{\Phi}_1^2 + \dot{\Phi}_2^2 + \mu_1(t, k) \Phi_1^2 + \mu_2(t, k) \Phi_2^2 \right] . \tag{3.94}$$

From this expression it is clear that the antisymmetric matrix does not influence the unboundedness from below of the Hamiltonian, instead such issues are encoded within the functions μ_i .

3.8 Appendix C: Mass eigenvalues for beyond Horndeski case

In section 3.4.3 we studied the mass eigenvalues of the EFTtoDE/MG in the presence of a dust fluid. We have presented the procedure and the results in the case of minimally coupled quintessence but refrained from showing general expressions due to their complexity. Here we present the mass eigenvalues for the beyond Horndeski theories which, when using the appropriate mapping, will yield the previously discussed quintessence results. Then, the eigenvalues are:

$$\begin{aligned}
 \mu_1 = & \left(-\frac{4F_1F_3 \left(F_2 \left(F_3\dot{F}_1 + F_1\dot{F}_3 \right) - 2F_1F_3\dot{F}_2 \right)^2 F_2^2}{3F_2^2 + F_1F_3} \right. \\
 & + \frac{4F_1F_3 \left(F_2 \left(F_3\dot{F}_1 + F_1\dot{F}_3 \right) - 2F_1F_3\dot{F}_2 \right)^2}{\frac{F_1F_3}{F_2^2} + 3} + (F_1F_2^2 (6F_1F_3 (3F_2^2 \\
 & + F_1F_3) \dot{a} \left(3\dot{F}_3F_2^2 + 6F_3 \left(\bar{\rho}_d - \dot{F}_2 \right) F_2 - 2F_3^2\dot{F}_1 \right) F_2^2 \\
 & + a \left(6F_3^3\dot{F}_1^2F_2^4 + F_1^2F_3 \left(\left(6F_3\ddot{F}_3 - 9\dot{F}_3^2 \right) F_2^3 + 12F_3 \left(\left(\dot{F}_2 - \bar{\rho}_d \right) \dot{F}_3 \right. \right. \\
 & + F_3 \left(\dot{\rho}_d - \ddot{F}_2 \right) \right) F_2^2 + 4F_3^2 \left(3\bar{\rho}_d \left(\dot{F}_2 - \bar{\rho}_d \right) + \dot{F}_1\dot{F}_3 - F_3\ddot{F}_1 \right) F_2 \\
 & - 8F_3^3\dot{F}_1\dot{F}_2 \right) F_2 + 2F_1^3F_3^2 \left(F_2\dot{F}_3 - 2F_3\dot{F}_2 \right)^2 + F_1 \left(9 \left(2F_3\ddot{F}_3 - 3\dot{F}_3^2 \right) F_2^6 \\
 & + 36F_3 \left(\left(\dot{F}_2 - \bar{\rho}_d \right) \dot{F}_3 + F_3 \left(\dot{\rho}_d - \ddot{F}_2 \right) \right) F_2^5 \\
 & - 12F_3^2 \left(3\bar{\rho}_d \left(\bar{\rho}_d - \dot{F}_2 \right) + F_3\ddot{F}_1 \right) F_2^4 + 4F_3^4\dot{F}_1^2F_2^2) \left. \right) / (a (3F_2^2 \\
 & + F_1F_3) (3F_2^2 + 2F_1F_3)) - (9F_2^6 \left(1 + \frac{1}{3\sqrt{\frac{F_2^4}{(3F_2^2 + 2F_1F_3)^2}}} \right) \\
 & \times \left(6F_1F_2F_3 (3F_2^2 + F_1F_3) \dot{a} \left(F_2F_3\dot{F}_1 + F_1 \left(2F_3 \left(\bar{\rho}_d - \dot{F}_2 \right) + F_2\dot{F}_3 \right) \right) \right. \\
 & + a \left(-3F_3^2\dot{F}_1^2F_2^4 + 2F_1F_3^2 \left(3F_2^2\ddot{F}_1 - F_3\dot{F}_1^2 \right) F_2^2 + F_1^2 \left(\left(6F_3\ddot{F}_3 \right. \right. \\
 & - 9\dot{F}_3^2 \right) F_2^3 + 12F_3 \left(\left(\dot{F}_2 - \bar{\rho}_d \right) \dot{F}_3 + F_3 \left(\dot{\rho}_d \right. \right. \\
 & - \ddot{F}_2 \right) \left. \left. \right) F_2^2 + 2F_3^2 \left(6\bar{\rho}_d \left(\dot{F}_2 - \bar{\rho}_d \right) - \dot{F}_1\dot{F}_3 + F_3\ddot{F}_1 \right) F_2 + 4F_3^3\dot{F}_1\dot{F}_2 \right) F_2 \\
 & + 2F_1^3F_3 \left(-2 \left(\bar{\rho}_d^2 - \dot{F}_2\bar{\rho}_d + \dot{F}_2^2 + F_2 \left(\ddot{F}_2 - \dot{\rho}_d \right) \right) F_3^2 \right. \\
 & \left. + F_2 \left(F_2\ddot{F}_3 - 2 \left(\bar{\rho}_d - 2\dot{F}_2 \right) \dot{F}_3 \right) F_3 - 2F_2^2\dot{F}_3^2 \right) \left. \right) / (a (3F_2^2 + F_1F_3))^2
 \end{aligned}$$

3 Stability conditions in the presence of matter

$$\begin{aligned}
& \times (3F_2^2 + 2F_1F_3)) \frac{1}{(16F_1^2F_2^4F_3^2)}, \\
\mu_2 = & \left(-\frac{4F_1F_3 \left(F_2 \left(F_3\dot{F}_1 + F_1\dot{F}_3 \right) - 2F_1F_3\dot{F}_2 \right)^2 F_2^2}{3F_2^2 + F_1F_3} \right. \\
& + \frac{4F_1F_3 \left(F_2 \left(F_3\dot{F}_1 + F_1\dot{F}_3 \right) - 2F_1F_3\dot{F}_2 \right)^2}{\frac{F_1F_3}{F_2^2} + 3} + \left. \left(9F_2^6 \left(1 + \frac{1}{3\sqrt{\frac{F_2^4}{(3F_2^2 + 2F_1F_3)^2}}} \right) \right) \right) \\
& \times \left(6F_1F_2F_3 (3F_2^2 + F_1F_3) \dot{a} \left(F_2F_3\dot{F}_1 + F_1 \left(2F_3 \left(\bar{\rho}_d - \dot{F}_2 \right) + F_2\dot{F}_3 \right) \right) \right) \\
& + a \left(-3F_3^2\dot{F}_1^2F_2^4 + 2F_1F_3^2 \left(3F_2^2\ddot{F}_1 - F_3\dot{F}_1^2 \right) F_2^2 \right. \\
& + F_1^2 \left(\left(6F_3\ddot{F}_3 - 9\dot{F}_3^2 \right) F_2^3 + 12F_3 \left(\left(\dot{F}_2 - \bar{\rho}_d \right) \dot{F}_3 + F_3 \left(\dot{\rho}_d - \ddot{F}_2 \right) \right) F_2^2 \right. \\
& + 2F_3^2 \left(6\bar{\rho}_d \left(\dot{F}_2 - \bar{\rho}_d \right) - \dot{F}_1\dot{F}_3 + F_3\ddot{F}_1 \right) F_2 + 4F_3^3\dot{F}_1\dot{F}_2 \left. \right) F_2 \\
& + 2F_1^3F_3 \left(-2 \left(\bar{\rho}_d^2 - \dot{F}_2\bar{\rho}_d + \dot{F}_2^2 + F_2 \left(\ddot{F}_2 - \dot{\rho}_d \right) \right) F_3^2 \right. \\
& + F_2 \left(F_2\ddot{F}_3 - 2 \left(\bar{\rho}_d - 2\dot{F}_2 \right) \dot{F}_3 \right) F_3 - 2F_2^2\dot{F}_3^2 \left. \right) \left. \right) / \left(a \left(3F_2^2 + F_1F_3 \right)^2 \right) \\
& \times \left(3F_2^2 + 2F_1F_3 \right) - \left(2F_2^2F_3 \left(6F_1F_3 \left(3F_2^2 + F_1F_3 \right) \dot{a} \left(9\dot{F}_3F_2^4 + 6F_1F_3\dot{F}_1F_2^2 \right) \right. \right. \\
& + F_1^2 \left(2F_3 \left(F_3\dot{F}_1 + 3F_2 \left(\dot{F}_2 - \bar{\rho}_d \right) \right) - 3F_2^2\dot{F}_3 \right) \left. \right) F_2^2 \\
& + a \left(-27F_3\dot{F}_1^2F_2^8 + 18F_1F_3 \left(3F_2^2\ddot{F}_1 - 2F_3\dot{F}_1^2 \right) F_2^6 + 18F_1^2F_3 \left(3F_2^2F_3\ddot{F}_1 \right. \right. \\
& - \dot{F}_1 \left(\dot{F}_3F_2^2 - 2F_3\dot{F}_2F_2 + F_3^2\dot{F}_1 \right) \left. \right) F_2^4 + 2F_1^3 \left(9 \left(\dot{F}_3^2 - F_3\ddot{F}_3 \right) F_2^4 \right. \\
& + 18F_3 \left(\bar{\rho}_d\dot{F}_3 + F_3(t) \left(\ddot{F}_2 - \dot{\rho}_d \right) \right) F_2^3 + 6F_3^2 \left(3\bar{\rho}_d^2 \right. \\
& - 3\dot{F}_2\bar{\rho}_d - 3\dot{F}_2^2 - \dot{F}_1\dot{F}_3 + 2F_3\ddot{F}_1 \left. \right) F_2^2 + 12F_3^3\dot{F}_1\dot{F}_2F_2 - 2F_3^4\dot{F}_1^2 \left. \right) F_2^2 \\
& + F_1^4F_3 \left(3 \left(\dot{F}_3^2 - 2F_3\ddot{F}_3 \right) F_2^2 + 12F_3 \left(\left(\bar{\rho}_d + \dot{F}_2 \right) \dot{F}_3 + F_3 \left(\ddot{F}_2 - \dot{\rho}_d \right) \right) F_2^2 \right. \\
& + 4F_3^2 \left(3 \left(\bar{\rho}_d + \dot{F}_2 \right) \left(\bar{\rho}_d - 2\dot{F}_2 \right) - \dot{F}_1\dot{F}_3 + F_3\ddot{F}_1 \right) F_2 + 8F_3^3\dot{F}_1\dot{F}_2 \left. \right) F_2 \\
& - 2F_1^5F_3^2 \left(F_2\dot{F}_3 - 2F_3\dot{F}_2 \right)^2 \left. \right) / \left(a \left(3F_2^2 + F_1F_3 \right)^2 \left(3F_2^2 + 2F_1F_3 \right) \right) \\
& / \left(16F_1^2F_2^4F_3^2 \right). \tag{3.95}
\end{aligned}$$