



Universiteit
Leiden
The Netherlands

Outcome of osteoarthritis and arthroplasty from patient perspective to molecular profiling.

Meessen, J.M.T.A.

Citation

Meessen, J. M. T. A. (2019, September 26). *Outcome of osteoarthritis and arthroplasty from patient perspective to molecular profiling*. Retrieved from <https://hdl.handle.net/1887/78663>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/78663>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/78663> holds various files of this Leiden University dissertation.

Author: Meessen, J.M.T.A.

Title: Outcome of osteoarthritis and arthroplasty from patient perspective to molecular profiling.

Issue Date: 2019-09-26

Patients who underwent total hip or knee arthroplasty are more physically active than the general Dutch population.

Jennifer Meessen
Wilfred Peter
Ron Wolterbeek
Suzanne Cannegieter
Claire Tilbury
Menno Bénard
Henrike van der Linden
Ron Onstenk
Rutger Tordoir
Stephan Vehmeijer
Suzan Verdegaal
Eric Vermeulen
Rob Nelissen
Thea Vliet Vlieland

Rheumatology International - 2017 37:219.



Abstract

Background: Total hip arthroplasty (THA) and total knee arthroplasty (TKA) bring relief of pain and functional disability to patients with end-stage osteoarthritis, however, the literature on their impact on patients' level of physical activity (PA) is scarce.

Methods: Cross-sectional study in patients who underwent THA/TKA surgery in the preceding 6–22 months and a random sample of persons aged >40 years from the Dutch general population, participating in a national survey. PA in minutes per week (min/week) and adherence to the Dutch recommendation for PA (NNGB yes/no) were measured by questionnaire to assess health-enhancing PA. Multivariable linear (total min/week) and logistic regression analyses (meeting recommendations PA), adjusting for confounders, were performed for THA and TKA separately.

Results: In total, 258 THA (62.3% female, aged 69.4 (9.1)) and 221 TKA (65.7% female, aged 69.5 (8.9)) patients and 4373 persons from the Dutch general population (51.4% female, aged 58.9 (11.6)) were included. The presence of THA was associated, after adjusting for age, sex, BMI education and musculoskeletal comorbidities, with more total min/week spent on PA (THA 13.8% increase, 95%CI 1.6–27.6%), whilst both TJA groups were associated with adhering to NNGB (THA: OR 1.79, 95%CI 1.26–2.56; TKA: OR 1.73, 95%CI 1.20–2.51).

Discussion: As this study used questionnaires to compare the PA of THA/TKA patients to the general population, some recall and selection bias might have been induced. After surgery, overall, TJA patients are more likely to adhere NNGB than a representative sample of persons >40 years from the Dutch general population.

Introduction

Worldwide, the numbers of patients undergoing total hip or total knee arthroplasty (THA or TKA) for hip or knee osteoarthritis (OA) are rapidly increasing. Overall, the outcomes are favourable, with a large majority of patients having less pain and improved physical functioning after surgery.^{1–4}

Although the benefits of THA and TKA are well documented for pain and function, relatively little is known on their impact on one specific aspect of physical functioning, i.e. physical activity (PA). Just like for any other individual, achieving and maintaining a sufficient level of PA is important for patients with hip and knee OA with respect to their potential general health benefits.

Moreover, in patients who undergo THA or TKA, PA may have an additional beneficial effect on the quality of the bone, which in turn may prevent complications such as early loosening.^{5–8} In addition, PA may have a positive effect on muscle strength and range of motion of the affected leg.⁹

With respect to the literature on PA after THA or TKA, de Groot et al. demonstrated in 84 THA and TKA patients that 6 months post-operatively PA levels as measured with an activity monitor did not significantly differ from the preoperative activity levels.¹⁰ Harding *et al*¹¹ found similar results when measuring PA by means of an accelerometer in 63 American THA and TKA patients before surgery and 6 months post-operatively. Kahn and Schwartzkopf¹² found no difference in PA as measured with an accelerometer between those on the waiting list for TKA and those who had TKA 2 years earlier. By using the patient-reported University of California at Los Angeles (UCLA) activity questionnaire, Baumann *et al*¹³ found that both THA and TKA patients were regularly active on moderate to high levels after on average 6–12 months after surgery. This finding is supported by Dahm *et al*¹⁴, who reported that 5.7 years after surgery TKA patients had an average physical activity score of 7.1 out of 10, with 10 being highly active. In contrast to these findings, Kahn and

Schwartzkopf¹² observed, using an accelerometer, that adherence to health-enhancing PA guidelines was only 5% in persons with TKA.

In all of these studies, a comparison with the general population was lacking. Two Dutch studies compared patient-reported physical activity in THA¹⁵ and TKA¹⁶ patients at 1–5 years postoperatively to that of age and gender-matched controls. It was found that in THA the proportion of persons reaching the Dutch Public Health Physical Activity guideline (the “Nederlandse Norm Gezond Bewegen”, NNGB) was similar to that of matched controls (51.2% (THA) vs 48.8% (controls))¹⁵, whereas in TKA patients the proportion of patients adhering to the guideline (54.5%) was significantly lower than that of the matched control population (63.7%).¹⁶ However, these studies did not take BMI into account, whilst BMI is one of the determinants of physical activity.¹⁷

Given the lack of knowledge on post-operative PA levels after total joint arthroplasty (TJA) compared to the general population, the aim of the present study was to compare the minutes of PA and proportion meeting the public health guidelines of THA and TKA patients to those of the general Dutch population. Moreover, factors other than TJA, possibly contributing to levels of physical activity, were also evaluated.

Methods

This cross-sectional, multicentre study concerned a comparison of PA levels of THA and TKA patients approximately one year after surgery with those of the general Dutch population. The data from the population of patients with THA and TKA were obtained from a study primarily aiming to make an inventory of the use of physical therapy and the presence of comorbidity¹⁸, whereas the data from the general population were obtained from the Dutch National Bureau of Statistics (in Dutch: Centraal Bureau voor de Statistiek, CBS).

Since the survey had to be filled in only once by patients, it was judged to fall outside the remit of the law for Medical Research Involving Human Subjects Act; MO [in Dutch; Wet medisch wetenschappelijk onderzoek met mensen (WMO)]. An exemption for medical ethical review was therefore given by the Medical Ethical Committee of the Leiden University Medical Center. The health monitoring conducted by the CBS commissioned by the Dutch Government also falls outside the remit of the WMO. The study was conducted in accordance with the Handbook for Good Clinical Research Practice of the World Health Organization¹⁹ and the Declaration of Helsinki principles.²⁰

Patients with THA or TKA

The patient data were obtained from a cross-sectional study performed in 2012, including patients who underwent THA or TKA for hip or knee OA in 2011 in four different hospitals in the Leiden region (Leiden University Medical Center in Leiden, Rijnland Hospital in Leiderdorp, Groene Hart Hospital in Gouda and Reinier de Graaf Hospital in Delft, the Netherlands). Patients receiving THA or TKA for reasons other than end-stage OA (such as fracture or rheumatoid arthritis) were excluded from the study, as well as patients undergoing revision surgery. Between July 2012 and October 2012, all patients operated in 2011 were approached by mail by their orthopaedic surgeon, resulting in a range of post-surgery time of 7–22 months. The orthopaedic surgeon sent all eligible persons an invitation letter, information leaflet, informed consent form, survey and pre-stamped return envelope. Patients who returned the envelope with a completed survey and signed informed consent were included in the study.

Data general population

Data from the Dutch general population were provided by the CBS and were derived from a nationwide survey on general health (Gezondheids-enquête).²¹ This questionnaire is annually administered to a representative sample of ± 8.000 Dutch inhabitants and is the prime health monitor tool of the Dutch government.²² The selection of participants is drawn from municipality registers. Persons living in institutionalized homes (e.g. nursing homes) are excluded. For the present study on physical activity, only data were selected from 2011, i.e. the same year as the data from patients with THA or TKA, and from respondents who were over 40 years of age, as none of the persons with arthroplasty was aged below 40.

Assessments

Included in both surveys were the following variables or questionnaires:

- Socio-demographic and basic health characteristics

Demographic variables included: age, gender and marital status (split into either married or not married). The height and weight of the patient were asked in order to calculate the body mass index (BMI). Smoking status (non-smoker, ex-smoker and smoker) and educational level (low (elementary school, lower secondary education), medium (secondary school or college) or high (higher secondary education or university)) were recorded.

- Physical activity (PA)

PA was assessed using the validated Dutch version of the short questionnaire to assess health (SQUASH).^{23,24} The SQUASH records the total amount of minutes per week (min/week) spent on PA in an average week in the past 12 months regarding eight different domains of active life: commuting, work activities, walking, cycling, gardening, odd jobs, household and sports. With the aid of the compendium of Ainsworth²⁵, PA can subsequently be categorized into light, moderate or vigorous intensity. Using this information, it is possible to define whether an individual adhered

to the Dutch Public Health recommendation (NNGB) for PA (30 min of moderate intensity PA on at least 5 days per week).²⁶

- Quality of life (QoL)

QoL of the persons with a THA/TKA was assessed with the Short Form 36 (SF36) questionnaire, whilst the QoL of the general Dutch population was assessed with the Short Form 12 (SF12).^{27,28} The SF36 outcomes of the THA/TKA patients were transformed to SF12 outcomes.

From the SF12, two summary scales were derived: the physical component scale (PCS) and the mental component scale (MCS). The higher the score on these scales, the better the physical or mental functioning.

- Comorbidity

The presence of comorbidity was assessed by means of a self-reported questionnaire of the CBS which comprised 19 different comorbidities.²⁹ For every comorbidity, the participants of the survey were asked to respond with either yes or no to the question “*Have you received any treatment for [condition] in the past year*”.

The included diseases were clustered into three groups:

- *Musculoskeletal comorbidities*: Severe back pain (including slipped disc), severe neck or shoulder pain, severe elbow wrist or hand pain, inflammatory arthritis or other joint conditions.
- *Non-musculoskeletal comorbidities*: Asthma or COPD (chronic obstructive pulmonary disease), (severe) cardiac disorder or coronary disease, arteriosclerosis (abdomen or legs), hypertension, (consequences of) stroke, severe bowel disorder, diabetes mellitus, migraine, psoriasis, chronic eczema, cancer and urine incontinence.
- *Sensory comorbidities*: Hearing impairments (group and face-to-face conversation), vision impairments (short and long distance) and dizziness in combination with falling.

Statistical analyses

The demographic and health characteristics of patients undergoing THA or TKA were each compared with those of the general Dutch population by means of two sample t-tests or Chi-square tests, where appropriate. Mann–Whitney tests were conducted to compare the min/week spent on PA for each of four different age groups (aged under 65, 65–69, 70–74 and 75+).

The min/week of PA was log-transformed to reach a normal distribution. Multivariable linear regression models were used to assess whether having had a joint replacement was associated with min/week spent on PA. Each analysis was done separately for THA versus the general Dutch population and TKA versus the general Dutch population. The antilog of the effect sizes (beta's) is reported for both the analyses with the 95% confidence interval (CI).

Multivariable logistic regression was used to assess the association between the presence of a joint replacement and adherence to the Dutch public health physical activity guideline. These results are presented as odds ratio (OR) with the 95% CI. The analyses were done separately for THA versus the general Dutch population and TKA and the general Dutch population.

All models (multivariable linear and multivariable logistic regression analyses) were constructed using a stepwise method. Potential confounders for the level of physical activity, i.e. sex, BMI, age and education level, were included in the models.

The determinants of minutes per week spent on activities categorized according to the three different levels of intensity of physical activity were determined for the arthroplasty groups and the Dutch population separately, by means of linear regression models including the variable of interest and correcting for age and sex. These analyses were performed including the variables age, sex, BMI, education, non-musculoskeletal comorbidities, musculoskeletal comorbidities, sensory comorbidities, MCS, PCS and time since surgery.

The level of statistical significance was set at $P < 0.05$, and analyses were performed using the SPSS statistical package (version 20.0, SPSS, Chicago, IL).

Results

Study population

Of the 545 THA and 465 TKA patients of the 4 hospitals who were invited to participate, 258 THA patients (response rate 47.3%) and 221 TKA patients (response rate 47.5%) completed the questionnaires. The selection of data from the general Dutch population from the year 2011 yielded 4373 surveys completed by people aged 40 years or older. Of those, 568 persons (13%) replied positively to the question “Have you received any treatment for osteo- or rheumatic arthritis in the past year?”.

The arthroplasty groups comprised statistically significantly more females, and the patients had a higher mean age and higher BMI than the general Dutch population. The PCS and MCS were statistically significantly lower in both the THA and the TKA groups than in the general population. There was no difference in the presence of sensory comorbidities between the arthroplasty groups and the general population. However, both musculoskeletal and non-musculoskeletal comorbidity were more present in the arthroplasty patients as compared to the general population (see also Table 1).

Table 1 - Characteristics of patients with Total Joint Arthroplasty and the general Dutch population

		General Dutch Population	Total Hip Arthroplasty	P*	Total Knee Arthroplasty	P*
Total		4373	258		221	
Female		2248 (51.4%)	159 (62.3%)	0.01	144 (65.7%)	<0.01
Age (year)		59 ± 11.6	69 ± 9.1	<0.01	70 ± 8.9	<0.01
Body Mass Index		26 ± 5.0	27 ± 4.1	<0.01	29 ± 5.0	<0.01
Education level	Low	653 (15.6%)	75 (37.3%)		72 (41.1%)	
	Medium	2391 (57.0%)	86 (42.8%)	<0.01	81 (46.3%)	<0.01
	High	1150 (27.4%)	40 (19.9%)		23 (12.6%)	
SF-12	PCS	53.9 ± 9.4	47.6 ± 11.2	<0.01	46.6 ± 10.8	<0.01
	MCS	44.1 ± 5.0	39.8 ± 5.2	<0.01	40.6 ± 5.1	<0.01
Comorbidities	≥ 1 Non-MSK	2465 (57.4%)	142 (72.8%)	<0.01	138 (84.1%)	<0.01
	≥1 MSK	1224 (27.9%)	93 (39.7%)	<0.01	84 (40.7%)	<0.01
	≥1 Sensory	400 (9.3%)	21 (8.4%)	0.736	21 (9.9%)	0.717

Variables reported as mean±SD or N(%).

*P - P-value for two sample T test or Chi² between hip/knee arthroplasty and the general population
MSK – Musculoskeletal comorbidities, PCS/MCS – Physical /Mental Component Score

Table 2 - Total minutes per week spent on physical activity by, age group for hip or knee arthroplasty or the general Dutch population

Gender	Aged	General Dutch population		Total hip arthroplasty			Total knee arthroplasty		
		N	Mean \pm SD	N	Mean \pm SD	P*	N	Mean \pm SD	P*
Men	<65	2521	2846 \pm 1543	32	2722 \pm 1252	0,692	21	2924 \pm 1468	0,959
	65-69	414	1976 \pm 1474	23	2315 \pm 1336	0,052	20	2079 \pm 1316	0,640
	70-74	313	1784 \pm 1330	20	2130 \pm 843	0,003	12	1968 \pm 1667	0,912
	≥ 75	372	1461 \pm 1209	21	1443 \pm 1065	0,840	22	2193 \pm 1477	0,018
Female	<65	2761	2845 \pm 1530	38	2745 \pm 1148	0,857	46	2883 \pm 1695	0,956
	65-69	401	2075 \pm 1412	34	2480 \pm 1908	0,364	25	2048 \pm 1539	0,764
	70-74	286	1967 \pm 1273	33	1867 \pm 1071	0,888	28	1944 \pm 2122	0,239
	≥ 75	525	1449 \pm 1173	53	1703 \pm 1453	0,350	45	1284 \pm 912	0,577
		N	Count (%)	N	Count (%)	P*	N	Count (%)	P*
Adherence to NGBB		4373	2954 (67,6%)	258	195 (75,6%)	0,007	221	161 (72,9%)	0,105

* P-value for Mann-Whitney or Chi² as compared to the general population

Table 2 shows the crude number of minutes spent per week on PA, stratified for age and gender. As can be seen in Figure 1, male arthroplasty patients spend more minutes per week physically active in the higher age groups as compared to the general Dutch population. The proportion of persons adhering to the NNGB guideline is in the arthroplasty groups higher than in the general Dutch population (THA 76%, TKA 73% and general Dutch population 68%).




Fig. 1 - Stratified representation of total minutes per week physical activity per age group and gender

Association between TJA and minutes per week spent on total physical activity

Univariately, both THA and TKA were significantly associated with more minutes per week physical activity when compared to the general population. As there were major differences between the groups, it was needed to correct for potential confounders. When correcting for age, gender, BMI, education and musculoskeletal comorbidities a statistically significant association was for THA, not TKA. With the adjustments, persons with a THA spend 13.8% more minutes per week on physical activity compared to the general population (see Table 3).

Table 3 – Regression analyses of min/week physical activity and adherence to NNGB.

	Total min/week Physical Activity					
	Total Hip Arthroplasty			Total Knee Arthroplasty		
	Beta	95% CI	P*	Beta	95% CI	P*
Univariate ^A	1.229	1.099-1.352	<0.001	1.324	1.186-1.483	<0.001
Multivariate 1 ^B	1.140	1.023-1.274	0.018	1.122	0.998-1.262	0.055
Multivariate 2 ^C	1.138	1.016-1.276	0.024	1.112	0.986-1.256	0.084
	Adherence to NNGB					
	Total Hip Arthroplasty			Total Knee Arthroplasty		
	OR	95% CI	P*	OR	95% CI	P*
Univariate ^A	1.487	1.111-1.989	0.008	1.289	0.952-1.745	0.101
Multivariate 1 ^B	1.750	1.243-2.465	0.001	1.750	1.219-2.512	0.002
Multivariate 2 ^C	1.789	1.253-2.556	0.001	1.731	1.195-2.507	0.004

A – Univariate analysis

B – adjusted for age, sex, BMI and education

C – adjusted for age, sex, BMI, education and musculoskeletal comorbidities.

Association between TJA and meeting public health recommendation

The multivariable logistic regression models showed that, adjusted for age, gender, BMI, education and muscular comorbidities, both THA and TKA patients had a significantly higher likelihood of meeting public health recommendations for healthy PA as compared to the general population (THA: OR 1.79 (95%CI 1.25–2.55); TKA: OR 1.73 (95%CI 1.20–2.51)). See also Table 3.

Determinants of physical activity

Regarding the determinants of physical activity in the general population, age, sex, PCS and education were found to be statistically significantly associated with total minutes per week spent on PA and per category of intensity (see Table 4).

In the general population, BMI was associated with the number of min/week of moderate and vigorous intensity PA, but not with light intensity PA. Within both the arthroplasty groups, it was found that age was a significant determinant of total min/week of PA and the min/week of light intensity PA. Within the THA group, age was also a determinant for the min/week of moderately intensive PA. Sex was associated with the min/week of moderate and vigorous intensity PA in both arthroplasty groups and also with the min/week of light intensity PA in the THA group.

In the general population, comorbidities were only found to be a determinant of total minutes per week of PA, but not of the min/week in the three categories of PA intensity. In THA patients only the presence of sensory comorbidities was associated with the total min/week of PA, whereas in TKA the presence of comorbidities was not associated with PA.

Although the association of a number of potential determinants with PA in the TKA and TJA groups did not reach statistical significance, overall the directions of the associations were similar to those within the general population (results not shown).

Table 4 – Variables statistically significantly associated with min/week physical activity after adjusting for age and sex.

		General Dutch population		Total hip arthroplasty		Total knee arthroplasty	
		Beta	P*	Beta	P*	Beta	P*
Total Physical Activity	Age	0.973	<0.001	0.966	<0.001	0.966	<0.001
	Sex ^A	0.953	0.037				
	BMI	0.995	0.022	0.960	0.005		
	Physical Component Score	1.016	<0.001				
	Low education ^B	0.774	<0.001				
	Non-musculoskeletal comorbidities ^C	1.076	0.002				
	MusculoskeletalS comorbidities ^C	1.052	0.046				
	Sensory comorbidities ^C	1.202	<0.001	1.517	0.028		
Light Physical Activity	Age	0.966	<0.001	0.604	<0.001	0.973	<0.001
	Sex ^A	0.748	<0.001	0.604	<0.001		
	BMI					0.759	0.044
	Physical Component Score	1.007	<0.001				
	Low education ^B	0.809	<0.001				
	Medium education ^B	0.834	<0.001				
Moderate Physical Activity	Age	0.991	<0.001	0.984	0.031		
	Sex ^A	1.084	0.020	1.596	0.01	1.538	0.009
	BMI	0.912	0.022			0.964	0.027
	Physical Component Score	1.016	<0.001				
	Mental Component Score	0.989	0.001				
	Low education ^B	1.180	0.006				
	Medium education ^B	1.384	<0.001				
Vigorous Physical Activity	Age	1.026	<0.001				
	Sex ^A	1.189	<0.001	1.489	0.005	1.432	0.022
	BMI	0.986	0.003				
	Physical Component Score	1.014	<0.001			1.021	0.027
	Mental Component Score					0.957	0.008
	Medium education ^B	1.151	0.002				

A – Females were reference

B – High education as reference

C - Affected as reference

Discussion

This study demonstrated that the presence of a THA was associated with more min/week spent on PA as well as better adherence to public health recommendations for PA (NNGB) when compared to the general population. TKA was found to only be associated with adhering to the NNGB when compared to the general Dutch population.

Overall, it seems the Dutch population spends more minutes per week on physical activity, but since the patient group differs from the general population the comparison between these groups should be adjusted. When adjusting for age, sex, BMI and education it is found that persons with THA do spend more minutes per week on physical activity and that persons with a THA and TKA are more likely to adhere to the Dutch guideline on physical activity, NNGB.

That TKA is associated with the NNGB but not to the minutes per week activity can be explained by the level of intensity of the physical activity performed. In the general population, more associations between potential determinants of physical activity and the actual numbers of PA reached statistical significance than in the arthroplasty groups. The lack of significance is probably due to the relatively small sample sizes in the arthroplasty groups, limiting the statistical power.

Our groups spent more min/week on PA than reported by two other Dutch studies (for THA in this study 2183 min/week PA, THA in Wagenmakers *et al*¹⁵ 1601 min/week, for TKA in this study 2153 min/week PA, TKA in Kersten *et al*¹⁶ 1347 min/week PA). In parallel, regarding the proportion of patients adhering to the Dutch recommendation for physical activity, the outcomes were more favourable in the present study (THA in this study 75.6% and THA in Wagenmakers *et al*¹⁵ 51.2%; TKA in this study 72.8% and TKA in Kersten *et al*¹⁶ 55%).

Both these latter two studies were done at 1–5 years post-surgery, and our study included patients within the first 22 months after surgery. As reported earlier by our group (Peter *et al*³⁰), 43.5% of the THA patients and 50.5% of the TKA patients had post-operative physiotherapy for more than 3 months. This implies that a vast

amount of our patients might still have intense training with aid of physiotherapists, motivating patients to adhere to the PA. As for the other two studies (Kersten, Wagenmakers), no data on prolonged post-operative physiotherapy are present, and thus, these patients might resume easier into their old, less active activity level.

A recent systematic review on physical activity after THA or TKA measured with accelerometers showed that the post-operative PA levels were lower in the arthroplasty groups as compared to healthy control participants.³¹ The differences in outcome could be because our sample of the general population might not be totally healthy and be less active than selected healthy persons. Also, as this study used a questionnaire whilst the systematic review concerned objective measures, participants might have caused some recall bias.

The general population in our study had an adherence rate to the Dutch PA of 67.5% which is comparable to reports from CBS published (Dutch adult population, 66% adhered to the Dutch public health physical activity guideline in 2012).³² The minutes per week spent on PA in our study was also consistent with the numbers reported by CBS (2589 min per week in 2012 and 2525 min per week in our study for overall physical activity for the Dutch population).³³

Factors we identified as influencing the level of PA of persons with hip or knee arthroplasty (BMI, increased age, physical component score) are in line with the findings in a systematic review by Stubbs *et al*³⁴ regarding PA in patients with hip or knee OA. The inverse association of BMI on the level of PA shows that it is an important factor, as well as age and gender, to include in any case–control study.¹⁷ Low-impact activities like walking or cycling seem to protect against function loss and experienced pain from OA, in contrast heavy load activities might be a risk factor for the development of osteoarthritis, but also early implant failure although debate exists on the latter.³⁵⁻³⁸ Since contradictory evidence exists on this topic, research into this field is necessary.

Current post-operative rehabilitation after a hip or knee arthroplasty is focussed at independent ambulation and regaining a normal walking pattern, which was deteriorated in the years before surgery due to the slowly progressing osteoarthritis. Secondary to this it aims at getting the patient physically active. As mentioned before, about half of our patients reported to receive physiotherapy for more than 3 months after surgery.³⁰ This might imply that these patients are more motivated to be active than the general population.

Another reason for the higher levels of PA in the arthroplasty groups might be the fact that PA is a risk factor for TJA.³⁹ As shown by de Groot *et al*¹⁰, the post-operative levels of PA did not significantly differ from preoperative levels, suggesting that PA levels of TJA were probably higher than those of the general population before surgery as well. Finally, the patients filling in the questionnaire knew that the subject of the study was PA, whilst the Dutch general population had to fill in an elaborate list of questions including all aspects of life, with only a subset on PA. Thus, the patients in our study might have overestimated their PA.

The limitations of this study are potential overestimation of outcome measures and recall bias, due to using the SQUASH questionnaire, more objective measures like accelerometers should be used in future studies. Furthermore, the preoperative levels of PA should be taken into account as an important confounder for outcome as well. Thus, more valid comparisons with the general population are possible.

Also, patients in our study who refused to fill in the questionnaires were not asked about their reasons as to why they declined to participate, and therefore, we have no information about any possible self-selection bias. In addition, the comorbidities of participants were all self-reported and we were unable to confirm the presence of comorbidities both in the general Dutch population and the arthroplasty groups.

The findings of this study give insights into the movement patterns of arthroplasty patients compared to the general Dutch population. Findings show that although a part of the arthroplasty patients adhere to the Dutch public health guideline, there is still a considerable group who should increase their PA levels.

Acknowledgements

The authors would like to thank all the participants of the original physiotherapy study and the Gezondheids Enquete. A special thanks to the onsite CBS data assistants (Martin Broxterman, Michael Vermaessen, Alfred Baven, Ivo Gorissen and Jan-Willem Bruggink) for their support and assistance.

References

1. Meftah M *et al.* (2012). Ten-year follow-up of a rotating-platform, posterior-stabilized total knee arthroplasty. *J Bone Joint Surg Am* 94(5):426–432.
2. Meding JB *et al.* (2012). Pain relief and functional improvement remain 20 years after knee arthroplasty. *Clin Orthop Relat Res* 470(1):144–9.
3. Kurtz S *et al.* (2007). Projections of primary and revision hip and knee arthroplasty in the US from 2005 to 2030. *J Bone Jt Surg Am* 89(4):780–785.
4. Wolf BR *et al.* (2012). Adverse outcomes in hip arthroplasty: long-term trends. *J Bone Jt Surg Am* 94(14):103.
5. Kuster MS (2002). Exercise recommendations after total joint replacement: a review of the current literature and proposal of scientifically based guidelines. *Sports Med* 32(7):433–445.
6. Trudelle-Jackson E and Smith SS (2004). Effects of a late-phase exercise program after total hip arthroplasty: a randomized controlled trial. *Arch Phys Med Rehabil* 85(7):1056–1062.
7. Cuschnaghan J *et al.* (2009). Long-term outcome following total knee arthroplasty: a controlled longitudinal study. *Ann Rheum Dis* 68(5):642–647.
8. Sharma L *et al.* (2003). Physical functioning over three years in knee osteoarthritis: role of psychosocial, local mechanical, and neuro muscular factors. *Arthritis Rheum* 48(12):3359–70.
9. Reeuwijk KG *et al.* (2010). Osteoarthritis of the hip or knee: which coexisting disorders are disabling? *Clin Rheumatol* 29(7):739–747.
10. de Groot IB *et al.* (2008). Small increase of actual physical activity 6 months after TJA. *Clin Orthop Relat Res* 466(9): 2201–2208.
11. Harding P *et al.* (2014). Do activity levels increase after total hip and knee arthroplasty? *Clin Orthop Relat Res* 472(5):1502–1511.
12. Kahn TL and Schwarzkopf R (2015). Does total knee arthroplasty affect physical activity levels? Data from the osteoarthritis initiative. *J Arthroplasty* 30(9):1521–1525.
13. Bauman S *et al.* (2007). Physical activity after total joint replacement: a cross-sectional survey. *Clin J Sport Med* 17(2):104–108.
14. Dahm DL *et al.* (2008). Patient-reported activity level after total knee arthroplasty. *J Arthroplasty* 23(3):401–407.
15. Wagenmakers R *et al.* (2008). Habitual physical activity behavior of patients after primary total hip arthroplasty. *Phys Ther* 88(9):1039–1048.
16. Kersten RF *et al.* (2012). Habitual physical activity after total knee replacement. *Phys Ther* 92(9):1109–1116.
17. Okeyo OD *et al.* (2009). Physical activity and dietary fat as determinants of body mass index in a cross-sectional correlational design. *East Afr J Public Health* 6(1):32–36.
18. Peter WF *et al.* (2015). The association between comorbidities and pain, physical function and quality of life following hip and knee arthroplasty. *Rheumatol Int* 35(7):1233–1241.
19. WHO (2005). Handbook for good clinical research practise (GCP): guidance for implementation.
20. WMA (2013). Declaration of Helsinki—ethical principles for medical research involving human subjects. *JAMA* 310(20):3.
21. van den Berg J and van der Wulp C (2003). Rapport van de Werkgroep Revisie POLS Gezondheidsenquête. Centraal Bureau voor de Statistiek. Retrieved from: <http://www.cbs.nl> on 9/11/15.
22. Rijksinstituut voor Volksgezondheid en Milieu (RIVM). Gezondheidsenquête. Retrieved from <https://bronnen.zorggegevens.nl> on 11/11/15.
23. Wagenmakers R *et al.* (2008). Reliability and validity of the short questionnaire to assess health-enhancing physical activity (SQUASH) in patients after THA. *BMC Musculoskeletal Disord* 9:141.
24. Wendel-Vos GC *et al.* (2003). Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 56(12):1163–1169.
25. Ainsworth BE *et al.* (2011). Compendium of physical activities: a second update of codes and MET values. *Med Sci Sports Exerc* 43(8):1575–81.
26. Volksgezondheid NK (2015). Nationaal Kompas Volksgezondheid: normen van lichamelijke (in)activiteit. Retrieved from: <http://www.nationaalkompas.nl/> on 11/11/15.
27. Aaronson NK *et al.* (1998). Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J Clin Epidemiol* 51(11):1055–1068.
28. Gandek B *et al.* (1998). Cross-validation of item selection and scoring for the SF-12 Health Survey in nine countries: results from the IQOLA Project. *J Clin Epidemiol* 51(11):1171–1178.
29. Sociaal Cultureel Planbureau. Onderzoek naar zorggebruik 2014–2017. Retrieved from <https://www.scp.nl/> on 10/11/2015.
30. Peter WF *et al.* (2016). The provision of preoperative and postoperative physical therapy in elderly people with hip and knee osteoarthritis undergoing primary joint replacement surgery. *Curr Orthop Pract* 27(2):10.
31. Arnold JB *et al.* (2016). Does physical activity increase after total hip or knee arthroplasty for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther* 46(6):431–442.
32. Nationaal Kompas Volksgezondheid (2012). Gezondheids determinanten - Fysieke Activiteit. Retrieved from <http://www.nationaalkompas.nl/> on 10/11/15.
33. Centraal Bureau voor de Statistiek. Statline Retrieved from <http://statline.cbs.nl/> on 20/9/15.
34. Stubbs B *et al.* (2015). What are the factors that influence physical activity participation in adults with knee and hip osteoarthritis? A systematic review of physical activity correlates. *Clin Rehabil* 29(1):80–94.
35. Fransen M *et al.* (2002). Therapeutic exercise for people with osteoarthritis of the hip or knee. A systematic review. *J Rheumatol* 29(8):1737–1745.
36. van Baar ME *et al.* (1999). Effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a systematic review of randomized clinical trials. *Arthritis Rheum* 42(7):1361–1369.
37. Vogel LA *et al.* (2011). Physical activity after total joint arthroplasty. *Sports Health* 3(5):441–450.
38. Golant A *et al.* (2010). Athletic participation after hip and knee arthroplasty. *Hosp Jt Dis* 68(2):76–83.
39. Wang Y *et al.* (2011). Is physical activity a risk factor for primary knee or hip replacement due to osteoarthritis? A prospective cohort study. *J Rheumatol* 38(2):350–357.