



Universiteit
Leiden
The Netherlands

Outcome of osteoarthritis and arthroplasty from patient perspective to molecular profiling.

Meessen, J.M.T.A.

Citation

Meessen, J. M. T. A. (2019, September 26). *Outcome of osteoarthritis and arthroplasty from patient perspective to molecular profiling*. Retrieved from <https://hdl.handle.net/1887/78663>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/78663>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/78663> holds various files of this Leiden University dissertation.

Author: Meessen, J.M.T.A.

Title: Outcome of osteoarthritis and arthroplasty from patient perspective to molecular profiling.

Issue Date: 2019-09-26

Frailty in end-stage hip or knee osteoarthritis: Validation of the Groningen Frailty Indicator (GFI) questionnaire

Jennifer Meessen
Claudia Leichtenberg
Claire Tilbury
Bart Kaptein
Lennard Koster
Eline Slagboom
Suzan Verdegaal
Ron Onstenk
Henrike van der Linden
Herman Kaptijn
Stephan Vehmeijer
Willem-Jan Marijnissen
Pieter-Jan Damen
Rob Nelissen
Thea Vliet Vlieland



Abstract

Background/Objective: Frailty is highly prevalent in the elderly, increasing the risk of poor health outcomes. The Groningen Frailty Indicator (GFI) is a 15-item validated questionnaire for the elderly. Its value in patients with end-stage hip or knee osteoarthritis (OA) has not yet been determined. This study assesses the validity of the GFI in this patient-group.

Methods: End-stage hip or knee OA patients completed the GFI (range 0-15, ≥ 4 is frail) before arthroplasty surgery. Convergent validity was determined by Spearman-rank correlation between the SF12 physical (PCS) and mental (MCS) component scores and the physical and mental GFI-domains, respectively. Discriminant validity was assessed by means of overall GFI-score and the pain-domain of the Hip/Knee Osteoarthritis Outcome Score (HOOS/KOOS).

Results: 3275 patients were included of whom 2957 (90.3%) completed the GFI. Mean GFI-scores were 2.78(2.41) and 2.28(1.99) in hip and knee OA-patients, respectively, with 570(35.9%) of hip and 344(24.1%) of knee patients considered frail. The convergent validity was moderate to strong (physical domain $R=-0.4$, mental domain $R=-0.6$) and discriminant validity low (R HOOS/KOOS-pain domain= -0.2), confirming the validity of the GFI-questionnaire in this population.

Conclusion: With 90% of participants completing the GFI, it is a feasible and valid questionnaire to assess frailty in end-stage hip and knee OA-patients. One-third (33.3%) of the patients undergoing hip arthroplasty and a quarter (24.1%) of those undergoing knee arthroplasty are frail. Whether this is associated with worse outcomes and can thus be used as a pre-operative predictor needs to be explored.

Introduction

Osteoarthritis (OA) is a degenerative joint disease which often leads to disability and pain. A highly effective treatment for end stage OA is arthroplasty surgery.^{1,2} Over 202,500 total hip and 402,100 total knee arthroplasties (THA and TKA) are performed annually in the United States of America alone, with the volume expected to increase up to 6-fold by 2030.³

At present, 83% of the patients receiving THA and 79% of patients receiving TKA are older than 60 years of age.⁴ As frailty is highly prevalent in the elderly, it is likely that a considerable proportion of patients undergoing THA or TKA are frail.⁵ Although there is not one definition for frailty, the most often used definitions include a combination of decrease of independence, strength, cognition, activity, energy, weight and walking speed.⁶⁻¹² Literature shows that there is considerable heterogeneity in the extent of frailty individuals may experience, with some persons accelerating fast while others are slowly progressing to higher levels of frailty.¹³ Within persons of the same age, also the onset of frailty differs per individual.¹⁴⁻¹⁷

It is generally acknowledged that frailty hampers the ability to resist stressors, leading to vulnerability for adverse outcomes after surgery.^{6, 16-19} As such, it is of importance to have more insight into frailty in the group of patients undergoing THA or TKA. As a first step into the exploration of the role of frailty in the outcomes of total joint surgery, an appropriate instrument for frailty is needed

The Groningen Frailty Indicator (GFI) is a frequently used questionnaire in the elderly to assess frailty. The advantage of the GFI is that it is a self-reported score, furthermore, this questionnaire has been validated specifically for elderly (mean age 81 years). In these elderly (both community dwelling and institutionalized), it was found that the GFI is feasible, reliable and valid.²⁰ However, it is not known yet how feasible the GFI is in a clinical setting as well as the validity of the GFI amongst the somewhat younger patients with end stage hip or knee OA waiting for arthroplasty surgery. Therefore, in this study we aimed to assess the feasibility and validity of the GFI as a tool to measure frailty in end stage hip or knee osteoarthritis patients scheduled to undergo arthroplasty surgery.

Methods

Study design

This study is part of the Longitudinal Leiden Orthopaedics Outcomes of Osteo-Arthritis study (LOAS). The LOAS study is an ongoing, multi-center, longitudinal prospective cohort study including patients undergoing primary total hip or knee arthroplasty (THA or TKA). Participants are recruited in 7 participating hospitals (the Leiden University Medical Center, Leiden; Alrijne Hospital, Leiden/Leiderdorp (former Diaconessenhuis and Rijnland Hospital); Groene Hart Hospital, Gouda; LangeLand Hospital, Zoetermeer; Reinier de Graaf Gasthuis, Delft; Albert Schweitzer Hospital, Dordrecht; Waterland Hospital, Purmerend). The LOAS study (Trial ID NTR3348) started in June 2012. The present study is only concerned with data gathered preoperatively from June 2012 – June 2016.²¹

Patients

All patients who were able to complete questionnaires in Dutch and who were 18 years or older were eligible for participation. Excluded were patients who did not provide informed consent, had insufficient Dutch language skills or of whom the physical or mental status did not allow participation. Eligible patients were informed about the study through written and oral information by their treating surgeon at the outpatient clinic. Only patients who agreed to be approached by the researcher received additional written information about the study by regular mail or e-mail, as well as a questionnaire, a stamped return envelope and a consent form.

Patients were included in the study once written informed consent was obtained according to the Declaration of Helsinki.²² For the purpose of the present analysis only data from patients who returned the preoperative questionnaire between the start of the study in June 2012 until June 2016 were included. Ethical approval was obtained by the Medial Ethics Committee of the Leiden University Medical Center (registration number P12.047) and funding was received from the Dutch Arthritis Foundation (LLP13).

The questionnaires were incorporated in current clinical setting of the included hospitals which all participate in the collection of patient reported outcome measures (PROMs) for the national Dutch Arthroplasty Register (LROI).

Assessments

Frailty: Frailty was assessed by the Groningen Frailty Indicator (GFI). This questionnaire consists of 15 questions covering several aspects of life, such as independence in daily tasks, involuntary weight loss, medication use, mental state, vision and hearing. Together these questions lead to a score between 0 and 15, a score of ≥ 4 is considered to be frail. The GFI is specifically directed to elderly persons both living at home as well as in institutions.^{20,23,24}

Overall health: Quality of life was measured using the validated Dutch version of the Short Form (SF)-12.²⁵ The SF-12 comprises 12 items on generic measurement of the overall health-related-quality of life. Scores range from 0 to 100, with 0 being lowest possible score and 100 the highest. From the SF12, 2 subscales can be calculated, the physical component score (PCS) and mental component score (MCS). These subdomains were assessed separately in the analyses.²⁶

Hip / Knee Symptoms: The Hip disability/Knee injury Osteoarthritis Outcome Score (HOOS/KOOS) questionnaires are validated questionnaires to measure the function of patients with end-stage osteoarthritis for hip or knee respectively.^{27,28} These questionnaires comprise five domains (activities of daily living, quality of life, sports, symptoms and pain). For the current study the validated Dutch version was used.^{29,30}

Statistical Analyses

Patient characteristics were analysed using descriptive statistics. Rates of patients who did not, partially or completely filled out the GFI were computed. Comparisons between patients who filled in the GFI completely and those who did not or partially were done by means of either Chi-Square tests for categorical variables and t-tests for continuous variables. In addition, for each GFI item the proportion of missing values was determined.

To explore determinants for completing the questionnaire a binary variable “completion of questionnaire” was constructed. This variable was used in a logistic

regression analysis to see if age, sex, BMI and comorbidities are of significant influence on the completion of the questionnaire.

The internal consistency of the GFI in this patient population was assessed by means of Cronbach's alpha, with an alpha of >0.7 being considered as good consistency [31]. Convergent validity of the GFI was determined by computing correlations between the physical domain of GFI (questions 1-9) and the PCS of the SF-12. The mental domain of the GFI (question 14 and 15) was correlated with the MCS of the SF-12. Correlations were computed using a Spearman rank correlation coefficients. As the corresponding subscales of the GFI and SF12 aim to measure similar constructs it was hypothesized that the correlation between the subscales of the GFI and SF12 will be high.

Discriminant validity of the questionnaire was assessed by correlating the physical domain of the GFI to the MCS and the mental domain of the GFI to the PCS. Also, a Spearman rank correlation analysis including the total GFI-score and pain as measured by the HOOS/KOOS questionnaire was performed. As the correlated constructs are conceptually different, we hypothesized the correlation between these domains would be low.

For those THA and TKA patients who completed the GFI the prevalence of frailty was calculated, based on the cut-off score of four.²⁴ The demographic variables of those assigned frail and those not designated as frail were compared by means of a t-test or Chi-square test, whichever was appropriate.

All analyses were performed with IBM SPSS statistics software version 23.

Results

Within the time frame of the present analysis 3275 patients with end-stage hip OA (N=1691) and knee OA (N=1584) were included in the cohort study. For both end stage hip and knee OA, 90.3% of the participants completed the questionnaire. In Table 1 the socio-demographic variables of patients returning the questionnaire that did and did not complete it fully were compared. In hip OA, those who did not fully complete the questionnaire were significantly older, whereas in knee OA those who did not complete the questionnaire fully were more often female and had a lower score on the HOOS/KOOS-activities of daily life domain. In both end stage hip and knee OA those who did not complete the questionnaire had a significantly lower score on the MCS.

Table 1 – Characteristics of patients with end stage OA who did and did not complete the Groningen Frailty Indicator questionnaire.

	End stage Hip OA			End stage Knee OA			
	GFI completed N=1527	GFI incomplete N=164	P*	GFI completed N=1430	GFI incomplete N=154	P*	
Female	925 (61.5%)	107 (67.3%)	0.155	911 (64.2%)	119 (77.3%)	0.001	
Age	67.8 ± 9.8	70.9 ± 9.4	<0.001	67.4 ± 8.9	67.6 ± 9.1	0.818	
BMI	27.2 ± 4.3	27.0 ± 5.4	0.529	29.4 ± 4.7	29.0 ± 4.4	0.373	
Not living alone	1187 (77.7%)	118 (71.9%)	0.097	1095 (76.5%)	115 (75.7%)	0.598	
Musculoskeletal Comorbidities	259 (17.8%)	29 (20.9%)	0.370	326 (24.1%)	39 (26.5%)	0.522	
Other Comorbidities	942 (70.7%)	80 (69.0%)	0.692	900 (74.7%)	85 (73.9%)	0.855	
SF12	Physical	32.2 ± 9.4	32.4 ± 9.2	0.821	32.3 ± 9.1	32.4 ± 9.7	0.918
	Mental	54.8 ± 9.9	52.9 ± 10.4	0.046	55.6 ± 9.4	54.0 ± 9.0	0.009
HOOS / KOOS	Pain	37.9 ± 18.6	39.8 ± 20.0	0.244	38.9 ± 17.6	36.4 ± 18.8	0.124
	Symptoms	39.8 ± 18.5	41.9 ± 20.6	0.252	43.7 ± 13.5	42.0 ± 12.4	0.178
	Activities of daily life	39.9 ± 19.2	41.8 ± 21.6	0.324	45.0 ± 18.2	40.8 ± 20.9	0.026
	Sport	18.1 ± 18.4	21.6 ± 21.7	0.200	10.7 ± 14.3	11.2 ± 15.5	0.852
	Quality of Life	33.4 ± 10.8	35.2 ± 12.1	0.083	33.6 ± 10.4	34.6 ± 11.8	0.327

* Characteristics of patients who completed and did not complete the GFI were tested by means of a T-test (normal distribution), Mann-Whitney test (not normal distribution) or Chi-square test (discrete variables).

On a total of 15 items, the median number of missing items for both joint locations was 0 (range 0 to 15), whereas the mean (SD) was 0.4 (1.9) (hip OA: 0.4 (2.0), knee OA: 0.3 (1.8)). Of the 164 patients with hip OA who did not complete all questions, 29 did not fill in any question whereas 99 missed only one question. Of the 154 patients with knee OA who did not complete all questions, 21 did not fill in any question and 102 persons had only one missing question.

Table 2 shows the percentage of missing values per question. Most frequently missed was question 15 “*How would you rate your physical fitness on a scale of 1 to 10?*” for both hip and knee (hip 4.4% missing, knee 4.2% missing). This was the only question with no predefined answering options; instead patients had to write down the number themselves. In addition, in patients with hip OA question 2 “*Are you able to walk independently outside?*” (2.8% missing) and question 3 “*Are you able to (un)dress yourself?*” (2.7% missing) were relatively often missing, while in knee OA patients question 6 “*Do you encounter problems in daily life because of impaired hearing?*” (2.6% missing) and question 2 “*Are you able to walk independently outside?*” (2.3% missing) were relatively often missing.

	Hip	Knee
1 Are you able to do groceries by yourself?	2.5%	1.9%
2 Are you able to walk independently outside?	2.8%	2.3%
3 Are you able to (un)dress yourself?	2.7%	2.2%
4 Are you able to use the bathroom by yourself?	2.7%	2.0%
5 Do you encounter problems in daily life because of impaired vision?	2.5%	2.6%
6 Do you encounter problems in daily life because of impaired hearing?	2.4%	1.8%
7 Did you unintentionally lose weight over the past 6 months?	2.4%	1.8%
8 Do you use 4 or more types of medication	2.7%	1.8%
9 Do you have any complaints of your memory?	2.1%	1.8%
10 Do you experience emptiness around you?	2.2%	1.8%
11 Do you miss the presence of other people around you?	2.4%	2.0%
12 Do you feel left alone?	2.7%	1.8%
13 Have you felt down or depressed lately?	2.5%	2.0%
14 Have you felt nervous or anxious lately?	2.5%	2.0%
15 How would you rate your physical fitness on a scale of 1-10 ?	4.4%	4.2%

To assess determinants for completing the GFI questionnaire a logistic regression model was build including age, sex, BMI, musculoskeletal and other comorbidities. Table 3 shows the odds ratio's associated to this model. It was found that age and sex are statistically significant determinants for completing the questionnaire in persons with end-stage OA of the lower limb corrected for BMI and comorbidities.

Table 3 – Odds ratio's for demographic characteristics associated with completing the GFI.

	Odds Ratio	95% CI	P-value
Age	0.981	0.966-0.997	0.020
Sex	1.497	1.100-2.038	0.010
BMI	1.006	0.974-1.039	0.714
Musculoskeletal comorbidities	0.946	0.661-1.354	0.762
Other comorbidities	0.890	0.644-1.230	0.481

Characteristics were included in logistic regression analysis to assess their association with completing the GFI questionnaire (yes/no).

Older age is, independent of gender, BMI and comorbidities, associated to lower odds for completing the questionnaire (OR: 0.98, P-value 0.020) while for gender it was found that, when correcting for age, BMI, musculoskeletal and other comorbidities, females have higher odds for completing the questionnaire as compared to males (OR: 1.50, P-value; 0.010). BMI and having musculoskeletal or other comorbidities were not statistically significant associated to the completing of the GFI questionnaire for persons with end-stage hip or knee OA.

The internal consistency of the GFI in patients scheduled to undergo arthroplasty was 0.69, just below the threshold of 0.7 of good internal consistency [31]. Regarding the validity of the GFI questionnaire the mental and physical domains of GFI were strongly to moderately correlated with the MCS of the SF12 (R = -0.59, P<0.001) and the PCS (R = -0.39, P<0.001), respectively, confirming the validity of the questionnaire. When performing cross-over analysis by correlating the mental domain of the GFI to the PCS of the SF-12 discriminatory validity was confirmed with a very weak correlation (R= -0.08; P<0.001).

In addition, the correlation of the physical domain of the GFI and MCS had a low correlation of $R = -0.28$ ($P < 0.001$). The correlation of the GFI with the HOOS/KOOS-pain score was, as hypothesized, low and also confirmed its discriminatory value to distinguish between pain and frailty ($R = -0.23$, $P < 0.001$).

Of the 2957 patients with end stage hip or knee OA who did complete the questionnaire, 853 (28.8%) were considered frail (a score of ≥ 4 on GFI). Patients with hip OA scored on average higher on the GFI (mean (SD) score: 2.78 (2.41) versus 2.28 (1.99)) and were more often considered frail as compared to persons with knee OA (33.3% versus 24.1%). Table 4 shows that frail persons were statistically significantly more often female, older and had a higher BMI as compared to those who are not frail. Also, frail persons scored statistically significantly lower on all scales of physical functioning of the HOOS/KOOS as well as on the physical and mental component scale of the SF-12 before arthroplasty surgery.

Table 4 – Comparison of demographic characteristics of frail and non-frail end stage OA patients

		Frailty as measured by GFI		
		Non frail N=2104	Frail N=853	P-value*
Joint	End stage hip OA	1018 (66.7%)	509 (33.3%)	< 0.001
	End stage knee OA	1086 (75.9%)	344 (24.1%)	
Female		1216 (58.4%)	620 (73.6%)	< 0.001
Age		67.07 ± 9.02	68.99 ± 9.97	< 0.001
BMI		28.07 ± 4.41	28.69 ± 5.14	0.002
Musculoskeletal comorbidities		351 (17.5%)	234 (29.4%)	< 0.001
Other comorbidities		1248 (68.1%)	594 (84.4%)	< 0.001
SF12	Physical	33.38 ± 9.52	29.33 ± 7.80	< 0.001
	Mental	58.33 ± 6.79	47.01 ± 11.06	< 0.001
Pain		40.56 ± 17.53	32.96 ± 18.45	< 0.001
Symptoms		43.05 ± 16.25	38.19 ± 16.34	< 0.001
HOOS / KOOS	Activities of daily life	45.36 ± 18.22	34.97 ± 18.51	< 0.001
	Sport	16.07 ± 17.54	10.64 ± 14.57	< 0.001
	Quality of Life	34.49 ± 10.79	31.06 ± 9.75	< 0.001

* Characteristics frail and non-frail patients were tested by means of a T-test (normal distribution), Mann-Whitney test (not normal distribution) or Chi-square test (discrete variables). A score of ≥ 4 was considered as frail.

Discussion

The GFI is a valid questionnaire to assess frailty in end stage hip or knee OA patients by means of a self-reported postal questionnaire. According to the GFI, using the cut-off of 4, about one-third of the patients undergoing THA and a quarter of the persons undergoing TKA are frail.

The feasibility of the use of the GFI within the current clinical setting for patients with end stage hip or knee OA is good, as 90% of the participants completed the questionnaire. In a study by Metzelthin et al. in older community dwelling persons showed that 77.4% of the persons completed the questionnaire.³²

Those who did not complete the questionnaire were more often male and older. The open question (question 15) was most often left empty, indicating that it is probably easier for patients to have closed questions with predefined answer options. Further research is needed to reconsider the format of this question aiming to obtain higher response rates.

Although the Cronbach's alpha of 0.69 is just below the threshold of good internal consistency of 0.7, it does indicate that the internal consistency of the GFI in our patient group is satisfactory and it is comparable to the alpha of 0.68 as found by Peters et al in home dwelling elderly in the Netherlands.^{20,31}

With respect to the convergent and discriminatory validity of the GFI for this specific patient group, the magnitude of the observed associations was in line with our hypotheses. Our convergent validity (range -0.6 -0.4) was comparable to the findings of Peters et al (range 0.4-0.61) [20]. The discriminatory validity in our patient group (range -0.08 - -0.3) was even stronger as compared to the elderly of Peters et al (range 0.08 - 0.5).²⁰

Significantly more patients with end stage hip OA were considered to be frail as compared to end stage knee OA (hip; 33%, knee; 24%, $P < 0.001$). However, both these numbers are lower as compared to the study of Peters *et al*²⁰ who found 60% of the independent living elderly in their study to be frail as measured by the GFI, but the average age in that study was 81 years, much higher than in the present study

(mean age 68 years). In a study among Romanian home-dwelling elderly (mean age 75), 75% of the participants were considered frail by the GFI.³³ These studies show that the presence of frailty shows wide variability depending on country, social status, diagnosis and age. The median and mean scores of the GFI in our patient group (2.00 and 2.54, respectively) were lower than the averages in independent living old persons found by Peters *et al*²⁰ (median 3) or reported by Metzehlhin *et al*²² and Drubbel *et al*²⁴ (means 3.8 and 3.2, respectively).

In both the latter studies the mean age was higher than in our study (77 and 73 years respectively). The lower frailty score in our patient groups can, apart from age, be explained by the fact that all patients were selected by an orthopaedic surgeon to receive arthroplasty surgery and were thus considered to be fit enough for major surgery.

The rates of persons with OA classified as being frail in our study are not easy to compare with other studies, as different methods to ascertain frailty were employed. Using Fried's Frailty Phenotype, Mandl *et al*²⁵ found that 8% of persons scheduled for knee arthroplasty were considered frail (although 17% reported difficulty with activities of daily life) with a similar rate found in men with hip osteoarthritis (8%) and in a study of persons with knee, hip or hand OA from 6 different European cohorts (10.2% considered frail).^{6,36,37}

A larger proportion, i.e. 22.4% of persons with hip or knee OA, was considered frail using Fried's Frailty Phenotype in a Brazilian study.³⁸ With the interpretation of these proportions it must be taken into account that the criteria of Fried's Frailty Phenotype are to be ascertained by a physician and do not include activities of daily life.⁶

Dent *et al*²⁹ have published an overview of the most commonly used frailty-questionnaires including, besides the GFI, three other self-reported frailty assessments: the Tilburg Frailty Index, the PRISMA-7 and the SPQ. However, none of these other three self-reported questionnaires have to our knowledge been used to assess the occurrence of frailty in persons with osteoarthritis.

Since a large proportion, about one third, of the patients scheduled to undergo major implant surgery are considered frail as scored by the self-reported GFI, the effects of frailty on their postoperative outcome should be assessed in future studies. This study has shown that the use of the GFI to discriminate between frail and non-frail total joint arthroplasty patients is appropriate.

Acknowledgements

We thank the patients and their relatives for their participation in the LOAS study, the students for their help in managing study logistics and the Reumafonds for the funding of this study (LLP13).

References

- Harris WH and Sledge CB (1990). Total hip and total knee replacement (1). *N Engl J Med*, 323(12): 801-7.
- Harris WH and Sledge CB (1990). Total hip and total knee replacement (2). *N Engl J Med*, 323(11): 725-31.
- Kurtz S, et al. (2007). Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am*, 89(4): 780-5.
- Landelijk Register Orthopedische Implantaten (LROI) (2015). Blik op Uitkomsten - Jaarrapportage LROI 2015.
- Santos-Eggimann B, et al. (2009). Prevalence of frailty in middle-aged and older community-dwelling Europeans living in 10 countries. *J Gerontol A Biol Sci Med Sci*, 64(6): 675-81.
- Fried LP, et al. (2001). Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*, 56(3): M146-56.
- de Vries NM, et al. (2011). Outcome instruments to measure frailty: a systematic review. *Ageing Res Rev*, 10(1): 104-14.
- Hamerman D. (1999). Toward an understanding of frailty. *Ann Intern Med*, 130(11): 945-50.
- Bales CW and Ritchie CS (2002). Sarcopenia, weight loss, and nutritional frailty in the elderly. *Annu Rev Nutr*, 22: 309-23.
- Levers MJ, et al. (2006). Factors contributing to frailty: literature review. *J Adv Nurs*, 56(3): 282-91.
- Rockwood K, et al. (1994). Frailty in elderly people: an evolving concept. *CMAJ*, 150(4): 489-95.
- Markle M and Browne G (2003). Conceptualizations of frailty in relation to older adults. *J Adv Nurs*, 44(1): 58-68.
- Gill TM, et al. (2006). Transitions between frailty states among community-living older persons. *Arch Intern Med*, 166(4): 418-23.
- Buchner DM and Wagner EH (1992). Preventing frail health. *Clin Geriatr Med*, 8(1): 1-17.
- Guralnik JM, et al. (2001). Progressive vs catastrophic loss of the ability to walk: implications for the prevention of mobility loss. *J Am Geriatr Soc*, 49(11): 1463-70.
- Fried LP, et al. (2004). Untangling the concepts of disability, frailty, and comorbidity: implications for improved targeting and care. *J Gerontol A Biol Sci Med Sci*, 59(3): 255-63.
- Fulop T, et al. (2010). Aging, frailty and age-related diseases. *Biogerontology*, 11(5): 547-63.
- Xue QL (2011). The frailty syndrome: definition and natural history. *Clin Geriatr Med*, 27(1): 1-15.
- Gobbens RJ, et al. (2012). Testing an integral conceptual model of frailty. *J Adv Nurs*, 68(9): 2047-60.
- Peters LL, et al. (2012). Measurement properties of the Groningen Frailty Indicator in home-dwelling and institutionalized elderly people. *J Am Med Dir Assoc*, 13(6): 546-51.
- Leichtenberg CS, et al. (2017). No associations between self-reported knee joint instability and radiographic features in knee osteoarthritis patients prior to Total Knee Arthroplasty: A cross-sectional analysis of the Longitudinal Leiden Orthopaedics Outcomes of Osteo-Arthritis study (LOAS) data. *Knee*, 24(4):816-823.
- World Medical Association (2013). Declaration of Helsinki - Ethical principles for medical research involving human subjects. *JAMA*, 310(20): 3.
- Peters LL, et al. (2015). Construct validity of the Groningen Frailty Indicator established in a large sample of home-dwelling elderly persons: Evidence of stability across age and gender. *Exp Gerontol*, 69: 129-41.
- Steverink N, et al. (2001). Measuring Frailty: Developing and testing the GFI (Groningen Frailty Indicator). *The Gerontologist*, 41(1): 236-237.
- Gandek B, et al. (1998). Cross-validation of item selection and scoring for the SF-12 Health Survey in nine countries: results from the IQOLA Project. International Quality of Life Assessment. *J Clin Epidemiol*, 51(11): 1171-8.
- Mols F, et al. (2009). Normative data of the SF-12 health survey with validation using postmyocardial infarction patients in the Dutch population. *Qual Life Res*, 18(4): 403-14.
- Nilsdotter AK, et al. (2003). Hip disability and osteoarthritis outcome score (HOOS)--validity and responsiveness in total hip replacement. *BMC Musculoskelet Disord*, 4: 10.
- Roos EM and Lohmander LS (2003). The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes*, 1:64.
- de Groot IB, et al. (2008). The Dutch version of the Knee Injury and Osteoarthritis Outcome Score: a validation study. *Health Qual Life Outcomes*, 6: 16.
- de Groot, IB, et al. (2007). Validation of the Dutch version of the Hip disability and Osteoarthritis Outcome Score. *Osteoarthritis Cartilage*, 15(1): 104-9.
- Field A. (2009). *Discovering Statistics using SPSS*. 3 ed. SAGE Publications Ltd.
- Metzelthin SF, et al. (2010). The psychometric properties of three self-report screening instruments for identifying frail older people in the community. *BMC Public Health*, 10: 176.
- Olariu M, et al. (2014). The psychometric qualities of the Groningen Frailty Indicator in Romanian community-dwelling old citizens. *Fam Pract*, 31(4): 490-5.
- Drubbel I, et al. (2013). Identifying frailty: do the Frailty Index and Groningen Frailty Indicator cover different clinical perspectives? a cross-sectional study. *BMC Fam Pract*, 14: 64.
- Mandl LA, et al. (2017). Is Frailty Associated with Adverse Events after Total Joint Arthroplasty? [abstract]. *Arthritis Rheumatol*, 69(10).
- Wise BL, et al. (2014). Frailty and hip osteoarthritis in men in the MrOS cohort. *J Gerontol A Biol Sci Med Sci*, 69(5): 602-8.
- Castell MV, et al. (2015). Osteoarthritis and frailty in elderly individuals across six European countries: results from the European Project on OsteoArthritis (EPOSA). *BMC Musculoskelet Disord*, 16: 359.
- Miguel C, et al. (2012). Frailty syndrome in the community-dwelling elderly with osteoarthritis. *Rev Bras Reumatol*, 52(3): 331-47.
- Dent E, et al. (2016). Frailty measurement in research and clinical practice: A review. *Eur J Intern Med*, 31: 3-10.