



Universiteit
Leiden
The Netherlands

The road to Insurmountability: Novel avenues to better target CC Chemokine Receptors

Ortiz Zacarías, N.V.

Citation

Ortiz Zacarías, N. V. (2019, December 4). *The road to Insurmountability: Novel avenues to better target CC Chemokine Receptors*. Retrieved from <https://hdl.handle.net/1887/81379>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/81379>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/81379> holds various files of this Leiden University dissertation.


Author: Ortiz Zacarías, N.V.

Title: The road to Insurmountability: Novel avenues to better target CC Chemokine Receptors

Issue Date: 2019-12-04


Chapter 3

Structure of CC Chemokine Receptor 2 with Orthosteric and Allosteric Antagonists



Yi Zheng, Ling Qin, Natalia V. Ortiz Zacarías, Henk de Vries, Gye Won Han, Martin Gustavsson, Marta Dabros, Chunxia Zhao, Robert Cherney, Percy Carter, Dean Stamos, Ruben Abagyan, Vadim Cherezov, Raymond C. Stevens, Adriaan P. IJzerman, Laura H. Heitman, Andrew Tebben, Irina Kufareva, Tracy M. Handel

Nature 2016, 540 (7633): 458-461



ABSTRACT

CC chemokine receptor 2 (CCR2) is one of 19 members of the chemokine receptor subfamily of human Class A G protein-coupled receptors. CCR2 is expressed on monocytes, immature dendritic cells and T cell subpopulations, and mediates their migration towards endogenous CC chemokine ligands such as CCL2.¹ CCR2 and its ligands are implicated in numerous inflammatory and neurodegenerative diseases² including atherosclerosis, multiple sclerosis, asthma, neuropathic pain, and diabetic nephropathy, as well as in cancer.³ These disease associations have motivated numerous preclinical studies and clinical trials⁴ (see ClinicalTrials.gov) in search of therapies that target the CCR2-chemokine axis. To aid drug discovery efforts,⁵ here we solve a structure of CCR2 in a ternary complex with an orthosteric (BMS-681⁶) and allosteric (CCR2-RA-[R]⁷) antagonist. BMS-681 inhibits chemokine binding by occupying the orthosteric pocket of the receptor in a previously unseen binding mode. CCR2-RA-[R] binds in a novel, highly druggable pocket that is the most intracellular allosteric site observed in Class A G protein-coupled receptors so far; this site spatially overlaps the G protein-binding site in homologous receptors. CCR2-RA-[R] inhibits CCR2 non-competitively by blocking activation-associated conformational changes and formation of the G protein-binding interface. The conformational signature of the conserved microswitch residues observed in double-antagonist-bound CCR2 resembles the most inactive G protein-coupled receptor structures solved so far. Like other protein-protein interactions, receptor-chemokine complexes are considered challenging therapeutic targets for small molecules, and the present structure suggests diverse pocket epitopes that can be exploited to overcome obstacles in drug design.

RESULTS AND DISCUSSION

A ternary complex between an engineered construct of human CCR2 isoform b (further referred to as CCR2-T4L or simply CCR2), an orthosteric antagonist BMS-681 (compound 13d in Carter *et al.*⁶), and an allosteric antagonist CCR2-RA-[R]⁷ was crystallized using the lipidic cubic phase (LCP) method,⁸ and the structure was determined to 2.8 Å resolution (Table S1 and Figure S1). Simultaneous addition of two compounds markedly stabilized detergent-solubilized CCR2-T4L compared with twice the concentration of each compound individually (Figure 1a), suggesting concurrent binding of CCR2-RA-[R] and BMS-681 to the receptor. The presence of both compounds was critical for crystallization.





Figure 1. Structure of a complex between CCR2, BMS-681 and CCR2-RA-[R] and comparison with other allosteric modulators of class A GPCRs. (a) Thermal denaturation curves demonstrate higher stability of CCR2-T4L in the presence of both BMS-681 and CCR2-RA-[R] compared with each compound individually. Data are representative of three independent experiments conducted on different days. (b) Overall view of double-antagonist-bound CCR2. (c, d) Structure viewed from the extracellular (c) and intracellular (d) side with simulated annealing omit maps of BMS-681 (c) and CCR2-RA-[R] (d) shown at 3σ. (e) CCR2-RA-[R] compared to other allosteric ligands crystallized with Class A GPCRs (PDB accession numbers 4MBS, 4XNV, 4PHU, and 4MQT). (f) CCR2-RA-[R] compared with the carboxy (C)-terminal helix of Gαs bound to the β₂-adrenergic receptor and transducin peptide bound to rhodopsin (PDB accession numbers 3SN6 and 4X1H).



In the structure, CCR2 adopts the canonical fold of class A G protein-coupled receptors (GPCRs) with seven transmembrane (TM) helices connected by three extracellular (EC) and three intracellular (IC) loops (Figure 1b). Both compounds are visible in the electron density (Figures 1b-d); BMS-681 binds in the extracellular orthosteric pocket (Figures 1b,c) while CCR2-RA-[R] is located more than 30 Å away (Figures 1b,d), in a site that is the most intracellular allosteric pocket observed in class A GPCRs so far (Figure 1e). The binding site of CCR2-RA-[R] spatially overlaps with the G protein-binding site in homologous receptors (Figure 1f). As for other chemokine receptors,⁹⁻¹² CCR2 is expected to have two conserved disulfide bonds in its extracellular domains, with Cys32-Cys277 connecting the amino (N) terminus to ECL3 (NT-ECL3), and Cys113-Cys190 connecting TM3 to ECL2. Electron density is apparent for the ECL2-TM3 disulfide bond but not for the N-terminal residues 1-36 or the NT-ECL3 disulfide bond (Figures 1b,c). Because the NT-ECL3 disulfide bond has been shown to be important for CCR2 signaling,¹³ its absence is unlikely to be an inherent feature of the receptor; instead, it might be caused by strain of the bond in the ligand-bound state of the receptor,¹⁴ possibly exacerbated by solvent exposure and radiation damage of the crystals.¹⁵

As with other chemokine receptors, the extracellular orthosteric pocket of CCR2 can be divided into a major and a minor subpocket, defined by helices III-VII and helices I-III and VII, respectively, and separated by residues Y120^{3,32} and E291^{7,39} (superscript indicates residue number according to Ballesteros-Weinstein nomenclature). BMS-681 binds predominantly in the minor subpocket (Figures 2a,b) and buries 366.3 Å² of surface area. The 6-trifluoromethyl quinazoline moiety protrudes between helices I and VII towards the lipid bilayer, while the tri-substituted cyclohexane packs against W98^{2,60}. The γ-lactam secondary exocyclic amine forms a hydrogen bond with the hydroxyl of T292^{7,40}, which is critical for binding of chemically related compounds such as BMS-558 (compound 22 in Cherney *et al.*¹⁶) and the Teijin lead series.^{17, 18} This amine is also within hydrogen-bonding distance from the backbone carbonyl of Q288^{7,36}. The carbonyl oxygen of the γ-lactam forms a hydrogen bond with Y49^{1,39}, which itself is hydrogen-bonded to the side chain of T292^{7,40}. The N1 nitrogen of the quinazoline is within 4 Å of the Q288^{7,36} side chain. The protonated tertiary amine on the cyclohexane ring is proximal to a structured water molecule in the binding site. Some CCR2 antagonists, particularly those containing a basic amine, are known to depend on the conserved E291^{7,39} in the receptor;¹⁹ however, no direct interaction is observed between E291^{7,39} and BMS-681. The receptor-bound, bioactive conformation of BMS-681 is strikingly similar to the crystallographic conformation of free BMS-681 (Figure 2c, Table S2), suggesting the absence of internal strain in the bound state.

BMS-681 engages several residues that are critical for CCR2 binding and/or activation of CCR2^{17, 18} including Y49^{1,39}, W98^{2,60}, Y120^{3,32}, and T292^{7,40}. Thus, it seems to directly compete with chemokine binding to the orthosteric pocket. Additionally, by inserting between

helices I and VII, BMS-681 may put strain onto residues C32-V37 connecting TM1 to ECL3, destabilize the conserved NT-ECL3 disulfide bond (absent in the structure), and prevent the N terminus and TM1 from adopting a productive chemokine binding conformation observed in homologous receptor-chemokine structures^{11, 12} (Figure S2).




Figure 2. Ligand binding sites and receptor interactions. (a, b) BMS-681 interactions with CCR2 viewed in three dimensions from the extracellular side (a) and in a two-dimensional schematic depiction (b). (c) Of the two conformers in the free BMS-681 structure, one is almost identical to the CCR2-complexed conformation. (d, e) CCR2-RA-[R] interactions viewed in three dimensions along the plane of the membrane and from the intracellular side (d), and in a two-dimensional depiction (e). In (b, e) polar and non-polar residue contacts are shown as blue and green, respectively. Bulk solvent and lipid are represented by blue and yellow shading, respectively.

On the opposite side of the receptor, CCR2-RA-[R] is caged by the intracellular ends of helices I-III and VI-VIII and buries 297.8 Å² of surface area. The inner hydrophobic part of the cage is made by V63^{1.53}, L67^{1.57}, L81^{2.43}, L134^{3.46}, A241^{6.33}, V244^{6.36}, I245^{6.37}, Y305^{7.53}, and F312^{8.50}, while the outer (cytosol-facing) polar part consists of T77^{2.39}, R138^{3.50}, G309^{5.47}, K311^{8.49}, and Y315^{8.53} (Figures 2d,e), as well as the backbones of engineered R237^{6.29} and K240^{6.32}. The binding pocket of CCR2-RA-[R] is highly enclosed and possesses a balanced combination of hydrophobic and polar features, all of which favors pocket ‘druggability’.⁵ Owing to the lack of a side-chain on G309^{8.47}, the hydroxyl and pyrrolone carbonyl groups of CCR2-RA-[R] can hydrogen-bond to the exposed backbone amides of E310^{8.48}, K311^{8.49}, and F312^{8.50} (Figures 2d,e). The acetyl group of the compound resides near the terminal amine of K311^{8.49}. The critical roles of V244^{6.36}, Y305^{7.53}, K311^{8.49}, and F312^{8.50} in CCR2-RA-[R] binding were established by an earlier mutagenesis study.²⁰ Because homologues of several residues in

the CCR2-RA-[R] binding pocket directly couple to the G protein in bovine rhodopsin²¹ and the β_2 adrenergic receptor (β_2 AR)²² structures (Figure S4), CCR2-RA-[R] appears to sterically interfere with G protein binding to CCR2.

The structure suggests an interesting symmetrical mechanism for the concurrent antagonistic action of the two compounds. BMS-681 interferes with chemokine binding directly and with G protein coupling indirectly, by stabilizing an inactive, presumably G protein-incompatible,⁶ conformation of the receptor. Conversely, CCR2-RA-[R] directly prevents G protein coupling and allosterically inhibits binding of the CCL2 chemokine,²³ which, like most GPCR agonists, requires an active, G protein-associated receptor for high affinity binding.²³ Bi-directional allosteric communication between the extra- and intracellular sides of the receptor is reminiscent of that previously observed in adenosine A_{2A} receptor (A_{2A} AR)²⁴ and β_2 AR²⁵ using allosteric inverse agonist antibodies/nanobodies that target the same epitope as CCR2-RA-[R]. Similar to these antibodies, CCR2-RA-[R] was previously shown to allosterically enhance, and to be allosterically enhanced by, binding of orthosteric antagonists,²³ demonstrating positive binding cooperativity.

We further characterized this cooperativity by studying the binding of BMS-681 to wild-type CCR2 and the crystallization construct CCR2-T4L using previously characterized radioactive probes [³H]-INCB3344 (orthosteric) and [³H]-CCR2-RA (allosteric).²³ In equilibrium competition binding assays on wild-type CCR2, both INCB3344 and CCR2-RA-[R] displaced their homologous radioligand with half-maximum inhibitory concentration (IC_{50}) values of 17 and 13 nM, respectively (Figures S4a,b and Table S3), comparable to previously reported values.²³ Compared to wild-type CCR2, the affinity of both antagonists towards CCR2-T4L was improved by approximately twofold, suggesting a slight engineering-related shift towards the inactive state. BMS-681 fully displaced [³H]-INCB3344 with nanomolar affinities for both constructs, but did not displace [³H]-CCR2-RA. Instead, at 1 μ M concentration it enhanced the binding of [³H]-CCR2-RA by >30% (Figures S4a,b and Table S3).

In kinetic radioligand experiments, the presence of BMS-681 also increased total binding of [³H]-CCR2-RA to both wild-type CCR2 and CCR2-T4L, with the increase as high as 62% in the case of CCR2-T4L (Figures S4c,d, and Table S4). BMS-681 (1 μ M) decreased the dissociation rate constant of [³H]-CCR2-RA, while producing a slight increase (wild-type CCR2) or no change (CCR2-T4L) in the observed association rate constants. Moreover, for CCR2-T4L, the presence of BMS-681 changed the biphasic dissociation profile of [³H]-CCR2-RA to monophasic, suggesting stabilization of the receptor population in a homogenous conformational state (Table S4). Along with the stability and equilibrium binding data, these results further corroborate the hypothesis that BMS-681 and CCR2-RA-[R] cooperatively stabilize a preferred inactive conformation of CCR2-T4L.




Figure 3. Crystallographic conformation of double-antagonist-bound CCR2 has pronounced structural signatures of an inactive state. Structures of active (US28, salmon), inactive (CCR5, grey), and more inactive (CCR2, blue) chemokine receptors viewed along the plane of the membrane (a) and across the membrane from the intracellular side (b-e). (b) Overlay of structures; arrows show the direction of activation-associated conformational changes; sticks show conserved Y^{5.58}, R^{3.50} and Y^{7.53}; the white mesh is CCR2-RA-[R]. (c-e) Detailed, single-receptor depictions of (b). (f) Although located ~30 Å apart, the orthosteric (BMS-681, magenta) and the allosteric (CCR2-RA-[R], yellow) ligands cooperate in stabilizing an inactive conformation of CCR2 through helix VII.

We next analyzed the structure of double-antagonist-bound CCR2-T4L to better understand this conformation. The plethora of existing class A GPCR structures suggests a conserved conformational signature of an active receptor state.²⁶ This signature involves increased separation between the intracellular end of helix VI and the rest of the TM bundle, an inward repositioning and rotation of helix VII, and concerted repacking of the highly conserved microswitches R^{3.50} (of the DR^{3.50}Y motif), Y^{5.58}, and Y^{7.53} (of the NPxxY^{7.53} motif) (Figures 3a,b) to form an intracellular binding interface for G protein. Furthermore, rather than adopting either an ‘on’ or ‘off’ state, receptors can occupy an ensemble of intermediate conformations.²⁷ The active state signature is fully represented in US28, the only agonist-bound chemokine receptor crystallized so far¹² (Figures 3a-c). By contrast, the double-antagonist-bound CCR2 structure appears to occupy the opposite end of the activation spectrum as it shares the conformational microswitch signatures of the most inactive GPCR structures observed thus far (Figures 3a-e).

As in the inactive CCR5-maraviroc complex,¹⁰ the intracellular ends of CCR2 helices III and VI are close together, and the conserved R^{3.50} interacts with D^{3.49} and T^{2.39}, effectively disrupting the G protein-binding pocket (Figures 3b,d,e). Similarly, in both CCR2 and CCR5 structures, the intracellular end of helix VII is in the inactive outward-facing conformation with Y^{7.53}

pointing towards helix II rather than the centre of the bundle. However, in CCR5, Y^{5.58} is oriented towards the centre of the bundle, whereas in the present CCR2 structure, it faces the lipid and is sterically blocked from approaching R^{3.50} and Y^{7.53} by F^{6.38} (Figures 3d,e). The net result of these interactions is that the crystallographically observed conformation of CCR2 appears to be even more inactive than that of CCR5 and most similar to dark rhodopsin²⁸ and Fab-bound β_2 AR.²⁹ Although receptor construct engineering appears to contribute to stabilization of this inactive state, the ligand binding and thermal denaturation data suggest that the concerted action of the two antagonists is also important. By directly interacting with the conserved activation microswitch residues, CCR2-RA-[R] is perfectly positioned to stabilize this inactive state: it sterically blocks Y^{7.53} from populating the active conformation and is propped against R^{3.50}, restricting its orientation away from the G protein interface (Figure 3b). Although located 30 Å away, BMS-681 appears to cooperate with CCR2-RA-[R] through their common interactions with helix VII, which moves outward on the intracellular side (opposite to its movement during activation) and inwards on the extracellular side (relative to CCR5 and US28) (Figure 3f).




Figure 4. Structural motifs exploited by small molecule antagonists of chemokine receptors. Receptor surface meshes are colored by polarity (cyan, polar; grey, nonpolar). (a) Modeled CCR2-CCL2 complex illustrates the extensive receptor-chemokine interface. (b-d) Structures of CCR2-BMS-681 (b), CCR5-maraviroc (PDB 4MBS) (c), and CXCR4-IT1t (PDB 3ODU) (d). Compounds utilize unique non-polar subpockets (yellow contours) within the open polar binding pockets of their target receptors. (e) The allosteric pocket possesses a balanced combination of hydrophobic and polar features, making it a promising target for drug development.

The CCR2 structure has general implications for the design of drugs targeting chemokine receptors as a family. As with most protein-protein interfaces, the orthosteric binding

pockets of chemokine receptors are large, wide open, and highly polar. Chemokines explore numerous hotspots within these pockets and their binding is additionally reinforced by the interaction with the flexible N termini of the receptors^{11, 12} (Figure 4a), collectively making for an extensive and versatile interaction that is conceptually difficult to inhibit with small molecules. The structure of CCR2 with BMS-681 and CCR2-RA-[R] extends the repertoire of ideas that can be used to overcome these obstacles. The binding mode of BMS-681 (Figure 4b) contrasts with both the binding mode of maraviroc to CCR5 (Figure 4c) where the ligand spans the major and the minor subpockets of the receptor, and that of IT1t to CXCR4 (Figure 4d) where the ligand is entirely accommodated in the minor subpocket. While occupying the minor subpocket of CCR2, BMS-681 protrudes between helices I and VII towards the lipid bilayer (Figure 4b) in an interaction facilitated by the trifluoromethyl group that is often present in CCR2 antagonists.³⁰ This interaction enables hydrophobic anchoring of BMS-681 to the otherwise polar and open binding site of CCR2; by doing so, it parallels the role of other unique non-polar subpockets exploited by crystallized small molecule antagonists of CCR5 and CXCR4 (Figures 4b-d). The novel subpocket explored by BMS-681 may have an additional advantage of disrupting the chemokine-compatible conformation of the receptor N terminus (Figure S2).

CCR2-RA-[R] demonstrates a previously unseen binding mode within an allosteric pocket on the intracellular side of CCR2. Although relatively small, this pocket has a desirable balance of polarity and hydrophobicity (Figures 2e, 4e). Homologous pockets may be present in other chemokine receptors, owing to a conserved G^{8.47}; in fact, compound binding in homologous regions has been indirectly demonstrated for CCR1 and CCR5, and directly for CCR4,³¹ CXCR1, and CXCR2.³² In most other receptors that have been crystallized thus far, the non-glycine residue at position 8.47 appears to both reduce the pocket volume and block access to the backbone amides of helix 8; consequently, the homologous pockets in these receptors may not be druggable although negative allosteric modulation with antibodies and nanobodies targeting the same region has been reported.^{24, 25} By simultaneously competing with G protein and blocking activation-related conformational changes, compound binding in the allosteric pocket seems a powerful way to antagonize the receptor. Therefore, for receptors where in which the allosteric pocket is druggable, targeting it with small molecules may open new avenues for GPCR drug discovery.

METHODS

Design and Expression of CCR2-T4L fusion constructs

The sequence of human CCR2 isoform b (Uniprot ID P41597-2) was engineered for crystallization by truncation of C-terminal residues 329-360 and by grafting T4 lysozyme (T4L) into the ICL3. In the process of construct optimization, the native CCR2 residues between L226^{5.62} and R240^{6.32} (L226^{5.62}-KTLRCRNEKKRH-R240^{6.32}) were removed and replaced with corresponding residues from the crystallized structure of M2 muscarinic acetylcholine receptor (PDB accession number 3UON, resulting amino acid sequence S226^{5.62}-RASKSRI-T4L-PPPSREK-K240^{6.32}). The presence of T4L in ICL3 is expected to prevent receptor activation; however the similar affinities of BMS-681 and CCR2-RA-[R] for both wild-type (WT) CCR2 and CCR2-T4L (Figures S4a,b and Table S3) suggest that the fusion construct is a good surrogate to WT CCR2 for understanding ligand recognition.

The CCR4-T4L coding sequence was cloned into a modified pFastBac1 vector (Invitrogen) with an HA signal sequence followed by a Flag tag at the N terminus and a PreScission protease site followed by a 10× His tag and another Flag tag at the C terminus. The receptor was expressed in *Spodoptera frugiperda* (Sf9) cells. High-titre recombinant baculovirus (>10⁹ viral particles/ml) was obtained using the Bac-to-Bac Baculovirus Expression System (Invitrogen) as previously described.¹¹ Sf9 cells at a cell density of (2–3) × 10⁶ cells/ml were infected with P1 virus at a multiplicity of infection of 5. Cells were harvested by centrifugation 48 h after infection and stored at –80 °C until use.

Purification of CCR2-T4L

Insect cell membranes were prepared by thawing frozen cell pellets in a hypotonic buffer containing 10 mM HEPES (pH 7.5), 10 mM MgCl₂, 20 mM KCl, and EDTA-free complete protease inhibitor cocktail tablets (Roche). Extensive washing of the raw membranes was performed by repeated douncing and centrifugation in the same hypotonic buffer (two or three times) and then in a high osmotic buffer containing 1.0 M NaCl, 10 mM HEPES (pH 7.5), 10 mM MgCl₂, 20 mM KCl, and EDTA-free complete protease inhibitor cocktail tablets (three or four times), thereby separating soluble and membrane associated proteins from integral transmembrane proteins. Stock solutions (40 mM) of BMS-681 and CCR2-RA-[R] were made in isopropanol. Washed membranes were resuspended into a buffer containing 50 μM BMS-681, 2 mg/ml iodoacetamide, and EDTA-free complete protease inhibitor cocktail tablets, and incubated at 4 °C for 1 h before solubilization. The membranes were then solubilized in 50 mM HEPES (pH 7.5), 400 mM NaCl, 1% (w/v) *n*-dodecyl-β-D-

maltopyranoside (DDM, Anatrace), 0.2% (w/v) cholesteryl hemisuccinate (CHS, Sigma) at 4 °C for 3 h. The supernatant was isolated by centrifugation at 50,000g for 30 min, and incubated in 20 mM HEPES (pH 7.5), 400 mM NaCl with TALON IMAC resin (Clontech) overnight at 4 °C. After binding, the resin was washed without addition of ligands with ten column volumes of Wash I Buffer (50 mM HEPES (pH 7.5), 400 mM NaCl, 10% (v/v) glycerol, 0.1% (w/v) DDM, 0.02% (w/v) CHS, 10 mM imidazole), followed by four column volumes of Wash II Buffer (50 mM HEPES (pH 7.5), 400 mM NaCl, 10% (v/v) glycerol, 0.02% (w/v) DDM, 0.01% (w/v) CHS, 50 mM imidazole). The protein was then eluted with three to four column volumes of Elution Buffer (50 mM HEPES (pH 7.5), 1 μ M BMS-681, 400 mM NaCl, 10% (v/v) glycerol, 0.02% (w/v) DDM, 0.01% (w/v) CHS, 250 mM imidazole). PD MiniTrap G-25 columns (GE Healthcare) were used to remove imidazole. The protein was then treated overnight with His-tagged PreScission protease to cleave the C-terminal His-tag and Flag-tag. PreScission protease and the cleaved C-terminal fragment were removed by binding to TALON IMAC resin for 2 h at 4 °C. The protein was collected as the TALON IMAC column flow-through. The protein was supplemented with 75 μ M each of BMS-681 and CCR2-RA-[R] before being concentrated to 30 mg/ml with a 100 kDa molecular mass cut-off Amicon centrifuge concentrator (Millipore). The estimated final compound concentrations were ~1–2 mM for both compounds.

Protein stability assays

The thermostability of CCR2-T4L was analyzed by a differential scanning fluorimetry assay adapted from previous publications³³ using a RotorGene Q 6-plex RT-PCR machine (Qiagen). Briefly, 1–5 μ g of protein was mixed with 3 μ M 7-diethylamino-3-(4'-maleimidylphenyl)-4-methylcoumarin (CPM) dye (2.5 mM stock in DMSO) in 25 mM HEPES pH 7.5, 400 mM NaCl, 0.02% DDM, 0.004% CHS, 10% glycerol, and indicated concentrations of compounds to a final volume of 20 μ l; samples were incubated for 5 min at room temperature and then heated gradually from 28 °C to 90 °C at a rate of 0.8 °C/min, with CPM fluorescence (excitation 365 nm, emission 460 nm) recorded every 1 °C. The melting temperature (T_m) was determined from the first derivative of the denaturation curve, using the Rotor-Gene Q – Pure Detection software (version 2.0.3).

Crystallization

Purified CCR2 in complex with BMS-681 and CCR2-RA-[R] was reconstituted into LCP by mixing with molten lipid using a mechanical syringe mixer.⁸ The protein–LCP mixture contained 40% (w/w) receptor solution, 54% (w/w) monoolein, and 6% (w/w) cholesterol.

Crystallization trials were performed in 96-well glass sandwich plates (Hampton research) using a Mosquito LCP robot (TTP Labtech) by dispensing 45 nl of protein-laden LCP and 800 nl of precipitant solution per well. Plates were incubated and imaged at 20 °C. Initial crystal hits were found from a precipitant condition containing 100 mM MES, pH 6.5, 30% (v/v) PEG400, 100 mM Li₂SO₄. After optimization, diffraction-quality crystals were obtained from 100 mM MES, pH 6.5, 30–32% (v/v) PEG400, 75–85 mM Li₂SO₄. Crystals usually grew to a maximum size of 60 µm × 10 µm × 10 µm in 1 week, and were harvested directly from the LCP matrix using MiTeGen micromounts and flash cooled in liquid nitrogen.

Data collection and structure determination

X-ray diffraction data were collected using a 10 µm collimated minibeam at a wavelength of 1.0332 Å with a Pilatus3 6M direct detector on the 23ID-D beamline (GM/CA CAT) of the Advanced Photon Source at the Argonne National Laboratory. Crystals were located and aligned by the rastering strategy.³⁴ Among the several hundred crystal samples screened, most crystals diffracted to 2.8–3.5 Å resolution when exposed to 0.3 s of unattenuated beam using 0.3° oscillations. A 93.1% complete data set at 2.80 Å resolution was obtained by merging data from 17 crystals, using XDS³⁵ and Aimless.³⁶ As the data showed anisotropy, the UCLA Diffraction Anisotropy Server (<http://services.mbi.ucla.edu/anisotryscale/>) was used to truncate the data to 3.0 Å along both *a** and *b** axes, and to 2.81 Å along the *c** axis. Initial phase information was obtained by molecular replacement with the program Phaser³⁷ using the receptor portion of the CCR5 structure (PDB accession number 4MBS) converted to polyalanines, and the T4L portion of the CXCR4 structure (PDB accession number 3ODU) as search models. The correct molecular replacement solution (translation function Z-score = 14.8) contained one CCR2-T4L molecule in the asymmetric unit. Refinement was performed with Phenix³⁸ followed by manual examination and rebuilding of the refined coordinates in the program COOT³⁹ using both $|2F_o| - |F_c|$ and $|F_o| - |F_c|$ maps, as well as omit maps. The final model included 295 residues (37–225 and 241–319) of the 360 residues of CCR2 and residues 2–161 of T4L plus 16 residues of two 8-residue linkers. The remaining N- and C-terminal residues were disordered and were not built. Strong electron density for one metal ion was observed. The identity of the ion was determined to be Zn²⁺ by X-ray fluorescence scans (Figure S5). The zinc ion is coordinated by a water molecule as well as side chains of H144^{3,56}, E238, and E1005. Data collection and refinement statistics are shown in Table S1.


Crystallization and structure determination of BMS-681

BMS-681 was dissolved in a minimal amount of CH_3CN and then 15% water was added. After standing overnight, the resulting crystals were collected. Data were obtained on a Bruker-AXS X8-Proteum Kappa goniometer and APEXII detector. Intensities were measured using Cu K α radiation ($\lambda = 1.5418 \text{ \AA}$) with the crystal kept at a constant temperature using an Oxford cryo system during data collection. Indexing and processing of the measured intensity data were performed with the SAINT-APEX2 (Bruker-AXS) program suite, structure solution with SHELXS-97, and structure refinement with SHELXL-97.

The derived atomic parameters (coordinates and temperature factors) were refined through full matrix least-squares. The function minimized in the refinements was $\sum_w (|F_o| - |F_c|)^2$. R is defined as $\sum ||F_o| - |F_c|| / \sum |F_o|$ while $R_w = [\sum_w (|F_o| - |F_c|)^2 / \sum_w |F_o|^2]^{1/2}$ where w is an appropriate weighting function based on errors in the observed intensities. Hydrogens were introduced in idealized positions with isotropic temperature factors, but no hydrogen parameters were varied. It should be noted that the refinement model illustrates disorder and partial occupancy factors of 'guest' solvent/water molecules within the crystalline lattice. The atomic positions of these disordered molecules were taken from the difference map analysis, which showed peaks of electron density of varying intensities at the refined positions representing the disordered solvent/water molecules. Data collection and refinement statistics are shown in Table S2.

Cell culture and transfections

Chinese hamster ovary (CHO) cells (provided by H. den Dulk, Leiden University, The Netherlands; originally obtained from and certified by American Type Culture Collection) were cultured in Dulbecco's Modified Eagle Medium/F-12 Nutrient Mixture (DMEM/F-12) supplemented with 10% (v/v) newborn calf serum, 50 IU/ml penicillin, and 50 $\mu\text{g}/\text{ml}$ streptomycin; they were maintained at 37°C and in 5% CO_2 . Cells were subcultured twice a week at a ratio of 1:30 to 1:50 by trypsinization. Transient transfection of CHO cells with WT CCR2 and CCR2-T4L constructs was performed using a polyethylenimine method, as described previously.²³ Briefly, CHO cells were grown on plates (diameter 15 cm) to around 50% confluence and then transfected with a DNA/polyethylenimine mixture containing 10 μg plasmid DNA—previously diluted in 150 mM NaCl solution—mixed with polyethylenimine solution (1 mg/ml) at a 1:6 DNA:polyethylenimine mass ratio. Before adding 1 ml of the transfection mixture to each plate, the culture medium of the cells was refreshed and the mixture incubated for 20 min at room temperature. Following transfection, cells were incubated for 48 h at 37°C and 5% CO_2 before membrane preparation. Twenty-four hours



after transfection, sodium butyrate was added to each plate at a final concentration of 3 mM to increase receptor expression. CHO cells were tested for mycoplasma contamination before use, the outcome of which was negative.

Membrane preparation

Membranes from CHO cells transiently expressing the WT CCR2 or CCR2-T4L were prepared as described previously.²³ Briefly, cells were detached from plates (diameter 15 cm) using 5 ml of phosphate-buffered saline and centrifuged for 5 min at 3000g. The membranes were separated from the cytosolic fractions by several centrifugation and homogenization steps. First, the pellets were resuspended and homogenized in ice-cold membrane buffer (50 mM Tris-HCl buffer, supplemented with 5 mM MgCl₂, pH 7.4) using an Ultra Thurrax Homogenizer (IKA-Werke, Staufen, Germany). Homogenized membranes were then centrifuged in an Optima LE-80 K ultracentrifuge (Beckman Coulter, Fullerton, California, USA) at 31,000g for 20 min at 4 °C. The final membrane pellet was resuspended also in ice-cold membrane buffer and aliquoted before storage. Membrane aliquots were stored at -80 °C and protein concentrations were measured using a standard BCA protein determination assay (Pierce Chemical Company, Rockford, Illinois, USA).

Radioligand binding assays

[³H]-INCB3344 (specific activity 32 Ci mmol⁻¹) and [³H]-CCR2-RA (specific activity 63 Ci mmol⁻¹) were custom-labeled by Vitrox (Placentia, California, USA). JNJ-27141491 was synthesized as described previously.⁴⁰ INCB3344 and CCR2-RA-[R] were synthesized in-house as described previously.^{7, 41}


All radioligand binding assays were performed at 25 °C in a 100 µl reaction volume containing assay buffer (50 mM Tris-HCl buffer (pH 7.4), 5 mM MgCl₂, 0.1% CHAPS) and 30 µg of membrane protein from CHO cells transiently expressing WT CCR2 or CCR2-T4L. For competition binding assays with [³H]-INCB3344, a concentration of 5 nM [³H]-INCB3344 was used, and non-specific binding was determined with 10 µM of unlabelled INCB3344. In the case of [³H]-CCR2-RA competition binding assays, a radioligand concentration of 3 nM was used and non-specific binding was determined with 10 µM of JNJ-27141491. In all cases, homologous or competition displacement assays were performed using six increasing concentrations of competing ligands. Kinetic experiments were also performed at 25 °C using 7 nM [³H]-CCR2-RA and 30 µg of membrane protein in a 100 µl reaction volume. For association experiments, CHO-CCR2 or CHO-CCR2-T4L membranes were added to the reaction at eight different time points, in the absence or presence of 1 µM BMS-681. For

dissociation experiments, membranes were first incubated with radioligand for 90 min; dissociation was then initiated by addition of 10 μ M of CCR2-RA-[R] at 12 different time points, in the presence or absence of 1 μ M BMS-681. More time points were used in the dissociation assays, to characterize the biphasic profile of [3 H]-CCR2-RA dissociation. In all cases, total radioligand binding did not exceed 10% of the total radioligand added to avoid ligand depletion. For all experiments, incubation was terminated by dilution with ice-cold wash buffer (50 mM Tris-HCl buffer (pH 7.4), 5 mM MgCl_2 , 0.05% CHAPS). Separation of bound from free radioligand was achieved by rapid filtration over a 96-well GF/B filter plate using a Perkin Elmer Filtermate-harvester (Perkin Elmer, Groningen, The Netherlands) and filter-bound radioactivity was determined in a Perkin Elmer 2450 Microbeta2 plate counter after addition of 25 μ l Microscint scintillation cocktail per well (Perkin-Elmer, Groningen, The Netherlands).

Statistical Methods

No statistical methods were used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.

All radioligand binding data were analyzed using Prism 6.0 and 7.0 (GraphPad Software, San Diego, California, USA). The pIC_{50} values were obtained by nonlinear regression analysis of competition displacement assays. Apparent association rate constants (k_{obs}) and maximum binding (B_{max} , used to calculate $\%B/B_{\text{control}}$) were determined by fitting the association data to a one-phase exponential association function. Dissociation rate constants were determined by fitting the dissociation data to a monophasic (k_{off}) or biphasic ($k_{\text{off, fast}}$ and $k_{\text{off, slow}}$) exponential decay model. All data shown represent means $\pm$ s.e.m. of at least three independent experiments performed in duplicate. An unpaired, two-tailed Student's *t*-test was used to compare differences in pIC_{50} as well as differences in kinetic parameters. Differences in binding enhancement (%Binding) in the absence (set at 100%) or presence of BMS-681 were analyzed using a one-way analysis of variance with Dunnett's post-hoc test. Significant differences are denoted as follows: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.



Data availability

The atomic coordinates and structure factors for the CCR2–BMS-681–CCR2-RA-[*R*] complex have been deposited in the Protein Data Bank under accession number 5T1A. The structure of free BMS-681 is deposited in the Cambridge Crystallographic Data Centre (<http://www.ccdc.cam.ac.uk/>) under accession number 1479580. All other data are available from the corresponding authors upon reasonable request.

REFERENCES

- Scholten, D. J.; Canals, M.; Maussang, D.; Roumen, L.; Smit, M. J.; Wijtmans, M.; de Graaf, C.; Vischer, H. F.; Leurs, R. Pharmacological modulation of chemokine receptor function. *Br J Pharmacol* **2012**, *165*, 1617-1643.
- O'Connor, T.; Borsig, L.; Heikenwalder, M. CCL2-CCR2 Signaling in Disease Pathogenesis. *Endocr Metab Immune Disord Drug Targets* **2015**, *15*, 105-118.
- Lim, S. Y.; Yuzhalin, A. E.; Gordon-Weeks, A. N.; Muschel, R. J. Targeting the CCL2-CCR2 signaling axis in cancer metastasis. *Oncotarget* **2016**, *7*, 28697-28710.
- Solari, R.; Pease, J. E.; Begg, M. "Chemokine receptors as therapeutic targets: Why aren't there more drugs?". *Eur. J. Pharmacol.* **2015**, *746*, 363-367.
- Cooke, R. M.; Brown, A. J.; Marshall, F. H.; Mason, J. S. Structures of G protein-coupled receptors reveal new opportunities for drug discovery. *Drug Discov Today* **2015**, *20*, 1355-1364.
- Carter, P. H.; Brown, G. D.; Cherney, R. J.; Batt, D. G.; Chen, J.; Clark, C. M.; Cvijic, M. E.; Duncia, J. V.; Ko, S. S.; Mandlekar, S.; Mo, R.; Nelson, D. J.; Pang, J.; Rose, A. V.; Santella, J. B., 3rd; Tebben, A. J.; Traeger, S. C.; Xu, S.; Zhao, Q.; Barrish, J. C. Discovery of a Potent and Orally Bioavailable Dual Antagonist of CC Chemokine Receptors 2 and 5. *ACS Med Chem Lett* **2015**, *6*, 439-444.
- Dasse, O.; Evans, J.; Zhai, H.-X.; Zou, D.; Kintigh, J.; Chan, F.; Hamilton, K.; Hill, E.; Eckman, J.; Higgins, P. Novel, acidic CCR2 receptor antagonists: lead optimization. *Lett. Drug. Des. Discov.* **2007**, *4*, 263-271.
- Caffrey, M.; Cherezov, V. Crystallizing membrane proteins using lipidic mesophases. *Nat Protoc* **2009**, *4*, 706-731.
- Wu, B.; Chien, E. Y. T.; Mol, C. D.; Fenalti, G.; Liu, W.; Katritch, V.; Abagyan, R.; Brooun, A.; Wells, P.; Bi, F. C.; Hamel, D. J.; Kuhn, P.; Handel, T. M.; Cherezov, V.; Stevens, R. C. Structures of the CXCR4 chemokine GPCR with small-molecule and cyclic peptide antagonists. *Science* **2010**, *330*, 1066-1071.
- Tan, Q.; Zhu, Y.; Li, J.; Chen, Z.; Han, G. W.; Kufareva, I.; Li, T.; Ma, L.; Fenalti, G.; Li, J. Structure of the CCR5 chemokine receptor-HIV entry inhibitor maraviroc complex. *Science* **2013**, *341*, 1387-1390.
- Qin, L.; Kufareva, I.; Holden, L. G.; Wang, C.; Zheng, Y.; Zhao, C.; Fenalti, G.; Wu, H.; Han, G. W.; Cherezov, V.; Abagyan, R.; Stevens, R. C.; Handel, T. M. Crystal structure of the chemokine receptor CXCR4 in complex with a viral chemokine. *Science* **2015**, *347*, 1117-1122.
- Burg, J. S.; Ingram, J. R.; Venkatakrishnan, A. J.; Jude, K. M.; Dukkupati, A.; Feinberg, E. N.; Angelini, A.; Waghray, D.; Dror, R. O.; Ploegh, H. L.; Garcia, K. C. Structural basis for chemokine recognition and activation of a viral G protein-coupled receptor. *Science* **2015**, *347*, 1113-1117.
- Montecarlo, F. S.; Charo, I. F. The amino-terminal domain of CCR2 is both necessary and sufficient for high affinity binding of monocyte chemoattractant protein 1. Receptor activation by a pseudo-tethered ligand. *J Biol Chem* **1997**, *272*, 23186-23190.
- Zhang, K.; Zhang, J.; Gao, Z. G.; Zhang, D.; Zhu, L.; Han, G. W.; Moss, S. M.; Paoletta, S.; Kiselev, E.; Lu, W.; Fenalti, G.; Zhang, W.; Müller, C. E.; Yang, H.; Jiang, H.; Cherezov, V.; Katritch, V.; Jacobson, K. A.; Stevens, R. C.; Wu, B.; Zhao, Q. Structure of the human P2Y12 receptor in complex with an antithrombotic drug. *Nature* **2014**, *509*, 115-118.
- Weik, M.; Ravelli, R. B.; Kryger, G.; McSweeney, S.; Raves, M. L.; Harel, M.; Gros, P.; Silman, I.; Kroon, J.; Sussman, J. L. Specific chemical and structural damage to proteins produced by synchrotron radiation. *Proc Natl Acad Sci U S A* **2000**, *97*, 623-628.
- Cherney, R. J.; Mo, R.; Meyer, D. T.; Nelson, D. J.; Lo, Y. C.; Yang, G.; Scherle, P. A.; Mandlekar, S.; Wasserman, Z. R.; Jezak, H.; Solomon, K. A.; Tebben, A. J.; Carter, P. H.; Decicco, C. P. Discovery of disubstituted cyclohexanes as a new class of CC chemokine receptor 2 antagonists. *J Med Chem* **2008**, *51*, 721-724.
- Berkhout, T. A.; Blaney, F. E.; Bridges, A. M.; Cooper, D. G.; Forbes, I. T.; Gribble, A. D.; Groot, P. H.; Hardy, A.; Ife, R. J.; Kaur, R.; Moores, K. E.; Shillito, H.; Willetts, J.; Witherington, J. CCR2: characterization of the antagonist binding site from a combined receptor modeling/mutagenesis approach. *J Med Chem* **2003**, *46*, 4070-4086.

18. Hall, S. E.; Mao, A.; Nicolaidou, V.; Finelli, M.; Wise, E. L.; Nedjai, B.; Kanjanapangka, J.; Harirchian, P.; Chen, D.; Selchau, V.; Ribeiro, S.; Schyler, S.; Pease, J. E.; Horuk, R.; Vaidehi, N. Elucidation of binding sites of dual antagonists in the human chemokine receptors CCR2 and CCR5. *Mol Pharmacol* **2009**, *75*, 1325-1336.
19. Cherney, R. J.; Nelson, D. J.; Lo, Y. C.; Yang, G.; Scherle, P. A.; Jezak, H.; Solomon, K. A.; Carter, P. H.; Decicco, C. P. Synthesis and evaluation of cis-3,4-disubstituted piperidines as potent CC chemokine receptor 2 (CCR2) antagonists. *Bioorg Med Chem Lett* **2008**, *18*, 5063-5065.
20. Zweemer, A. J.; Bunnik, J.; Veenhuizen, M.; Miraglia, F.; Lenselink, E. B.; Vilums, M.; de Vries, H.; Gibert, A.; Thiele, S.; Rosenkilde, M. M.; IJzerman, A. P.; Heitman, L. H. Discovery and mapping of an intracellular antagonist binding site at the chemokine receptor CCR2. *Mol. Pharmacol.* **2014**, *86*, 358-368.
21. Blankenship, E.; Vahedi-Faridi, A.; Lodowski, D. T. The high-resolution structure of activated opsin reveals a conserved solvent network in the transmembrane region essential for activation. *Structure* **2015**, *23*, 2358-2364.
22. Rasmussen, S. G.; DeVree, B. T.; Zou, Y.; Kruse, A. C.; Chung, K. Y.; Kobilka, T. S.; Thian, F. S.; Chae, P. S.; Pardon, E.; Calinski, D.; Mathiesen, J. M.; Shah, S. T.; Lyons, J. A.; Caffrey, M.; Gellman, S. H.; Steyaert, J.; Skiniotis, G.; Weis, W. I.; Sunahara, R. K.; Kobilka, B. K. Crystal structure of the β_2 adrenergic receptor-Gs protein complex. *Nature* **2011**, *477*, 549-555.
23. Zweemer, A. J.; Nederpelt, I.; Vrieling, H.; Hafith, S.; Doornbos, M. L.; de Vries, H.; Abt, J.; Gross, R.; Stamos, D.; Saunders, J.; Smit, M. J.; IJzerman, A. P.; Heitman, L. H. Multiple binding sites for small-molecule antagonists at the CC chemokine receptor 2. *Mol. Pharmacol.* **2013**, *84*, 551-561.
24. Hino, T.; Arakawa, T.; Iwanari, H.; Yurugi-Kobayashi, T.; Ikeda-Suno, C.; Nakada-Nakura, Y.; Kusano-Arai, O.; Weyand, S.; Shimamura, T.; Nomura, N.; Cameron, A. D.; Kobayashi, T.; Hamakubo, T.; Iwata, S.; Murata, T. G-protein-coupled receptor inactivation by an allosteric inverse-agonist antibody. *Nature* **2012**, *482*, 237-240.
25. Staus, D. P.; Strachan, R. T.; Manglik, A.; Pani, B.; Kahsai, A. W.; Kim, T. H.; Wingler, L. M.; Ahn, S.; Chatterjee, A.; Masoudi, A.; Kruse, A. C.; Pardon, E.; Steyaert, J.; Weis, W. I.; Prosser, R. S.; Kobilka, B. K.; Costa, T.; Lefkowitz, R. J. Allosteric nanobodies reveal the dynamic range and diverse mechanisms of G-protein-coupled receptor activation. *Nature* **2016**, *535*, 448-452.
26. Katritch, V.; Cherezov, V.; Stevens, R. C. Structure-function of the G protein-coupled receptor superfamily. *Annu Rev Pharmacol Toxicol.* **2013**, *53*, 531-556.
27. Manglik, A.; Kim, T. H.; Masureel, M.; Altenbach, C.; Yang, Z.; Hilger, D.; Lerch, M. T.; Kobilka, T. S.; Thian, F. S.; Hubbell, W. L.; Prosser, R. S.; Kobilka, B. K. Structural insights into the dynamic process of β_2 -adrenergic receptor signaling. *Cell* **2015**, *161*, 1101-1111.
28. Palczewski, K.; Kumasaka, T.; Hori, T.; Behnke, C. A.; Motoshima, H.; Fox, B. A.; Le Trong, I.; Teller, D. C.; Okada, T.; Stenkamp, R. E.; Yamamoto, M.; Miyano, M. Crystal structure of rhodopsin: a G protein-coupled receptor. *Science* **2000**, *289*, 739-745.
29. Rasmussen, S. G.; Choi, H. J.; Rosenbaum, D. M.; Kobilka, T. S.; Thian, F. S.; Edwards, P. C.; Burghammer, M.; Ratnala, V. R.; Sanishvili, R.; Fischetti, R. F.; Schertler, G. F.; Weis, W. I.; Kobilka, B. K. Crystal structure of the human β_2 adrenergic G-protein-coupled receptor. *Nature* **2007**, *450*, 383-387.
30. Pease, J.; Horuk, R. Chemokine receptor antagonists. *J. Med. Chem.* **2012**, *55*, 9363-9392.
31. Andrews, G.; Jones, C.; Wreggett, K. A. An intracellular allosteric site for a specific class of antagonists of the CC chemokine G protein-coupled receptors CCR4 and CCR5. *Mol. Pharmacol.* **2008**, *73*, 855-867.
32. Nicholls, D. J.; Tomkinson, N. P.; Wiley, K. E.; Brammall, A.; Bowers, L.; Grahames, C.; Gaw, A.; Meghani, P.; Shelton, P.; Wright, T. J.; Mallinder, P. R. Identification of a putative intracellular allosteric antagonist binding-site in the CXC chemokine receptors 1 and 2. *Mol. Pharmacol.* **2008**, *74*, 1193-1202.
33. Alexandrov, A. I.; Mileni, M.; Chien, E. Y. T.; Hanson, M. A.; Stevens, R. C. Microscale fluorescent thermal stability assay for membrane proteins. *Structure* **2008**, *16*, 351-359.
34. Cherezov, V.; Hanson, M. A.; Griffith, M. T.; Hilgart, M. C.; Sanishvili, R.; Nagarajan, V.; Stepanov, S.; Fischetti, R. F.; Kuhn, P.; Stevens, R. C. Rastering strategy for screening and centring of microcrystal samples of human membrane proteins with a sub-10 μm size X-ray synchrotron beam. *J R Soc Interface* **2009**, *6 Suppl 5*, S587-S597.

35. Kabsch, W. Xds. *Acta Crystallogr D Biol Crystallogr* **2010**, *66*, 125-132.
36. Winn, M. D.; Ballard, C. C.; Cowtan, K. D.; Dodson, E. J.; Emsley, P.; Evans, P. R.; Keegan, R. M.; Krissinel, E. B.; Leslie, A. G.; McCoy, A.; McNicholas, S. J.; Murshudov, G. N.; Pannu, N. S.; Pottterton, E. A.; Powell, H. R.; Read, R. J.; Vagin, A.; Wilson, K. S. Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol Crystallogr* **2011**, *67*, 235-242.
37. McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R. J. Phaser crystallographic software. *J Appl Crystallogr* **2007**, *40*, 658-674.
38. Adams, P. D.; Afonine, P. V.; Bunkoczi, G.; Chen, V. B.; Davis, I. W.; Echols, N.; Headd, J. J.; Hung, L. W.; Kapral, G. J.; Grosse-Kunstleve, R. W.; McCoy, A. J.; Moriarty, N. W.; Oeffner, R.; Read, R. J.; Richardson, D. C.; Richardson, J. S.; Terwilliger, T. C.; Zwart, P. H. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **2010**, *66*, 213-221.
39. Emsley, P.; Lohkamp, B.; Scott, W. G.; Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **2010**, *66*, 486-501.
40. Doyon, J.; Coesemans, E.; Boeckx, S.; Buntinx, M.; Hermans, B.; Van Wauwe, J. P.; Gilissen, R. A.; De Groot, A. H.; Corens, D.; Van Lommen, G. Discovery of potent, orally bioavailable small-molecule inhibitors of the human CCR2 receptor. *ChemMedChem* **2008**, *3*, 660-669.
41. Brodmerkel, C. M.; Huber, R.; Covington, M.; Diamond, S.; Hall, L.; Collins, R.; Leffert, L.; Gallagher, K.; Feldman, P.; Collier, P.; Stow, M.; Gu, X.; Baribaud, F.; Shin, N.; Thomas, B.; Burn, T.; Hollis, G.; Yeleswaram, S.; Solomon, K.; Friedman, S.; Wang, A.; Xue, C. B.; Newton, R. C.; Scherle, P.; Vaddi, K. Discovery and pharmacological characterization of a novel rodent-active CCR2 antagonist, INCB3344. *J Immunol* **2005**, *175*, 5370-5378.
42. Chen, V. B.; Arendall, W. B., 3rd; Headd, J. J.; Keedy, D. A.; Immormino, R. M.; Kapral, G. J.; Murray, L. W.; Richardson, J. S.; Richardson, D. C. MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr D Biol Crystallogr* **2010**, *66*, 12-21.

SUPPORTING INFORMATION




Figure S1. CCR2-T4L crystals and crystal packing. (a-c) Crystal packing of CCR2-T4L. CCR2 is a blue ribbon with ECL2 colored red and T4L yellow. The unit cell is shown as a green box. CCR2-T4L molecules are arranged in a type I packing with hydrophilic stacking mediated by T4L and T4L-ECL2 interactions along axis c. (a) Crystal packing in the ac plane. CCR2 makes abundant hydrophobic contacts with its neighbor via an interface mediated by antiparallel helix IV–helix VI interactions related by a screw axis along axis a. (b) Crystal packing in the bc plane. Contacts between receptors and T4L involve ECL2 and the intracellular surface of CCR2 including helix VIII. Direct contacts between T4L are along axis b. One layer of CCR2-T4L molecules at the very top of the stacking column is omitted for clarity. (c) Crystal packing in the ab plane. There are no direct interactions between T4L along axis a. (d) Crystals of CCR2-T4L in the LCP bolus. Average crystals grew to 60 μm × 10 μm × 10 μm before harvesting.




Figure S2. BMS-681 binding may disrupt a chemokine-recognizing conformation of the CCR2 N terminus and helix I. (a) Model of CCR2-CCL2 built by homology from the structure of CXCR4-vMIP-II¹¹ suggests that a productive chemokine-compatible conformation of the receptor requires re-orientation of the N terminus from almost parallel to almost perpendicular to the membrane plane, and formation of an extra helical turn in helix I to bring it closer to helix VII and ECL3. (b) Binding of BMS-681 may disrupt this chemokine-compatible conformation by inserting between helices I and VII.

	TM1-ICL1-TM2								TM3-ICL2								TM5-ICL3-TM6								TM7-H8										
CCR2 residue	63	66	67	71	72	75	77	81	134	138	141	142	144	145	148	149	150	225	228	229	236	237	240	241	244	245	248	305	308	309	310	311	312	315	
CCR2-RA-[R] contacts	•		•					•	•														•	•	•	•	•	•	•	•	•	•	•	•	
CCR2	V	I	L	K	K	C	T	D	I	L	R	A	I	H	A	A	L	K	I	T	L	K	K	R	A	V	I	I	Y	V	G	E	K	F	Y
bRho	T	V	T	.	.	T	L	N	.	.	.	V	V	K	P	N	F	R	L	.	V	.	A	E	V	M	V	M	.	M	N	K	Q	.	C
β ₂ AR	A	I	E	R	.	N	I	.	.	I	.	.	S	P	Y	Q	S	.	V	E	A	L	.	K	.	T	L	.	R	S	P	D	.	A	
Gat-CT contacts (bRho)							•			•	•	•	•					•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
Gas-CT contacts (β ₂ AR)																•																			

Figure S3. CCR2-RA-[R] directly binds to CCR2 residues that are homologous to those involved in G protein coupling in other GPCRs. Partial alignment of intracellular regions of CCR2 and homologous regions in bovine Rho (bRho) and β_2 adrenergic receptor (β_2 AR), alongside profile of contacts that CCR2-RA-[R], the $G\alpha_t$ C-terminal peptide,²¹ and $G\alpha_s$ C terminus²² make with the three respective receptors. Contacts are shown by circles above and below the alignment, with circle area indicative of contact strength. Backbone and side-chain contacts are gray and black, respectively. Assuming structural homology between the CCR2-G protein interface and at least one of the bRho- $G\alpha_t$ and β_2 AR- $G\alpha_s$ interfaces, several residue positions seem to be involved in binding both CCR2-RA-[R] and the C terminus of the G protein.




Figure S4. Equilibrium binding and binding kinetics of BMS-681 and CCR2-RA-[R] with WT CCR2 and CCR2-T4L. (a, b) Displacement of [^3H]-INCB3344 (5 nM, a) and [^3H]-CCR2-RA (3 nM, b) from WT CCR2 and CCR2-T4L in CHO cells by increasing concentrations of unlabelled INCB3344, CCR2-RA-[R] and BMS-681. (c, d) Association and (e, f) dissociation of 7 nM [^3H]-CCR2-RA from CHO cell membranes transiently expressing WT CCR2 (c, e) or CCR2-T4L (d, f) at 25°C, in the absence or presence of 1 μM BMS-681. Figures represent normalized and combined data from three independent experiments performed in duplicate, with results presented as mean $\pm$ S.E.M percentage of specific [^3H]-CCR2-RA binding.

Figure S5. A Zn^{2+} binding site was identified by X-ray fluorescence emission analysis of the CCR2-T4L-BMS-681-CCR2-RA-[R] crystals. (a) View of the Zn^{2+} ion at an interface formed by CCR2 helices III and VI and the N terminus of T4L. The Zn^{2+} ion is coordinated by side chains of H144/3.56 (from WT receptor), E238/3.0 (from the engineered part of the receptor), and E1005 (from T4L) as well as a structured water. (b) Background fluorescence signal of an empty MiTeGen micromount is low, indicating the absence of metal ion. Excitation at 12 keV results in a peak at 11.7 keV (owing to the incidence beam). (c) X-ray fluorescence emission signal from a wide fluorescence scan of the CCR2-T4L crystal. The fluorescence peaks at 8.60 keV and 9.53 keV correspond to X-ray emission lines $\text{K}\alpha$ (8.64 keV) and $\text{K}\beta$ (9.57 keV) and indicate the presence of Zn^{2+} bound to CCR2-T4L. (d) A zoomed-in view of the X-ray fluorescence emission signal from (c).




Table S1. Data collection and refinement statistics (molecular replacement)

	CCR2-T4L-BMS-681-CCR2-RA-[R] ^a
Data collection ^b Wavelength (Å)	1.03319
Space group	P2 ₁ 2 ₁ 2 ₁
Unit cell parameters a,b,c (Å)	59.19 64.69 169.90
Number of reflections measured	82,111
Number of unique reflections	15,550
Resolution (Å)	48-2.8 (2.95-2.8)
R _{merge} (%)	22.5(101)
R _{plim} (%)	12.8(88.4)
Mean I/s(I)	6.9(0.8)
Completeness (%)	93.1(66.6)
Redundancy	5.3(1.8)
Refinement	
Resolution (Å)	25-2.81 (3.0, 3.0, 2.81)
Number of reflections (test set)	14515 (746)
R _{work} /R _{free}	0.233/0.274 (0.319/0.392)
Number of atoms	3,580
CCR2	2,215
T4L	1,243
BMS-681	35
CCR2-RA-[R]	24
Monoolein	25
Sulfate	20
Water	17
Zn	1
Mean overall B value (Å ²)	41.4
Wilson B	40.4
Protein	41.5
Ligands	41.3
Water	22.9
Root mean square deviation	
Bond lengths (Å)	0.003
Bond angles (°)	0.85
Ramachandran plot statistics ^c (%)	
Favored regions	97.1
Allowed regions	2.9
Disallowed regions	0

^aDiffraction data from 17 crystals were merged into a complete data set

^bHighest resolution shell statistics are shown in parentheses

^cAs defined in MolProbity.⁴²

Table S2. Small Molecule (BMS-681) X-ray data collection and refinement.

Empirical formula	C26 H36 F3 N5 O3.58
Formula weight	532.80
Temperature	173(2) K
Wavelength	1.54178 Å
Crystal system	Tetragonal
Space group	P4 ₃ 2 ₁ 2
Unit cell dimensions	a = 20.4436(4) Å α = 90°. b = 20.4436(4) Å β = 90°. c = 28.9325(7) Å γ = 90°.
Volume	12092.1(4) Å ³
Z	16
Density (calculated)	1.171 Mg/m ³
Absorption coefficient	0.768 mm ⁻¹
F(000)	4522
Crystal size	0.46 x 0.18 x 0.16 mm ³
Theta range for data collection	2.65 to 58.78°.
Resolution range	16.7 to 0.9 Å
Index ranges	-22 ≤ h ≤ 22, -21 ≤ k ≤ 22, -31 ≤ l ≤ 14
Reflections collected	108743
Independent reflections	8518 [R(int) = 0.1259]
Completeness to theta = 58.78°	98.6 %
Absorption correction	None
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	8518 / 22 / 713
Goodness-of-fit on F ²	1.058
Final R indices [I>2sigma(I)]	R1 = 0.0770, wR2 = 0.2087
R indices (all data)	R1 = 0.0860, wR2 = 0.2178
Absolute structure parameter; Flack(x)	0.1(2)
Absolute structure parameter; Hooft(y), P3true	0.03(5), 1.000
Largest diff. peak and hole	0.543 and -0.405 e.Å ⁻³

Table S3. Displacement of specific [³H]-INCB3344 (5 nM) and [³H]-CCR2-RA (3 nM) binding from CCR2 constructs transiently expressed on CHO cells.

Construct	[³ H]-INCB3344 displacement by INCB-3344	[³ H]-INCB3344 displacement by BMS-681	[³ H]-CCR2-RA displacement by CCR2-RA-[R]	[³ H]-CCR2-RA enhancement by BMS-681
	pIC ₅₀ ± S.E.M (IC ₅₀ , nM)			%Binding
WT CCR2	7.8 ± 0.0 (17)	8.1 ± 0.0 (8)	7.9 ± 0.0 (13)	134 ± 3% ^a **
CCR2-T4L	8.1 ± 0.1* (8)	8.6 ± 0.1** (3)	8.2 ± 0.0** (6)	157 ± 13% ^a ****

Values represent mean ± S.E.M of at least three independent experiments performed in duplicate. ^aPercentage of [³H]-CCR2-RA (3 nM) binding in presence of BMS-681 (1 μM). Values higher than 100% represent binding enhancement compared with the 100% control without BMS-681. Differences in pIC₅₀ values between constructs were analyzed using a Student's t-test, with significant differences noted as follows: **P* < 0.05, ***P* < 0.01. Differences in percentage Binding in the absence (100%) and presence of BMS-681 were analyzed using a one-way analysis of variance with Dunnett's post-hoc test, with significant differences noted as follows: ***P* < 0.01, *****P* < 0.0001.

Table S4. Observed association and dissociation rate constants of [³H]-CCR2-RA (7 nM) on membranes from CHO cells transiently expressing WT CCR2 and CCR2-T4L, in the absence or presence of 1 μM BMS-681.

	CHO-CCR2		CHO-CCR2-T4L	
	Control	+ 1 μM BMS-681	Control	+ 1 μM BMS-681
k _{obs} (min ⁻¹)	0.031 ± 0.002	0.038 ± 0.003*	0.015 ± 0.003	0.015 ± 0.001
% B/B _{control} ^a	100 ± 0.0	135 ± 2.0****	100 ± 0.0	162 ± 8.4**
k _{off,fast} (min ⁻¹)	0.089 ± 0.015	0.069 ± 0.012*	0.077 ± 0.013	
k _{off,slow} (min ⁻¹)	0.016 ± 0.005	0.012 ± 0.004	0.010 ± 0.003	0.049 ± 0.003 ^b
%fast	70 ± 10	71 ± 11	69 ± 8	N/A ^b

Values represent mean ± S.E.M of three independent experiments performed in duplicate. ^aThe percentage of maximum binding in the absence (B_{control}) or presence (B) of BMS-681 (1 μM). ^bFor CHO-CCR2-T4L only, dissociation kinetics of [³H]-CCR2-RA (7 nM) in the presence of BMS-681 (1 μM) fitted best with a monophasic exponential decay model, resulting in a single k_{off} value, as shown in the table. Thus for CHO-CCR2-T4L, the statistical significance between k_{off} measurements with and without BMS-681 could not be calculated. Statistical significance was analyzed using a Student's t-test, with significant differences versus control noted as follows: **P* < 0.05, ***P* < 0.01, *****P* < 0.0001

