




REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY

ISSN 0034 – 365 X | E-ISSN 2337 – 8824 | Accredited 10/E/KPT/2019



REINWARDTIA

A JOURNAL ON TAXONOMIC BOTANY, PLANT SOCIOLOGY AND ECOLOGY

Vol. 18 (2): 51 – 133, December 10, 2019

Chief Editor

Kartini Kramadibrata (Mycologist, Herbarium Bogoriense, Indonesia)

Editors

Dedy Darnaedi (Taxonomist, Herbarium Bogoriense, Indonesia)

Tukirin Partomihardjo (Ecologist, Herbarium Bogoriense, Indonesia)

Joeni Setijo Rahajoe (Ecologist, Herbarium Bogoriense, Indonesia)

Marlina Ardiyani (Taxonomist, Herbarium Bogoriense, Indonesia)

Himmah Rustiami (Taxonomist, Herbarium Bogoriense, Indonesia)

Lulut Dwi Sulistyaningsih (Taxonomist, Herbarium Bogoriense, Indonesia)

Eka Fatmawati Tihuraa (Morphologist, Herbarium Bogoriense, Indonesia)

Topik Hidayat (Taxonomist, Indonesia University of Education, Indonesia)

Eizi Suzuki (Ecologist, Kagoshima University, Japan)

Jun Wen (Taxonomist, Smithsonian Natural History Museum, USA)

Barry J. Conn (Taxonomist, School of Life and Environmental Sciences, The University of Sydney, Australia)

David G. Frodin (Taxonomist, Royal Botanic Gardens, Kew, United Kingdom)

Graham Eagleton (Wagstaffe, NSW, Australia)

Secretary

Ruslan Bukhori

Layout

Liana Astuti

Illustrators

Wahyudi Santoso

Anne Kusumawaty

Correspondence on editorial matters and subscriptions for Reinwardtia should be addressed to:

HERBARIUM BOGORIENSE, BOTANY DIVISION,
RESEARCH CENTER FOR BIOLOGY– INDONESIAN INSTITUTE OF SCIENCES
CIBINONG SCIENCE CENTER, JLN. RAYA JAKARTA – BOGOR KM 46,
CIBINONG 16911, P.O. Box 25 CIBINONG
INDONESIA

PHONE (+62) 21 8765066; Fax (+62) 21 8765062

E-MAIL: reinwardtia@mail.lipi.go.id

<http://e-journal.biologi.lipi.go.id/index.php/reinwardtia>

Cover images: *Dinochloa glabra* Widjaja & Ervianti, *spec. nov.* A. Culm sheath. B. Leaves. C. Leaf sheath. D. Inflorescence (1. Floret. 2. Palea. 3. Lemma. 4. Glume (a, b, c). 5. Lodicule (a, b, c). 6. Anthers. 7. Stigma. 8. Fruit). From *Widjaja EAW 8864* (BO), drawing by Wahyudi Santoso (BO).

The Editors would like to thank all reviewers of volume 18(2):

Abdul Latiff Mohamad, Universiti Kebangsaan Malaysia (UKM), Bangi, Selangor, Malaysia

Andrew Powling, School of Biological Sciences, University of Portsmouth, United Kingdom

Barry J. Conn, School of Life and Environmental Sciences, The University of Sydney, Australia

Hans Joachim Esser, Botanische Staatssammlung München, Germany

Martin Dancak, Faculty of Science Palacky University, Czech Republic

Sumitra Salam, Nambol L. Sanoi College, Bishnupur, Manipur, India

Wong Khoon Meng, Herbarium Singapore, Singapore Botanic Gardens, 1 Cluny Road, Singapore

LECTOTYPIFICATION AND AMENDED DESCRIPTION OF *PHYLLANTHUS* (PHYLLANTHACEAE) SPECIES DESCRIBED BY KOORDERS FROM SULAWESI, INDONESIA

Received July 22, 2019; accepted September 11, 2019

RODERICK W. BOUMAN

Institute of Biology Leiden, Leiden University, PO Box 9505, 2300 RA Leiden, The Netherlands. Hortus botanicus Leiden, Leiden University, PO Box 9500, 2300 RA Leiden, The Netherlands. Naturalis Biodiversity Center, PO Box 9517, 2300 RA Leiden, The Netherlands. Email: r.w.bouman@hortus.leidenuniv.nl

PAUL J.A. KEßLER

Institute of Biology Leiden, Leiden University, PO Box 9505, 2300 RA Leiden, The Netherlands. Hortus botanicus Leiden, Leiden University, PO Box 9500, 2300 RA Leiden, The Netherlands. Email: p.j.a.kessler@hortus.leidenuniv.nl

PETER C. VAN WELZEN

Institute of Biology Leiden, Leiden University, PO Box 9505, 2300 RA Leiden, The Netherlands. Naturalis Biodiversity Center, PO Box 9517, 2300 RA Leiden, The Netherlands. Email: peter.vanwelzen@naturalis.com

ABSTRACT

BOUMAN, R. W., KEßLER, P. J. A. & VAN WELZEN, P. C. 2019. Lectotypification and amended description of *Phyllanthus* (Phyllanthaceae) species described by Koorders from Sulawesi, Indonesia. *Reinwardtia* 18(2): 97–103. — Two species of *Phyllanthus* collected and described by Koorders during his travels on the island of Sulawesi (Indonesia) are lectotypified, descriptions amended and their taxonomic affinity is discussed. *Phyllanthus mindorensis* was found to be too similar to *P. celebicus* and is placed in the synonymy of the latter. A key is provided to the species of *Phyllanthus* on Sulawesi.

Key words: Celebes, *Eriococcus*, Euphorbiaceae *s.l.*, Koorders, Phyllanthaceae, *Phyllanthus*, Sulawesi, taxonomy.

ABSTRAK

BOUMAN, R. W., KEßLER, P. J. A. & VAN WELZEN, P. C. 2019. Lektotipikasi dan deskripsi ulang jenis *Phyllanthus* (Phyllanthaceae) yang dipertelakan Koorders dari Sulawesi, Indonesia. *Reinwardtia* 18(2): 97–103. — Dua jenis *Phyllanthus* asal Sulawesi (Indonesia) yang dikoleksi dan dipertelakan oleh Koorders pada waktu perjalanan beliau di pulau Sulawesi disajikan lektotipifikasinya, dideskripsi ulang dan didiskusikan kekerabatan taksonominya. *Phyllanthus mindorensis* sangat menyerupai *P. celebicus* oleh karena itu dijadikan sebagai sinonim. Disajikan pula kunci jenis-jenis *Phyllanthus* Sulawesi.

Kata kunci: Celebes, *Eriococcus*, Euphorbiaceae *s.l.*, Koorders, Phyllanthaceae, *Phyllanthus*, Sulawesi, taksonomi.

INTRODUCTION

The flora of Sulawesi represents an interesting biodiversity hotspot that borders several biogeographical zones, with the Sunda shelf to the west, the Sahul shelf to the east and the Philippines to the north (Stelbrink *et al.*, 2012). It is the largest island of Wallacea, a biogeographic region that also includes the Moluccas, the Lesser Sunda Islands (Dickerson, 1928), and botanically usually also the Philippines (van Welzen *et al.*, 2011). While this island has become better explored recently, the flora remains understudied and many taxa did not receive any taxonomic treatment for some time. The enumeration of Euphorbiaceae for Central Malesia by Airy Shaw (1982) lists ten species of *Phyllanthus* L. for Sulawesi (Table 1), but this was only based on a limited number of collections. Airy Shaw (1982) made no redescription of the species and did not treat the

Wallacean islands extensively like he did for Borneo (Airy Shaw, 1975) and Papua New Guinea (Airy Shaw, 1980). Several species are probably still undiscovered and it is important that an adequate comparison can be made between those previously described from the island. Koorders (1898) reported two new species of *Phyllanthus* (*P. celebicus* Koord. and *P. minahassae* Koord.) in his travel account of the island, but only included a brief description of their habit with no mention of flower morphology. During the preparations for a new classification of the genus *Phyllanthus* several taxonomic problems were identified, often concerning rare species (Bouman *et al.*, 2018). This included the species of Koorders (1898), which could not yet be placed in any subgeneric group of *Phyllanthus* (Bouman *et al.*, 2018).

During a recent visit to the Herbarium Bogoriense (BO) on Java by the first author, the

Table 1. Species of *Phyllanthus* in Sulawesi, compiled from Robinson (1909), Airy Shaw (1982) and supplemented by records from the L herbarium. Species are listed by subgenus following Bouman *et al.* (2018). *Phyllanthus mindorensis* was listed by Airy Shaw (1982) and is treated here as synonym.

Subgenus	Species
<i>Eriococcus</i> (Hassk.)Croizat & Metcalf	<i>Phyllanthus buxifolius</i> (Blume) Müll.Arg. <i>Phyllanthus celebicus</i> Koord. <i>Phyllanthus lamprophyllus</i> Müll.Arg. <i>Phyllanthus minahassae</i> Koord. <i>Phyllanthus trichosporus</i> Adelb.
<i>Macraea</i> (Wight)Jean F.Brunel	<i>Phyllanthus lancifolius</i> Merr. <i>Phyllanthus samarensis</i> Müll.Arg. <i>Phyllanthus virgatus</i> G.Forst.
<i>Gomphidium</i> (Baill.)G.L.Webster	<i>Phyllanthus tenuirhachis</i> J.J.Sm.
<i>Kirganelia</i> (A.Juss.)Kurz	<i>Phyllanthus reticulatus</i> Poir.
<i>Emblica</i> (Gaertn.)Kurz	<i>Phyllanthus urinaria</i> L.
<i>Swartziani</i> (G.L.Webster)Ralim. & Petra Hoffm.	<i>Phyllanthus amarus</i> Schumach. & Thonn.
<i>Afroswartziani</i> Ralim. & Petra Hoffm.	<i>Phyllanthus debilis</i> Klein ex. Willd.

types of the Koorders' species could be studied. Here, we place these species in *Phyllanthus* subgenus *Eriococcus* (Hassk.) Croizat & Metcalf section *Eriococcus* and expand the descriptions of both species. Affinities to, and differences with, other species are discussed and a provisional key to the known *Phyllanthus* species of Sulawesi is provided.

TAXONOMIC TREATMENT

Both species discussed here are placed in *Phyllanthus* subgenus *Eriococcus* section *Eriococcus* based on the morphology of the staminate flowers. The staminate flowers in both species consist of four sepals in a cross shape with fimbriate margins, four disc glands and two connate stamens with horizontally dehiscing anthers. This is consistent with *Phyllanthus* subgenus *Eriococcus* section *Eriococcus* (see Müller, 1866) and both species are classified here in this taxon. Subgenus *Eriococcus* section *Emblicastrum* Müll.Arg., which is represented by *P. lamprophyllus* Müll.Arg. on Sulawesi, differs in the usually upright orientated sepals with entire margins, thicker leaves and the presence of a tubular style. No species of section *Eriococcus* is endemic to Papua New Guinea (Bouman *et al.*, 2018) and the closest affinity of both species are similar *Phyllanthus* species of the Philippines.

Roughly ten species of section *Eriococcus* occur on the Philippines and they are very similar in staminate flower and leaf morphology. Unfortunately many of these have been rarely collected. Differences for identification with the Philippine species are discussed below in the notes under the species, but they undoubtedly represent the closest relation to the species of Sulawesi within subgenus *Eriococcus*. A provisional key for *Phyllanthus* in Sulawesi is provided. Information for the key was derived from herbarium specimens, Luo *et al.* (2011), Robinson (1909) and Verwijs *et al.* (in press). All acronyms for herbaria follow Thiers (2019, continuously updated).

Note

Phyllanthus subgenus *Kirganelia* is represented by *P. reticulatus* Poir. on Sulawesi, though there is some discussion on how to differentiate it from *P. microcarpus* (Benth.)Müll.Arg. In the flora of Thailand (Chantaranothai, 2007), they are distinguished based on the presence or absence of indumentum while Luo *et al.* (2011) distinguish them based on habit and floral characters. Specimens from Sulawesi seen for this study were characterized by emergent styles and sometimes bisexual inflorescences while there were both pubescent and glabrous forms present. This conforms to the definition Luo *et al.* (2011) for *P. reticulatus* and it is treated as such here.

Key to the *Phyllanthus* species of Sulawesi

1. Branching non-phyllanthoid (lamine leaves on main axes present, lateral branches subtended by leaves and not deciduous); stamens 3, filaments free; fruits capsules — subgenus *Macraea* 2
1. Branching phyllanthoid (leaves on main axes reduced to cataphylls, lateral branchlets bear laminate leaves and are deciduous); stamens 2, 3 or 5; filaments free or connate (or in whorls); fruits capsules or berries..... 4
2. Prostrate or erect herbs or subshrubs, up to 1 m high, axes glabrous; pistillate pedicel 3–9 mm long *P. virgatus*
2. Erect shrubs to trees, up to 2 m high, axes mostly pubescent; pistillate pedicel 8–50 mm long..... 3
3. Leaf blades mostly ovate-elliptic, 9–79 mm long, apex acuminate; staminate sepals 1.1–1.4 × 0.5–0.8 mm; pistillate pedicel 8–50 mm long..... *P. lancifolius*
3. Leaf blades elliptic, 7–24 mm long, apex acute to obtuse or rounded to retuse; staminate sepals 1.2–1.6 × 0.6–0.8 mm; pistillate pedicel 8–10 mm long..... *P. samarensis*
4. Herbs (or only woody at the base)..... 5
4. Shrubs to trees..... 7
5. Pistillate inflorescences at basal part of lateral branchlets; ovary warted; seeds with transverse ridges— subgenus *Emblica*..... *P. urinaria*
5. Pistillate inflorescences at distal part of lateral branchlets; ovary smooth; seeds smooth or with longitudinal striae..... 6
6. Leaf blades oblong, apex rounded, upper side green; inflorescences mostly bisexual; staminate flowers with 5 sepals— subgenus *Swartziani*..... *P. amarus*
6. Leaf blades ovate, apex acute, upper side dark green; inflorescences unisexual; staminate flowers with 6 sepals— subgenus *Afroswartziani*..... *P. debilis*
7. Staminate flowers with 5 sepals, stamens 5, fused in 2 whorls; stigmas entire; fruits berries— subgenus *Kirganelia*..... *P. reticulatus*
7. Staminate flowers with 4 or 6 sepals, stamens 2 or 3, filaments free or connate in one whorl; stigmas bifid or entire (not seen in *P. minahassae*); fruits capsules..... 8
8. Leaves usually symmetric, blade elongated elliptic-ovate, longer than 5 cm; sepals 6 in both flowers of both sexes; stamens 3, filaments free, anthers dehiscent vertically — subgenus *Gomphidium* *P. tenuirhachis*
8. Leaves usually basally asymmetric, blade elliptic or ovate; shorter than 4.5 cm; sepals in staminate flowers 4, pistillate flowers with 5 or 6 sepals; stamens 2, filaments connate, anthers dehiscent horizontally— subgenus *Eriococcus*..... 9
9. Leaves coriaceous; staminate sepals erect, tubular, margins entire..... 10
9. Leaves membranous to slightly coriaceous; staminate sepals spreading, margins fimbriate, dentate or lacinate..... 11
10. Leaf blades up to 4 cm long; filaments of stamens thickened at apex; ovary 4–8-locular; style absent, stigma almost entire with only a bilobed fold at apex..... *P. buxifolius*
10. Leaf blades up to 1.2 cm long; filaments of stamens slender; ovary 3-locular; style present, tube of *ca.* 2.2 mm, stigmas entire..... *P. lamprophyllus*
11. Branchlets 8–10 cm long, glabrous; capsules glabrous, smooth; seeds *ca.* 1.8 mm high *P. trichosporus*
11. Branchlets 17–40 cm long, pubescent; capsules verrucate or smooth and puberulous; seeds *ca.* 2–4.8 mm high..... 12
12. Leaf blades shorter than 22 mm; sepals red to brown; pistillate sepals not enlarged in fruit, shorter than 4 mm; fruit pedicel up to 3.5 cm long..... 1. *P. celebicus*
12. Leaf blades longer than 25 mm; sepals pale green; pistillate sepals enlarged in fruit, up to 16 mm long; fruit pedicel 3–4 cm long..... 2. *P. minahassae*

Species descriptions of *P. celebicus* and *P. minahassae*

1. PHYLLANTHUS CELEBICUS Koord.

Phyllanthus celebicus Koord., Meded. Lands Plantentuin 19: 588, 627 (Koorders 1898). — Lectotype (designated here): Indonesia, Sulawesi, Minahasa province (Menado), near Pinomorongan, near Kajoewatoe, *H.S. Koorders 16949* (fieldnumber 1917) (lecto BO (BO129623); isolecto L (L.2246433, L.2246434).

Phyllanthus mindorensis C.B.Rob., Philipp. J. Sci., C 4: 82 (Robinson 1909). — Type: *L.M. Merritt 5370* (not traced); paratype: Philippines, Mindoro, *L.M. Merritt 8789* (K (K001056684)); paratype *L.M. Merritt 8606* (not traced). **Syn. nov.**

Shrubs, 0.5–1 m high, monoecious, all axes puberulent to pubescent with dark brown short stiff hairs; branching phyllanthoid; branchlets terete, 17–26 cm long, bearing 32–56 leaves, internodes 2–3 mm long. *Cataphyllary stipules* triangular, ca. 1.5 × 2 mm, membranous, caducous, base plane, tending to fuse with cataphyll, margin entire, apex acute. *Cataphylls* triangular, 1–1.5 × ca. 1 mm, membranous, caducous, margin entire, apex acute. *Stipules* triangular, ca. 1 × 0.5 mm, caducous, membranous, base plane, margin brittle, apex acute. *Leaves* distichous; sessile to petiole 0.5–1 mm long, glabrous; blade elliptic, asymmetric, 10–21 × 5–7.5 mm, 2–3.2 times longer than wide, membranous to slightly coriaceous, base cuneate, margin slightly thickened and slightly revolute, apex rounded, mucronate, mucro 0.1–0.2 mm long, glabrous except for some hairs on basal part of lower side, upper side darker green; venation pinnate, midrib prominent, flat on both sides, lateral veins 4–6 per side, indistinct. *Inflorescences* axillary fascicles, usually unisexual and originating from small brachyblasts, up to 4 staminate flowers together near basal part of branchlets, pistillate flowers solitary on distal part of branchlets. *Staminate flowers* ca. 1.1 mm in diameter closed, 2.2–3.5 mm when open; pedicel 4.5–5 mm long, terete, glabrous, thin; sepals 4, ovate, 1–1.5 × 0.8–1.1 mm, spreading, red-brown, margin dentate to lacinate, apex acute, midrib indistinct; disc glands 4, alternating with sepals, obovate, 0.2–0.3 × 0.3–0.6 mm, indented from thecae; stamens 2, 0.4–0.5 mm long, filaments and connectives connate, filaments ca. 0.2 mm high, anthers slightly stipitate resulting in cross-shaped connective, apically extended for ca. 0.1 mm, thecae oblong, ca. 0.3 × 0.2 mm, hanging above disc gland, dehiscing horizontally via slits. *Pistillate flowers*: pedicel ca. 16 mm long, terete, pubescent, thin; sepals 5, ovate, ca. 1.5 × 1.1 mm, red-brown, margin fimbriate to lacinate, apex

acute, midrib indistinct; disc annular, strongly star-shaped, ca. 0.8 mm in diameter at shortest point, with large oblong lobes alternating with sepals, lobes 0.3–0.7 mm long and ca. 0.2 mm wide, slightly thickened at the end; ovary globose, ca. 1.3 mm wide, ca. 1.1 mm high, puberulous; style absent; stigmas 3, bifid for entire length, ca. 0.2 mm long, curved upwards. *Fruits* capsules, only dehisced remains left; pedicel terete, 1.8–3.5 cm long, greenish, puberulous, rarely glabrous, smooth; sepals not enlarged in fruit, columella triangular, ca. 1.2 mm long. *Seeds* trigonous, ca. 2 × 1 mm, covered with transverse striae that break up epidermis, striae radiating from hilum.

Distribution. Philippines (Luzon, Mindoro) and Sulawesi (Tenga, Utara, Minahassa, Pangkadyeu).

Habitat. This species was collected from fertile turf and dry riverbeds, but it is known only from few specimens in Sulawesi. In the Philippines it was collected also from coastal areas. Altitude: 10–600 m a.s.l. This species was found with flowers and fruits in February till June, but more specimens could expand on this as we saw now collections from later in the year than June.

Etymology. This species is named after the area where it was found as Celebes is the name formerly used for Sulawesi.

Notes. The collection *Koorders 16949* stored at Bogor herbarium (BO) is selected as lectotype, while additional specimens are stored in the Naturalis Biodiversity Center (L). The material at Bogor bears Koorders' handwriting and notes on comparisons with other species, making it likely that it was used in the first description by him (Koorders, 1898). *Phyllanthus celebicus* agrees in almost all characters with *P. mindorensis* C.B.Rob., which was first described for the Philippines (Robinson, 1909) and its distribution was expanded by Airy Shaw (1982) to include Sulawesi. Airy Shaw noted on material of *P. celebicus* that it was very similar to *P. mindorensis* and differences between the Koorders specimens and the description by Robinson (1909) only include minor qualitative differences. Pistillate flowers were dissected from specimens from the Philippines and Sulawesi previously assigned to *P. mindorensis* and these were found to have similarly lobed discs, but with slightly shorter lobes (ca. 0.3 vs 0.6–0.7 mm). Taking into account the similarities in vegetative and floral characters and the variation shown by specimens from Sulawesi, we decided to synonymize *P. mindorensis* with *P. celebicus*.

Specimens examined. Indonesia, Sulawesi *W.H. de Vries s.n.* (L). Indonesia, Sulawesi, Minahassa,

Manembo-Nembo *A.H.G. Alston 16549* (L). Indonesia, Sulawesi, Road Palu - Donggala, near Loli *M.M.J. van Balgooy 2976* (BO, L). Indonesia, Sulawesi, Minahasa province (Menado), near Pinomorong, near Kajoewatoe, *H.S. Koorders 16949* (BO, L). Indonesia, Sulawesi, Tengah, Palu *G.J. de Joncheere 1024* (L). Indonesia, Sulawesi, Utara, Tondano *E.A. Forsten* (L). Philippines, Luzon, Batangas prov., Mt. Lobo, *PNH (M.D. Sulit) 15718* (L). Philippines, Luzon, Mt Arayat, *FB (H.M. Curran) 19333* (L). Philippines, Mindoro, Mt. Yagaw, *PNH (H.C. Conklin) 18674* (L). Philippines, Rizal Prov., *BS (M. Ramos) 731* (L). Philippines, Luzon, *BS (M. Ramos) 8194* (L).

2. P. MINAHASSAE Koord.

Phyllanthus minahassae Koord., Meded. Lands Plantentuin 19: 588, 627 (Koorders 1898). — Lectotype (designated here): Indonesia, Sulawesi, Minahasa province, near camp Totok, Rata Totok, *S.H. Koorders 16954* (fieldnumber 2595) (lecto BO (BO-1310079); isolecto L (L2059450)).

Shrubs, 1–2 m high, monoecious?, all axes pubescent with dark brown short stiff hairs; branching phyllanthoid; branchlets terete, 20–40 cm long, bearing 30–52 leaves, internodes 2–3 mm long. *Cataphyllary stipules* and *cataphylls* caducous, not seen. *Stipules* triangular, *ca.* 1.5 × 0.3 mm, persistent, membranous, pubescent, base plane, margin brittle, apex acute. *Leaves* distichous; petiole *ca.* 1.8 mm long, pubescent; blade elliptic, asymmetric, 2.6–4.1 × 1–1.7 cm, 2.4–3.4 times longer than wide, membranous to slightly coriaceous, pubescent, base cuneate, margin slightly thickened and flat, pubescent, apex acute-rounded to obtuse, mucronate, mucro 0.1–0.2 mm long, puberulous at base on lower side, glabrous on upper side, upper side darker green than lower side; venation pinnate, midrib flat on either side, lateral veins 4–6, indistinct. *Inflorescences* axillary fascicles, usually unisexual, staminate flowers up to 6 together, at basal part of branchlets, pistillate flowers solitary on distal part of branchlets. *Staminate flowers ca.* 1.5 mm in diameter when closed, 3.3–4 mm when open; pedicel 1.3–4 mm long, terete, pubescent, thin; sepals 4, ovate, 1.5–1.6 × *ca.* 1 mm, spreading, outside covered in short hairs, pale green, margin fimbriate and brittle, apex acute to attenuate, midrib indistinct; disc glands 4, alternating with sepals, oblong, 0.2–0.4 × 0.4–0.6 mm, indented from thecae; stamens 2, *ca.* 0.4–0.5 mm high, filaments and connectives connate, staminal column *ca.* 0.2 mm high, anthers slightly stipitate resulting in cross shaped connective, apically extended for 0.1–0.2 mm, thecae oblong, *ca.* 0.3 × 0.2 mm, hanging above disc gland, dehiscing horizontally via slits. *Pistillate flowers* not seen,

information derived from fruit; sepals 6, elliptic, enlarged (only in fruit?) to 14–16 × 5–6 mm, outside covered in short hairs, pale green, margin entire, pubescent, apex acute to obtuse, midrib indistinct; disc, ovary and stigmas not seen. *Fruits* capsules (already dehisced, only fragments on type), estimated at *ca.* 6 mm in diameter and *ca.* 6 mm high, minutely verrucate; pedicel terete, 3–4 cm long, puberulous; columella triangular, 4.8–6 mm long. *Seeds* trigonous, 4.2–4.8 × *ca.* 2.2 mm, covered with transverse striae that break up epidermis, striae radiating from hilum.

Distribution. Known from two collections by Koorders from the same area in Sulawesi, Minahasa province.

Habitat. The label information of the type showed it to be common in forests on fertile volcanic sand. It was found at *ca.* 200 m asl. Flowering and fruiting is in March, but this is only based on the type specimen and one additional collection around the same time.

Etymology. Named after the province where it was found on Sulawesi.

Notes. The lectotype is here selected from two collections by Koorders, both bear his handwriting, but only the specimen designated here as lectotype possesses remains of fruits and seeds. The staminate flowers were described from the other collection (*Koorders 16954*; BO1310078), but this specimen is in poorer condition, has less information on the label and has only dehisced fruits without seeds. *Phyllanthus minahassae* is probably closely related to *P. celebicus*, but it has larger leaves and very distinct pistillate sepals (at least in fruit). Large pistillate sepals are also found in the Indian species, *P. macrocalyx* Müll. Arg. (also subgenus *Eriococcus*) and some species of subgenus *Gomphidium* (Baill.) G.L. Webster (McPherson & Schmid 1991), but there they actually enclose the fruit in development (Naveen Kumar *et al.*, 2015).

Specimens examined. Indonesia, Sulawesi, Minahasa province, near camp Totok, Rata Totok *H.S. Koorders 16924* (fieldnumber 2430) (BO, L). Indonesia, Sulawesi, Minahasa province, near camp Totok, Rata Totok *S.H. Koorders 16954* (fieldnumber 2595) (BO, L).

DISCUSSION

With the rather brief description of *P. celebicus* and *P. minahassae* by Koorders (1898), these two species could not be confidently assigned to any subgroup within *Phyllanthus*. By examining material collected by Koorders including his notes on the labels, we were able to place both species

in subgenus *Eriococcus*. This subgenus is distributed in Asia (and one species in Australia) and its 4-merous staminate flower is a consistent character that can be used to identify them (Bouman *et al.*, 2018). However, while the staminate flower offers more information on subgeneric placement, the pistillate flower is often more useful for species identification as found here for *P. celebicus* and *P. minahassae*. Subgenus *Eriococcus* is mainly known from mainland Asia (see Bouman *et al.*, 2018), but only includes few species from Western Malesia (e.g. *P. acutissimus* Miq., *P. kinabaluicus* Airy Shaw, *P. singalensis* (Miq.) Müll.Arg.), which often have symmetric leaves (pers. obs.). Therefore, the species treated here are most likely closely related to other species from the Philippines, which also have asymmetric leaves that appear quite similar. Some species from the Philippines and Sulawesi of section *Eriococcus* seem to form a complex, all with usually pubescent axes, strongly asymmetric leaves and usually red flowers with fimbriate sepals (some exceptions occur). Other species that form this complex, aside from those already mentioned, include *P. laciniatus* C.B.Rob., *P. leytenensis* Elmer, *P. sibuyanensis* Elmer, and perhaps *P. blancoanus* Müll.Arg. (see Robinson 1909). A thorough revision or phylogenetic study with sampling of these taxa might help to improve some of the species delimitations. This group of species could lend support to the inclusion of the Philippines in Wallacea (see van Welzen *et al.*, 2011). While not many species of *Phyllanthus* occur on Sulawesi, the two species treated here already highlight the interesting flora found on the island and the need to study it further.

ACKNOWLEDGEMENTS

This work was done as part of the PhD research of the first author, funded by the Hortus botanicus Leiden and Leiden University. Travel funding granted by the Alberta Mennega foundation to visit herbaria in Indonesia and the Flora Malesiana symposium in Brunei is gratefully acknowledged. The directors and keepers of BO and L are thanked for use of their collections. Deby Arifiani and Michael Mambrasar are thanked for their help at BO herbarium with locating and studying the Koorders collections. Aninda Wibowo and Eka Iskandar helped with the abstract in Indonesian, for which we are very grateful. The second author thanks the Leiden University Fund (LUF) for their support of the chair Botanical Gardens and Botany of Southeast Asia. The last author thanks the Treub Maatschappij, the Society for the Advancement of Research in the Tropics, for their support of the Ornstein chair in Tropical Plant Biogeography.

REFERENCES

- AIRY SHAW, H. K. 1975. The Euphorbiaceae of Borneo. *Kew Bull., Addit. Ser.* 4: 1–224.
- AIRY SHAW, H. K. 1980. The Euphorbiaceae of New Guinea. *Kew Bull., Addit. Ser.* 8: 3–230.
- AIRY SHAW, H. K. 1982. The Euphorbiaceae of Central Malesia (Celebes, Moluccas, Lesser Sunda Is.). *Kew Bull.* 37: 1–40.
- BOUMAN, R. W., KEßLER, P. J. A., TELFORD, I. R. H., BRUHL, J. J. & VAN WELZEN, P. C. 2018. Subgeneric delimitation of the plant genus *Phyllanthus* L. (Phyllanthaceae). *Blumea* 63: 167–198.
- CHANTARANOTHAI, P. 2007. *Phyllanthus*. In: P. C. VAN WELZEN & K. CHAYAMARIT (Eds.). *Flora of Thailand* 8(2). The Forest Herbarium, Bangkok. Pp. 305–592.
- DICKERSON, R. E. 1928. *Distribution of life in the Philippines*. Bureau of Sciences, Manila.
- KOORDERS, H. S. 1898. Verslag eener botanische dienstreis door de Minahassa. *Meded. Lands Plantentuin* 19: 1–716.
- LUO, S., ESSER, H.-J., ZHANG, D., & RENNER S. S. 2011. Nuclear ITS sequences help disentangle *Phyllanthus reticulatus* (Phyllanthaceae), an Asian species not occurring in Africa, but introduced to Jamaica. *Syst. Bot.* 36: 99–104.
- MCPHERSON, G. & SCHMID, M. 1991. *Flore de la Nouvelle Calédonie et Dépendances 17: Euphorbiaceae*. Muséum National d'Histoire Naturelle, Paris.
- MÜLLER, J. 1866. Euphorbiaceae excl. Euphorbieae. In: A. L. P. P. DE CANDOLLE (Ed.), *Prodromus Systematis Naturalis Regni Vegetabilis* 15, 2: Victor Masson, Paris. Pp. 189–1286.
- NAVEEN KUMAR, V. V., PRABHUKUMAR, K. M., BHAVADAS, N. & SUNIL, C. N. 2015. Rediscovery of *Phyllanthus macrocalyx* Müll.Arg. (Phyllanthaceae), a rare endemic species of Western Ghats, India. *Curr. Sci.* 109: 33–35.
- ROBINSON, C. B. 1909. Philippine Phyllanthinae. *Philipp. J. Sci., C.* 4: 71–106.
- STELBRINK, B., ALBRECHT, C., HALL, R. & RINTELEN, T. VON. 2012. The biogeography of Sulawesi revisited: is there evidence for a vicariant origin of taxa on Wallace's "anomalous island"? *Evolution* 66: 2252–2271.
- THIERS, B. 2019. Continuously updated. Index Herbariorum: A Global Directory of public Herbaria and associated Staff. New York Botanical Garden's Virtual Herbarium, New York. Available from <http://sweetgum.nybg.org/science/ih/> [Accessed 10 July 2019].

VAN WELZEN, P. C., PARNELL, J. A. N. & SLIK, J. W. F. 2011. Wallace's line and plant distributions: two or three phytogeographical areas and where to group Java?. *Biol. J. Linn. Soc.* 103: 531–545.

VERWIJS, J. I. M., BOUMAN, R. W. & VAN WELZEN, P. C. in press. A taxonomic revision of *Phyllanthus* subgenus *Macraea* (Phyllanthaceae). *Blumea* in press.

INSTRUCTION TO AUTHORS

Scope. *Reinwardtia* is a scientific regular journal on plant taxonomy, plant ecology and ethnobotany published in June and December. Manuscript intended for a publication should be written in English.

Titles. Titles should be brief, informative and followed by author's name and mailing address in one-paragraphed.

Abstract. English abstract followed by Indonesian abstract of not more than 250 words. Keywords should be given below each abstract.

Manuscript. Manuscript is original paper and represent an article which has not been published in any other journal or proceedings. The manuscript of no more than 36 pages by using Times New Roman 11, MS Word for Windows of A4 with double spacing, submitted to the editor through <reinwardtia@mail.lipi.go.id>. New paragraph should be indented in by 5 characters. For the style of presentation, authors should follow the latest issue of *Reinwardtia* very closely. Author(s) should send the preferred running title of the article submitted. Every manuscript will be sent to two blind reviewers.

Identification key. Taxonomic identification key should be prepared using the aligned couplet type.

Nomenclature. Strict adherence to the International Code of Nomenclature is observed, so that taxonomic and nomenclatural novelties should be clearly shown. English description for new taxon proposed should be provided and the herbaria where the type specimens area deposited should be presented. Name of taxon in taxonomic treatment should be presented in the long form that is name of taxon, author's name, year of publication, abbreviated journal or book title, volume, number and page.

Map/line drawing illustration/photograph. Map, line drawing illustration, or photograph preferably should be prepared in landscape presentation to occupy two columns. Illustration must be submitted as original art accompanying, but separated from the manuscript. The illustration should be saved in JPG or GIF format at least 350 pixels. Legends or illustration must be submitted separately at the end of the manuscript.

References. Bibliography, list of literature cited or references follow the Harvard system as the following examples.

- Journal : KRAENZLIN, F. 1913. *Cyrtandraceae* novae Philippinenses I. *Philipp. J. Sci.* 8: 163–179.
MAYER, V., MOLLER, M., PERRET, M. & WEBER, A. 2003. Phylogenetic position and generic differentiation of *Epithemateae* (Gesneriaceae) inferred from plastid DNA sequence data. *American J. Bot.* 90: 321–329.
- Proceedings : TEMU, S. T. 1995. Peranan tumbuhan dan ternak dalam upacara adat “Djoka Dju” pada suku Lio, Ende, Flores, Nusa Tenggara Timur. In: NASUTION, E. (Ed.). *Prosiding Seminar dan Lokakarya Nasional Etnobotani II*. LIPI & Perpustakaan Nasional. Pp. 263–268. (In Indonesian).
SIMBOLON, H. & MIRMANTO, E. 2000. Checklist of plant species in the peat swamp forests of Central Kalimantan, Indonesia. In: IWAKUMA, T., INOUE, T., KOHYAMA, T., OSAKI, M., SIMBOLON, H., TACHIBANA, H., TAKAHASHI, H., TANAKA, N., YABE, K. (Eds.). *Proceedings of the International Symposium on: Tropical Peatlands*. Pp. 179 – 190.
- Book : RIDLEY, H. N. 1923. *Flora of the Malay Peninsula* 2. L. Reeve & Co. Ltd, London.
- Part of Book : BENTHAM, G. 1876. *Gesneriaceae*. In: BENTHAM, G. & HOOKER, J. D. (Eds.). *Genera Plantarum* 2. Lovell Reeve & Co., London. Pp. 990–1025.
- Thesis : BAIRD, L. 2002. *A Grammar of Kéo: An Austronesian Language of East Nusantara*. Australian National University, Canberra. [PhD. Thesis].
- Website : <http://www.nationaalherbarium.nl/fmcollectors/k/KostermansAJGH.html>. (Accessed 15 February 2012).



Reinwardtia

Published by Herbarium Bogoriense, Botany Division, Research Center for Biology,
Indonesian Institute of Sciences

Address: Jln. Raya Jakarta-Bogor Km. 46 Cibinong 16911, P.O. Box 25 Cibinong

Telp. (+ 62) 21 8765066; Fax (+62) 21 8765062

Email: reinwardtia@mail.lipi.go.id

***REINWARDTIA* Author Agreement Form**

Title of article :

Name of Author(s) :

I/We hereby declare that:

- My/Our manuscript was based on my/our original work.
- It was not published or submitted to other journal for publication.
- I/we agree to publish my/our manuscript and the copyright of this article is owned by Reinwardtia.
- We have obtained written permission from copyright owners for any excerpts from copyrighted works that are included and have credited the sources in our article.

Author signature (s)

Date

Name

REINWARDTIA

Vol. 18. No. 2. 2019

CONTENTS

- INA ERLINAWATI, NI PUTU SRI ASIH, AGUNG KURNIAWAN & YUZAMMI. Studies on the Araceae of the Lesser Sunda Islands II: New record for *Scindapsus hederaceus* Miq. in Bali 51
- RISMITA SARI, MIFTAHUL HUDA, RATNA SUSANDARINI & INGGIT PUJI ASTUTI. *Rafflesia hasseltii* Suringar (Rafflesiaceae): A new record to Kalimantan, Indonesia 65
- ROMITA DEVI NGANGBAM, NAOREM PREMITA DEVI, MAIBAM HARIPRIYA DEVI & P. K. SINGH. Rediscovery of *Aldrovanda vesiculosa* L. (Droseraceae), an endangered plant, from Manipur in India after six decades, with studies on micromorphology and physico-chemistry of water 71
- MOHD NORFAIZAL GHAZALLI, AMIN ASYRAF TAMIZI, MUHAMAD IKHWANUDDIN MAT ESA, EDWARD ENTALAI BESI, DOME NIKONG, ANUAR RASYIDI MOHD NORDIN & AHMAD ZAKI ZAINI. The systematic significance of leaf epidermal micromorphology of ten *Nepenthes* species (Nepenthaceae) from Peninsular Malaysia 81
- RODERICK W. BOUMAN, PAUL J. A. KEBLER & PETER C. VAN WELZEN. Lectotypification and amended description of *Phyllanthus* (Phyllanthaceae) species described by Koorders from Sulawesi, Indonesia 97
- SANATOMBI DEVI YUMKHAM, NAOREM PREMITA DEVI, SANDHYARANI DEVI KHOMDRAM & MAYANGLAMBAM ROMA DEVI. *Trichodesma kumareum* (Boraginaceae), a new species from North East India 105
- DITA ERVIANTI, ELIZABETH A. WIDJAJA & AGUNG SEDAYU. New species of climbing and scrambling bamboo from Sulawesi, Indonesia 115
- SOEJATMI DRANSFIELD. Book review: The Spectacular Indonesian Bamboos 133

Reinwardtia is an accredited Journal (10/E/KPT/2019)

<http://e-journal.biologi.lipi.go.id/index.php/reinwardtia>

Herbarium Bogoriense
Botany Division
Research Center for Biology – Indonesian Institute of Sciences
Cibinong Science Center
Jln. Raya Jakarta – Bogor, Km 46
Cibinong 16911, P.O. Box 25 Cibinong
Indonesia

