



Universiteit  
Leiden  
The Netherlands

## **The evolution of chemical diversity in plants : pyrrolizidine alkaloids and cytochrome P450s in *Jacobaea***

Chen, Y.

### **Citation**

Chen, Y. (2020, January 29). *The evolution of chemical diversity in plants : pyrrolizidine alkaloids and cytochrome P450s in Jacobaea*. Retrieved from <https://hdl.handle.net/1887/83487>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/83487>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/83487> holds various files of this Leiden University dissertation.

**Author:** Chen, Y.

**Title:** The evolution of chemical diversity in plants : pyrrolizidine alkaloids and cytochrome P450s in *Jacobaea*

**Issue Date:** 2020-01-29

---

# Chapter 6

## Summary and conclusions

## Summary and conclusions

Plants produce an astonishing variety of secondary metabolites (SMs) which are thought to play vital roles in the fitness of plants through ecological interactions (Wink, 2003; Moore *et al.*, 2014). SMs enable plants to deal with antagonists (e.g. herbivores, pathogens, neighboring plants) and mutualists (e.g. pollinators, predators/parasitoids against herbivores, mycorrhizal fungi, rhizobium and other beneficial bacteria) as well as abiotic factors (e.g. UV light, drought, frost) (Kessler and Halitschke, 2007). The most characteristic features of SMs are their striking chemical diversity and inter- or intraspecific variation. So far, an estimated 200,000 SMs have been isolated and identified from plants (Kessler and Kalske, 2018). SMs are assigned to different compound classes such as glucosinolates, alkaloids, terpenoids and flavonoids. Within each chemical class, both qualitative and quantitative SM diversity are common in plants. For instance, in *Arabidopsis thaliana* 34 different glucosinolates have been identified, showing 20-fold difference in total concentration in leaves of different ecotypes (Kliebenstein *et al.*, 2001). As yet, it is poorly understood how SM diversity comes about and why it is maintained in nature (Moore *et al.*, 2014).

In this thesis we focus on SMs involved in plant protection against herbivores. Aiming at a better understanding of mechanisms behind SM diversity, researchers have put much effort in studying the distribution patterns of SMs under particular phylogenetic frameworks (Wink, 2003; Wink, 2008; Pelsler *et al.*, 2005; Mint Evolutionary Genomics Consortium, 2018). At the level of a particular class of compounds (e.g. quinolizidine alkaloids, glucosinolates) there is a strong phylogenetic signal although some exceptions exist. This suggests that once the ability to produce the basic structures of these classes of compounds has evolved it is conserved through evolutionary time and not often convergent evolution takes place. In contrast the distributions of different chemical modifications of these structures on phylogenetic trees are often random. That is to say particular SMs within a class lack phylogenetic signals and phylogenetic distances are not correlated with differences in SM bouquets (Pelsler *et al.*, 2005; Mint Evolutionary Genomics Consortium, 2018; Chapter 2). This suggests that such modifications are not conserved and can rapidly appear or disappear on an evolutionary time scale. An evolutionary explanation would be that the basic structures that helps to protect plants against a broad set of herbivores comes about once and is conserved and that further evolutionary fine-tuning is dependent on the selective effects of the specific set of generalist and specialist herbivores each plant species is confronted with at a particular moment in evolutionary time. At the molecular level, the fact that new basic structures do not rapidly evolve in many plant families suggests that the enzymatic changes needed to create them are relatively complex. The further modification of newly evolved basic structures involves relative simple chemical reactions such as oxidation that are carried out by more general enzymes that are already present in the plant. The multiple functions of such an enzyme are subsequently reduced to single function by duplication of the underlying gene allowing optimizing the catalyzation for the new function (Hughes, 1994; DePristo, 2007). Examples

are the large gene families of cytochrome P450s (CYPs) of which the members show large similarity and that rapidly evolve (Bak *et al.*, 2006; Frey *et al.*, 2009; Chapter 3). Apparently this enables fast evolution, loss of phylogenetic signal and rapid evolutionary fine-tuning of their efficacy. The process of evolutionary tinkering is most likely in addition speeded up by genetic changes in the expression of CYPs. The investigation of the diversity of both SM profiles and the CYP family in plants may allow insights into the evolution of SM pathways that coordinate the respective enzymes.

The pyrrolizidine alkaloids (PAs) of *Jacobaea* were used as the model system in this thesis. PAs are a class of SMs with typical great diversity that are constitutively formed in plants containing them and are thought to mediate the interactions between plants and herbivores (Hartmann, 1999). The *Jacobaea* species (26 species and formerly a part of *Senecio* species, Asteraceae) all produce PAs (Pelser *et al.*, 2005; Langel *et al.*, 2011) but the composition and concentration are often species-specific (Soldaat *et al.*, 1996; Hartmann and Dierich, 1998; Langel *et al.*, 2011). Nevertheless, individual PAs seem to have random distributions on the phylogenetic tree lacking phylogenetic signals (Pelser *et al.*, 2005). Two hypotheses can be used to explain the random occurrences of PAs in *Jacobaea* species, i.e. (i) convergent evolution where the ability of plant species to produce particular PAs evolved several times, (ii) differential gene regulation of the PA pathway where all the plant species possess the machinery to produce those PAs but do not all express them. In order to figure out which hypothesis is correct, the occurrence and expression of genes underlying SM pathways need to be investigated.

So far, more than 400 PAs have been found (Chou and Fu, 2006), of which more than 100 PAs are macrocyclic senecionine-type (Hartmann and Witte, 1995; Langel *et al.*, 2011). Most of our current knowledge of PA biosynthetic diversification has come from the studies of the 12-membered macrocyclic senecionine-type PAs in the Senecioneae (Langel *et al.*, 2011). The primary product senecionine *N*-oxide synthesized in roots (Hartmann and Toppel, 1987; Hartmann *et al.*, 1988; Hartmann *et al.*, 1989) undergoes structural transformations in a position-specific and stereoselective manner resulting in the rearrangement of the skeletal structure and oxidative modifications thereof in shoots (Hartmann and Dierich, 1998; Pelser *et al.*, 2005). The enzymes responsible for these processes have not been identified. CYPs catalyze a wide range of regiospecific, stereospecific and irreversible oxidation steps in plant SM biosynthesis (Renault *et al.*, 2014), thus playing an important role in the evolution of chemical diversity. Given the common oxygenated site-specific modifications of PAs derived from the primary PA senecionine *N*-oxide, CYPs are good candidates for involvement in PA biosynthesis.

In this thesis the following questions were proposed: what are the distribution patterns of PAs in *Jacobaea* species and are these patterns related to their phylogenetic relationships? How does this PA diversity come about? Are CYPs involved in PA biosynthesis? Following these questions, the experimental chapters of this thesis can be divided into two sections: (i)

the evolution of PA diversity among and within *Jacobaea* species (Chapter 2), (ii) a candidate gene approach targeting CYPs for involvement in PA diversity (Chapter 3-5).

### 1. The evolution of PA diversity among and within *Jacobaea* species

In chapter 2, aiming to understand the mechanism behind chemical diversity, PA patterns of *Jacobaea* species were studied in a phylogenetic context. The presences and concentrations of 80 PAs in eight to ten week old leaves of 17 *Jacobaea* species including different individuals and populations grown in a controlled chamber were analyzed by LC-MS/MS. A great diversity of PA profiles was observed with the numbers of PAs ranging from 21 to 59 per species, and with the total PA concentrations ranging from 32.9 to 3835.7  $\mu\text{g/g}$  dry weight. These profiles were largely confirmed to be species-specific both qualitatively and quantitatively, which is in line with previous findings (Soldaat *et al.*, 1996; Hartmann and Dierich, 1998; Langel *et al.*, 2011). Jacobine-like PAs, senecionine-like PAs and otosenine-like PAs contributed more to the classification of different *Jacobaea* species than other PA structural groups. PA patterns from different populations within some *Jacobaea* species (e.g. *J. alpina* and *J. paludosa*) differed from each other even surpassing differences between species. The phylogeny of 17 *Jacobaea* species were reconstructed using 11 chloroplast regions and three nuclear DNA genes to trace the evolution of PA diversity at species level. With ancestral state reconstruction of the occurrences of individual PAs, complex evolutionary patterns for almost all PAs were found showing that the occurrence of virtually all PAs evolved several times, which is in agreement with the findings of Pelsler *et al.* (2005). Two different measures, Blomberg's  $K$  (Blomberg, 2003) and Pagel's  $\lambda$  (Pagel, 1999), were used to evaluate the correlations between quantitative PA traits and phylogenetic relationships, and significant phylogenetic signals for nine out of 80 PAs only under  $\lambda$  statistics were found. Given the common intraspecific PA diversity found in *Jacobaea* species (this thesis; Witte *et al.*, 1992; Macel *et al.*, 2004), it can be assumed that this high PA diversity is due to the regulation of PA biosynthesis genes in plants as a life strategy to meet their different biological needs rather than the evolutionary gains and losses of particular PA biosynthesis genes.

### 2. A candidate gene approach targeting CYPs for involvement in PA diversity

Testing our hypothesis that PA diversity in *Jacobaea* species is the result of the regulation of PA biosynthesis genes requires the genes underlying PA biosynthesis. CYPs have formed a large gene family in plants and have often been found to be involved in SM pathways as oxidative enzymes. As diversification of PAs is often through site-specific oxidation (Hartmann and Dierich, 1998; Pelsler *et al.*, 2005), CYPs are likely to be involved in PA biosynthesis. Yet, no public CYP database of *Jacobaea* species is available at the start of the studies described in this thesis. Therefore, a systematical study of CYPs was performed with regard to their diversity and evolution in *J. vulgaris* and *J. aquatica* as described in Chapter 3. In total, 221 (classified into eight clans and 38 families) and 157 (classified into eight clans and 35 families) full-length CYPs were retrieved from *de novo* assembled transcriptomes of *J.*

*vulgaris* and *J. aquatica*, respectively. Based on KEGG annotation, the CYPs assigned as being SM metabolic pathway enzymes were all from the CYP71 clan but no CYPs were assigned as being involved in alkaloid pathways. This does not necessarily mean that they are not involved in alkaloid biosynthesis since the current KEGG database does not contain information about PA biosynthetic enzymes. Phylogenetic analyses of the six largest CYP families (CYP71, CYP76, CYP706, CYP82, CYP93 and CYP72) from the two *Jacobaea* species, two other members of the Asteraceae, *Helianthus annuus* and *Lactuca sativa* and the outgroup of *Arabidopsis thaliana* were performed. The phylogenetic trees showed strong lineage-specific expansion of CYPs, suggesting that the evolution of CYPs has been very fast even within the Asteraceae family. Only CYPs of the closely related species *J. vulgaris* and *J. aquatica* were found often in pairs, confirming a close relationship in evolutionary history of these two species. The studies described in Chapter 3 provide a CYP database for future exploration of their functions, including possible involvement in PA biosynthesis and PA diversity.

Chapter 4 describes an attempt to identify candidate CYPs that may be involved in PA biosynthesis based on the association between metabolic and transcriptomic profiles between different *Jacobaea* samples grown under controlled conditions. *J. aquatica* and four groups of F<sub>2</sub> hybrids of a cross between *J. vulgaris* and *J. aquatica* with PA contrasts were used for constitutive PA contrasts, especially in jacobine-like and erucifoline-like PAs. A methyl jasmonate treatment on tissue culture plants of *J. vulgaris* were performed to induce increase of erucifoline-like PAs. In total, 44 PAs were detected by LC-MS/MS and PA profiles of different *Jacobaea* samples were compared separately in the constitutive and induced groups by summing up concentrations of PAs containing the same site-specific oxidative modifications which might be catalyzed by a CYP enzyme: 15,20-epoxidation, 12,13-epoxidation or 19-hydroxylation, 18-hydroxylation, 13,19-dehydrogenation and 8-oxidation. RNA sequencing was performed separately for the constitutive and induced groups to analyze the expression of CYPs which may be involved in PA oxidative conversions. In total, 33 and 27 CYP candidate genes were sieved out for the constitutive and induced PA conversions, respectively. Most of these candidate enzymes were from the CYP71 clan without known functions. There were 11 CYP subfamilies found both in the constitutive and induced groups, where three subfamilies (CYP72A, CYP706E, CYP82Q) may be responsible for the formation of erucifoline-like PAs which contain both 12,13-epoxidation and 19-hydroxylation.

Chapter 5 describes functional tests of eight CYP candidates for involvement in the PA biosynthesis pathway using heterologous expression in yeast and *in vitro* enzyme assays using microsomal membrane preparations presumably containing the expressed CYPs. None of the eight CYP enzyme preparations showed PA conversion with the selected PA substrates (senecionine/integerrimine, seneciophylline, jacobine, erucifoline or a PA mixture as well as the respective *N*-oxides) based on *in vitro* enzyme assays using extracted microsomal membranes. The reasons of the negative results might be lack of efficient expression, and/or requirement for more optimal reaction conditions or better detection method. More likely is

that the tested eight CYPs are not enzymes responsible for PA biosynthesis, or at least do not act on the tested substrates.

### 3. Discussion and conclusions

*Jacobaea* species have been frequently used to study their PA diversity (Vrieling *et al.*, 1993; Hartmann and Dierich, 1998; Macel *et al.*, 2004; Cheng *et al.*, 2011). Compared with previous studies, we studied both qualitative and quantitative PA variation at more levels including among species, among populations and among individuals at the same time. Our results revealed stronger evidence that PA patterns of *Jacobaea* species are indeed species-specific both in concentrations and compositions, though for some species PA profiles also differed largely between populations. It is the first study showing that the distribution pattern of concentrations among and within *Jacobaea* species was highly similar to that of compositions, implying the same or closely related mechanisms behind quantitative and qualitative variation of PAs.

The occurrence of individual PAs was studied at species level and limited phylogenetic signals were found. A reasonable assumption is that all *Jacobaea* species possess the machinery for PA production and that differences in PA occurrence are due to differential gene regulation. Based on the fact that PA diversity is largely created by oxidation reactions, CYPs, a major class of oxidative enzymes in plants, were chosen as an entry for identification of PA biosynthesis genes. Based on current scientific literature, the curated CYP database developed in this thesis is the first in *Jacobaea* species. A gene-to-metabolite approach was used to identify CYP candidates possibly involved in PA biosynthesis and the functions of eight candidate CYP enzymes was checked but this did not lead to any indication for their involvement in PA biosynthesis. The majority of candidate gene identified in Chapter 4 still needs to be tested. There is a possibility that the enzymes underlying PA biosynthesis are not CYPs but other oxidative enzymes such as flavin-dependent monooxygenases or peroxidases (Burton, 2003), for which gene-to-metabolite correlations can also be retrieved from the data described in Chapter 4.

### References

- Blomberg SP, Garland T, Ives AR. 2003. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* 57:717-745.
- Böttcher F, Adolph R-D, Hartmann T. 1993. Homospermidine synthase, the first pathway-specific enzyme in pyrrolizidine alkaloid biosynthesis. *Phytochemistry* 32:1373-1384.
- Burton SG. 2003. Oxidizing enzymes as biocatalysts. *Trends in Biotechnology* 21:543-549.
- Cheng D, Hirk H, Mulder PPJ, Vrieling K, Klinkhamer PGL. 2011. Pyrrolizidine alkaloid variation in shoots and roots of segregating hybrids between *Jacobaea vulgaris* and *Jacobaea aquatica*. *New Phytologist* 192: 1010-1023.



- Chou MW, Fu PP. 2006. Formation of DHP-derived DNA adducts in vivo from dietary supplements and Chinese herbal plant extracts containing carcinogenic pyrrolizidine alkaloids. *National Center for Toxicological Research* 22:321-327.
- Hartmann T, Dierich B. 1998. Chemical diversity and variation of pyrrolizidine alkaloids of the senecionine type: biological need or coincidence? *Planta* 206: 443-451.
- Hartmann T, Ehmke A, Eilert U, von Borstel K, Theuring C. 1989. Site of synthesis, translocation and accumulation of pyrrolizidine alkaloid *N*-oxides in *Senecio vulgaris* L. *Planta* 177:98-107.
- Hartmann T, Sander H, Adolph R, and Toppel G. 1988. Metabolic links between the biosynthesis of pyrrolizidine alkaloids and polyamines in root cultures of *Senecio vulgaris*. *Planta* 175:82-90.
- Hartmann T, Toppel G. 1987. Senecionine *N*-oxide, the primary product of pyrrolizidine alkaloid biosynthesis in root cultures of *Senecio vulgaris*. *Phytochemistry* 26:1639-1643.
- Hartmann T, Witte L. 1995. Chemistry, biology and chemoeology of the pyrrolizidine alkaloids. In Pelletier SW (ed) *Alkaloids: chemical and biological perspective*. Oxford: Pergamon Vol. 9, pp.155-233.
- Hartmann T. 1999. Chemical ecology of pyrrolizidine alkaloids. *Planta* 207:483-495.
- Kessler A, Halischke R. 2007. Specificity and complexity: the impact of herbivore-induced plant response on arthropod community structure. *Current Opinion in Plant Biology* 10:409-414.
- Kessler A, Kalske A. 2018. Plant secondary metabolite diversity and species interactions. *Annual Review Ecology, Evolution, and Systematics* 49:115-138.
- Kliebenstein DJ, Kroymann J, Brown P, Figuth A, Pedersen D, Gershenzon J, Mitchell-Olds T. 2001. Genetic control of natural variation in *Arabidopsis* glucosinolate accumulation. *Plant Physiology* 126:811-825.
- Langel D, Ober D, Pelser PB. 2011. The evolution of pyrrolizidine alkaloid biosynthesis and diversity in the Senecioneae. *Phytochemistry Reviews* 10:3-74.
- Liu Q, Beyraghdar Kashkooli A, Manzano D, Pateraki I, Richard L, Kolkman P, Lucas MF, Guallar V, de Vos RCH, Franssen MCR, van der Krol A, Bouwmeester H. 2018. Kauniolide synthase is a P450 with unusual hydroxylation and cyclization-elimination activity. *Nature Communications* 9:4657.
- Macel M, Vrieling K, Klinkhamer PG. 2004. Variation in pyrrolizidine alkaloid patterns of *Senecio jacobaea*. *Phytochemistry* 65:865-873.
- Mint Evolutionary Genomics Consortium. 2018. Phylogenomic mining of the mints reveals multiple mechanisms contributing to the evolution of chemical diversity in Lamiaceae. *Molecular Plant* 11:1084-1096.
- Moore BD, Andrew RL, Külheim C, Foley WJ. 2014. Explaining intraspecific diversity in plant secondary metabolites in an ecological context. *New Phytologist* 201:733-750.
- Pagel M. 1999. Inferring the historical patterns of biological evolution. *Nature* 401:877-884.
- Pelser PB, de Vos H, Theuring C, Beuerle T, Vrieling K, Hartmann T. 2005. Frequent gain and loss of pyrrolizidine alkaloids in the evolution of *Senecio* section *Jacobaea* (Asteraceae). *Phytochemistry* 66:1285-1295.
- Raunio H, Kuusisto M, Juvonen RO, Pentikäinen OT. 2015. Modeling of interactions between xenobiotics and cytochrome P450 (CYP) enzymes. *Frontiers in Pharmacology* 6:123.
- Renault H, Bassard JE, Hamberger B, Werck-Reichhart D. 2014. Cytochrome P450-mediated metabolic engineering: current progress and future challenges. *Current Opinion in Plant Biology* 19:27-34.
- Soldaat LL, Boutin JP, Derridj S. 1996. Species-specific composition of free amino acids on the leaf surface of four *Senecio* species. *Journal of Chemical Ecology* 22: 1-12.

- Vrieling K, de Vos H, van Wijk CAM. 1993. Genetic analysis of the concentrations of pyrrolizidine alkaloids in *Senecio jacobaea*. *Phytochemistry* 32:1141-1144.
- Wink M. 2003. Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry* 64:3-19.
- Wink M. 2008. Plant secondary metabolism: diversity, function and its evolution. *Natural Product Communications* 3:1205-1216.
- Witte L, Ernst L, Adam H, Hartmann T. 1992. Chemotypes of two pyrrolizidine alkaloid-containing *Senecio* species. *Phytochemistry* 31:559-565.
- Hughes AL. 1994. The evolution of functionally novel proteins after gene duplication. *Proceedings of the Royal Society of London, Series B: Biological Sciences* 256:119-124.
- DePristo MA. 2007. The subtle benefits of being promiscuous: adaptive evolution potentiated by enzyme promiscuity. *HFSP Journal* 1:94-98.
- Bak S, Paquette SM, Morant M, Vinther Moranth A, Saito S, Bjarnholt N, Zagrobelny M, Jorgensen K, Osmani S, *et al.* 2006. Cyanogenic glycosides: a case study for evolution and application of cytochromes P450. *Phytochemistry Reviews* 5:309-329.
- Frey M, Schullehner K, Dick R, Fiesselmann A, Gierl A. 2009. Benzoxazinoid biosynthesis, a model for evolution of secondary metabolic pathways in plants. *Phytochemistry* 70:1645-165