



Universiteit
Leiden
The Netherlands

Patterns and predictors of burn scar outcome in the first 12 months after burn: The patient's perspective

Rashaan, Z.M.; Kwa, K.A.A.; Wal, B.A. van der; Tuinebreijer, W.E.; Zuijlen, P.P.M. van; Breederveld, R.S.

Citation

Rashaan, Z. M., Kwa, K. A. A., Wal, B. A. van der, Tuinebreijer, W. E., Zuijlen, P. P. M. van, & Breederveld, R. S. (2019). Patterns and predictors of burn scar outcome in the first 12 months after burn: The patient's perspective. *Burns*. doi:10.1016/j.burns.2019.03.025


Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/119915>

Note: To cite this publication please use the final published version (if applicable).

AUTHOR QUERY FORM

| | | |
|---|---|---|
|  ELSEVIER | Journal: JBUR Article Number: 5807 | Please e-mail your responses and any corrections to: E-mail: |
|---|---|---|

Dear Author,


Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the ‘Q’ link to go to the location in the proof.

| Location in article | Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof |
|--|--|
| Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 | <p>Please check the presentation of all the tables, and correct if necessary.</p> <p>Please provide the significance of bold values in Table 3.</p> <p>Your article is registered as a regular item and is being processed for inclusion in a regular issue of the journal. If this is NOT correct and your article belongs to a Special Issue/Collection please contact e.wileman@elsevier.com immediately prior to returning your corrections.</p> <p>Please confirm that the provided email is the correct address for official communication, else provide an alternate e-mail address to replace the existing one, because private e-mail addresses should not be used in articles as the address for communication.</p> <p>Author names have been tagged as given names and surnames (surnames are highlighted in teal color). Please confirm if they have been identified correctly.</p> <p>Highlights must be provided as 3-5 bullet points, each bullet point having a maximum of 125 characters. Please rephrase the following highlights or provide new highlights.</p> <p>Please check the hierarchy of section headings, and correct if necessary.</p> <p>Please provide the volume, issue or page range for Ref. [11].</p> <div style="border: 1px solid black; padding: 10px; margin-top: 20px;"> <p>Please check this box or indicate your approval if you have no corrections to make to the PDF file <input style="float: right; width: 40px; height: 25px; margin-left: 10px;" type="checkbox"/></p> </div> |

Thank you for your assistance.

Available online at www.sciencedirect.com**ScienceDirect**journal homepage: www.elsevier.com/locate/burns

1 Highlights

Patterns and predictors of burn scar outcome in the first 12 months post-burn: The patient's perspective

Burns xxx (2019) xxx-xxx

Z.M. Rashaan*, K.A.A. Kwa, M.B.A. van der Wal, W.E. Tuinebreijer, P.P.M. van Zuijlen, R.S. Breederveld


- The POSAS patient total and individual item scores showed a statistically significant improvement of the scar quality in the first 12 months post-burn, except for relief.
- Sex, age, depth of the wound, percentage of TBSA and flame burns were predictors of various POSAS patient items at 3, 6 and 12 months post-burn.
- The effect of the predictors was not the same on the individual POSAS patient items.

UNCORRECTED PROOF

Q6

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/burns

1 Q3 Patterns and predictors of burn scar outcome in the 2 first 12 months post-burn: The patient's perspective

3 Q4 Z.M. Rashaan^{a,b,*}, K.A.A. Kwa^{a,b}, M.B.A. van der Wal^e,
4 W.E. Tuinebreijer^b, P.P.M. van Zuijlen^{c,d}, R.S. Breederveld^{a,b}

5 ^a Department of Surgery, Leiden University Medical Centre, Albinusdreef 2, 2333 ZA Leiden, Leiden, The Netherlands

6 ^b Burn Centre and Department of Surgery, Red Cross Hospital, Vondellaan 13, 1942 LE, Beverwijk, The Netherlands

7 ^c Burn Centre and Department of Plastic and Reconstructive Surgery, Red Cross Hospital, Vondellaan 13, 1942 LE,
8 Beverwijk, The Netherlands

9 ^d Department of Plastic and Reconstructive Surgery and MOVE Research Institute, VU University of Amsterdam, Van
10 der Boechorststraat 7, 1081 BT, Amsterdam, The Netherlands

11 ^e Association of Dutch Burn Centres, Zeestraat 27-29, 1941 AJ, Beverwijk, The Netherlands

ARTICLE INFO

Article history:

Accepted 26 March 2019

Available online xxx

Keywords:

Burn wounds

Burn scars

POSAS

Scar quality

Predictors and patterns

ABSTRACT

Objective: This study aimed to provide insight into the patterns and factors that predict burn scar outcomes at 3, 6 and 12 months post-burn.

Methods: The Patient and Observer Scar Assessment Scale (POSAS) was used to assess the scar formation of each patient. Structural equation modelling was used. The predictor variables used in this study were sex, three age categories, TBSA, depth of the wound and cause of the burn.

Results: The POSAS patient total and individual item scores demonstrated a statistically significant decrease in the first 12 months post-burn, except for the relief item. Male patients had a lower total and items scores (better scar quality) for pain and pruritus compared with female patients. Full thickness burns had a higher scores for pruritus, pliability, thickness and relief compared to the partial-thickness burns. Ages younger than 5 years, higher TBSA values and flame burns were predictors of various POSAS items at 3 and 6 months post-burn. **Conclusion:** The POSAS patient total and individual item scores demonstrated a statistically significant improvement in the scar quality in the first 12 months post-burn, except for the relief. Sex, age, depth of the wound, the percentage of TBSA and flame burns were predictors of various POSAS patient items at 3, 6 and 12 months post-burn.

© 2019 Elsevier Ltd and ISBI. All rights reserved.

1. Introduction

Burn scars have extensive impacts on burn patients in terms of quality of life, functional impairment and physiological

problems [1-3]. Thus, the optimal management of burn scars requires more insight into the factors that influence the severity of burn scars.

To date, sex, age, skin type, location, bacterial colonisation, time to wound healing, type of graft, multiple surgical

Abbreviations: CFI, comparative fit index; POSAS, the Patient and Observer Scar Assessment Scale; SEM, structural equation modelling; LGM, latent growth curve model; RMSEA, the root mean square error of approximation.

* Corresponding author.

E-mail addresses: z.rashaan@vmc.nl (Z.M. Rashaan), kkwa@rkz.nl (K.A.A. Kwa), mbavdwal@hotmail.com (M.B.A. van der Wal), wetuineb@knmng.nl (W.E. Tuinebreijer), pvanzuijlen@rkz.nl (P.P.M. van Zuijlen), rbreederveld@rkz.nl (R.S. Breederveld).
<https://doi.org/10.1016/j.burns.2019.03.025>

0305-4179/© 2019 Elsevier Ltd and ISBI. All rights reserved.

procedures, burn severity and the skin being subjected to stretching have been found to be risk factors for hypertrophic scarring [4-8]. The impacts of burn scars not only entail the appearance of the scar but also involve of its accompanying symptoms. Up to 47% of patients experience pain that is associated with their burn scars [9]. In addition, pruritus was found to still be present in 67% of the burn patients at two years post-burn [10]. It should be noted that different burn scar assessment strategies were used in these studies, and these studies were often limited by the lack of an appropriate tool for evaluating scar outcomes.

Currently, the Patient and Observer Scar Assessment Scale (POSAS) is widely used to assess scar quality [11]. The POSAS consists of observer and patient components and has been found to be a reliable and valid instrument for the assessment of burn scars [12,13]. The POSAS patient scale by Draaijers et al. (version 1.0) incorporates scores for the following six items by using a 10-point rating scale: pain, itch, color, pliability, thickness and relief [12,14]. A high score indicates a worse scar quality. There is a paucity of research investigating the changes in the POSAS scores after burns [15]. Van der Wal et al. described that full thickness wounds and a higher percentage of TBSA were significant predictors of a higher POSAS score, whereas the aetiology and age of the patient had no influence on the scar quality [16]. In addition, POSAS assessment a three months post-burn found to be predictive of final scar quality at twelve months post-burn [17].

The purpose of the present study was to describe the influence of predictors on changes in POSAS patient scores at 3, 6 and 12 months post-burn.

2. Materials and methods

2.1. Recruitment and study population

This retrospective study was performed at the burn centre outpatient clinic at the Red Cross Hospital, Beverwijk in the Netherlands between June 2004 and December 2009. This study was conducted in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The POSAS questionnaire is a standard part of each routine follow-up visit of each of the burn patients in the outpatient clinic at 3, 6 and 12 months at our specialized burn centre. The data of the patients who were admitted to the burn centre and who were subsequently seen at the outpatient clinic at 3, 6 and 12 months post-burn were included in the analysis. In this consecutive sample, the patients who participated in clinical trials for wound or scar treatments were excluded from the study. The parents or caregivers were asked to fill in the POSAS patient component for patients who were under the age of 5 years. Baseline characteristics such as sex, age at the time of burn, the percentage of total body surface area (TBSA), burn depth (partial or full thickness) and the cause of the burn wound (flame or scald) were collected. At our institution, patients with full-thickness burns were operated (skin grafting). Mixed burns (partial and full-thickness) were conservatively treated for approximately 10-14 days. Burn wounds of $>3\text{ cm}^2$ that

were not yet healed, were considered for skin grafting procedures. Partial-thickness burns were treated with topical antiseptics or hydrofibre dressings. This treatment algorithm was chosen because wound healing that takes more than three weeks to complete, is considered to be a risk factor for hypertrophic scar formation [18]. Patients were categorized into the following three age-groups: **<5 years, 5-18 years and >18 years**. The cut-off value of 5 years was chosen because of two reasons. First, the epidemiology of burn wounds tends to be different between children <5 years and older children. In general, scald burns were more common in children who were younger than 5 years compared with older children [19,20]. Second, the POSAS patient scores of this age category were completed by the caregivers, which may influence the outcomes compared with older children who completed the POSAS patient scores on their own. The study location at three months post-burn was defined as the most apparent part of the scar according to the patient. Standard treatment consisted of silicones or pressure garments depending on the location and scar activity. If there was a significant functional impairment during ADL, then there was an indication of reconstruction surgery during the first 12 months post-burn. After 12 months post-burn, an operation was indicated for both functional impairment and esthetical reasons.

2.2. The POSAS

To the best of our knowledge, there is conflicting data in the literature concerning the analysis of the POSAS patient scores. Van der Wal et al. found that the POSAS patient questionnaire was unidimensional. Therefore, the individual and sum of the items of the POSAS patient scores could be used for statistical analysis [21]. Conversely, de Jong et al. found that the POSAS patient questionnaire was multidimensional. Therefore, the only individual POSAS patients scores could be used for statistical analysis [13]. In this study, we used both the individual and sum of the POSAS patient scores for statistical analysis. If the patient was unable to answer the questionnaire, e.g. in the case of children <5 years or in the case of mentally impaired patients, then the caretaker was asked to score the items.

2.3. Study model and statistical analyses

Structural equation modelling (SEM) was performed using the IBM SPSS statistical package AMOSTM 24 [22]. We applied a latent growth curve model (LGM), which was a special application of the SEM with several advantages. Latent growth curve modelling in AMOS was able to accommodate irregularly spaced measurements at the three time points (3, 6 and 12 months post-burn) in our data [23]. In addition, the use of LGM made it possible to assess the fit of the model to the data and to effectively compute the maximum likelihood estimates in our dataset, which was not completed at all three of the time points (Appendix B). The Inter-individual differences in the changes over time were assessed, and group-level statistics such as the mean change rates and mean intercepts were provided. The LGM accounts for the of change (slope curve analysis) at the individual level (patient) and at the group level

(for instance, the depth of the burn wound, sex, etc.). The fit of the LGM was tested. The absolute and comparative fit indices were calculated.

The following predictor variables were entered into the models: sex, age <5 years, age 5-18 years and age >18 years, the percentage of TBSA, depth of the wound and cause of the burn. Our model was based on our earlier study that used the POSAS patient scale to study the influence of time-invariant predictors (such as sex, the percentage of TBSA, wound depth and age categories) on the POSAS scale in the same group of patients [16]. The three different intercept estimates represented the patients' total scores at 3, 6 or 12 months. The time moment of the intercept was dependent on how the time values were coded (0, 1, 3; -1, 0, 2 or -3, -2, 0). The slope estimates represented the patients' rates of change between 3, 6 and 12 months post-burn. Positive intercepts indicated higher POSAS scores at 3, 6 and 12 months post-burn, which thus indicated a worse scar quality compared to that of the reference group. Significant negative slopes in the POSAS scores indicated a slower rate of change in the presented predictor category compared to that in the reference category (for example, flame burns compared to the reference category scald burns).

The correlations between the intercepts and slopes were calculated. A positive value indicated a high initial score at 3 months post-burn with a greater rate of change, whereas a negative correlation indicated a high initial score at 3 months post-burn with a lower rate of change.

The LGM was investigated in a model for the total score and was individually investigated in a model for the six items that were incorporated in the POSAS patient scale, both with and without predictors. The intercept estimate can be interpreted as the influence of the predictors on the POSAS patient scores at 3, 6 and 12 months post-burn. The positive intercepts implied higher POSAS scores compared to the reference category. The slope estimate can be interpreted as the influence of the predictors on the changes in the POSAS scores over time. Positive slopes indicate higher degree of change over time compared to the reference category. A detailed description of the study model and statistical analyses can be found in Appendix D.

3. Results

3.1. Baseline characteristics

A total of 284 children and 190 adult patients were included in this study. The patients' characteristics are shown in Table 1. There were no statistically significant differences in the total TBSA ($p=0.99$, independent t -test), full-thickness burns ($p=0.30$, independent t -test), or surgeries on the evaluated scars ($p=0.53$, chi-square test) that were observed between the groups of patients who completed all three evaluations ($n=157$) and the patients who completed one or two of the evaluations moments ($n=317$).

3.2. The fit indices for the different models

The fit indices for the different models are presented in Appendix B. The fit indices for the model with the total score and the six predictors (Appendix A) revealed the following results: The minimum discrepancy (CMIN) was 6.751 with

Table 1 – Patient characteristics.

| Characteristic | <18 years | ≥18 years |
|--------------------------------------|----------------|------------------|
| Number of patients (%) | 284 (60) | 190 (40) |
| Sex, n (%) | | |
| - Male | 186 (64.5) | 103 (54.2) |
| - Female | 98 (34.5) | 87 (45.8) |
| Age at burn, median in years (range) | 2.5 (0.7-17.8) | 43.2 (18.6-85.6) |
| TBSA, median (range) | | |
| - Total | 7 (0.5-76) | 7.3 (0.5-85) |
| - Full thickness | 1 (0.5-75) | 3 (0.5-60) |
| Cause of the burn (%) | | |
| - Scald | 172 (60.6) | 26 (13.7) |
| - (Flash)flame | 70 (24.6) | 115 (60.5) |
| - Contact | 19 (6.7) | 15 (7.9) |
| - Oil/ fat | 20 (7.0) | 27 (14.2) |
| - Chemical | 0 (0) | 6 (3.2) |
| - Electricity | 3 (1.1) | 0 (0) |
| - Other | 0 (0) | 1 (0.5) |
| Treatment of evaluated scar | | |
| - Conservative treatment, n (%) | 86 (30.3) | 31 (16.3) |
| - Surgery (skin grafting), n (%) | 198 (69.7) | 159 (83.7) |
| Evaluated, n (%) | | |
| - At 3 months post burn | 224 (78.9) | 135 (71.1) |
| - At 6 months post burn | 205 (72.2) | 122 (64.2) |
| - At 12 months post burn | 156 (55.3) | 76 (40.0) |
| Total evaluations, n (%) | | |
| - One evaluation completed | 76 (26.8) | 81 (42.6) |
| - Two evaluations completed | 101 (35.6) | 71 (37.4) |
| - Three evaluations completed | 97 (34.2) | 38 (20.0) |

TBSA: total body surface area.

7 degrees of freedom (df) and a p -value of 0.455. The comparative fit index (CFI) was 1.00. The root mean square error of approximation (RMSEA) was 0.0001 with a confidence interval of 0.0001-0.055. These values of the fit indices agree with a good-to-perfect fit with the total score and the six predictors. All of the models that evaluated the six individual items had a perfect fit. The model with the total score without the six predictors had a moderate fit, and the models with the items of thickness or relief and without the six predictors had a poor fit.

3.3. Patterns of change in the POSAS patient scores

The parameter estimates for the intercept and slopes of the model that evaluated the separate total POSAS patient scale scores and the separate 6 items without the 6 predictors are shown in Table 2. The parameter estimates for the total POSAS scores obtained from the predictor models are presented in Table 2 and Appendix A. Pain had the lowest separate intercept score, which implied that pain had the lowest item score out of the six items in the POSAS at 3 months post-burn. The total score and all of the items (except relief) had significant negative slopes, which implied that the rates of change in the scores showed a decreasing trend. The covariances between the predictor variables of the total POSAS patient scale scores are shown in Appendix C.

3.4. Sex

Male patients had lower total POSAS patient scores at 3, 6 and 12 months post-burn, with no significant difference in the rate

Q1

192
193
194
195
196
197
198
199
200

201

202
203
204
205
206
207
208
209
210
211
212
213
214

215

216
217

Table 2 – Estimates of the intercepts, slopes and covariances between intercepts and slopes of the total scores and items pain, pruritus, color, pliability, thickness and relief without predictors.

| POSAS patient scale | | Intercept | | | | Slope | | | | Covariances | | | |
|---------------------|-----------|-----------|------|-------|--------|----------|------|-------|--------|-------------|------|-------|-------|
| | | Estimate | SE | CR | P | Estimate | SE | CR | p | Estimate | SE | CR | p |
| Total score | 3 months | 29.18 | 0.55 | 53.25 | <0.001 | -2.02 | 0.25 | -8.25 | <0.001 | -2.86 | 4.56 | -0.63 | 0.531 |
| | 6 months | 27.16 | 0.47 | 57.75 | <0.001 | | | | | -1.13 | 2.94 | -0.38 | 0.701 |
| | 12 months | 23.12 | 0.65 | 35.60 | <0.001 | | | | | 2.34 | 8.83 | 0.26 | 0.791 |
| Items: | | | | | | | | | | | | | |
| Pain | 3 months | 2.38 | 0.10 | 22.45 | <0.001 | -0.17 | 0.04 | -4.26 | <0.001 | -0.07 | 0.14 | -0.47 | 0.637 |
| | 6 months | 2.21 | 0.09 | 25.01 | <0.001 | | | | | 0.01 | 0.09 | 0.07 | 0.942 |
| | 12 months | 1.87 | 0.11 | 16.75 | <0.001 | | | | | 0.15 | 0.27 | 0.55 | 0.581 |
| Pruritus | 3 months | 4.54 | 0.13 | 33.81 | <0.001 | -0.49 | 0.06 | -8.09 | <0.001 | -0.67 | 0.26 | -2.55 | 0.011 |
| | 6 months | 4.05 | 0.11 | 36.05 | <0.001 | | | | | -0.10 | 0.18 | -0.54 | 0.592 |
| | 12 months | 3.07 | 0.15 | 19.92 | <0.001 | | | | | 1.05 | 0.51 | 2.06 | 0.039 |
| Color | 3 months | 6.94 | 0.11 | 64.43 | <0.001 | -0.54 | 0.06 | -9.56 | <0.001 | -0.52 | 0.21 | -2.52 | 0.012 |
| | 6 months | 6.40 | 0.09 | 74.02 | <0.001 | | | | | -0.17 | 0.14 | -1.27 | 0.204 |
| | 12 months | 5.32 | 0.14 | 39.26 | <0.001 | | | | | 0.52 | 0.39 | 1.32 | 0.187 |
| Pliability | 3 months | 5.79 | 0.13 | 44.66 | <0.001 | -0.47 | 0.07 | -7.07 | <0.001 | -0.40 | 0.30 | -1.37 | 0.172 |
| | 6 months | 5.33 | 0.11 | 50.73 | <0.001 | | | | | -0.10 | 0.19 | -0.52 | 0.600 |
| | 12 months | 4.40 | 0.16 | 27.59 | <0.001 | | | | | 0.50 | 0.58 | 0.87 | 0.383 |
| Thickness | 3 months | 5.26 | 0.13 | 39.84 | <0.001 | -0.31 | 0.06 | -5.06 | <0.001 | 0.08 | 0.28 | 0.29 | 0.770 |
| | 6 months | 4.96 | 0.11 | 43.61 | <0.001 | | | | | 0.10 | 0.18 | 0.52 | 0.601 |
| | 12 months | 4.34 | 0.16 | 26.80 | <0.001 | | | | | 0.12 | 0.57 | 0.21 | 0.832 |
| Relief | 3 months | 5.08 | 0.13 | 39.10 | <0.001 | -0.09 | 0.06 | -1.34 | 0.179 | -0.35 | 0.29 | -1.19 | 0.236 |
| | 6 months | 5.00 | 0.11 | 47.33 | <0.001 | | | | | -0.07 | 0.19 | -0.37 | 0.715 |
| | 12 months | 4.82 | 0.16 | 30.83 | <0.001 | | | | | 0.48 | 0.57 | 0.85 | 0.397 |

SE, standard error; CR, critical ratio.

of change when compared to female patients. (Table 3) The male patients had lower pain scores at 3 and 6 months post-burn, with an equal rate of change compared to females. Men tended to have lower itch scores at 3 and 6 months post-burn. Nevertheless, the changes in the scores over time were comparable. (Table 4A) Male patients had higher POSAS scores for relief at 3 and 6 months post-burn, with lower pliability scores at 6 and 12 months post-burn. However, the changes in the scores were comparable to those observed in female patients (Table 4B).

3.5. Wound depth

Patients with full thickness burns had higher POSAS patient total scores at 3 months post-burn and a lower rate of change during the first 12 months post-burn compared to patients with partial thickness burns. The total POSAS scores for full thickness and partial thickness burns showed no difference at 12 months post-burn (Table 3). Pruritus scores at 3 months were significantly higher in patients with full thickness burns than those in patients with partial thickness burns. The rate of change in the pruritus scores was significantly lower in patients with full thickness burns (Table 4A). Finally, patients with full thickness burns had significantly higher POSAS scores for pliability, thickness and relief at 3 and 6 months post-burn compared with patients with partial thickness burns (Table 4B).

3.6. Age

There was no significant difference in the total POSAS scores between younger patients or patients who were older than

5 years. However, patients who were younger than 5 years had significantly lower pruritus scores at 12 months post-burn and lower rates of change compared to older patients. (Table 4A) Patients aged below 5 years had higher scar color, pliability and thickness scores at 3 and 6 months post-burn, while patients older than 18 years had a higher scar color scores at 12 months post-burn and a greater change in scores than the younger patients (Table 4B). Patients older than 18 years had higher pain scores at 3, 6 and 12 months post-burn than younger patients, but groups of patients had equal rates of change. (Table 4A)

3.7. Aetiology and percentage of TBSA

The covariances between the predictor variables of the total POSAS patient score are shown in Appendix A and Appendix C. No effects of the percentage of TBSA or cause of burn were found on the total POSAS patient scale. (Table 3) Patients with flame burns generally had significantly higher color scores at 3 and 6 months post-burn. (Table 4A) Patients with a higher percentage of TBSA had higher POSAS score for relief at 3 and 6 months post-burn. (Table 4B) Pruritus scores at 6 and 12 months post-burn were higher in patients with a higher percentage of TBSA values.

4. Discussion

The change in the POSAS patient scale scores was studied between 3 and 6 months post-burn and between 6 and 12 months post-burn. The POSAS patient total score and all of the item scores showed a statistically significant decline in these two time periods, except for the relief item. The greatest

Table 3 – Regression weights and p-values of the POSAS patient scores and the predictors TBSA, burn depth, age category, sex and cause of burn.

| POSAS patient scale total score | | Estimate | SE | CR | p |
|---------------------------------|------------------------|----------|-------|--------|--------|
| Predictors | | | | | |
| Sex: male | Intercept at 3 months | -3.327 | 1.138 | -2.922 | 0.003 |
| | Intercept at 6 months | -3.204 | 0.973 | -3.292 | <0.001 |
| | Intercept at 12 months | -2.959 | 1.332 | -2.222 | 0.026 |
| | Slope | 0.122 | 0.504 | 0.243 | 0.808 |
| Depth: full thickness | Intercept at 3 months | 3.543 | 1.283 | 2.762 | 0.006 |
| | Intercept at 6 months | 1.997 | 1.097 | 1.820 | 0.069 |
| | Intercept at 12 months | -1.095 | 1.501 | -0.730 | 0.466 |
| | Slope | -1.546 | 0.568 | -2.722 | 0.006 |
| Age < 5 years | Intercept at 3 months | 3.130 | 1.664 | 1.881 | 0.060 |
| | Intercept at 6 months | 1.673 | 1.423 | 1.176 | 0.240 |
| | Intercept at 12 months | -1.242 | 1.942 | -0.640 | 0.522 |
| | Slope | -1.458 | 0.735 | -1.984 | 0.047 |
| Age > 18 years | Intercept at 3 months | 0.649 | 1.443 | 0.450 | 0.653 |
| | Intercept at 6 months | 1.229 | 1.234 | 0.996 | 0.319 |
| | Intercept at 12 months | 2.388 | 1.689 | 1.414 | 0.157 |
| | Slope | 0.580 | 0.639 | 0.907 | 0.364 |
| Cause: flame burns | Intercept at 3 months | 1.006 | 1.490 | 0.675 | 0.499 |
| | Intercept at 6 months | 0.840 | 1.272 | 0.661 | 0.509 |
| | Intercept at 12 months | 0.509 | 1.719 | 0.296 | 0.767 |
| | Slope | -0.166 | 0.651 | -0.255 | 0.799 |
| TBSA | Intercept at 3 months | 0.024 | 0.044 | 0.552 | 0.581 |
| | Intercept at 6 months | 0.041 | 0.037 | 1.107 | 0.268 |
| | Intercept at 12 months | 0.076 | 0.051 | 1.486 | 0.137 |
| | Slope | 0.017 | 0.019 | 0.893 | 0.372 |

SE, standard error; CR, critical ratio. Reference categories were female sex, partial thickness burns, age 5-18 years, scald burns. TBSA was a continuous variable in the model.

decline was observed during the longer time period between 6 and 12 months post-burn. The pain item scale presented the lowest decline score, and the color item exhibited the highest decline score. Therefore, the pain and color items had the lowest and highest influences on the total POSAS score, respectively. The low pain scores could be the result of effective medication for pain and/or the result of real low pain values in patients after 3 months post-burn. The high color values represent the importance of color for the patient assessment of his or her scars. Patients with the highest total and item scores presented the lowest changes during the 3 and 6 months post-burn, thus leading to the lowest decline in the total score.

In our study, a strong effect of sex was observed on the total POSAS patient score. Male patients had a better scar quality, which was caused by lower score of pruritus and pain, as well as a better score for pliability and relief compared to the scar quality in female patients. Various studies have demonstrated higher pain-related symptoms in women compared with men [26-28]. Sex role beliefs, pain coping strategies, pain-related expectations and even hormonal factors may possibly explain the difference in pain experience between males and females [29]. In line with our study, two studies observed higher itch intensity scores in women compared to men, although this phenomenon is not well understood [10,30]. Higher pliability

and relief scores in the female group in our study could possibly be explained by the differences in body images between males and females. In general, women have a more negative body image compared to men [31-33]. Dyer et al. observed that women with scars that resulted from accidents or surgeries reported a more negative body image [33].

Patients with full thickness burns had higher total POSAS scores, which were caused by higher scores for the pruritus, pliability, thickness and relief items. Other studies have also described higher itching scores for full thickness burns and grafted wounds [10,30,34]. An increase in both mediators and neuronal damage are thought to contribute to pruritus symptoms in full thickness burns [35]. In our study, pruritus diminished after 3 months post-burn; a finding that has been previously described in other studies [10,16]. Higher POSAS scores for pliability, thickness and relief are explained by the loss of epidermal and dermal structures.

Previous studies have found that the age of the patient does not influence scar behavior [6,16,36]. Our results are consistent with these reports when considering the total POSAS score. However, this is not the case when looking at the separate items. Patients who were aged below 5 years had significantly higher scores for color, pliability and thickness at 3 and 6 months post-burn, and these patients also had significantly less pruritus at 12 months post-burn. The fact that caretakers

Table 4A – Regression weights and *p*-values of the items pain, pruritus and color of the POSAS patient scale from the predictors TBSA, burn depth, age category, sex and cause of burn.

| Items POSAS patient scale | | Pain | | Pruritus | | Color | |
|---------------------------|---------------------|----------|----------|----------|----------|----------|----------|
| Predictors | | Estimate | <i>p</i> | Estimate | <i>p</i> | Estimate | <i>p</i> |
| Sex: male | Intercept 3 months | -0.730 | <0.001 | -0.676 | 0.015 | -0.181 | 0.419 |
| | Intercept 6 months | -0.598 | <0.001 | -0.614 | 0.009 | -0.039 | 0.830 |
| | Intercept 12 months | -0.335 | 0.124 | -0.489 | 0.118 | 0.246 | 0.375 |
| | Slope | 0.132 | 0.117 | 0.062 | 0.611 | 0.143 | 0.217 |
| Depth: full thickness | Intercept 3 months | 0.125 | 0.572 | 0.756 | 0.016 | 0.249 | 0.324 |
| | Intercept 6 months | 0.090 | 0.631 | 0.374 | 0.156 | 0.100 | 0.624 |
| | Intercept 12 months | 0.019 | 0.939 | -0.392 | 0.266 | -0.200 | 0.524 |
| | Slope | -0.035 | 0.708 | -0.383 | 0.005 | -0.150 | 0.250 |
| Age < 5 years | Intercept 3 months | -0.070 | 0.807 | 0.069 | 0.866 | 1.031 | 0.002 |
| | Intercept 6 months | -0.071 | 0.771 | -0.316 | 0.355 | 0.660 | 0.012 |
| | Intercept 12 months | -0.072 | 0.822 | -1.084 | 0.017 | -0.082 | 0.839 |
| | Slope | 0.000 | 0.997 | -0.384 | 0.031 | -0.371 | 0.027 |
| Age > 18 years | Intercept 3 months | 1.282 | <0.001 | -0.480 | 0.175 | 0.008 | 0.978 |
| | Intercept 6 months | 1.330 | <0.001 | -0.311 | 0.294 | 0.360 | 0.115 |
| | Intercept 12 months | 1.427 | <0.001 | -0.009 | 0.983 | 1.065 | 0.003 |
| | Slope | 0.049 | 0.648 | 0.169 | 0.275 | 0.352 | 0.016 |
| Cause: flame burns | Intercept 3 months | 0.303 | 0.240 | -0.065 | 0.858 | 0.951 | 0.001 |
| | Intercept 6 months | 0.313 | 0.150 | -0.027 | 0.929 | 0.527 | 0.025 |
| | Intercept 12 months | 0.333 | 0.236 | 0.049 | 0.903 | -0.320 | 0.371 |
| | Slope | -0.010 | 0.926 | 0.038 | 0.810 | -0.424 | 0.004 |
| TBSA | Intercept 3 months | 0.001 | 0.854 | 0.020 | 0.067 | 0.001 | 0.900 |
| | Intercept 6 months | 0.002 | 0.801 | 0.023 | 0.011 | 0.005 | 0.491 |
| | Intercept 12 months | 0.002 | 0.808 | 0.029 | 0.016 | 0.012 | 0.254 |
| | Slope | 0.000 | 0.948 | 0.003 | 0.506 | 0.004 | 0.405 |

Reference categories were female sex, partial thickness burns, age 5-18 years, scald burns. TBSA was a continuous variable in the model.

321 completed the questionnaires for the patients under 5 years
 322 old may have contributed to the differences in the outcomes
 323 between the age groups. We did not find any studies that
 324 reported the influence of age on color change in burn scars.

325 Furthermore, it should be noted that different studies have
 326 described a negative association between age and hypertrophic
 327 scar formation [37]. This finding is supported by the
 328 decreased proliferation, reepithelization and inflammatory
 329 responses that are observed during wound healing, as well as
 330 the slower epidermal turnover and the different remodeling
 331 phase that are observed in aged individuals [7,37,38]. However,
 332 the present study did not investigate hypertrophic scar
 333 formation. Finally, patients who were above 18 years had
 334 higher pain scores at 3, 6 and 12 months post-burn compared
 335 to patients who were below 18 years.

336 The percentage of TBSA was a predictor for the pruritus,
 337 thickness and relief item scores. The effect of the percentage
 338 of TBSA on pruritus has been well described in various
 339 studies. However, there are conflicting data on the effect of
 340 the percentage of TBSA on the duration of pruritus. Van Loey
 341 et al. described a higher TBSA to be a risk factor for pruritus at
 342 3 months post-burn [10]. The scar tissue modulation and
 343 nerve density which are thought to be highest in the first
 344 6 months post-burn could explain this effect. However, in
 345 line with other studies, we found the effect of the percentage

346 of TBSA to be significant even at 12 months post-burn [30,39].
 347 Furthermore, the effect of full thickness burns and the
 348 percentage of TBSA on itching is different than the effect of
 349 full thickness burns on pain. Pain scores were observed to be
 350 the lowest of all the scored items on the POSAS patient scale.
 351 This could be caused by a different mechanism or by a better
 352 treatment for pain.

353 Scald injuries are more often observed in patients who are
 354 under 5 years, whereas fire/flame burns are observed more
 355 often in older patients. Additionally, more males than females
 356 are admitted to burn centres. Full thickness burns and burns
 357 with a higher percentage of TBSA tend to occur more often in
 358 patients who are older than 18 years. Flame burns are more
 359 often deep dermal or full-thickness burns. Overall, our data are
 360 corroborated by the findings of various epidemiological
 361 studies [19,40].

362 Our study had several limitations. First, the age-related
 363 findings of the patients who were under 5 years should be
 364 interpreted with caution, given that the care givers completed
 365 the questionnaires. Second, no sample size calculation was
 366 performed, given the large number of included patients and
 367 given that the data were retrospectively collected. However, a
 368 sample size calculation could still be relevant, based on the
 369 amount of missing data. Third, the extent of the influence of
 370 the excluded patients on the results of the current study is

Table 4B – Regression weights and p-values of the items pliability, thickness and relief of the POSAS patient scale from the predictors TBSA, burn depth, age category, sex and cause of burn.

| Items POSAS patient scale | | Pliability | | Thickness | | Relief | |
|---------------------------|---------------------|------------|--------|-----------|-------|----------|--------|
| Predictors | | Estimate | p | Estimate | p | Estimate | p |
| Sex: male | Intercept 3 months | -0.393 | 0.138 | -0.136 | 0.617 | -0.520 | 0.051 |
| | Intercept 6 months | -0.545 | 0.012 | -0.269 | 0.252 | -0.435 | 0.045 |
| | Intercept 12 months | -0.847 | 0.011 | -0.537 | 0.114 | -0.264 | 0.421 |
| | Slope | -0.151 | 0.262 | -0.134 | 0.294 | 0.085 | 0.528 |
| Depth: full thickness | Intercept 3 months | 1.151 | <0.001 | 0.797 | 0.009 | 1.076 | <0.001 |
| | Intercept 6 months | 0.682 | 0.005 | 0.463 | 0.080 | 0.863 | <0.001 |
| | Intercept 12 months | -0.254 | 0.497 | -0.204 | 0.595 | 0.438 | 0.238 |
| | Slope | -0.468 | 0.002 | -0.334 | 0.020 | -0.213 | 0.162 |
| Age < 5 years | Intercept 3 months | 1.333 | <0.001 | 0.953 | 0.016 | 0.574 | 0.141 |
| | Intercept 6 months | 0.799 | 0.012 | 0.787 | 0.022 | 0.453 | 0.153 |
| | Intercept 12 months | -0.267 | 0.580 | 0.453 | 0.360 | 0.209 | 0.662 |
| | Slope | -0.533 | 0.007 | -0.167 | 0.369 | -0.122 | 0.536 |
| Age > 18 years | Intercept 3 months | 0.492 | 0.143 | 0.026 | 0.940 | -0.031 | 0.928 |
| | Intercept 6 months | 0.307 | 0.264 | 0.018 | 0.951 | 0.147 | 0.592 |
| | Intercept 12 months | -0.062 | 0.882 | 0.004 | 0.993 | 0.503 | 0.228 |
| | Slope | -0.185 | 0.280 | -0.007 | 0.964 | 0.178 | 0.299 |
| Cause: flame burns | Intercept 3 months | 0.277 | 0.425 | -0.514 | 0.147 | -0.025 | 0.942 |
| | Intercept 6 months | 0.318 | 0.262 | -0.302 | 0.325 | -0.153 | 0.589 |
| | Intercept 12 months | 0.400 | 0.349 | 0.123 | 0.780 | -0.408 | 0.336 |
| | Slope | 0.041 | 0.813 | 0.212 | 0.196 | -0.127 | 0.465 |
| TBSA | Intercept 3 months | 0.004 | 0.667 | 0.021 | 0.042 | 0.026 | 0.010 |
| | Intercept 6 months | 0.006 | 0.508 | 0.015 | 0.094 | 0.023 | 0.006 |
| | Intercept 12 months | 0.008 | 0.541 | 0.003 | 0.821 | 0.016 | 0.200 |
| | Slope | 0.001 | 0.827 | -0.006 | 0.212 | -0.003 | 0.517 |

Reference categories were female sex, partial thickness burns, age 5-18 years, scald burns. TBSA was a continuous variable in the model.

371 unknown, because no data of the excluded patients were
 372 recorded. Fourth, there are conflicting data on whether the
 373 POSAS score is a unidimensional instrument. Therefore, the
 374 scores of the individual items could be summed into a total
 375 score [13,21]. In theory, the POSAS patient questionnaire is
 376 based on a formative model in which the individual items of
 377 the POSAS patient score are causal indicators of the scar
 378 quality. A formative questionnaire could consist of more than
 379 one dimension. Thus the individual items could be summed to
 380 a final score, for example as is done for the Apgar score. Finally,
 381 the included study predictors were obtained from the available
 382 literature, whereas no systematic search was performed. As a
 383 result, there may be predictors that are not included in the
 384 current study, which may be relevant in the context of changes
 385 in the POSAS scores at 3, 6 and 12 months post-burn.

386 **5. Conclusion**

387 This retrospective study, the POSAS patient total and
 388 individual item scores demonstrated a statistically significant
 389 improvement in the first 12 months post-burn, except for the
 390 relief item. Furthermore, sex, age, depth of the wound,
 391 percentage of TBSA and flame burns were predictors of

various POSAS patient items at 3, 6 and 12 months post-burn. 392
 However, the effect of these predictors was not the same for 393
 the individual POSAS patient items. 394

Source of funding 395

This work was conducted without external financial support. 396

Conflict of interest 397

None of the authors have any potential conflicts of interest to 398
 disclose. 399

Updated references 400

[24,25]. 401

Acknowledgements 402

None. 403

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.burns.2019.03.025>.

REFERENCES

- [1] Stavrou, Weissman O, Tessone A, Zilinsky I, Holloway S, Boyd J, et al. Health related quality of life in burn patients—a review of the literature. *Burns* 2014;40(5):788-96.
- [2] Falder, Browne A, Edgar D, Staples E, Fong J, Rea S, et al. Core outcomes for adult burn survivors: a clinical overview. *Burns* 2009;35(5):618-41.
- [3] van Baar E, Essink-Bot ML, Oen IM, Dokter J, Boxma H, van Beeck EF. Functional outcome after burns: a review. *Burns* 2006;32(1):1-9.
- [4] Gangemi N, Gregori D, Berchialla P, Zingarelli E, Cairo M, Bollero D, et al. Epidemiology and risk factors for pathologic scarring after burn wounds. *Arch Facial Plast Surg* 2008;10(2):93-102.
- [5] Bombaro M, Engrav LH, Carrougher GJ, Wiechman SA, Faucher L, Costa BA, et al. What is the prevalence of hypertrophic scarring following burns? *Burns* 2003;29(4):299-302.
- [6] Deitch A, Wheelahan TM, Rose MP, Clothier J, Cotter J. Hypertrophic burn scars: analysis of variables. *J Trauma* 1983;23(10):895-8.
- [7] Butzelaar, Ulrich MM, Mink van der Molen AB, Niessen B, Beelen RH. Currently known risk factors for hypertrophic skin scarring: a review. *J Plast Reconstr Aesthet Surg* 2016;69(2):163-9.
- [8] Lawrence W, Mason ST, Schomer K, Klein MB. Epidemiology and impact of scarring after burn injury: a systematic review of the literature. *J Burn Care Res* 2012;33(1):136-46.
- [9] Li-Tsang W, Lau JC, Chan CC. Prevalence of hypertrophic scar formation and its characteristics among the Chinese population. *Burns* 2005;31(5):610-6.
- [10] Van Loey E, Bremer M, Faber AW, Middelkoop E, Nieuwenhuis MK. Itching following burns: epidemiology and predictors. *Br J Dermatol* 2008;158(1):95-100.
- [11] Mundy R, Miller HC, Klassen AF, Cano SJ, Pusic AL. Patient-reported outcome instruments for surgical and traumatic scars: a systematic review of their development, content, and psychometric validation. *Aesthetic Plast Surg* 2016.
- [12] Draaijers J, Tempelman FR, Botman YA, Tuinebreijer WE, Middelkoop E, Kreis RW, et al. The patient and observer scar assessment scale: a reliable and feasible tool for scar evaluation. *Plast Reconstr Surg* 2004;113(7):1960-5 discussion 6-7.
- [13] DeJong M, Phillips M, Edgar DW, Wood FM. Patient opinion of scarring is multidimensional: an investigation of the POSAS with confirmatory factor analysis. *Burns* 2017;43(1):58-68.
- [14] The patient and observer scar assessment scale (POSAS).
- [15] Tyack, Wasiak J, Spinks A, Kimble R, Simons M. A guide to choosing a burn scar rating scale for clinical or research use. *Burns* 2013;39(7):1341-50.
- [16] van der Wal B, Vloemans JF, Tuinebreijer WE, van de Ven P, van Unen E, van Zuijlen PP, et al. Outcome after burns: an observational study on burn scar maturation and predictors for severe scarring. *Wound Repair Regen* 2012;20(5):676-87.
- [17] Goei, van der Vlies CH, Tuinebreijer WE, van Zuijlen PPM, Middelkoop E, van Baar ME. Predictive validity of short term scar quality on final burn scar outcome using the Patient and Observer Scar Assessment Scale in patients with minor to moderate burn severity. *Burns* 2017;43(4):715-23.
- Cubison C, Pape SA, Parkhouse N. Evidence for the link between healing time and the development of hypertrophic
- [18] scars (HTS) in paediatric burns due to scald injury. *Burns* 2006;32(8):992-9.
- [19] Dokter, Vloemans AF, Beerthuisen GI, van der Vlies CH, Boxma H, Breederveld R, et al. Epidemiology and trends in severe burns in the Netherlands. *Burns* 2014;40(7):1406-14.
- [20] Vloemans F, Dokter J, van Baar ME, Nijhuis I, Beerthuisen GI, Nieuwenhuis MK, et al. Epidemiology of children admitted to the Dutch burn centres. Changes in referral influence admittance rates in burn centres. *Burns* 2011;37(7):1161-7.
- [21] van der Wal B, Tuinebreijer WE, Bloemen MC, Verhaegen PD, Middelkoop E, van Zuijlen PP. Rasch analysis of the Patient and Observer Scar Assessment Scale (POSAS) in burn scars. *Qual Life Res* 2012;21(1):13-23.
- [22] Arbuckle. AMOSTM 22 user's guide. IBM; 2013.
- [23] Byrne M. Structural equation modeling with AMOS. Basic concepts, applications, and programming. Third ed. New York & London: Taylor & Francis Group; 2016.
- [24] DiLalla F. A structural equation modeling overview for medical researchers. *J Dev Behav Pediatr* 2008;29(1):51-4.
- [25] Byrne M, Lam WW, Fielding R. Measuring patterns of change in personality assessments: an annotated application of latent growth curve modeling. *J Pers Assess* 2008;90(6):536-46.
- [26] Andersson I, Ejlertsson G, Leden I, Rosenberg C. Chronic pain in a geographically defined general population: studies of differences in age, gender, social class, and pain localization. *Clin J Pain* 1993;9(3):174-82.
- [27] Forgyas G, Rzewnicki R, Ober AJ, Forgyas DK. Headache in college students: a comparison of four populations. *Headache* 1993;33(4):182-90.
- [28] Sternbach A. Pain and 'hassles' in the United States: findings of the Nuprin pain report. *Pain* 1986;27(1):69-80.
- [29] Fillingim B. Sex, gender, and pain: women and men really are different. *Curr Rev Pain* 2000;4(1):24-30.
- [30] Carrougher J, Martinez EM, McMullen KS, Fauerbach JA, Holavanahalli RK, Herndon DN, et al. Pruritus in adult burn survivors: postburn prevalence and risk factors associated with increased intensity. *J Burn Care Res* 2013;34(1):94-101.
- [31] Cash F, Morrow JA, Hrabosky JI, Perry AA. How has body image changed? A cross-sectional investigation of college women and men from 1983 to 2001. *J Consult Clin Psychol* 2004;72(6):1081-9.
- [32] Smith E, Thompson JK, Raczynski JM, Hilner JE. Body image among men and women in a biracial cohort: the CARDIA Study. *Int J Eat Disord* 1999;25(1):71-82.
- [33] Dyer, Mayer-Eckhard L, White AJ, Alpers GW. The role of scar origin in shaping men's body image. *Am J Mens Health* 2015;9(2):115-23.
- [34] Kuipers C, Bremer M, Braem L, Goemanne AS, Middelkoop E, van Loey NE. Itch in burn areas after skin transplantation: patient characteristics, influencing factors and therapy. *Acta Derm Venereol* 2015;95(4):451-6.
- [35] Goutos, Dziewulski P, Richardson PM. Pruritus in burns: review article. *J Burn Care Res* 2009;30(2):221-8.
- [36] Schwanholt A, Ridgway CL, Greenhalgh DG, Staley MJ, Gaboury TJ, Morress CS, et al. A prospective study of burn scar maturation in pediatrics: does age matter? *J Burn Care Rehabil* 1994;15(5):416-20.
- [37] Mahdavian Delavary, van der Veer M, Ferreira JA, Niessen FB. Formation of hypertrophic scars: evolution and susceptibility. *J Plast Surg Hand Surg* 2012;46(2):95-101.
- [38] Stevenson, Thornton J. Effect of estrogens on skin aging and the potential role of SERMs. *Clin Interv Aging* 2007;2(3):283-97.
- [39] Willebrand, Low A, Dyster-Aas J, Kildal M, Andersson G, Ekselius L, et al. Pruritus, personality traits and coping in long-term follow-up of burn-injured patients. *Acta Derm Venereol* 2004;84(5):375-80.
- [40] Smolle, Cambiaso-Daniel J, Forbes A, Wurzer P, Hundeshagen G, Branski LK, et al. Recent trends in burn epidemiology worldwide: a systematic review. *Burns* 2016.