

Agency for Healthcare Research and Quality (AHRQ) (R01HS018721).

Author Contributions: Both authors participated in the conception, drafting, and approval of the manuscript.

Sponsor's Role: The Pittsburgh Claude D. Pepper Center and Agency for Healthcare Research and Quality played no role in the conception or preparation of the manuscript.

REFERENCES

1. Caljouw MAA, van den Hout WB, Putter H et al. Effectiveness of cranberry capsules to prevent urinary tract infections in vulnerable older persons: A double-blind randomized placebo-controlled trial in long-term care facilities. *J Am Geriatr Soc* 2014;62:103–110.
2. Nicolle LE, Mayhew WJ, Bryan L. Prospective randomized comparison of therapy and no therapy for asymptomatic bacteriuria in institutionalized elderly women. *Am J Med* 1987;83:27–33.
3. Ouslander JG, Schapira M, Schnelle JF et al. Does eradicating bacteriuria affect the severity of chronic urinary incontinence in nursing home residents? *Ann Intern Med* 1995;122:749–754.
4. Nace DA, Drinka PJ, Crnich CJ. Clinical uncertainties in the approach to long term care residents with possible urinary tract infection. *J Am Med Dir Assoc* 2014;15:133–139.
5. AGS Choosing Wisely Workgroup. American Geriatrics Society identifies five things that healthcare providers and patients should question. *J Am Geriatr Soc* 2013;61:622–631.
6. Vance J. AMDA-choosing wisely. *J Am Med Dir Assoc* 2013;14:639–641.
7. Loeb M, Bentley DW, Bradley S et al. Development of minimum criteria for the initiation of antibiotics in residents of LTCFs: Results of a consensus conference. *Infect Control Hosp Epidemiol* 2001;22:120.
8. Stone ND, Ashraf MS, Calder J et al. Surveillance definitions of infections in long term care facilities: Revisiting the McGeer criteria. *Infect Control Hosp Epidemiol* 2012;33:965–977.
9. Drinka P. Limiting the impact of antibiotic resistance in nursing homes. *J Am Med Dir Assoc* 2010;11:537–539.
10. D'Agata E, Loeb MB, Mitchell SL. Challenges in assessing nursing home residents with advanced dementia for suspected urinary tract infections. *J Am Geriatr Soc* 2013;61:62–66.

RESPONSE TO DAVID NACE AND PAUL DRINKA

To the Editor: We appreciate the thoughtful comments of Drs. Nace and Drinka concerning our article.^{1,2} They highlighted our definitions of clinical and strict urinary tract infection (UTI) and concluded that, by using the clinical definition, it is not possible to infer any benefit from cranberry capsules in the prevention of UTI.

We agree that, for many studies, the appropriate criterion standard for diagnosing UTI is detection of the pathogen in the presence of clinical symptoms. A less-rigorous definition can easily lead to overdiagnosis and false conclusions, but residents in long-term care facilities (LTCF) are a vulnerable population, mostly with severe cognitive impairment, multiple chronic comorbidities, functional disabilities, and urinary incontinence. Signs and symptoms of UTI are frequently absent,³ and differentiating asymptomatic from symptomatic UTI in older persons with dementia is difficult and challenging.^{4,5} The use of the criterion standard for diagnosing UTI is not suitable for LTCF residents and would lead to substantial underdiagnosis. As a result, there is no criterion standard in diagnosing UTI in LTCF populations, and most clinical criteria to ascertain UTI in LTCF residents are based on consensus.^{6–8}

To make research in real-world LTCF populations possible, we have chosen a clinical UTI definition. Our clinical

definition is a broad and practical definition, following clinical practice guidelines for LTCF residents^{9,10} and based on the experience of the elderly care physician and nursing staff, which is consistent with a previous study.⁵ Experienced staff can achieve even better diagnostic precision than urine culture.¹¹ There is also recent evidence that mic-turition-related signs and symptoms are predictive of UTI.⁴ Although our clinical definition is different from the strict definition, it closely reflects clinical care in LTCF and adds knowledge to practice guidelines to assist physicians in making decisions. Moreover, our cost-effectiveness analysis showed the relevance of our clinical definition, because clinical UTIs were followed by a significant deterioration in quality of life, survival, care dependency, and costs.¹²

Our study concludes that, for elderly LTCF residents with high UTI risk, taking cranberry capsules twice daily results in a lower incidence of clinical UTI. In daily practice, prevention with cranberry will also reduce antibiotic prescription, including inappropriate prescriptions.

Monique A. A. Caljouw, MSc
Department of Public Health and Primary Care, Leiden
University Medical Center, Leiden, the Netherlands

Wilbert B. van den Hout, PhD
Department of Medical Decision Making, Leiden
University Medical Center, Leiden, the Netherlands

Hein Putter, PhD
Department of Medical Statistics, Leiden University
Medical Center, Leiden, the Netherlands

Wilco P. Achterberg, MD, PhD
Herman J. M. Cools, MD, PhD
Jacobijn Gussekloo, MD, PhD
Department of Public Health and Primary Care, Leiden
University Medical Center, Leiden, the Netherlands

ACKNOWLEDGMENTS

Conflict of Interest: The authors do not have financial or other kind of personal conflicts with this report.

Author Contributions: All authors participated in the conception, drafting, and approval of the report.

Sponsor's Role: None.

REFERENCES

1. Nace DA, Drinka PJ. Cranberry capsules reducing the incidence of what? *J Am Geriatr Soc* 2014;62:1616–1617.
2. Caljouw MAA, van den Hout WB, Putter H et al. Effectiveness of cranberry capsules to prevent urinary tract infections in vulnerable older persons: A double-blind randomized placebo-controlled trial in long-term care facilities. *J Am Geriatr Soc* 2014;62:103–110.
3. D'Agata E, Loeb MB, Mitchell SL. Challenges in assessing nursing home residents with advanced dementia for suspected urinary tract infections. *J Am Geriatr Soc* 2013;61:62–66.
4. Buhr GT, Genao L, White HK. Urinary tract infections in long-term care residents. *Clin Geriatr Med* 2011;27:229–239.
5. Rowe TA, Juthani-Mehta M. Diagnosis and management of urinary tract infection in older adults. *Infect Dis Clin North Am* 2014;28:75–89.
6. Loeb M, Bentley DW, Bradley S et al. Development of minimum criteria for the initiation of antibiotics in residents of long-term-care facilities: Results of a consensus conference. *Infect Control Hosp Epidemiol* 2001;22:120–124.

7. McGeer A, Campbell B, Emori TG et al. Definitions of infection for surveillance in long-term care facilities. *Am J Infect Control* 1991;19:1–7.
8. Juthani-Mehta M, Tinetti M, Perrelli E et al. Interobserver variability in the assessment of clinical criteria for suspected urinary tract infection in nursing home residents. *Infect Control Hosp Epidemiol* 2008;29:446–449.
9. Went P, Achterberg W, Bruggink R et al. Richtlijn Urineweg-Infecties. [Guideline urinary tract infections]. Utrecht, the Netherlands: Verenoo, Dutch Association of Elderly Care Physicians, 2006.
10. High KP, Bradley SF, Gravenstein S et al. Clinical practice guideline for the evaluation of fever and infection in older adult residents of long-term care facilities: 2008 update by the Infectious Diseases Society of America. *J Am Geriatr Soc* 2009;57:375–394.
11. Schmiemann G, Knieh E, Gebhardt K et al. The diagnosis of urinary tract infection: A systematic review. *Dtsch Arztebl Int* 2010;107:361–367.
12. van den Hout WB, Caljouw MAA, Putter H et al. Cost-effectiveness of cranberry capsules to prevent urinary tract infections in long-term care facilities: Economic evaluation alongside a randomized controlled trial. *J Am Geriatr Soc* 2014;62:111–116.

REVIEWING THE SAFETY OF LORATADINE FOR ELDERLY ADULTS: A POTENTIAL SHORTCOMING OF THE 2012 BEERS CRITERIA

To the Editor: Anticholinergic drugs produce a variety of adverse effects, including dry mouth and eyes, constipation, blurred vision, rapid heart rate, dizziness, sedation, confusion, delirium, hallucinations, and cognitive impairment.^{1,2} Furthermore, anticholinergic toxicity has been reported as a common problem in elderly adults, and anticholinergic drug use is closely associated with serious negative outcomes in this population, such as risk of falls, behavioral symptoms (including agitation), and high mortality.^{1,2} Anticholinergic drugs are often mentioned in explicit criteria for inappropriate medication use in older adults, such as the Beers criteria.^{1,3} In 2012, the American Geriatrics Society revised these criteria and included loratadine on the list of potentially inappropriate medication for elderly adults owing to its strong anticholinergic properties; these drugs were to be avoided in cases of lower urinary tract symptoms, benign prostatic hyperplasia, chronic constipation, and cognitive impairment and delirium.³

Loratadine is a second-generation H1 antihistamine, as are levocabastine, azelastine, bilastine, desloratadine, ebastine, cetirizine, fexofenadine, levocetirizine, and rupatadine; the characteristics of second-generation H1 antihistamines make them more effective and safer than first-generation H1 antihistamines (e.g., diphenhydramine, hydroxyzine, promethazine, clemastine, triprolidine). Moreover, first-generation H1 antihistamines are associated with undesirable sedation and anticholinergic side effects.^{4–6} Under the umbrella term “sedation” is a range of conditions including somnolence, impaired concentration, and poor learning ability.⁵ Central nervous system (CNS) and cardiac toxicity have been the most serious adverse effects associated with H1 antihistamines. Cardiac toxicity is rare but is of considerable concern because of the associated risk of death. CNS toxicity produced by first-generation H1 antihistamines is widespread, but even during the years when first-generation H1 antihistamines were widely used, case reports of cardiac toxicity were uncommon, and epidemiological studies involving large sample sizes identified only a few cases of H1 antihistamine-associated ventricular arrhythmias.⁷ Thereafter, sev-

eral second-generation H1 antihistamines, the use of which was devoid of any associated cardiac toxicity and significant CNS toxicity,^{5,7} became the H1 antihistamines of choice.⁷

Loratadine is considered a nonsedating antihistamine. At recommended doses (10 mg/d), no significant differences between loratadine and placebo for any measure of cognitive or psychomotor performance, mood, or sedation were observed.^{5,6} In contrast, other performance studies that used higher doses of loratadine (20 and 40 mg) showed significant performance impairment and sedation in some tests (e.g., choice reaction time, adaptive tracking, digit-symbol substitution) in comparison with placebo.⁵ In light of the evidence demonstrating its safety, the inclusion of loratadine in the list of potentially inappropriate medications solely on the basis of the studies that the 2012 revised Beers criteria reference is not justified. Those three studies, which the American Geriatrics Society cited,^{8–10} do not provide any evidence of loratadine being an antihistamine with strong anticholinergic properties. One of these studies⁹ gave loratadine 2 points on the Anticholinergic Risk Scale, indicating that it entails intermediate risk; another study⁸ gave loratadine 0 points, which means that it has no known anticholinergic properties. Moreover, a recent systematic review of anticholinergic risk scales in older adults¹ did not mention that loratadine has strong anticholinergic properties or the potential for serious adverse effects in elderly adults. That review¹ cited two of the three studies that the 2012 Beers criteria referenced;^{8,9} the third study,¹⁰ also used as a reference by the American Geriatrics Society, was not included in the systematic review, probably because it was a narrative review that did not provide any evidence of loratadine possessing strong anticholinergic properties.

That review¹ has great clinical implications because the summarized studies describe different drugs with anticholinergic properties. Although this study was not a systematic review of randomized clinical trials, the data can be applied to clinical practice. The findings of this review provide insights into a broad issue that could have not been addressed through clinical trials. The finding that loratadine is safe for elderly adults is of great consequence to health professionals concerned with providing treatment for people of this age group. In addition, it highlights a probable shortcoming of the Beers criteria, which are currently used as a basis for supporting prescription for elderly adults.³

Márcio Galvão Oliveira, BPharm, PhD
Instituto Multidisciplinar em Saúde, Universidade Federal da Bahia, Bahia, Brazil

Welma Wildes Amorim, MD
Curso de Medicina, Universidade Estadual do Sudoeste da Bahia, Bahia, Brazil

Alfredo José Rodrigues-Neto
Instituto Multidisciplinar em Saúde, Universidade Federal da Bahia, Bahia, Brazil

ACKNOWLEDGMENTS

Conflict of Interest: The editor in chief has reviewed the conflict of interest checklist provided by the authors and