



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Best Practice & Research Clinical Gastroenterology



13

Improving the outcome of fistulising Crohn's disease



Ilse Molendijk, MD, PhD Student^{a,1}, Koen C.M.J. Peeters, MD, PhD, GE-surgeon^{b,2}, Coen I.M. Baeten, MD, PhD, GE-surgeon^{c,3}, Roeland A. Veenendaal, MD, PhD, Gastroenterologist, Head of Department^{a,4}, Andrea E. van der Meulen-de Jong, MD, PhD, Gastroenterologist, Head of IBD Unit^{a,*}

^a Dept. of Gastroenterology and Hepatology (C4-14), Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, The Netherlands

^b Dept. of Surgery (K6), Leiden University Medical Center, P.O. Box 9600, 2300 RC Leiden, The Netherlands

^c Dept. of Surgery, Groene Hart Hospital, P.O. Box 1098, 2800 BB Gouda, The Netherlands

A B S T R A C T

Keywords:

Crohn's disease
Perianal fistulas
Classical treatment
Potential future therapies
Research targets

Fistulas are a frequent manifestation of Crohn's disease (CD) and can result in considerable morbidity. Approximately 35% of all patients with CD will experience one fistula episode during their disease course of which 54% is perianal. The major symptoms of patients with perianal fistulas are constant anal pain, the formation of painful swellings around the anus and continuous discharge of pus and/or blood from the external fistula opening. The exact aetiology of perianal fistulas in CD patients remains unclear, but it is thought that a penetrating ulcer in the rectal mucosa caused by active CD forms an abnormal passage between the epithelial lining of the rectum and the perianal skin. Genetic, microbiological and immunological factors seem to play important roles in this process. Although the incidence of perianal fistulas in patients with CD is quite high, an effective treatment is not yet

* Corresponding author. Tel.: +31 71 526 2915; fax: +31 71 524 8115.

E-mail addresses: i.molendijk@lumc.nl (I. Molendijk), k.c.m.j.peeters@lumc.nl (K.C.M.J. Peeters), coen.baeten@ghz.nl (C.I.M. Baeten), r.a.veenendaal@lumc.nl (R.A. Veenendaal), ae.meulen@lumc.nl (A.E. van der Meulen-de Jong).

¹ Tel.: +31 71 526 5719; fax: +31 71 524 8115.

² Tel.: +31 71 526 2309; fax: +31 71 526 6750.

³ Tel.: +31 182 505420; fax: +31 182 505519.

⁴ Tel.: +31 71 526 2915; fax: +31 71 524 8115.

discovered. In this review all available medical and surgical therapies are discussed and new treatment options and research targets will be highlighted.

© 2014 Elsevier Ltd. All rights reserved.

Introduction

Fistulas in Crohn's disease (CD) are a major problem which can result in considerable morbidity. Approximately 35% of all CD patients have at least one fistula episode. A quarter of the fistulas is between two parts of intestine, 9% is rectovaginal and 13% is different, including fistulas between the intestine and bladder and around a stoma [1]. In case of an enterovesical fistula there may be recurrent polymicrobial urinary tract infections, pneumaturia and faecaluria. When a rectovaginal fistula develops, dyspareunia, malodorous vaginal discharge and recurring episodes of vaginitis can occur. Perianal fistulas are the most common type of fistulas in fistulising CD. The cumulative incidence of perianal fistulas was estimated at 23%–26% after 20 years of CD [1,2]. Patients with perianal fistulas can present with symptoms such as constant anal pain or pain after defecation, (painful) swelling around the anus, continuous (malodorous) discharge of pus and/or blood from the external opening with skin irritation around the anus, fever and even incontinence [3]. In 20–45% of the CD patients a perianal fistula developed before or at the time of diagnosis CD [1,2]. Patients with colonic and active rectal disease have more frequently perianal fistulas compared to patients with isolated ileal or ileocolonic disease [1,2,4–6]. Male gender, age at diagnosis of CD and smoking are other risk factors although data are conflicting [1,2,5,7–9]. The formation of perianal fistulas in CD is based on the presence of a penetrating ulcer in the rectal or anal mucosa resulting in an abnormal granulating connection between the epithelial lining of the rectum or anal canal and the perianal skin [10,11]. However, most perianal fistulas are cryptoglandular fistulas (90%) and are not associated with CD. They originate from the intersphincteric anal glands due to a local infection with abscess formation [12]. Normally the internal sphincter is a barrier for bacterial overgrowth, only chronic infection, CD inflammation or local




Fig. 1. Anatomy of the muscles surrounding the rectum and anal canal. A fistula tract through the lower third of the external muscle is classified as a low perianal fistula; fistulas in the two-thirds above as high perianal fistulas. Figure reproduced with permission from Bemelman from van Koperen et al [93].

trauma can cause abscesses beyond this barrier in the intersphincteric space. When such an abscess increases it will usually drain in two ways: it can either drain through the intersphincteric space downwards to form an intersphincteric fistula erupting into the perianal skin or it can overcome the external sphincteric plane into the ischiorectal fossa resulting in a transsphincteric perianal fistula. In




Fig. 2. Parks classification of perianal fistulas. (a) An intersphincteric fistula situated between the internal and external sphincter and a transsphincteric fistula passing the internal and external sphincter into the ischiorectal fossa. (b) A suprasphincteric fistula passing through the external sphincter above the puborectalis muscle into the ischiorectal fossa and an extrasphincteric fistula with its origin above the puborectalis draining through the pelvic floor into the ischiorectal fossa without passing the internal or external sphincter. Figure reproduced with permission from Schouten from Schouten et al [94].

CD these abscesses are often a complex delta of channels and patients might present with large or multiple abscesses. The exact aetiology of perianal fistulising CD remains unclear, however genetic, microbiological and immunological factors seem to play important roles. The risk haplotype of the carnitine/organic cation transporter (OCTN) on the IBD5 locus (5q31) is associated with penetrating and perianal CD due to altered bacteria killing resulting in inflammation [13]. Furthermore, diminished clearance of intracellular pathogens by autophagy caused by a specific polymorphism in the immunity-related GTPase family M (IRGM) gives an increased risk of penetrating and perianal CD [14]. It is hypothesized that microbiota also contribute to the development of perianal fistulas as faecal diversion leads to long-term improvement [15]. Especially gram-positive microorganisms are present in CD perianal fistulas [16]. The likelihood of spontaneous healing of perianal fistulas is very low as the rectum, where the luminal opening of perianal fistulas in CD is most frequently located, functions as a reservoir for faeces that is pushed into the luminal opening of the fistula resulting in a continuous contamination. More importantly, epithelialization of the fistula tract in patients with CD hinders fistula closing. Although a range of potent drugs and advanced surgical techniques are available nowadays, the treatment of perianal fistulising CD remains challenging. In this review both classical and future treatments for perianal CD will be discussed.

Anatomy (Fig. 1)

The anal canal is approximately 2–4 cm long and is closely related to both the internal and external sphincter that both play an important role in remaining faecal continence. The internal sphincter (inner muscular layer) is a continuation of the circular smooth muscular layer of the rectum and becomes thicker at the anal canal. The external sphincter is formed from the pelvic floor musculature. Because the external sphincter is longer than the internal sphincter it covers the internal sphincter to the anoderm. The surface of the anal canal is formed by the mucosa of the rectum with columnar epithelium (endoderm) and the mucosa of the skin with squamous epithelium (ectoderm). The junction of these two surfaces is the linea dentata or pectinate line. Above the pectinate line the mucosa is folded into columns: the columns and crypts of Morgagni with ducts to the anal glands. These anal glands penetrate into the intersphincteric space.

Classification

Nowadays, the relative simple Parks classification [17] identifying four types of perianal fistulas (intersphincteric, transsphincteric (Fig. 2(a)), suprasphincteric and extrasphincteric (Fig. 2(b))) is often used. This Parks classification is also useful for CD, although more (complicated) tracts can occur in a CD patient. Furthermore, this system does not include other perianal manifestations of CD (e.g. abscesses or strictures). In clinical practice the 'simple' or 'complex' classification combining physical inspection of the perianal area and endoscopy to determine rectal inflammation, is mostly used [6,11]. A simple fistula has its origin below the pectinate line (superficial, low intersphincteric or low transsphincteric) and has a single external opening without evidence of a perianal abscess, a rectovaginal fistula or an anorectal stricture. A complex fistula is high intersphincteric, high transsphincteric, extrasphincteric or suprasphincteric and may have multiple external openings. They may be associated with a rectovaginal fistula, an anorectal stricture, active rectal disease at endoscopy and pain or fluctuation suggesting a perianal abscess.

Diagnosis

Knowledge about the exact route and internal opening of the fistula, its relation to the sphincters and the presence of abscesses, is crucial in the management of perianal fistulising CD. Inspection of the perianal area is the first step in this process. The fistula disease activity can be quantified with the Perianal Disease Activity Index (PDAI) [18] and comprises five categories: discharge, pain/restriction activities, restriction of sexual activity, type of perianal disease and the degree of induration (all 0–4 points). Pelvic magnetic resonance imaging (MRI) (Fig. 3) is accurate in determining the exact route of the fistula, differentiating between a fibrotic and septic fistula and locating abscesses [19–22]. Introducing an endoanal coil receiver results in more detailed images of the location of the internal opening,




Fig. 3. Pelvic magnetic resonance imaging (MRI) of a perianal fistula. (a) Transversal image of a perianal fistula (arrow) after introduction of an endoanal coil receiver and (b) a coronal image of a perianal fistula (arrows). The fistula tract shows a high signal intensity indicating active disease.

the extent of the fistula tract and its relation with the sphincters [23,24]. Anorectal endoscopic ultrasound (EUS) requires expertise, but can be equivalent to pelvic MRI and is less expensive and time consuming [25,26]. Examination under anaesthesia (EUA) has the advantage of the possibility of concomitant drainage of abscesses and placement of non-cutting setons. The presence of related abnormalities is not detectable with digital examination (e.g. high abscesses and sinuses). In addition, MRI predicts patient outcome better than solely EUA [27,28]. The random combination of two of three methods (MRI, EUS and EUA) resulted in a 100% correct classification of perianal fistulising CD [29].

Description of perianal fistulas with older methods such as fistulography and computerized tomography (CT) is obsolete, because using these techniques it is not possible to distinguish between a fibrotic and septic fistula. Furthermore, small secondary tracts may not be visible with fistulography.

Independent of the diagnostic method used, proctosigmoidoscopy should be performed to assess whether the rectum and/or sigmoid is inflamed since active inflammation influences therapy choice and outcome.

Treatment

Treatment of perianal fistula is often indicated. Apart from the inconvenience of discharge from an untreated fistula, development of abscesses, sepsis, incontinence and carcinoma [30,31] have been described. Although a range of medical and surgical options is available nowadays, the treatment of perianal fistulas is still challenging. Achieving complete closure of the fistula tract is a long process with in many cases multiple relapses. Spontaneous closure of complex fistula in CD is rare, though it has been reported that a simple transsphincteric fistula has a spontaneous closure rate up to 50% [32]. Nevertheless the chance of recurrence is up to 60% after two years [33].

Classical medical treatment

Antibiotics

Antibiotics such as ciprofloxacin and metronidazole are broadly used as first-line treatment of perianal fistulas; however, only one randomized, double-blind, placebo-controlled pilot study was

published to evaluate the safety and efficacy of ciprofloxacin and metronidazole as a treatment for active perianal fistulising CD. No significant differences were found after ten weeks of treatment [34]. In addition, re-exacerbations were common after discontinuation of these treatments [34,35]. Monotherapy with antibiotics is therefore not considered to induce complete healing of perianal fistulas.

Immunosuppressants

An open-label study from 2003 showed that perianal fistulas treated with the immunosuppressant azathioprine in combination with eight weeks of antibiotics responded significantly more often than fistulas treated with antibiotics alone [36]. In addition, immunosuppression with azathioprine and 6-mercaptopurine solely has been shown to be effective in the treatment of perianal fistulas (54% response in comparison to 21% in placebo group) [37]. In patients with previous failure or intolerance to 6-mercaptopurine methotrexate seemed an effective option with a 44–56% response rate [38,39]. However, the recurrence rate was high when the dose of methotrexate was tapered or changed to oral administration.

Anti-TNF α agents

Several trials clearly demonstrated the benefit of the anti-tumour necrosis factor alpha (anti-TNF α) agents infliximab, adalimumab and certolizumab pegol for the induction and maintenance of remission in perianal fistulising disease [40–43]. However, a meta-analysis reported no statistically significant difference in the relative risk of fistulas remaining unhealed with anti-TNF α agents versus placebo [44]. On exclusion of short-term follow-up results, the effect of anti-TNF α on fistula healing became significant. Unfortunately, in a retrospective analysis the five-year probability of recurrence of a perianal fistula that was initially healed with infliximab therapy, was estimated at 40.1% [45]. When infliximab was combined with ciprofloxacin a trend to higher response compared to infliximab alone was observed [46]. In the ADAFI trial [47] patients with perianal fistulising CD were treated with adalimumab monotherapy or adalimumab plus ciprofloxacin. After 12 weeks, the treatment with ciprofloxacin was stopped. At that point, a significant higher response and remission rate was reported in the combination group. However, at week 24 no significant difference in fistula healing between the two treatment groups was found. Another possibility is to inject anti-TNF α locally in the fistula tract. In three studies a small number of patients were open-label treated with multiple local injections of infliximab without any reported adverse events of the treatment [48–50]. The remission rate varied among the studies with a sustained remission between 36.4% [49] and 87.5% [50] after approximately one year. However, in the latter study, local injection of infliximab was combined with fistulectomy. Local injections with adalimumab as a treatment for perianal fistulising CD also appeared to be safe [51,52] with remission rates of 75–77.8% [52,53]. No relapse was observed after a mean follow-up time of 17.5 months in one study [52]. Because the results of locally injected anti-TNF α agents are encouraging, it would be worth it to set up randomized controlled studies to evaluate the efficacy of these local treatments for perianal fistulising CD.

5-Aminosalicylic acid, corticosteroids, cyclosporine and tacrolimus

There is no role for 5-aminosalicylic acid and corticosteroids in the treatment of perianal fistulising CD [35]. The same applies to intravenous cyclosporine and oral tacrolimus: clinical response is seen, but transition to oral treatment or stopping the drugs is associated with a high relapse rates [54–56].

Classical surgical treatment

Before elaborating on the optimal surgical treatment technique, it has to be stated that a conservative surgical approach is warranted in most cases: aggressive surgery may result in outcomes that are worse than the CD itself. Faecal incontinence is a feared complication and may occur even after partial sphincter division. Concomitant proctitis has to be taken into account, and therefore, optimal medical therapy to control disease activity is paramount before embarking on surgery.

Fistulotomy and non-cutting setons

The most simple and classical surgical treatment for perianal fistulas is to open the fistula tract widely by fistulotomy and to let the wound heal by granulation (Fig. 4). Fistulotomy in case of simple superficial fistula is successful in up to 80–100% of the cases [11]. Fistulotomy is not preferred in case of a trans- and extra-sphincteric fistula as a part of the sphincters are cut during surgery. For these fistulas a non-cutting seton for initial drainage can best be placed (Fig. 5). This seton will drain the fistula tract and reduce the local inflammation. If the inflammation has diminished, the seton can be removed. However, the majority of the patients need additional surgical therapy, especially when optimal medical treatment does not appear to prevent disease recurrence. Therefore, seton drainage is often a bridge to a more definite surgical treatment.

Mucosal advancement flap (MAF)

When rectal inflammation is limited, the creation of an MAF to cover the internal opening is a good option. This technique can also be applied in case of a rectovaginal fistula with success rates of 54%–71% in two retrospective series [57,58]. The mucosa and submucosa and even sometimes the muscle is mobilized and then advanced over the internal opening. This technically demanding procedure is known to be successful in experienced hands in even up to 71% of CD patients [59]. The majority of the MAF-reports are however limited to patients with cryptoglandular disease, obscuring its clinical value in CD patients. Incontinence may occur in 13% and 9% respectively [60].

Fibrin glue

The success rates of sealing the fistula tract with a mix of both fibrin and thrombin are varying. Fibrin can be retrieved from autologous blood but commercial fibrin adhesives are available as well. Optimistic reports on patients with cryptoglandular disease disclose success rates from 68% to 85% at one year [61,62]. Results in CD are less favourable: a randomised trial including 77 patients with moderate disease were randomised between observation after seton removal and fibrin glue administration [63]. After eight weeks, 38% of the glue patients experienced disease remission whereas only 16% did in the observation group.




Fig. 4. Fistulotomy of a superficial perianal fistula. (a) The fistula is explored with a probe to find both openings and (b) is opened to heal by granulation. Figure reproduced with permission from Bemelman from van Koperen et al [93].



Fig. 5. Non-cutting seton in a perianal fistula to assure drainage and to promote fibrosis of the fistula tract.

Fistula plug

Fistula plugs consist of inert porcine intestinal submucosa that is known to avoid inflammatory reaction after implantation due to its inert nature. After three months the plug is populated with patient's endogenous cells [64,65]. In patients with CD healing rates were 54% [66] without affecting faecal continence [67]. A prospective study was conducted on 73 patients with anorectal fistulas of differing aetiologies. Only eight CD patients were included of which four patients (50%) were successfully managed by plug insertion [61]. Fistula plugs were compared with MAF in two randomised studies, however no CD patients were included. One trial reported poor healing rates in patients with plug insertion (29%) being not statistically different from patients who received an advancement flap (48%) [68]. The other trial was stopped early due dramatic performance of the plug (only 20% success), being much worse than the advancement flap group (88%) [69]. Plug protrusion shortly after surgery is the predominant cause of treatment failure.

Ligation of the intersphincteric fistula tract (LIFT)

This rather new technique was launched by Rojanasakul in 2009 [70] and consists of dissection between the internal and external sphincter up to the level of the fistula tract. There, the fistula is ligated and the rest of the external tract is curetted [71]. The (theoretical) advantage of LIFT over MAF is the complete preservation of sphincter function. LIFT is an emerging technique that has proved to be effective in 57–83% of patients with cryptoglandular disease who had previous unsuccessful treatment [72–74]. There is no randomised comparison with other techniques yet. A small series of 15 CD patients was recently published showing an LIFT healing rate of 67% at 12 months of follow-up without any development of faecal incontinence [75].

Video assisted anal fistula treatment (VAAFT)

VAAFT is a novel technique that has hardly been studied. In only one study VAAFT was applied in CD patients [76]. In 11/13 included patients VAAFT was successfully completed: visualization of the fistula tract and/or side tracts was done using a fistuloscope after which the internal opening was correctly localized under direct vision with irrigation. Additionally, MAFs were created in all patients and in four of 11 patients also faecal diversion was performed. Success rate after nine months was 82% (9/11 patients) with no deterioration in continence. Although promising, this technique still has to prove its value in future studies to come.

Faecal diversion

Only if other options have proved to be ineffective, the construction of a stoma can ameliorate symptoms related to perianal fistulas [77]. Patients should know that many of these stomas, although often considered as a temporary measure, turn out to be definite [78–80].

Proctectomy

If all treatments fail, proctectomy may be considered. These patients suffer from debilitating abscess formation, colonic disease, complex high fistulas and/or anal stenosis. Unfortunately, also proctectomy has the risk of bad wound healing and perianal sinus development in almost 50% of the cases [81].

Potential future therapies

Vedolizumab

Recently Sandborn et al [82] reported the results of the first induction and maintenance trials of vedolizumab, an $\alpha 4\beta 7$ -integrin antibody. Although the number of patients with fistulas at baseline in this trial was quite low (165/1115; 14.8%) and the number of patients available for evaluation at week 52 even lower ($n = 57$), vedolizumab every eight weeks resulted in a significant higher closure rate (41.2%) compared to placebo (11.1%) ($p = 0.03$) [82].

Mesenchymal stromal cells

A new experimental approach to the treatment of perianal fistula in CD is cellular therapy with mesenchymal stromal cells (MSCs). MSCs are non-haematopoietic precursors of connective tissue cells with immunomodulatory and tissue regenerative properties, making them a potential therapeutic option for inflammatory disorders, including fistulising CD. Several studies indicate soluble factors released by MSCs, such as COX-2-dependent prostaglandin E2 and nitric oxide, to be the main mechanism of MSC-mediated immune suppression. Others believe that cell–cell contact is needed to achieve inhibition of for instance T cell proliferation [83–85]. In the past years several reports on clinical trials using MSCs derived from bone marrow or adipose tissue as a treatment for perianal fistulising CD have been published [86–89]. A phase I trial in which nine fistulas were injected with autologous MSCs derived from adipose tissue, demonstrated 75% healed fistulas after eight weeks without the occurrence of serious adverse events [86]. This trial was followed by a phase II study of the same group [87] in which they included 49 adult patients with complex cryptoglandular ($n = 35$) and CD ($n = 14$) perianal fistulas. Treatment consisted of either local application of fibrin glue or fibrin glue plus 20 million autologous MSCs derived from adipose tissue. Evaluation took place eight weeks after the treatment and in case of no healing, a second dose of fibrin glue or fibrin glue plus 40 million MSCs was injected. Although the majority of the patients had perianal fistulas based on cryptoglandular disease, overall healing of the fistulas was observed in 16% of the patients who received only fibrin glue versus 71% of the patients who received fibrin glue with additional MSC treatment. An Italian group treated ten patients with refractory fistulising CD every four weeks with a local injection of 20 million autologous MSCs derived from bone marrow as long as the autologous MSCs were available [88]. In 70% of the patients, fistulas closed completely without any serious adverse events. Furthermore, rectal mucosal healing was observed and PDAI was improved. The first paper on the treatment of perianal fistulising CD with allogeneic MSCs was published last year [89]. Patients were treated locally with 20 million MSCs derived from adipose tissue per fistula. If the fistula was not healed after 12 weeks, another 40 million cells were injected. Although the investigators and patients were not blinded, 69.2% of the patients showed a reduction in the number of draining fistulas after 24 weeks. In more than half of the patients complete closure of the treated fistula was achieved and in 30% of the patients complete healing of all fistulas was observed after 24 weeks. However, all these studies were not powered for efficacy analysis, so double-blind randomized controlled trials with a sufficient number of patients are needed in order to prove the actual efficacy of MSCs in the treatment of fistulising CD.

Research targets to improve outcome

Targeting immune cells and genetics

Further unravelling and understanding of predisposing genes and the presence of immune cells and their function in CD fistulas are needed to elucidate the next steps in improving outcome of fistulising CD.

Targeting microbiota

Antibiotics seem partially effective in the treatment of perianal fistulising CD suggesting that microbiota contribute to the development of perianal fistulas. Supporting this hypothesis, faecal diversion leads to long-term improvement of this disease [15]. However, until now only one paper is published on microorganisms found in perianal fistulising CD [16]. It could well be that the bacteria that colonize perianal fistulas in CD are not sensitive to metronidazole or ciprofloxacin. In addition, patients carrying a variant of the nucleotide-binding oligomerization domain 2 (NOD2)/caspase recruitment domain 15 (CARD15) tend to respond less to metronidazole or ciprofloxacin than patients with NOD2/CARD15 wild-type [90]. The feasibility of individualized antibiotic therapy should be investigated to improve the outcome of perianal fistulising CD.

Targeting myofibroblasts

Gut myofibroblasts are important players in intestinal tissue damage via the release of matrix metalloproteinases (MMPs) and intestinal wound healing via tissue inhibitors of metalloproteinases (TIMPs). Important for wound healing is the ability of myofibroblasts to migrate. In CD fistulas their migration ability is impaired because of an increased release of TNF α by myofibroblasts. In addition, MMP activity is increased in CD patients resulting in tissue damage and thereby formation of fistulas. Anti-TNF agents enhance migration of myofibroblasts and the increase the production of TIMP-1 resulting in the healing of fistulas by reducing MMP activity [15,91,92]. However, recurrence rates after anti-TNF treatment with consecutive healing are high [45]. Local injections of TIMP-1 could possibly improve outcome and therefore be a target for future investigation.

Targeting epithelial mesenchymal transition (EMT)

On the other hand, the diminished migration capacity of myofibroblasts in CD fistulas may induce EMT to repair mucosal defects by forming a new epithelial layer at the site of tissue damage. However, epithelial cells that undergo EMT become mobile because they lack tight intercellular adhesions, and could therefore be crucial in the pathogenesis of CD fistula development. Evidence for EMT was observed near the luminal origin of enteroenteric, enterovesical, enterocutaneous and perianal fistulas [92], suggesting that EMT-interference could be a novel treatment approach for all types of CD fistulas.

Summary

Although the medical treatment and surgical approaches are improving, the high recurrence rate of perianal fistulas indicates that there is an ongoing need for effective therapies for CD patients with perianal fistulas. Perianal fistulas in CD require specialised and dedicated gastroenterologists and surgeons to optimise treatment outcome. Surgery is often mandatory, but mostly not successful enough and may even deteriorate functional outcome. The increasing number of surgical techniques calls for well-designed clinical trials to establish their true clinical relevance in the treatment of perianal fistulising CD. Apart from surgical expertise a specialized IBD gastroenterology department is necessary to improve patient outcome. In addition, new medical treatment options such as vedolizumab and mesenchymal stromal cells are tested for safety and efficacy in the treatment of perianal fistulising CD. Future targets to improve outcome include genetics, present immune cells, microbiota, myofibroblasts and EMT. For now, a multimodal approach with drugs to keep luminal disease in

remission and to eliminate septic foci together with surgical treatment to reduce symptoms with preservation of faecal continence is advised.

Practice points

- Knowledge about the type, location and route of the fistula and the presence of related abscesses is crucial in the management of perianal fistulising CD and should be obtained with MRI, EUS and/or EUA;
- Spontaneous closure of complex fistula in CD is rare;
- A multimodal approach with both drugs and surgery is recommended in the treatment of complex perianal fistulising CD;
- Achieving complete closure of the fistula tract is a long process with in many cases multiple relapses.

Research agenda

- The safety and efficacy of new therapeutic options for perianal fistulising CD need to be established;
- Future targets to improve outcome include genetics, microbiota, myofibroblasts and EMT;
- New surgical techniques (LIFT, VAAFT) to eradicate perianal fistulas need to be verified in patients with CD.

Conflict of interest

Authors confirm that there are no conflicts of interest.

Acknowledgement

This work was supported by Digest Science Foundation, 2009. Authors confirm that Digest Science Foundation had no role in the collection, analysis and interpretation of data and in the writing of the manuscript.

References

- *[1] Schwartz DA, Loftus Jr EV, Tremaine WJ, Panaccione R, Harmsen WS, Zinsmeister AR, et al. The natural history of fistulizing Crohn's disease in Olmsted County, Minnesota. *Gastroenterology* 2002 Apr;122(4):875–80.
- [2] Hellers G, Bergstrand O, Ewerth S, Holmstrom B. Occurrence and outcome after primary treatment of anal fistulae in Crohn's disease. *Gut* 1980 Jun;21(6):525–7.
- [3] Kamm MA, Ng SC. Perianal fistulizing Crohn's disease: a call to action. *Clin Gastroenterol Hepatol* 2008 Jan;6(1):7–10.
- [4] Lockhart-Mummery HE. Symposium. Crohn's disease: anal lesions. *Dis Colon Rectum* 1975 Apr;18(3):200–2.
- [5] Ingle SB, Loftus Jr EV. The natural history of perianal Crohn's disease. *Dig Liver Dis* 2007 Oct;39(10):963–9.
- [6] Bell SJ, Williams AB, Wiesel P, Wilkinson K, Cohen RC, Kamm MA. The clinical course of fistulating Crohn's disease. *Aliment Pharmacol Ther* 2003 May 1;17(9):1145–51.
- [7] Lapidus A, Bernell O, Hellers G, Lofberg R. Clinical course of colorectal Crohn's disease: a 35-year follow-up study of 507 patients. *Gastroenterology* 1998 Jun;114(6):1151–60.
- [8] Devaraj B, Khabbazi S, Cosman BC. Recent smoking is a risk factor for anal abscess and fistula. *Dis Colon Rectum* 2011 Jun;54(6):681–5.
- [9] Cosnes J, Cattan S, Blain A, Beaugerie L, Carbonnel F, Parc R, et al. Long-term evolution of disease behavior of Crohn's disease. *Inflamm Bowel Dis* 2002 Jul;8(4):244–50.
- *[10] Van AG, Dignass A, Reinisch W, van der Woude CJ, Sturm A, De VM, et al. The second European evidence-based consensus on the diagnosis and management of Crohn's disease: special situations. *J Crohns Colitis* 2010 Feb;4(1):63–101.

- [11] Sandborn WJ, Fazio VW, Feagan BG, Hanauer SB. AGA technical review on perianal Crohn's disease. *Gastroenterology* 2003 Nov;125(5):1508–30.
- [12] Parks AG. Pathogenesis and treatment of fistula-in-ano. *Br Med J* 18-2-1961;1(5224):463–9.
- [13] Vermeire S, Pierik M, Hlavaty T, Claessens G, van SN, Joossens S, et al. Association of organic cation transporter risk haplotype with perianal penetrating Crohn's disease but not with susceptibility to IBD. *Gastroenterology* 2005 Dec;129(6):1845–53.
- [14] Latiano A, Palmieri O, Cucchiara S, Castro M, D'Inca R, Guariso G, et al. Polymorphism of the IRGM gene might predispose to fistulizing behavior in Crohn's disease. *Am J Gastroenterol* 2009 Jan;104(1):110–6.
- [15] Tozer PJ, Whelan K, Phillips RK, Hart AL. Etiology of perianal Crohn's disease: role of genetic, microbiological, and immunological factors. *Inflamm Bowel Dis* 2009 Oct;15(10):1591–8.
- [16] West RL, van der Woude CJ, Endtz HP, Hansen BE, Ouwedijk M, Boelens HA, et al. Perianal fistulas in Crohn's disease are predominantly colonized by skin flora: implications for antibiotic treatment? *Dig Dis Sci* 2005 Jul;50(7):1260–3.
- [17] Parks AG, Gordon PH, Hardcastle JD. A classification of fistula-in-ano. *Br J Surg* 1976 Jan;63(1):1–12.
- [18] Irvine EJ. Usual therapy improves perianal Crohn's disease as measured by a new disease activity index. McMaster IBD Study Group. *J Clin Gastroenterol* 1995 Jan;20(1):27–32.
- [19] Beets-Tan RG, Beets GL, van der Hoop AG, Kessels AG, Vliegen RF, Baeten CG, et al. Preoperative MR imaging of anal fistulas: does it really help the surgeon? *Radiology* 2001 Jan;218(1):75–84.
- [20] Lunniss PJ, Armstrong P, Barker PG, Reznek RH, Phillips RK. Magnetic resonance imaging of anal fistulae. *Lancet* 1992 Aug 15;340(8816):394–6.
- [21] Haggert PJ, Moore NR, Shearman JD, Travis SP, Jewell DP, Mortensen NJ. Pelvic and perineal complications of Crohn's disease: assessment using magnetic resonance imaging. *Gut* 1995 Mar;36(3):407–10.
- [22] Barker PG, Lunniss PJ, Armstrong P, Reznek RH, Cottam K, Phillips RK. Magnetic resonance imaging of fistula-in-ano: technique, interpretation and accuracy. *Clin Radiol* 1994 Jan;49(1):7–13.
- [23] DeSouza NM, Hall AS, Puni R, Gilderdale DJ, Young IR, Kmiot WA. High resolution magnetic resonance imaging of the anal sphincter using a dedicated endoanal coil. Comparison of magnetic resonance imaging with surgical findings. *Dis Colon Rectum* 1996 Aug;39(8):926–34.
- [24] Stoker J, Hussain SM, van KD, Elevelt AJ, Lameris JS. Endoanal coil in MR imaging of anal fistulas. *Am J Roentgenol* 1996 Feb;166(2):360–2.
- [25] Orsoni P, Barthet M, Portier F, Panuel M, Desjeux A, Grimaud JC. Prospective comparison of endosonography, magnetic resonance imaging and surgical findings in anorectal fistula and abscess complicating Crohn's disease. *Br J Surg* 1999 Mar;86(3):360–4.
- [26] Buchanan GN, Halligan S, Bartram CI, Williams AB, Tarroni D, Cohen CR. Clinical examination, endosonography, and MR imaging in preoperative assessment of fistula in ano: comparison with outcome-based reference standard. *Radiology* 2004 Dec;233(3):674–81.
- [27] Van BB, Grandin C, Kartheuser A, Hoang P, Mahieu P, Detry R, et al. MRI of complicated anal fistulae: comparison with digital examination. *J Comput Assist Tomogr* 1994 Jan;18(1):87–90.
- [28] Spencer JA, Chapple K, Wilson D, Ward J, Windsor AC, Ambrose NS. Outcome after surgery for perianal fistula: predictive value of MR imaging. *Am J Roentgenol* 1998 Aug;171(2):403–6.
- [29] Schwartz DA, Wiersma MJ, Dudiak KM, Fletcher JG, Clain JE, Tremaine WJ, et al. A comparison of endoscopic ultrasound, magnetic resonance imaging, and exam under anesthesia for evaluation of Crohn's perianal fistulas. *Gastroenterology* 2001 Nov;121(5):1064–72.
- [30] Thomas M, Bienkowski R, Vandermeer TJ, Trostle D, Cagir B. Malignant transformation in perianal fistulas of Crohn's disease: a systematic review of literature. *J Gastrointest Surg* 2010 Jan;14(1):66–73.
- [31] Ogawa H, Haneda S, Shibata C, Miura K, Nagao M, Ohnuma S, et al. Adenocarcinoma associated with perianal fistulas in Crohn's disease. *Anticancer Res* 2013 Feb;33(2):685–9.
- [32] Halme L, Sainio AP. Factors related to frequency, type, and outcome of anal fistulas in Crohn's disease. *Dis Colon Rectum* 1995 Jan;38(1):55–9.
- [33] Makowiec F, Jehle EC, Starlinger M. Clinical course of perianal fistulas in Crohn's disease. *Gut* 1995 Nov;37(5):696–701.
- [34] Thia KT, Mahadevan U, Feagan BG, Wong C, Cockeram A, Bitton A, et al. Ciprofloxacin or metronidazole for the treatment of perianal fistulas in patients with Crohn's disease: a randomized, double-blind, placebo-controlled pilot study. *Inflamm Bowel Dis* 2009 Jan;15(1):17–24.
- [35] Vavricka SR, Rogler G. Fistula treatment: the unresolved challenge. *Dig Dis* 2010;28(3):556–64.
- [36] Dejaco C, Harter M, Waldhoer T, Miehsler W, Vogelsang H, Reinisch W. Antibiotics and azathioprine for the treatment of perianal fistulas in Crohn's disease. *Aliment Pharmacol Ther* 2003 Dec;18(11–12):1113–20.
- [37] Pearson DC, May GR, Fick GH, Sutherland LR. Azathioprine and 6-mercaptopurine in Crohn disease. A meta-analysis. *Ann Intern Med* 1995 Jul 15;123(2):132–42.
- [38] Mahadevan U, Marion JF, Present DH. Fistula response to methotrexate in Crohn's disease: a case series. *Aliment Pharmacol Ther* 2003 Nov 15;18(10):1003–8.
- [39] Soon SY, Ansari A, Yaneza M, Raouf S, Hirst J, Sanderson JD. Experience with the use of low-dose methotrexate for inflammatory bowel disease. *Eur J Gastroenterol Hepatol* 2004 Sep;16(9):921–6.
- [40] Sands BE, Anderson FH, Bernstein CN, Chey WY, Feagan BG, Fedorak RN, et al. Infliximab maintenance therapy for fistulizing Crohn's disease. *N Engl J Med* 2004 Feb 26;350(9):876–85.
- [41] Present DH, Rutgeerts P, Targan S, Hanauer SB, Mayer L, Van Hogezaand RA, et al. Infliximab for the treatment of fistulas in patients with Crohn's disease. *N Engl J Med* 1999 May 6;340(18):1398–405.
- [42] Schreiber S, Lawrance IC, Thomsen OO, Hanauer SB, Bloomfield R, Sandborn WJ. Randomised clinical trial: certolizumab pegol for fistulas in Crohn's disease – subgroup results from a placebo-controlled study. *Aliment Pharmacol Ther* 2011 Jan;33(2):185–93.
- [43] Colombel JF, Schwartz DA, Sandborn WJ, Kamm MA, D'Haens G, Rutgeerts P, et al. Adalimumab for the treatment of fistulas in patients with Crohn's disease. *Gut* 2009 Jul;58(7):940–8.

- *[44] Ford AC, Sandborn WJ, Khan KJ, Hanauer SB, Talley NJ, Moayyedi P. Efficacy of biological therapies in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol* 2011 Apr;106(4):644–59 [quiz].
- [45] Bouguen G, Siproudhis L, Gizard E, Wallenhorst T, Billioud V, Bretagne JF, et al. Long-term outcome of perianal fistulizing Crohn's disease treated with infliximab. *Clin Gastroenterol Hepatol* 2013 Aug;11(8):975–81.
- [46] West RL, van der Woude CJ, Hansen BE, Felt-Bersma RJ, van Tilburg AJ, Drapers JA, et al. Clinical and endosonographic effect of ciprofloxacin on the treatment of perianal fistulae in Crohn's disease with infliximab: a double-blind placebo-controlled study. *Aliment Pharmacol Ther* 2004 Dec;20(11–12):1329–36.
- [47] Dewint P, Hansen BE, Verhey E, Oldenburg B, Hommes DW, Pierik M, et al. Adalimumab combined with ciprofloxacin is superior to adalimumab monotherapy in perianal fistula closure in Crohn's disease: a randomised, double-blind, placebo controlled trial (ADAFI). *Gut* 2014 Feb;63(2):292–9.
- [48] Poggioli G, Laureti S, Pierangeli F, Rizzello F, Ugolini F, Gionchetti P, et al. Local injection of infliximab for the treatment of perianal Crohn's disease. *Dis Colon Rectum* 2005 Apr;48(4):768–74.
- [49] Asteria CR, Ficari F, Bagnoli S, Milla M, Tonelli F. Treatment of perianal fistulas in Crohn's disease by local injection of antibody to TNF-alpha accounts for a favourable clinical response in selected cases: a pilot study. *Scand J Gastroenterol* 2006 Sep;41(9):1064–72.
- [50] Alessandrini L, Kohn A, Cosentino R, Marrollo M, Papi C, Monterubbianesi R, et al. Local injection of infliximab in severe fistulating perianal Crohn's disease: an open uncontrolled study. *Tech Coloproctol* 2011 Dec;15(4):407–12.
- [51] Poggioli G, Laureti S, Pierangeli F, Bazzi P, Coscia M, Gentilini L, et al. Local injection of adalimumab for perianal Crohn's disease: better than infliximab? *Inflamm Bowel Dis* 2010 Oct;16(10):1631.
- [52] Tonelli F, Giudici F, Asteria CR. Effectiveness and safety of local adalimumab injection in patients with fistulizing perianal Crohn's disease: a pilot study. *Dis Colon Rectum* 2012 Aug;55(8):870–5.
- [53] Maggi L, Capone M, Giudici F, Santarlasci V, Querci V, Liotta F, et al. CD4+CD161+ T lymphocytes infiltrate Crohn's disease-associated perianal fistulas and are reduced by anti-TNF-alpha local therapy. *Int Arch Allergy Immunol* 2013; 161(1):81–6.
- [54] Present DH, Lichtiger S. Efficacy of cyclosporine in treatment of fistula of Crohn's disease. *Dig Dis Sci* 1994 Feb;39(2): 374–80.
- [55] Sandborn WJ, Present DH, Isaacs KL, Wolf DC, Greenberg E, Hanauer SB, et al. Tacrolimus for the treatment of fistulas in patients with Crohn's disease: a randomized, placebo-controlled trial. *Gastroenterology* 2003 Aug;125(2):380–8.
- [56] Egan LJ, Sandborn WJ, Tremaine WJ. Clinical outcome following treatment of refractory inflammatory and fistulizing Crohn's disease with intravenous cyclosporine. *Am J Gastroenterol* 1998 Mar;93(3):442–8.
- [57] Hull TL, Fazio VW. Surgical approaches to low anovaginal fistula in Crohn's disease. *Am J Surg* 1997 Feb;173(2):95–8.
- [58] Kodner IJ, Mazon A, Shemesh EI, Fry RD, Fleshman JW, Birnbaum EH. Endorectal advancement flap repair of rectovaginal and other complicated anorectal fistulas. *Surgery* 1993 Oct;114(4):682–9.
- [59] Joo JS, Weiss EG, Nogueras JJ, Wexner SD. Endorectal advancement flap in perianal Crohn's disease. *Am Surg* 1998 Feb; 64(2):147–50.
- *[60] Soltani A, Kaiser AM. Endorectal advancement flap for cryptoglandular or Crohn's fistula-in-ano. *Dis Colon Rectum* 2010 Apr;53(4):486–95.
- [61] Cintron JR, Park JJ, Orsay CP, Pearl RK, Nelson RL, Abcarian H. Repair of fistulas-in-ano using autologous fibrin tissue adhesive. *Dis Colon Rectum* 1999 May;42(5):607–13.
- [62] Park JJ, Cintron JR, Orsay CP, Pearl RK, Nelson RL, Sone J, et al. Repair of chronic anorectal fistulae using commercial fibrin sealant. *Arch Surg* 2000 Feb;135(2):166–9.
- [63] Grimaud JC, Munoz-Bongrand N, Siproudhis L, Abramowitz L, Senejoux A, Vitton V, et al. Fibrin glue is effective healing perianal fistulas in patients with Crohn's disease. *Gastroenterology* 2010 Jun;138(7):2275–81.
- [64] Champagne BJ, O'Connor LM, Ferguson M, Orangio GR, Schertzer ME, Armstrong DN. Efficacy of anal fistula plug in closure of cryptoglandular fistulas: long-term follow-up. *Dis Colon Rectum* 2006 Dec;49(12):1817–21.
- [65] Badylak SF. The extracellular matrix as a scaffold for tissue reconstruction. *Semin Cell Dev Biol* 2002 Oct;13(5):377–83.
- [66] Garg P, Song J, Bhatia A, Kalia H, Menon GR. The efficacy of anal fistula plug in fistula-in-ano: a systematic review. *Colorectal Dis* 2010 Oct;12(10):965–70.
- *[67] O'Riordan JM, Datta I, Johnston C, Baxter NN. A systematic review of the anal fistula plug for patients with Crohn's and non-Crohn's related fistula-in-ano. *Dis Colon Rectum* 2012 Mar;55(3):351–8.
- [68] van Koperen PJ, Bemelman WA, Gerhards MF, Janssen LW, van Tets WF, van Dalsen AD, et al. The anal fistula plug treatment compared with the mucosal advancement flap for cryptoglandular high transsphincteric perianal fistula: a double-blinded multicenter randomized trial. *Dis Colon Rectum* 2011 Apr;54(4):387–93.
- [69] Ortiz H, Marzo J, Ciga MA, Oteiza F, Armendariz P, de MM. Randomized clinical trial of anal fistula plug versus endorectal advancement flap for the treatment of high cryptoglandular fistula in ano. *Br J Surg* 2009 Jun;96(6):608–12.
- [70] Rojanasakul A. LIFT procedure: a simplified technique for fistula-in-ano. *Tech Coloproctol* 2009 Sep;13(3):237–40.
- [71] Schouten WR, Gosselink MP, Thijsse S, van OR. Intersphincteric ligation of perianal fistula. *Ned Tijdschr Geneesk* 2013; 157(43):A6505.
- [72] Ooi K, Skinner I, Croxford M, Faragher I, McLaughlin S. Managing fistula-in-ano with ligation of the intersphincteric fistula tract procedure: the Western Hospital experience. *Colorectal Dis* 2012 May;14(5):599–603.
- [73] Bleier JL, Moloo H, Goldberg SM. Ligation of the intersphincteric fistula tract: an effective new technique for complex fistulas. *Dis Colon Rectum* 2010 Jan;53(1):43–6.
- [74] Shanwani A, Nor AM, Amri N. Ligation of the intersphincteric fistula tract (LIFT): a sphincter-saving technique for fistula-in-ano. *Dis Colon Rectum* 2010 Jan;53(1):39–42.
- [75] Gingold DS, Murrell ZA, Fleshner PR. A prospective evaluation of the ligation of the intersphincteric tract procedure for complex anal fistula in patients with Crohn disease. *Ann Surg*; 2013 Dec 26 [Epub ahead of print].
- [76] Schwandner O. Video-assisted anal fistula treatment (VAAFT) combined with advancement flap repair in Crohn's disease. *Tech Coloproctol* 2013 Apr;17(2):221–5.
- *[77] Kasperek MS, Glatzle J, Temeltcheva T, Mueller MH, Koenigsrainer A, Kreis ME. Long-term quality of life in patients with Crohn's disease and perianal fistulas: influence of fecal diversion. *Dis Colon Rectum* 2007 Dec;50(12):2067–74.

- [78] Galandiuk S, Kimberling J, Al-Mishlab TG, Stromberg AJ. Perianal Crohn disease: predictors of need for permanent diversion. *Ann Surg* 2005 May;241(5):796–801.
- [79] Mueller MH, Geis M, Glatzle J, Kasparek M, Meile T, Jehle EC, et al. Risk of fecal diversion in complicated perianal Crohn's disease. *J Gastrointest Surg* 2007 Apr;11(4):529–37.
- [80] Yamamoto T, Allan RN, Keighley MR. Effect of fecal diversion alone on perianal Crohn's disease. *World J Surg* 2000 Oct;24(10):1258–62.
- [81] Yamamoto T, Bain IM, Allan RN, Keighley MR. Persistent perineal sinus after proctocolectomy for Crohn's disease. *Dis Colon Rectum* 1999 Jan;42(1):96–101.
- [82] Sandborn WJ, Feagan BG, Rutgeerts P, Hanauer S, Colombel JF, Sands BE, et al. Vedolizumab as induction and maintenance therapy for Crohn's disease. *N Engl J Med* 2013 Aug 22;369(8):711–21.
- [83] Gonzalez MA, Gonzalez-Rey E, Rico L, Buscher D, Delgado M. Adipose-derived mesenchymal stem cells alleviate experimental colitis by inhibiting inflammatory and autoimmune responses. *Gastroenterology* 2009 Mar;136(3):978–89.
- [84] Duffy MM, Pindjakova J, Hanley SA, McCarthy C, Weidhofer GA, Sweeney EM, et al. Mesenchymal stem cell inhibition of T-helper 17 cell-differentiation is triggered by cell-cell contact and mediated by prostaglandin E2 via the EP4 receptor. *Eur J Immunol* 2011 Oct;41(10):2840–51.
- [85] Ren G, Zhang L, Zhao X, Xu G, Zhang Y, Roberts AI, et al. Mesenchymal stem cell-mediated immunosuppression occurs via concerted action of chemokines and nitric oxide. *Cell Stem Cell* 2008 Feb 7;2(2):141–50.
- [86] Garcia-Olmo D, Garcia-Arraz M, Herreros D, Pascual I, Peiro C, Rodriguez-Montes JA. A phase I clinical trial of the treatment of Crohn's fistula by adipose mesenchymal stem cell transplantation. *Dis Colon Rectum* 2005 Jul;48(7):1416–23.
- [87] Garcia-Olmo D, Herreros D, Pascual I, Pascual JA, Del-Valle E, Zorrilla J, et al. Expanded adipose-derived stem cells for the treatment of complex perianal fistula: a phase II clinical trial. *Dis Colon Rectum* 2009 Jan;52(1):79–86.
- [88] Ciccocioppo R, Bernardo ME, Sgarella A, Maccario R, Avanzini MA, Ubezio C, et al. Autologous bone marrow-derived mesenchymal stromal cells in the treatment of fistulising Crohn's disease. *Gut* 2011 Jun;60(6):788–98.
- [89] de la Portilla F, Alba F, Garcia-Olmo D, Herreras JM, Gonzalez FX, Galindo A. Expanded allogeneic adipose-derived stem cells (eASCs) for the treatment of complex perianal fistula in Crohn's disease: results from a multicenter phase I/IIa clinical trial. *Int J Colorectal Dis* 2013 Mar;28(3):313–23.
- [90] Angelberger S, Reinisch W, DeJaco C, Miehsler W, Waldhoer T, Wehkamp J, et al. NOD2/CARD15 gene variants are linked to failure of antibiotic treatment in perianal fistulating Crohn's disease. *Am J Gastroenterol* 2008 May;103(5):1197–202.
- [91] Di SA, Pender SL, Jackson CL, Prothero JD, Gordon JN, Picariello L, et al. Functional modulation of Crohn's disease myofibroblasts by anti-tumor necrosis factor antibodies. *Gastroenterology* 2007 Jul;133(1):137–49.
- [92] Bataille F, Rohrmeier C, Bates R, Weber A, Rieder F, Brenmoehl J, et al. Evidence for a role of epithelial mesenchymal transition during pathogenesis of fistulae in Crohn's disease. *Inflamm Bowel Dis* 2008 Nov;14(11):1514–27.
- [93] van Koperen PJ, Horsthuis K, Bemelman WA, Stoker J, Slors JF. Perianal fistulas: developments in the classification and diagnostic techniques, and a new treatment strategy. *Ned Tijdschr Geneesk* 2008 Dec 20;152(51–52):2774–80.
- [94] Schouten WR, Zimmerman DD, Meuwissen SG. Gastrointestinal surgery and gastroenterology. XIII. Classification and diagnosis of perianal fistulas. *Ned Tijdschr Geneesk* 2001 Jul 21;145(29):1398–402.