



Universiteit
Leiden
The Netherlands

Effect of Vagus Nerve Integrity on Short and Long-Term Efficacy of Antireflux Surgery

Rijn, S. van; Rinsma, N.F.; Herwaarden-Lindeboom, M.Y.A. van; Ringers, J.; Gooszen, H.G.; Rijn, P.J.J. van; ... ; Masclee, A.A.M.

Citation

Rijn, S. van, Rinsma, N. F., Herwaarden-Lindeboom, M. Y. A. van, Ringers, J., Gooszen, H. G., Rijn, P. J. J. van, ... Masclee, A. A. M. (2016). Effect of Vagus Nerve Integrity on Short and Long-Term Efficacy of Antireflux Surgery. *American Journal Of Gastroenterology*, 111(4), 508-515. doi:10.1038/ajg.2016.42

Version: Not Applicable (or Unknown)
License: [Leiden University Non-exclusive license](#)
Downloaded from: <https://hdl.handle.net/1887/117199>

Note: To cite this publication please use the final published version (if applicable).

Effect of Vagus Nerve Integrity on Short and Long-Term Efficacy of Antireflux Surgery

S. van Rijn, MD^{1,2}, N.F. Rinsma, MD¹, M.Y.A. van Herwaarden-Lindeboom, MD, PhD³, J. Ringers, MD⁴, H.G. Gooszen, MD, PhD⁵, P.J.J. van Rijn, MD⁶, R.A. Veenendaal, MD, PhD⁷, J.M. Conchillo, MD, PhD¹, N.D. Bouvy, MD, PhD² and Adrian A.M. Masclee, MD, PhD¹

- OBJECTIVES:** Vagus nerve injury is a feared complication of antireflux surgery (ARS) that may negatively affect reflux control. The aim of the present prospective study was to evaluate short-term and long-term impact of vagus nerve injury, evaluated by pancreatic polypeptide response to insulin-induced hypoglycemia (PP-IH), on the outcome of ARS.
- METHODS:** In the period from 1990 until 2000, 125 patients with gastroesophageal reflux disease (GERD) underwent ARS at a single center. Before and 6 months after surgery, vagus nerve integrity testing (PP-IH), 24-h pH-monitoring, gastric emptying, and reflux-associated symptoms were evaluated. In 2014, 14–25 years after surgery, 110 patients were contacted again for evaluation of long-term symptomatic outcome using two validated questionnaires (Gastrointestinal Symptom Rating Scale (GSRS) and GERD-Health Related Quality of Life (HRQL)).
- RESULTS:** Short-term follow-up: vagus nerve injury (PP peak ≤ 47 pmol/l) was observed in 23 patients (18%) 6 months after fundoplication. In both groups, a comparable decrease in reflux parameters and symptoms was observed at 6-month follow-up. Postoperative gastric emptying was significantly delayed in the vagus nerve injury group compared with the vagus nerve intact group. Long-term follow-up: patients with vagus nerve injury showed significantly less effective reflux control and a higher re-operation rate.
- CONCLUSIONS:** Vagus nerve injury occurs in up to 20% of patients after ARS. Reflux control 6 months after surgery was not affected by vagus nerve injury. However, long-term follow-up showed a negative effect on reflux symptom control and re-operation rate in patients with vagus nerve injury.

Am J Gastroenterol 2016; 111:508–515; doi:10.1038/ajg.2016.42; published online 15 March 2016

INTRODUCTION

Gastroesophageal reflux disease (GERD) is a common health problem that affects 10–20% of individuals in Western society (1). Antireflux surgery (ARS) is indicated for patients with documented GERD and chronic symptoms refractory to proton-pump inhibitors (PPIs) or not willing to take lifelong PPIs. The aim of ARS is an optimal long-term control of reflux symptoms and reflux signs, without or with minimal side effects.

Despite improvements in surgical techniques such as laparoscopic antireflux surgery (2,3), postoperative complications still occur (4–9). Vagus nerve injury is a well-known complication of ARS. Abdominal vagus function cannot be measured or evaluated

directly. Data on the prevalence of vagus nerve dysfunction after antireflux surgery are therefore based on the presence of symptoms such as diarrhea, nausea, and vomiting (6,10,11). In the past, an indirect method has been developed to measure vagus (dys)function by measuring the response of plasma pancreatic polypeptide (PP) secretion to insulin-induced hypoglycemia, the insulin hypoglycemia (IH)–PP test (12,13).

In a previous study by our group, vagus nerve injury measured by the IH-PP test was observed in 10% of GERD patients after laparoscopic hemi-fundoplication (14). In that study, no significant differences were observed in the short-term postoperative outcome—that is, symptom control, reflux recurrence, and delayed

¹Division of Gastroenterology–Hepatology, NUTRIM Research Institute, Maastricht University Medical Center, Maastricht, The Netherlands; ²Department of General Surgery, Maastricht University Medical Center, Maastricht, The Netherlands; ³Department of Paediatric Surgery, Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht, The Netherlands; ⁴Department of General Surgery, Leiden University Medical Center, Leiden, The Netherlands; ⁵Department of OR/Evidence Based Surgery, Radboud University Medical Center Nijmegen, Nijmegen, The Netherlands; ⁶Department of General Surgery, Lange Land Hospital Zoetermeer, Zoetermeer, The Netherlands; ⁷Department of Gastroenterology and Hepatology, Leiden University Medical Center, Leiden, The Netherlands. **Correspondence:** Adrian A.M. Masclee, MD, PhD, Division of Gastroenterology–Hepatology, NUTRIM Research Institute, Maastricht University Medical Center, P. Debyeilaan 25, 6229 HX Maastricht, The Netherlands. E-mail: a.masclee@mumc.nl

Received 5 August 2015; accepted 24 January 2016

gastric emptying between patients with or without vagus nerve injury. It should, however, be noted that the number of patients in that study (14) as in other studies assessing vagus nerve injury after antireflux surgery was small, and follow-up of patients was only short (14–16).

Data on the impact of vagotomy on outcome of antireflux surgery have been obtained by Oelschlager *et al.* These investigators reported on the effect of intended vagus nerve injury on the outcome of antireflux surgery. During a mean follow-up period of 18 months, vagus nerve damage did not negatively influence the outcome of antireflux surgery, nor that of gastric emptying (17).

It should be beyond any doubt that vagus nerve dysfunction measured by PP response to IH (IH-PP) truly results from injury during ARS and not from other causes. Therefore, the test should be performed both before and after surgery. Up to now, vagus nerve function in ARS has not been systematically and prospectively evaluated in a large cohort.

Aim of our study was to evaluate both short- and long-term impact of vagus nerve injury as evaluated by the IH-PP test on reflux symptoms, reflux control, and quality of life after ARS. We hypothesize that ARS-related vagus nerve injury will have a negative outcome on symptoms, reflux control, gastric emptying, and quality of life both after short- and long-term follow-up. In the nineties of the past century, as part of clinical work-up, we prospectively evaluated vagus nerve function by PP response to IH in patients undergoing ARS at the Leiden University Medical Center. Only data from patients with vagus nerve testing both before and after ARS have been evaluated. At 6 months after surgery, objective and subjective reflux parameters were measured. In 2014, a follow-up of this prospective cohort for long-term efficacy has been performed with questionnaires on reflux symptoms and quality of life.

METHODS

In the period from 1990 until 2000, out of a larger group, 125 patients with GERD underwent ARS at the Leiden University Medical Centre with detailed evaluation on the effect of ARS on several parameters. Patients underwent esophageal manometry, 24-h pH-monitoring, gastric emptying, and vagus nerve integrity tests both prior and 6 months after surgery as clinical evaluation. Symptoms were evaluated using questionnaires. In 2014, up to 25 years after surgery patients were contacted again and two validated questionnaires evaluating typical and atypical GERD symptoms were sent. Of the initial 125 patients 110 were eligible for follow-up (Figure 1).

Operation

Several surgical procedures were performed as procedure of choice from 1990 until 2000. The selection of the procedure was based on time-dependent surgeon's preference, not on patient characteristics. Initially, in the early nineties, the open Belsey Mark IV procedure was standard (18). After laparoscopic surgery became available at the Leiden University Medical Centre in the year 1994, the laparoscopic Nissen (19), and later on the Toupet




Figure 1. Cohort flow diagram.

fundoplication (20), became the procedure of choice based on literature data. In all patients, surgery was indicated based on documented GERD and symptoms refractory to medical therapy including PPIs.

Short-term follow-up

24-h pH-monitoring. We performed 24-h ambulatory intra-esophageal pH-monitoring as previously described (4). Reflux parameters used in this study were the percentage of time with a pH level <4 for the total recording time, for time in the upright, and the supine position. Abnormal acid exposure time was defined as the percentage of time with pH <4 for total time $\geq 4.0\%$, upright time $\geq 4.4\%$, and supine time $\geq 1.2\%$ (21).

Esophageal manometry. Esophageal body and lower esophageal sphincter (LES) pressure recordings were obtained by a polyvinyl assembly with a 5-mm outer diameter incorporating a 6-cm-long sleeve sensor and seven side holes as described previously (9). The outcome variable used in this study was the LES mean pressure measured in mm Hg.

Gastric emptying. Gastric emptying was measured using a radionuclide scintigraphy with a dual-isotope method as previously described (14). Patients underwent gastric emptying testing pre- and postoperatively. Data of gastric emptying for solids were analyzed by determining lag phase, gastric emptying, and gastric half emptying time. The retention for liquids was calculated at 15, 30, 60, and 90 min. Gastric emptying for solids was considered normal if values for lag phase ranged from 13 to 31 min and for emptying rate from 36 to 79% per hour. Gastric emptying for solids was considered delayed if the emptying rate was <35% per hour.

Vagus nerve integrity. As previously described, vagus nerve integrity was measured indirectly by the response of PP to insulin-

Table 1. Grading system to combine severity and frequency of reflux symptoms such as heartburn, fullness, and nausea

	Severe	Moderate	Mild	Absent
Daily	Grade 3	Grade 2	Grade 1	Grade 1
Weekly	Grade 2	Grade 1	Grade 1	Grade 1
Monthly	Grade 1	Grade 1	Grade 1	Grade 0
Infrequent	Grade 0	Grade 0	Grade 0	Grade 0

induced hypoglycemia (14). After an overnight fast, patients were administered a bolus of 0.1 U/kg of insulin (Actrapid, Novo Nordisk Farma BV, Bagsvaerd, Denmark) intravenously. After administration of insulin, blood samples were drawn at regular intervals during 150 min to measure PP and glucose levels. During the first 90 min, blood was drawn with 10 min time intervals. At 90 min, a standard test meal was given to the patients, and blood was withdrawn with 15-min intervals up to 150 min. PP levels were measured by a specific radioimmunoassay, as described previously (22). Nadir blood glucose levels in response to administration of insulin were <2.5 mmol/l in all patients. A peak increment in plasma PP ≤ 47 pmol/l was considered to be compatible with truncal vagotomy. Cutoff values calculated for this IH-PP test were based on data obtained from patients who underwent intentional vagotomy in comparison to a healthy control group (14). All participants in this study underwent vagus nerve integrity testing before and 6 months after surgery.

Symptoms. GERD symptoms were scored by a standardized symptom questionnaire before and 6 months after operation. Symptoms included heartburn, dysphagia, diarrhea, sensation of fullness, abdominal distention, vomiting, nausea, and belching. Severity and frequency of symptoms were combined according to a scoring system as shown in **Table 1**.

Long-term follow-up

In May 2014, two validated GERD-symptom questionnaires, the GERD-Health Related Quality of Life (HRQL) and Gastrointestinal Symptom Rating Scale (GSRS) symptom questionnaire, were sent out to the initial cohort.

We used these quantitative methods to evaluate symptom severity in GERD. Subjects were asked for severity and frequency of gastrointestinal symptoms (burning, bloating, belching, sensation of abdominal fullness, nausea, and pain) 14–25 years after surgery. The questionnaires assess typical and atypical GERD symptoms and patients' quality of life. In the GERD-HRQL questionnaire, patients were asked to indicate on a scale from 0 to 5 which number reflects best the severity of the symptom asked for in the question. The questionnaire has a minimum score of 0 and a maximum score of 45 (23). The GSRS questionnaire has a scale from 0 to 3 for symptom severity and a scale from 0 to 4 for symptom frequency for every question with a minimum of 0 and maximum of 12 points per question (24). All these questions are anchored at the high end with the most negative or lowest intensity feelings

(e.g., extremely unpleasant, not at all) and with opposing terms at the low end (e.g., extremely pleasant, very high and extreme). The survey also contained questions regarding the use of antireflux medication postoperatively and on re-operation.

Statistical analysis

All data are presented as mean and s.e.m. for normally distributed data and as median and interquartiles for abnormally distributed data. Comparisons were performed using the Wilcoxon signed-ranked test, the Mann–Whitney *U* test, the χ^2 test, and a dependent or an independent *t*-test as appropriate. $P < 0.05$ was regarded as significant. All statistical analyses were performed using commercially available computer software (IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.)

RESULTS

Baseline characteristics

The vagus nerve injury group contained 23 patients (11 males, mean age 51 (range: 21–77) years, mean follow-up 19 (range: 14–25) years). The vagus nerve intact group consisted of 102 patients (49 males, mean age 48 (range: 19–73) years, mean follow-up 19 (range: 14–24) years). There were no statistically significant differences between these groups with respect to age, gender, and duration of follow-up. Before surgery, patients underwent gastric emptying testing, esophageal manometry, 24-h pH-monitoring, and vagus nerve integrity testing. Preoperative characteristics with respect to symptoms, acid exposure time, LES pressure, vagus nerve integrity testing, or gastric emptying were not significantly different between the vagus nerve injury and the vagus nerve intact group. The results of the preoperative work-up are listed in **Table 2**.

Short-term outcome

Vagus nerve integrity. Before surgery, none of the 125 patients had evidence of vagus nerve dysfunction. The plasma PP peak increase in response to insulin-induced hypoglycemia had a median value of 143 (100–193) pmol/l. After surgery the median value in the vagus nerve injury group was 17 (5–31) pmol/l compared with 113 (83–170) pmol/l in the vagus nerve intact group (**Figure 2**). Postoperative plasma PP response was compatible with truncal vagotomy (PP peak increase ≤ 47 pmol/l) in 23 patients. With respect to the surgical procedure, 47 patients underwent Nissen fundoplication (13% vagus nerve injury), 47 patients underwent Toupet hemi-fundoplication (17% vagus nerve injury), and 31 underwent BM IV fundoplication (29% vagus nerve injury) No significant difference in the distribution of vagus nerve injury was seen among the type of operations. We divided the patient population into two groups: one group with and one group without vagus nerve injury.

Gastric emptying. Solids

Gastric emptying of solids increased significantly after ARS for patients in the vagus nerve intact group. In the vagus nerve injury group, gastric emptying of solids did not change significantly after

Table 2. Outcomes of 24-h pH-monitoring, esophageal manometry, and gastric emptying in patients with and without vagus nerve injury before and 6 months after antireflux surgery

	Vagus nerve injury			Vagus nerve intact			P ¹	P ²
	Pre	6 months	P	Pre	6 months	P		
<i>pH monitoring^a</i>								
Total acid exposure (%)	9.7 (4.2–12.2)	2.0 (0.3–5.6)	<0.01	8.0 (5.2–14.9)	1.1 (0.3–3.3)	<0.001	NS	NS
Upright acid exposure time (%)	10.2 (4.2–16.4)	3.0 (0.5–5.0)	<0.01	8.6 (5.8–12.1)	1.3 (0.6–3.6)	<0.001	NS	NS
Supine acid exposure time (%)	6.2 (1.7–19.3)	0.0 (0.0–4.6)	0.01	7.1 (1.1–13.9)	0.1 (0.0–1.4)	<0.001	NS	NS
<i>Manometry^b</i>								
Mean LES pressure (mmHg)	11 (9–14)	16 (12–18)	0.01	12 (8–16)	18 (13–22)	<0.001	NS	NS
<i>Gastric emptying^c</i>								
Gastric lag phase (min)	13 (9–17)	11 (7–21)	NS	17 (8–26)	8 (4–14)	<0.001	NS	<0.05
Gastric emptying rate (%/h)	38 (27–49)	33 (26–43)	NS	37 (29–45)	47 (38–57)	<0.001	NS	<0.001
Gastric half emptying time (min)	84 (75–136)	100 (78–147)	NS	97 (75–120)	69 (58–87)	<0.001	NS	0.001
<i>Liquid retention (%) at^c</i>								
15 min	43 (37–47)	48 (30–54)	NS	53 (40–69)	51 (36–60)	NS	NS	NS
30 min	26 (10–37)	28 (21–45)	NS	35 (22–40)	31 (22–38)	NS	NS	NS
60 min	13 (2–26)	17 (14–28)	NS	16 (10–22)	14 (9–18)	NS	NS	<0.05
90 min	6 (0–14)	13 (8–20)	NS	8 (2–13)	7 (2–11)	NS	NS	<0.01

LES, lower esophageal sphincter; NS, not significant.

P¹ Comparison of baseline characteristics of both groups. P² comparison of postoperative outcomes of both groups. Presented as median (25–75%).

^apH-monitoring n=19/63.

^bManometry n=21/72.

^cGastric emptying n=13/51.




Figure 2. Difference between vagus nerve injury and the vagus nerve intact group 6 months after surgery for the hypoglycemia-induced pancreatic polypeptide test. At time 0 min, insulin (0.1 U/kg) was administered.

ARS compared with preoperatively. The groups were not significantly different with respect to emptying parameters in the preoperative state; however, after ARS lag phase, emptying rate and half emptying time were significantly delayed in the patients with vagus nerve injury when compared with the vagus nerve intact group (Figure 3).

Liquids

No differences in gastric emptying of liquids were observed apart from a minor but statistically significant difference in reten-

tion of liquids at 60 and 90 min postoperative between vagus nerve injury and vagus nerve intact patients (Table 2).

24 h pH-monitoring and esophageal manometry

ARS significantly reduced acid exposure time and increased the mean LES pressure 6 months after the procedure (Table 2). No statistically significant differences were found between both groups.

Reflux symptoms

Prior and 6 months after ARS, patients filled in a standard symptom questionnaire. The prevalence of symptoms after ARS was higher in the vagus nerve injury group. However, this did not reach statistical significance (Figure 4).

Long-term outcome

Responders. Of the 110 patients who received questionnaires via a letter, completed questionnaires were received from 71 patients (65%). Thirty-four patients did not respond, four cases were excluded from the study because of an incorrect address, and 1 patient did not fill out the questionnaires correctly (Figure 1). Of these 71 patients, 13 (18%) had vagus nerve injury, and 58 (82%) had an intact vagus nerve.

Heartburn—GERD-HRQL. For the assessment of reflux symptoms >10 years after surgery, the GERD-HRQL questionnaire was




Figure 3. Comparison between vagus nerve injury and vagus nerve intact group for the difference in gastric emptying rate (%/h; left panel) and gastric half emptying time (min; right panel) 6 months post-surgery. * $P < 0.001$.




Figure 4. Difference in symptomatic outcome 6 months after surgery between the vagus nerve injury and the vagus nerve intact group.

used. Comparison between groups showed higher scores in the vagus nerve injury group (Table 3).

Dysphagia, regurgitation, gas-bloating, and cough. Severity and frequency of postoperative dysphagia, regurgitation, gas-bloating, and cough were assessed for both groups using the GSRS questionnaire. Similar results for all symptoms were shown in both groups (Table 3).

Satisfaction. For the assessment of general satisfaction, the GERD-HRQL was used. Three categories for satisfaction, “satisfied”, “neutral”, and “unsatisfied”, were described at the end of the questionnaire. Of the total group of 71 patients that responded 14–25 years after their antireflux surgery, 62% was satisfied, 23% was neutral, and 15% was unsatisfied. In the vagus nerve injury group 33.3% of patients were satisfied, 33.3% neutral and 33.3% unsatisfied. In the vagus nerve intact group 68% were satisfied, 21% neutral and 11% unsatisfied. A statistical significant difference was found between the two groups for the satisfaction rate (Table 3).

Proton-pump inhibitor usage. Of the 71 patients, 42% ($n=30$) were using a PPI after fundoplication. The other 58% ($n=41$) had not used PPIs since surgery. In the vagus nerve injury group, 7 patients (54%) restarted the PPI use; in the vagus nerve

Table 3. Long-term outcome of antireflux surgery in patients with and without vagus nerve injury

	Vagus nerve injury ($n=13$)	Vagus nerve intact ($n=58$)	<i>P</i>
<i>GERD-HRQL questionnaire</i>			
Total score	13 (5–25)	3(0–9)	<0.001
Satisfaction rate (%)	33	68	<0.05
PPI use (%)	54	41	NS
Re-operation rate (%)	54	16	0.007
<i>GSRS questionnaire</i>			
Heartburn	2(0–6)	1(0–3)	NS
Regurgitation	1(0–8)	0(0–2)	NS
Abdominal distention	3(1–6)	1(0–6)	NS
Dysphagia	2(0–5)	1(0–6)	NS
Coughing	2(0–6)	2(0–6)	NS

HRQL, Health Related Quality of Life; GERD, gastroesophageal reflux disease; GSRS, Gastrointestinal Symptom Rating Scale; NS, not significant; PPI, proton-pump inhibitor.

intact group, 24 patients (41%) had recommenced taking PPI (Table 3).

Re-operation rate. Re-operation rate in the vagus nerve injury group was significantly higher: 7 of 13 patients (54%) underwent re-operation. Three patients were operated because of reflux recurrence, 3 patients because of dysphagia, and 1 patient had severe gastroparesis for which a BII resection of the stomach was performed. In patients without vagus nerve injury, 9 out of 58 (16%) patients underwent re-operation. Six patients were operated because of reflux recurrence, 1 patient because of dysphagia, 1 because of a slipped fundoplication, and 1 because of a rotated fundoplication. This difference in the re-operation rate between the two groups was statistically significant (Table 3). In addition, no significant difference in distribution of re-operation rate was seen among the different type of surgical procedures (Belsey Mark IV, laparoscopic Nissen, laparoscopic Toupet).

DISCUSSION

In this short- and long-term follow-up of a prospective cohort, we evaluated the impact of vagus nerve injury on reflux control after antireflux surgery. The prevalence of vagus nerve injury tested by IH-PP was 20% in patients who underwent ARS. Our short-term data (6 months) indicate that symptom control and reflux control were not different among patients with and without vagus nerve injury. Postoperative gastric emptying was significantly delayed in the vagus nerve injury vs. vagus intact group. Concerning the data at 20-year follow-up, a significantly worse outcome with respect to heartburn symptoms, satisfaction rate, and re-operation rate was seen in patients with vagus nerve injury. Our results indicate that vagus nerve injury during antireflux surgery negatively affects long-term reflux control.

Using the IH-PP test, we observed a high prevalence of vagus nerve injury after ARS. In a previous study, we reported on vagus nerve injury after partial fundoplication. In that study, the prevalence of vagus nerve injury after ARS was 10% (14). It should be taken into account that the number of patients in that study was much smaller and that we evaluated only one single surgical technique and therefore that the study may not give an accurate estimate of the prevalence of vagus nerve injury after ARS in general—that is, after all types of ARS. In the present study, three different surgical techniques (BM IV, Toupet, and Nissen) have been employed and a large group of patients was included. In 23 patients, the results were compatible with vagus nerve injury: 13% (6/47) in the Nissen group, 18% (8/47) in the Toupet group, and 29% (9/31) in the BM IV fundoplication group. We did not observe significant differences between surgical procedures with respect to the prevalence of vagus nerve injury.

Risk factors for vagus nerve injury after ARS have not been studied systematically. One may assume that the risk of vagus nerve injury is both procedure and surgeon related as operating and manipulating near the branches of the vagus nerve may easily lead to direct or indirect injury. An intrathoracic approach (BM IV) is technically more challenging and therefore more prone to vagus nerve injury. As mentioned earlier, we did not observe significant differences in vagus nerve dysfunction between the surgical procedures that have been employed in this study, although the percentage of patients with vagus nerve injury after BM IV was higher than after the two other procedures.

Other factors that potentially may increase a risk of vagus nerve injury during ARS include the presence of a large para-esophageal or axial hiatal hernia, previous abdominal surgery, severe esophagitis causing adhesions, poor visibility of the anatomy in the gastroesophageal region, anatomical variations of the vagus nerve, and limited experience with the procedure of the surgeon. Patients included in this study underwent antireflux surgery as a primary procedure and no para-esophageal hiatal hernias were present. Furthermore, all patients were operated in a high volume center by surgeons experienced with the types of operations employed in this study.

The mechanisms by which vagus nerve injury negatively affects the outcome after ARS are not fully understood. The vagus nerve is

known to have an important role in the motor and sensory regulatory functions of the esophagus and stomach. Vagus nerve injury may impair these functions and thereby lead to upper gastrointestinal symptoms.

Previous studies have established an association between vagus nerve injury after ARS and impaired gastric motility, resulting in delayed gastric emptying or even gastroparesis (25–27). In more detail, impairment of the gastric accommodation of the proximal stomach, hypomotility of the antrum (with loss of trituration and retention of solids), and loss of pyloric relaxation (with stasis) have been associated with vagotomy and conditions after gastric surgery (28–34).

Longer retention of food in the stomach leads to higher intra-gastric pressures and thereby may promote reflux even in the presence of a restored esophago-gastric barrier. Recently, Gourcerol *et al.* showed that, in GERD patients with delayed gastric emptying, the number of postprandial reflux events measured with pH-impedance monitoring was increased, with a longer bolus clearance time and more proximal esophageal extension of the refluxate. This was also true for reflux symptoms in patients with delayed gastric emptying compared with patients with normal gastric emptying (35). Rebecchi *et al.* evaluated the association between gastric emptying with long-term reflux control in GERD patients after ARS. These authors showed that delayed gastric emptying was strongly associated with poor reflux control (36). Taken together, these findings indicate that delayed gastric emptying contributes to GERD and that it may negatively influence the effect of ARS.

We observed that gastric emptying improved for solids but not for liquids after ARS in the vagus nerve intact group, whereas it remained unchanged both for liquids and solids in the vagus nerve injury group at 6 months after surgery.

Delayed gastric emptying is seen in up to 40% of patients with GERD (37). Fundoplication is known to accelerate gastric emptying (14,38). In ARS, part of the fundus is utilized for creating the fundoplication and thereby reducing postprandial gastric volume and gastric capacity. This results in a higher proximal gastric tone and reduces duration of postprandial relaxation of the proximal stomach, leading to propulsion of food into the antrum with acceleration in gastric emptying (38–40). Acceleration of gastric emptying after fundoplication may positively influence the antireflux effect of the procedure. The delay in liquid emptying at 60 and 90 min seen after ARS in the vagus nerve injury group vs. vagus nerve intact group is not in line with literature data. It has been indicated that vagotomy leads to a reduction in fundus accommodation and body of the stomach and thereby an acceleration of liquid emptying (26,41–43). In our study, testing of solid and liquid gastric emptying was conducted simultaneously, with potential interference of the solid and liquid components.

We confirmed findings from previous studies and extended there upon: vagus nerve injury did not negatively affect outcomes 6 months after ARS despite the persistence of delayed gastric emptying (14,15,17). However, during long-term follow-up, a worse outcome of reflux control was found in the vagus nerve

injury group. A possible explanation for the difference in control of reflux symptoms during short- and long-term follow-up is not readily available. One option is that the newly created gastroesophageal valve may gradually lose efficacy with respect to its reflux barrier over time (44). Increasing age with loss of compliance and elasticity of tissue may lead to reduced efficacy of the gastroesophageal valve.

The re-operation rate differed significantly between both groups. Over 50% of patients in the vagus nerve injury group have been re-operated. Most patients underwent re-operation because of recurrence of reflux. In one patient from the vagus nerve injury group, severe gastroparesis had developed. This was so debilitating that a partial resection of the stomach had to be performed. Despite high re-operation rates, symptoms associated with GERD persisted in this patient group. The findings of our study suggest that patients who have recurrent reflux symptoms following ARS might benefit from vagal integrity testing before deciding on additional treatment, as a normal outcome of gastric emptying testing does not appear to exclude vagal nerve injury. Therefore, we believe careful and objective evaluation of symptoms and measurement of gastric emptying and PP-IH testing after ARS is warranted before considering re-operation in these patients.

Some limitations to this study should be addressed. The long-term symptomatic outcome cannot be directly compared with the 6-month data because other types of questionnaires were used. Furthermore, not all patients underwent all tests such as gastric emptying test, esophageal manometry, and 24-h pH-monitoring pre- and 6 months after surgery.

The strength of our study is the evaluation of vagus nerve integrity by IH-PP measurements both before and after fundoplication in all 125 patients so that we are certain that the outcome of the test when pointing to vagus nerve dysfunction was related to the surgical procedure. To our best knowledge, our study is the largest to date reporting on vagus nerve dysfunction after ARS with long-term results.

In conclusion: vagus nerve injury occurs in up to 20% of patients who undergo antireflux surgery. Vagus nerve injury does not negatively affect short-term reflux control 6 months after antireflux surgery. However, long-term follow-up showed a negative effect on symptom control and a significantly higher re-operation rate in patients with vagus nerve injury. Preserving the main trunks of the vagus nerve is therefore of utmost importance in order to maintain vagus nerve integrity and contribute to reflux control after ARS.

CONFLICT OF INTEREST

Guarantor of the article: Adrian A.M. Masclee, MD, PhD.

Specific author contributions: S. van Rijn: acquisition, analysis and interpretation of data, and drafting of manuscript. N.F. Rinsma: analysis of data and drafting of manuscript. M.Y.A. van Herwaarden-Lindeboom, J. Ringers, P.J.J. van Rijn, H.G. Gooszen, and R.A. Veenendaal: data acquisition. J.M. Conchillo: Approval of final manuscript. N.D. Bouvy: interpretation of data. A.A.M. Masclee: study concept, interpretation of data, drafting and approval of final manuscript, and overall study supervision.

Financial support: This study was conducted without funding.

Potential competing interests: None.

Study Highlights

WHAT IS CURRENT KNOWLEDGE

- ✓ Antireflux surgery (ARS) is an accepted alternative therapy for patients with symptoms refractory to proton-pump inhibitors (PPIs) or not willing to take lifelong PPIs.
- ✓ Vagus nerve injury is a feared complication of ARS that may negatively affect reflux control.

WHAT IS NEW HERE

- ✓ The prevalence of vagus nerve injury after ARS is 20%.
- ✓ At short-term follow-up, gastric emptying was significantly delayed in the vagus nerve injury group vs. the vagus nerve intact group.
- ✓ At short-term follow-up, reflux control was similar between both groups.
- ✓ Long-term follow-up showed significantly less reflux control and a higher re-operation rate in the vagus nerve injury group.

REFERENCES

1. Dent J, El-Serag HB, Wallander MA *et al.* Epidemiology of gastro-oesophageal reflux disease: a systematic review. *Gut* 2005;54:710–7.
2. Dallemagne B, Weerts JM, Jehaes C *et al.* Laparoscopic Nissen fundoplication: preliminary report. *Surg Laparosc Endosc* 1991;1:138–43.
3. Dallemagne B, Perretta S. Twenty years of laparoscopic fundoplication for GERD. *World J Surg* 2011;35:1428–35.
4. Wang YR, Dempsey DT, Richter JE. Trends and perioperative outcomes of inpatient antireflux surgery in the United States, 1993–2006. *Dis Esophagus* 2011;24:215–23.
5. Broeders JA, Mauritz FA, Ahmed Ali U *et al.* Systematic review and meta-analysis of laparoscopic Nissen (posterior total) versus Toupet (posterior partial) fundoplication for gastro-oesophageal reflux disease. *Br J Surg* 2010;97:1318–30.
6. Papasavas P. Functional problems following esophageal surgery. *Surg Clin North Am* 2005;85:525–38.
7. Broeders JA, Sportel IG, Jamieson GG *et al.* Impact of ineffective oesophageal motility and wrap type on dysphagia after laparoscopic fundoplication. *Br J Surg* 2011;98:1414–21.
8. Kessing BF, Broeders JA, Vinke N *et al.* Gas-related symptoms after antireflux surgery. *Surg Endosc* 2013;27:3739–47.
9. Lindeboom MY, Ringers J, Straathof JW *et al.* The effect of laparoscopic partial fundoplication on dysphagia, esophageal and lower esophageal sphincter motility. *Dis Esophagus* 2007;20:63–8.
10. Kozarek RA, Low DE, Raltz SL. Complications associated with laparoscopic anti-reflux surgery: one multispecialty clinic's experience. *Gastrointest Endosc* 1997;46:527–31.
11. Trus TL, Bax T, Richardson WS *et al.* Complications of laparoscopic paraesophageal hernia repair. *J Gastrointest Surg* 1997;1:221–7. discussion 228
12. Schwartz TW. Pancreatic polypeptide: a hormone under vagal control. *Gastroenterology* 1983;85:1411–25.
13. Schwartz TW, Holst JJ, Fahrenkrug J *et al.* Vagal, cholinergic regulation of pancreatic polypeptide secretion. *J Clin Invest* 1978;61:781–9.
14. Lindeboom MY, Ringers J, van Rijn PJ *et al.* Gastric emptying and vagus nerve function after laparoscopic partial fundoplication. *Ann Surg* 2004;240:785–90.
15. DeVault KR, Swain JM, Wentling GK *et al.* Evaluation of vagus nerve function before and after antireflux surgery. *J Gastrointest Surg* 2004;8:883–8. discussion 888–889.
16. Straathof JW, Ringers J, Masclee AA. Prospective study of the effect of laparoscopic Nissen fundoplication on reflux mechanisms. *Br J Surg* 2001;88:1519–24.
17. Oelschlagel BK, Yamamoto K, Woltman T *et al.* Vagotomy during hiatal hernia repair: a benign esophageal lengthening procedure. *J Gastrointest Surg* 2008;12:1155–62.

18. Skinner DB, Belsey RH. Surgical management of esophageal reflux and hiatus hernia. Long-term results with 1,030 patients. *J Thorac Cardiovasc Surg* 1967;53:33–54.
19. Nissen R. [A simple operation for control of reflux esophagitis]. *Schweiz Med Wochenschr* 1956;86:590–2.
20. Toupet A. [Technic of esophago-gastroplasty with phrenogastropexy used in radical treatment of hiatal hernias as a supplement to Heller's operation in cardiospasm]. *Mem Acad Chir (Paris)* 1963;89:384–9.
21. Masclee AA, de Best AC, de Graaf R *et al*. Ambulatory 24-hour pH-metry in the diagnosis of gastroesophageal reflux disease. Determination of criteria and relation to endoscopy. *Scand J Gastroenterol* 1990;25:225–30.
22. Lamers CB, Diemel CM, van Leer E *et al*. Mechanism of elevated serum pancreatic polypeptide concentrations in chronic renal failure. *J Clin Endocrinol Metab* 1982;55:922–6.
23. Velanovich V. The development of the GERD-HRQL symptom severity instrument. *Dis Esophagus* 2007;20:130–4.
24. Allen CJ, Parameswaran K, Belda J *et al*. Reproducibility, validity, and responsiveness of a disease-specific symptom questionnaire for gastro-esophageal reflux disease. *Dis Esophagus* 2000;13:265–70.
25. Camilleri M, Parkman HP, Shafi MA *et al*. Clinical guideline: management of gastroparesis. *Am J Gastroenterol* 2013;108:18–37.quiz 38.
26. Jamieson GG, Maddern GJ, Myers JC. Gastric emptying after fundoplication with and without proximal gastric vagotomy. *Arch Surg* 1991;126:1414–7.
27. Olinde AJ, Maher JW, McGuigan JE *et al*. The effect of fundoplication with or without proximal gastric vagotomy on gastric emptying and serum gastrin. *Am Surg* 1985;51:690–2.
28. Shafi MA, Pasricha PJ. Post-surgical and obstructive gastroparesis. *Curr Gastroenterol Rep* 2007;9:280–5.
29. Behrns KE, Sarr MG. Diagnosis and management of gastric emptying disorders. *Adv Surg* 1994;27:233–55.
30. Miyano Y, Sakata I, Kuroda K *et al*. The role of the vagus nerve in the migrating motor complex and ghrelin- and motilin-induced gastric contraction in suncus. *PLoS One* 2013;8:e64777.
31. Fich A, Neri M, Camilleri M *et al*. Stasis syndromes following gastric surgery: clinical and motility features of 60 symptomatic patients. *J Clin Gastroenterol* 1990;12:505–12.
32. Azpiroz F, Malagelada JR. Gastric tone measured by an electronic barostat in health and postsurgical gastroparesis. *Gastroenterology* 1987;92:934–43.
33. Bredenoord AJ, Chial HJ, Camilleri M *et al*. Gastric accommodation and emptying in evaluation of patients with upper gastrointestinal symptoms. *Clin Gastroenterol Hepatol* 2003;1:264–72.
34. Lagoo J, Pappas TN, Perez A. A relic or still relevant: the narrowing role for vagotomy in the treatment of peptic ulcer disease. *Am J Surg* 2014;207:120–6.
35. Gourcerol G, Benanni Y, Boueyre E *et al*. Influence of gastric emptying on gastro-esophageal reflux: a combined pH-impedance study. *Neurogastroenterol Motil* 2013;25:800–e634.
36. Rebecchi F, Allaix ME, Giaccone C *et al*. Gastric emptying as a prognostic factor for long-term results of total laparoscopic fundoplication for weakly acidic or mixed reflux. *Ann Surg* 2013;258:831–6.discussion 836–837.
37. Gonlachanvit S, Maurer AH, Fisher RS *et al*. Regional gastric emptying abnormalities in functional dyspepsia and gastro-oesophageal reflux disease. *Neurogastroenterol Motil* 2006;18:894–904.
38. Vu MK, Straathof JW, v d Schaar PJ *et al*. Motor and sensory function of the proximal stomach in reflux disease and after laparoscopic Nissen fundoplication. *Am J Gastroenterol* 1999;94:1481–9.
39. Vu MK, Ringers J, Arndt JW *et al*. Prospective study of the effect of laparoscopic hemifundoplication on motor and sensory function of the proximal stomach. *Br J Surg* 2000;87:338–43.
40. Lindeboom MY, Vu MK, Ringers J *et al*. Function of the proximal stomach after partial versus complete laparoscopic fundoplication. *Am J Gastroenterol* 2003;98:284–90.
41. Van Hee R, Mistiaen W, Block P. Gastric emptying of liquids after highly selective vagotomy for duodenal ulcer. *Hepatogastroenterology* 1989;36:92–6.
42. Stadaas JO. Effect of vagotomy on gastric motility. *Scand J Gastroenterol Suppl* 1976;42:85–8.
43. Gleysteen JJ, Burdeshaw JA, Hallenbeck GA. Gastric emptying of liquids after different vagotomies and pyloroplasty. *Surg Gynecol Obstet* 1976;142:41–8.
44. Robinson B, Dunst CM, Cassera MA *et al*. 20 years later: laparoscopic fundoplication durability. *Surg Endosc* 2014;29:2520–4.