

Higher functionality of extracellular vesicles isolated using size-exclusion chromatography compared to ultracentrifugation

Mol, E.A.; Goumans, M.J.; Doevendans, P.A.; Sluijter, J.P.G.; Vader, P.

Citation

Mol, E. A., Goumans, M. J., Doevendans, P. A., Sluijter, J. P. G., & Vader, P. (2017). Higher functionality of extracellular vesicles isolated using size-exclusion chromatography compared to ultracentrifugation. *Nanomedicine: Nanotechnology, Biology And Medicine, 13*(6), 2061-2065. doi:10.1016/j.nano.2017.03.011

Version:	Not Applicable (or Unknown)
License:	Leiden University Non-exclusive license
Downloaded from:	https://hdl.handle.net/1887/94814

Note: To cite this publication please use the final published version (if applicable).





BASIC SCIENCE

Nanomedicine: Nanotechnology, Biology, and Medicine 13 (2017) 2061–2065

Short Communication

Nanotechnology, Biology, and Medicine

nanomedjournal.com

Higher functionality of extracellular vesicles isolated using size-exclusion chromatography compared to ultracentrifugation

Emma A. Mol, MSc^a, Marie-José Goumans, PhD^b, Pieter A. Doevendans, PhD, MD^{a,c,d}, Joost P.G. Sluijter, PhD^{a,c,d}, Pieter Vader, PhD^{e,*}

^aDepartment of Cardiology, Laboratory of Experimental Cardiology, University Medical Center Utrecht, Utrecht, the Netherlands

^bDepartment of Molecular Cell Biology, Leiden University Medical Center, Leiden, the Netherlands

^cUMC Utrecht Regenerative Medicine Center, University Medical Center, Utrecht, the Netherlands

^dNetherlands Heart Institute (ICIN), Utrecht, the Netherlands

^eLaboratory of Clinical Chemistry and Haematology, University Medical Center Utrecht, Utrecht, the Netherlands

Received 12 January 2017; accepted 19 March 2017

Abstract

Extracellular vesicles (EVs) are nano-sized, lipid bilayer-enclosed particles involved in intercellular communication. EVs are increasingly being considered as drug delivery vehicles or as cell-free approach to regenerative medicine. However, one of the major challenges for their clinical application is finding a scalable EV isolation method that yields functional EVs. Although the golden standard for EV isolation is ultracentrifugation (UC), a recent study suggested that isolation using size-exclusion chromatography (SEC) yielded EVs with more intact biophysical properties. Whether this also leads to differences in functionality remained to be investigated. Therefore, we investigated possible differences in functionality of cardiomyocyte progenitor cell-derived EVs isolated using UC and SEC. Western blot analysis showed higher pERK/ERK ratios in endothelial cells after stimulation with SEC-EVs compared to UC-EVs, indicating that SEC-EVs bear higher functionality. Therefore, we propose to use SEC-EVs for further investigation of EVs' therapeutic potential. Further optimization of isolation protocols may accelerate clinical adoption of therapeutic EVs.

© 2017 Elsevier Inc. All rights reserved.

Key words: Extracellular vesicles; Exosomes; Ultracentrifugation; Size-exclusion chromatography; Functionality

Extracellular vesicles (EVs) are nano-sized endogenous messengers containing a plethora of biological cargo including proteins and RNA, reflecting the content of the secreting cell. By mediating intercellular communication, EVs can influence

The authors declare no conflicts of interest.

E-mail address: pvader@umcutrecht.nl (P. Vader).

recipient cell behavior, and affect physiological and pathological processes.¹⁻³ For this reason, EVs are increasingly being considered for therapeutic purposes, including cell-free approaches for regenerative medicine and drug delivery.⁴⁻⁶ The interest in using EVs for cardiac therapy increased after it became clear that the beneficial effects of stem cell therapy after a myocardial infarction (MI) were mainly due to paracrine actions.⁷ EVs were identified to be the major component of the stem cell secretome responsible for the observed increase in cardiac function.⁸ Therefore, using EVs as an off-the-shelf therapeutic may circumvent some of the drawbacks of cell based therapy, such as cell survival, retention, rejection, and the use of replicating cells.

One of the major challenges for implementation of EVs as therapeutics is the development of a scalable, reproducible, and standardizable isolation method that results in an acceptable yield of EVs. To date, the most common EV isolation method is differential ultracentrifugation (UC). This method relies on

http://dx.doi.org/10.1016/j.nano.2017.03.011 1549-9634/© 2017 Elsevier Inc. All rights reserved.

Please cite this article as: Mol EA, et al, Higher functionality of extracellular vesicles isolated using size-exclusion chromatography compared to ultracentrifugation. *Nanomedicine: NBM* 2017;13:2061-2065, http://dx.doi.org/10.1016/j.nano.2017.03.011

Funding sources: EAM is supported by the Project SMARTCARE-II of the BioMedical Materials Institute, co-funded by the ZonMw-TAS program (#116002016), the Dutch Ministry of Economic Affairs, Agriculture and Innovation and the Netherlands CardioVascular Research Initiative (CVON): the Dutch Heart Foundation, Dutch Federations of University Medical Centers, the Netherlands Organization for Health Research and Development, and the Royal Netherlands Academy of Sciences. PV is funded by a VENI fellowship (#13667) from the Netherlands Organisation for Scientific Research (NWO).

^{*}Corresponding author at: Department of Clinical Chemistry and Haematology, University Medical Center Utrecht, Utrecht 3584 CX, the Netherlands.



Figure 1. Schematic overview of EV isolation protocols used in this study. Abbreviations: CM = conditioned medium, SEC = size-exclusion chromatography, UC = ultracentrifugation.

sedimentation at high speed for separating EVs from other (extra)cellular components. Although the UC protocol is relatively straightforward, it is also time consuming, and may yield aggregated EVs after pelleting.^{9,10} Furthermore, UC isolation results in low and operator-dependent yields and EVs can be damaged due to shearing forces, as a result of centrifugation at high speeds.^{11,12} An additional method for EV isolation, based on ultrafiltration and size-exclusion chromatography (SEC) to separate EVs from other media components, was recently suggested by Nordin et al.¹³ EVs isolated using chromatography (SEC-EVs) are more intact than EVs isolated using UC (UC-EVs), likely due to the absence of centrifugation at high speeds. Whether this also leads to differences in functionality of SEC- and UC-EVs remained to be investigated.

Methods

A detailed version of the Methods is available online under Supplementary Material.

Results

In order to investigate whether the isolation protocol affects EV functionality, EVs derived from cardiomyocyte progenitor cells (CPCs) were isolated using UC and SEC. CPCs are being intensively investigated for cardiac-related therapies, and CPC-derived EVs have previously been shown to bear pro-angiogenic properties.¹⁴⁻¹⁶ A schematic representation of the UC and SEC isolation protocols used in this study is shown in Figure 1.

First, EV yield was compared by quantification of EV protein content and number of particles, as shown in Figure 2, *A*. No significant differences in total EV protein or particle yield between UC and SEC were found. This is in contrast with previous observations showing that SEC isolation results in a higher EV yield compared to UC isolation.¹³ This may be explained by variation between cell types or due to slight differences in UC or SEC isolation procedure (e.g., rotor/filter type or pore size). Next, UC-EVs and SEC-EVs were characterized based on size distribution, morphology, and the presence or absence of protein markers. Figure 2, B shows a representative size distribution profile of UC-EVs and SEC-EVs based on Nanoparticle Tracking Analysis. SEC-EVs had a smaller size distribution with the highest peak at approximately 90 nm, compared to a broader size range for UC-EVs, peaking at approximately 100 nm. Transmission electron microscopy analysis showed no major morphological differences between UC-EVs and SEC-EVs, as both preparations contained both smaller and larger vesicles (Figure 2, C). Western blot analysis revealed that both UC-EVs and SEC-EVs were enriched for EV marker proteins Alix and CD63, but not TSG101 (Figure 2, C). Although the expected band for Alix (96 kDa) was present for both UC-EVs and SEC-EVs, an extra band at 90 kDa was observed in the UC-EV preparation. The presence of double bands for Alix may be explained by differential phosphorylation status, as Alix is known to have multiple phosphorylation sites.¹⁷ Why this second band was exclusively found in UC-EVs remains unclear, but might suggest a different vesicle sub-class or activation status. The endoplasmatic reticulum protein calnexin was only detected in the cell lysate, confirming the absence of contamination with other membrane compartments in EVs. β-Actin was found in similar levels in EVs and cell lysate.

CPC-derived EVs have previously been shown to stimulate migration of human microvascular endothelial cells (HMECs) in a scratch wound assay.¹⁴ As the mitogen-activated protein kinase1/2 (MAPK1/2)–extracellular signal-regulated kinase1/2 (ERK1/2) pathway is known to play an important role in cell survival, migration and angiogenesis during wound healing,¹⁸⁻²⁰ EV-induced ERK1/2 phosphorylation was used as a read-out to evaluate the possible differences in functionality of UC-EVs and SEC-EVs. To investigate the functionality of EV preparations, HMECs were stimulated with UC-EVs and SEC-EVs (Figure 3, *A*).

Due to a lack of consensus in the EV-field on the most accurate method for EV quantification, HMECs were stimulated with both equal amounts of UC-EV and SEC-EV protein (Figure 3, *B*), and equal numbers of EV particles (Figure 3, *C*). Levels of phosphorylated ERK1/2 and total ERK1/2 were determined using Western blotting, after which pERK/ERK ratios were calculated. A dose-dependent increase in ERK phosphorylation was observed for



Figure 2. Characterization of UC-EVs and SEC-EVs. (A) EV yield as determined by microBCA analysis (EV protein, upper panel) and Nanoparticle Tracking Analysis (NTA) (EV particle number, lower panel). Statistical analysis was performed using an unpaired Student's *t* test. (B) Size distribution profile of UC-EVs and SEC-EVs as determined by NTA. (C) Transmission electron microscopy pictures of UC-EVs and SEC-EVs. Scale bar = 100 nm. (D) Western blot of UC-EVs, SEC-EVs and cell lysate (CL). Abbreviations: CL = cell lysate, MW = molecular weight, SEC = size-exclusion chromatography, UC = ultracentrifugation.

SEC-EVs, as treatment with 3 µg SEC-EVs led to a higher pERK/ ERK ratio compared to 1 µg SEC-EVs (2.1 ± 0.3 for 3 µg SEC-EVs vs 1.2 ± 0.2 for 1 µg SEC-EVs). Moreover, stimulation with 3 µg SEC-EVs resulted in a higher pERK/ERK ratio compared to stimulation with 3 µg of UC-EVs (2.1 ± 0.3 for SEC-EVs vs 1.0 ± 0.2 for UC-EVs). The same trend was observed after adding equal numbers of EV particles to HMECs (Figure 3, *C*). Stimulation of HMECs with 6.10^{10} SEC-EVs resulted in higher pERK/ERK ratio compared to 6.10^{10} UC-EVs (3.56 ± 1.29 for SEC particles vs 1.42 ± 0.24 for UC particles). These results show that CPC-derived SEC-EVs have higher functionality compared to UC-EVs.

Discussion

The striking difference in functionality between UC-EVs and SEC-EVs may result from the high shear forces that are applied during high speed centrifugation during UC isolation. These may

detrimentally affect signaling molecules on the EV surface, thereby preventing UC-EVs to activate, bind to or be taken up by recipient cells. Indeed, UC-EVs have previously been described to appear ruptured when studied using transmission electron microscopy and fluorescence correlation spectroscopy.¹³ Furthermore, the size distribution profile of SEC-EVs also differed from UC-EVs, as SEC-EVs were found to be smaller in size compared to UC-EVs. The apparent larger size may be the result of aggregation or fusion of EVs during UC, as also suggested by others.^{9,11-13} Alternatively, UC isolation may enrich for larger EVs that sediment more efficiently. Whether and how this contributes to EV functionality remains to be investigated. Furthermore, as characteristics of isolated EVs vary between cell types, differences in functionality between SEC-EVs and UC-EVs may be cell type-dependent. This needs to be addressed in future studies.

In this study, we used induction of ERK phosphorylation as an outcome parameter to assess EV functionality, as activation of HMECs via pERK has been shown to be indicative for the



Figure 3. Assessment of UC-EV and SEC-EV functionality. (A) HMECs were stimulated with EVs for 30 min, after which phosphorylated ERK1/2 and total ERK1/2 protein levels were determined using Western blotting. (B) Stimulation of HMECs with equal amounts of UC-EV and SEC-EV protein. (C) Stimulation with equal numbers of UC-EV and SEC-EV particles. Abbreviations: ERK = extracellular signal-regulated kinase, SEC = size-exclusion chromatography, UC = ultracentrifugation.

angiogenic potential of EVs.¹⁵ One could argue that SEC-EVs may be contaminated with other soluble materials that affect ERK phosphorylation. Although we cannot exclude this completely, Western blot analyses as well as protein and particle number measurements indicate that SEC isolation allows for EV preparations with similar purity as UC (Figure 1, A, D). Additionally, CPC-EVs may affect other processes involved in cardiac disease, as treatment with CPC-EVs has been shown to result in increased cardiac function after MI in mice by enhancing angiogenesis, as well as reducing cardiomyocyte apoptosis.¹⁶ Whether SEC-EVs display increased functionality for cardiac repair in vivo, as well as for other therapeutic strategies, therefore remains to be investigated.

In conclusion, one of the major challenges for developing EV therapeutics is finding a scalable isolation method that yields EVs with high functionality. Although previous reports already suggested that EV function might be affected by the isolation procedure, to our knowledge, we are the first to show that EV isolation technique can actually affect their functionality. SEC isolation results in more functional EVs compared to UC isolation, which is especially important when developing EVs as therapeutics.

Acknowledgments

The authors thank Sander van de Weg (UMC Utrecht) for excellent assistance with electron microscopy.

Appendix A. Supplementary data

Supplementary data to this article can be found online at http://dx.doi.org/10.1016/j.nano.2017.03.011.

References

- Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 2007;9(6):654-9.
- Ratajczak J, Miekus K, Kucia M, Zhang J, Reca R, Dvorak P, et al. Embryonic stem cell-derived microvesicles reprogram hematopoietic progenitors: evidence for horizontal transfer of mRNA and protein delivery. *Leukemia* 2006;**20**(5):847-56.

- Skog J, Würdinger T, van Rijn S, Meijer DH, Gainche L, Sena-Esteves M, et al. Glioblastoma microvesicles transport RNA and proteins that promote tumour growth and provide diagnostic biomarkers. *Nat Cell Biol* 2008;10(12):1470-6.
- Sluijter JP, Verhage V, Deddens JC, Van Den Akker F, Doevendans PA. Microvesicles and exosomes for intracardiac communication. *Cardiovasc Res* 2014;**102**(2):302-11.
- Haney MJ, Klyachko NL, Zhao Y, Gupta R, Plotnikova EG, He Z, et al. Exosomes as drug delivery vehicles for Parkinson's disease therapy. J Control Release 2015;207:18-30.
- Julich H, Willms A, Lukacs-Kornek V, Kornek M. Extracellular vesicle profiling and their use as potential disease specific biomarker. *Front Immunol* 2014;5:413 1-6.
- Timmers L, Lim SK, Hoefer IE, Arslan F, Lai RC, van Oorschot AA, et al. Human mesenchymal stem cell-conditioned medium improves cardiac function following myocardial infarction. *Stem Cell Res* 2011;6(3):206-14.
- Lai RC, Arslan F, Lee MM, Sze NS, Choo A, Chen TS, et al. Exosome secreted by MSC reduces myocardial ischemia/reperfusion injury. *Stem Cell Res* 2010;4(3):214-22.
- Linares R, Tan S, Gounou C, Arraud N, Brisson AR. High-speed centrifugation induces aggregation of extracellular vesicles. *J Extracell Vesicles* 2015;4:29509-15.
- Bobrie A, Colombo M, Krumeich S, Raposo G, Théry C. Diverse subpopulations of vesicles secreted by different intracellular mechanisms are present in exosome preparations obtained by differential ultracentrifugation. *J Extracell Vesicles* 2012;1:18397-408.
- Lamparski HG, Metha-Damani A, Yao JY, Patel S, Hsu DH, Ruegg C, et al. Production and characterization of clinical grade exosomes derived from dendritic cells. *J Immunol Methods* 2002;**270**(2):211-26.

- Taylor DD, Shah S. Methods of isolating extracellular vesicles impact down-stream analyses of their cargoes. *Methods* 2015;87:3-10.
- Nordin JZ, Lee Y, Vader P, Mäger I, Johansson HJ, Heusermann WS, et al. Ultrafiltration with size-exclusion liquid chromatography for high yield isolation of extracellular vesicles preserving intact biophysical and functional properties. *Nanomedicine* 2015;11(4):879-83.
- Vrijsen KR, Sluijter JP, Schuchardt MW, van Balkom BW, Noort WA, Chamuleau SA, et al. Cardiomyocyte progenitor cell-derived exosomes stimulate migration of endothelial cells. *J Cell Mol Med* 2010;14(5):1064-70.
- Vrijsen KR, Maring JA, Chamuleau SA, Verhage V, Mol EA, Deddens JC, et al. Exosomes from cardiomyocyte progenitor cells and mesenchymal stem cells stimulate angiogenesis via EMMPRIN. *Adv Healthc Mater* 2016;5(19):2555-65.
- Barile L, Lionetti V, Cervio E, Matteucci M, Gherghiceanu M, Popescu LM, et al. Extracellular vesicles from human cardiac progenitor cells inhibit cardiomyocyte apoptosis and improve cardiac function after myocardial infarction. *Cardiovasc Res* 2014;103(4):530-41.
- Watson DC, Bayik D, Srivatsan A, Bergamaschi C, Valentin A, Niu G, et al. Efficient production and enhanced tumor delivery of engineered extracellular vesicles. *Biomaterials* 2016;105:195-205.
- Ballif BA, Blenis J. Molecular mechanisms mediating mammalian mitogen-activated protein kinase (MAPK) kinase (MEK)-MAPK cell survival signals. *Cell Growth Differ* 2001;12(8):397-408.
- Eliceiri BP, Klemke R, Strömblad S, Cheresh DA. Integrin alphavbeta3 requirement for sustained mitogen-activated protein kinase activity during angiogenesis. *J Cell Biol* 1998;140(5):1255-63.
- Alavi A, Hood JD, Fraustro R, Stupack DG, Cheresh DA. Role of Raf in vascular protection from distinct apoptotic stimuli. *Science* 2003;**301**(5629):94-6.