

Technical comment to “*Aggregatibacter actinomycetemcomitans*-induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis.”

One-sentence summary:

The link between rheumatoid arthritis and exposure to a bacterial toxin was not found in a population of RA patients from the Netherlands

Mikhail Volkov¹, Jacqueline Dekkers¹, Bruno G. Loos², Sergio Bizzarro², Thomas W.J. Huizinga¹, Helle A. Praetorius³, René E.M. Toes¹, Diane van der Woude¹

1. Department of Rheumatology, Leiden University Medical Center, Leiden 2333 ZA, the Netherlands
2. Department of Periodontology, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam and Vrije Universiteit, Amsterdam 1081 HV, the Netherlands
3. Department of Biomedicine, Aarhus University, Aarhus 8000, Denmark

In a recent publication in *Science Translational Medicine*, Konig et al. described a potential explanation for the link between periodontal infection and rheumatoid arthritis (RA)(1). They identified a specific periodontitis-associated bacterium: *Aggregatibacter actinomycetemcomitans* (Aa), which can dysregulate the activity of citrullinating enzymes in neutrophils via its lytic toxin (leukotoxin A: LtxA). The generated citrullinated autoantigens are the target of a key autoantibody system in RA: anti-citrullinated protein antibodies (ACPA). Furthermore, the authors report that the effect of the most important genetic risk factor for RA, the human leukocyte antigen-DRB1 shared epitope (HLA SE) alleles, was limited to RA patients who had been exposed to Aa, as determined by seropositivity to LtxA in a set of 194 RA patients. Based on these findings, the authors hypothesized that LtxA may be a key factor in the initiation of the RA-specific anti-citrullinated protein immune response in genetically predisposed individuals. In light of the crucial implications of this hypothesis for the current thinking regarding RA development, we aimed to replicate these findings.

To this end, we focused on two main questions: 1) Is the increased exposure to Aa, as measured by the presence of anti-LtxA-antibodies, specific for RA or also present in other forms of inflammatory arthritis? 2) Can we replicate the finding that the association between HLA SE alleles and ACPA-positive RA is limited to the anti-LtxA-positive subset? To answer these questions, we established an enzyme-linked immunosorbent assay (ELISA) against purified LtxA. The purification of LtxA was performed according to the method described by Reinholdt et al. (2), which is the same method carried out from the same clone of the strain of Aa (HK921; JP2 positive) in the same laboratory as in the manuscript by König et al. (1). A hemolytic assay confirmed that the activity of LtxA corresponded to the normal range of freshly purified LtxA. To confirm that the presence of antibodies against LtxA was indeed a reflection of exposure to the bacterium Aa, we first tested serum samples from periodontitis patients (free from any chronic diseases such as RA) whose subgingival lesions had been tested for the presence of Aa by culturing techniques (3). As depicted in Figure 1A, the concentrations of anti-LtxA-antibodies were clearly increased in Aa positive patients [median concentration (interquartile range (IQR): 8853 (5544; 14147))] versus Aa negative patients [median concentration (IQR): 764 (523; 4636)], confirming the specificity of the ELISA (Mann-Whitney U-test p-value < 0.001). Serial dilutions of a serum mix from 3 strongly positive RA patients were used as a standard, and the lowest point of the linear part of the standard curve (2000 AU/ml) was defined as the cut-off for positivity. With this cut-off, the specificity of our assay in Aa-negative periodontitis patients was 75%.

We then tested sera from 594 patients participating in the Leiden Early Arthritis Clinic with various different diagnoses, including RA according to the 1987 ACR criteria (4). Furthermore, we also measured anti-LtxA concentrations and positivity in a group of 156 healthy controls (without chronic illnesses) from the Leiden area. Figure 1B depicts the anti-LtxA concentrations in these groups and illustrates that anti-LtxA antibodies could be found in a substantial proportion of RA patients, but also in patients with other forms of arthritis. The specificity of the assay in the healthy controls was 78.2%. Next, we investigated whether among RA patients, anti-LtxA-antibodies were preferentially present within the ACPA-positive group, and whether there was an association with the HLA SE alleles. As depicted in Figure 1C, neither of these associations could be found in our cohort. Furthermore, as can be seen in Table 1, the association between HLA SE alleles and anti-CCP-positive RA was similar

among RA patients positive and negative for anti-LtxA. Therefore, the effect of the HLA SE alleles appears not to be confined to the patient group positive for anti-LtxA antibodies. In summary, in this large cohort of arthritis patients, anti-LtxA antibodies were not specifically associated with RA, and among RA patients, there was no association with the presence of ACPA or HLA SE alleles, in contrast to the findings of Konig et al. (1). A possible explanation for these divergent results could be that the anti-LtxA assay used here may have differed from the assay in the previous publication. However, the essential constituent (LtxA) and setup of the assay was derived from the same source. Furthermore, we performed various controls, such as analysis of periodontitis patients, the results of which support the validity of our assay.

It is also possible that differences in proportions of positive patients reported in the original study by Konig et al. (1) and in our study could be due to differences in patient populations between the United States and the Netherlands. It seems likely that differences in living environment, genetic background, and referral strategy exist between these two countries. However, because Aa is a prevalent microorganism causing periodontitis in both countries (5,6), it appears unlikely that population differences can explain the contrasting findings regarding the possible role of Aa.

Although microbial influences may well be important in the development of RA, our results do not support a key role of exposure to LtxA originating from the periodontal pathogen Aa in linking the effect of the HLA SE alleles and periodontal disease to anti-citrullinated protein autoimmunity in RA. The hypothesis of a causal link between periodontitis/periodontitis-associated microorganisms and RA seems appealing, but it is also possible that RA patients are more frequently anti-LtxA-positive simply because they have more periodontitis, as is known from epidemiological studies (7,8). We hope our findings can contribute to the discussion about the origins of autoimmunity in RA, and look forward to other replication experiments regarding these intriguing observations.

References and notes

1. M. F. Konig *et al.*, Aggregatibacter actinomycetemcomitans-induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis. *Science translational medicine* **8**, (2016).

2. J. Reinholdt *et al.*, Monodisperse and LPS-free *Aggregatibacter actinomycetemcomitans* leukotoxin: Interactions with human beta(2) integrins and erythrocytes. *Biochimica Et Biophysica Acta-Proteins and Proteomics* **1834**, 546-558 (2013).
3. S. Bizzarro, U. Van der Velden, B. G. Loos, Local disinfection with sodium hypochlorite as adjunct to basic periodontal therapy: a randomized controlled trial. *Journal of Clinical Periodontology* **43**, 778-788 (2016).
4. D. P. de Rooy, M. P. van der Linden, R. Knevel, T. W. Huizinga, A. H. van der Helm-van Mil, Predicting arthritis outcomes--what can be learned from the Leiden Early Arthritis Clinic? *Rheumatology.(Oxford)* **50**, 93-100 (2011).
5. M. Montecchi *et al.*, Microbiological Distribution of Six Periodontal Pathogens Between Untreated Italian and Dutch Periodontal Patients. *Oral Health Prev Dent.* **14**, 329-37 (2016).
6. C. Chen, T. Wang, W. Chen, Occurrence of *Aggregatibacter actinomycetemcomitans* serotypes in subgingival plaque from United States subjects. *Mol Oral Microbiol.* **25**, 207-14 (2010).
7. N. Pischon *et al.*, Association among rheumatoid arthritis, oral hygiene, and periodontitis. *J Periodontology* **79**, 979-86 (2008).
8. P. de Pablo, T. Dietrich, T.E. McAlindon, Association of periodontal disease and tooth loss with rheumatoid arthritis in the US population. *J Rheumatology* **35**, 70-6 (2008).

Acknowledgements: We thank Gerrie Stoeken-Rijsbergen for performing a part of the experiments on the samples from the periodontitis patients.

Funding: These studies were supported by a scientific training bursary of the European League Against Rheumatism (M.V.), an NWO-ZonMW VENI grant from the Netherlands Organization for Scientific Research (D.W.) and the Dutch Arthritis Foundation.

Author contributions: M.V. and J.D. performed the experiments and the data analysis. B.G.L. and S.B. contributed the sera from periodontitis patients. H.A.P. contributed the leukotoxin A. T.W.J.H. and R.T. designed the study and aided in the interpretation of the data. D.W. supervised the experiments, aided in the interpretation of the data and wrote the manuscript. All authors were involved in drafting the manuscript and approved the final version to be published.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: Anonymized data can be requested from the authors.

Table 1

Table 1 The association of SE alleles with anti-CCP based on exposure to LtxA in patients with RA. Anti-LtxA, anti-leukotoxin antibodies as determined by ELISA; Anti-CCP, anti-cyclic citrullinated peptide antibody, cutoff for positivity >20 U; SE, HLA-DRB1 shared epitope allele

	Anti-LtxA-negative RA (n = 143)			P	Anti-LtxA-positive (n = 189)			P
	SE- negative (n = 44)	SE- positive (n = 99)	OR		SE- negative (n = 66)	SE- positive (n = 123)	OR	
Anti-CCP positivity, %	34	70	4,45	0.0001	29	68	5,33	<0.0001

Figure legend

Figure 1. Anti-LtxA-antibody concentrations in various patient groups.

(A) Serum antibodies to LtxA were measured in periodontitis patients with and without Aa infection (n=16 per group) confirmed by culturing.

(B) Distribution of anti-LtxA antibodies in sera of 594 patients suffering from early arthritis and 156 controls. Concentrations of anti-LtxA antibodies in the serum of each individual are shown.

(C) Serum antibodies to LtxA were measured in RA patients stratified for the presence of anti-CCP2 antibodies, classified as ACPA positive (n=214) and ACPA negative (n=155) RA patients (left panel). Presence of anti-LtxA antibodies is shown in sera of RA patients carrying HLA-DRB1 shared epitope alleles, classified as SE positive (n=243) and SE negative (n=117) RA patients (right panel).

OA: inflammatory osteoarthritis, SpA: spondyloarthritis with peripheral arthritis, PsA: psoriatic arthritis, Sarc: sarcoidosis. Red lines indicate the median for each group, error bars indicate interquartile ranges. The dashed line indicates the cut-off for positivity. The number of patients per group and percentage of patients positive according to the cut-off are shown underneath. Mann-Whitney test was used for statistical comparison of antibody concentrations between RA patients and other subgroups, ACPA positive and ACPA negative, SE positive and SE negative patients. Significant differences are indicated by corresponding p-values.