



Universiteit
Leiden
The Netherlands

Palladium-induced granulomas analysed with inductively coupled plasma mass spectrometry

Marsidi, N.; Beijnen, J.H.; Zuuren, E.J. van

Citation

Marsidi, N., Beijnen, J. H., & Zuuren, E. J. van. (2018). Palladium-induced granulomas analysed with inductively coupled plasma mass spectrometry. *Contact Dermatitis*, 79(1), 41-+. doi:10.1111/cod.12979

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/75715>

Note: To cite this publication please use the final published version (if applicable).

Palladium-induced granulomas analysed with inductively coupled plasma mass spectrometry

Nick Marsidi¹  | Jos H. Beijnen² | Esther J. van Zuuren¹

¹Department of Dermatology, Leiden University Medical Centre, Leiden, The Netherlands

²Department of Clinical Pharmacology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

Correspondence

Dr Nick Marsidi, Department of Dermatology, Leiden University Medical Centre, Albinusdreef 2, Leiden, 2300 RC, The Netherlands.
Email: n.marsidi@lumc.nl

KEYWORDS: allergic contact, case report, foreign body, granuloma, ICP-MS, palladium

CASE REPORT

A 28-year-old female was referred to our dermatology clinic because of persistent swelling of the earlobes several months after ear piercing. Clinical examination showed symmetrical nodes on both earlobes (Figure 1). Histology showed epithelioid granulomas with a lymphocytic infiltrate, as seen in sarcoidosis and foreign body reactions. There were no further signs of sarcoidosis (normal chest X-ray and normal angiotensin-converting enzyme findings) or foreign material. Patch testing with the European baseline series and a dental series (including various metals) was performed. Positive reactions to nickel sulfate 5% pet. [+ on day (D) 2 and D3] and palladium chloride 1% pet. (+ on D3) were observed. Four weeks after the patch test, a persistent reaction on the patient's back remained at the palladium test site. A biopsy showed epithelioid granulomas similar to those previously seen in the excised nodes.

Inductively coupled plasma mass spectrometry (ICP-MS) (ICP-quadrupole-MS, Varian 810-MS) detected palladium (^{105}Pd) in all of our skin samples (thickness, 4 μm) and showed a 3-fold higher signal than that from skin samples of control patients. The content of ^{105}Pd in each sample was semiquantitatively determined to be 0.6 ng of ^{105}Pd per sample.

DISCUSSION

Similar cases have been described in the literature, in which granulomas have been found in the earlobe with a positive reaction to

palladium after patch testing.¹⁻⁴ The presence of palladium has been confirmed in earrings, and is a likely source of this reaction.⁴ The authors described the combination of these findings as an allergic contact granuloma, but the skin biopsies have never been tested for the presence of metals.

This raises the question of whether contact sensitization alone is enough to cause granulomas, as the clinical relevance of palladium hypersensitivity remains uncertain. Tillman et al. investigated the wearing of palladium-coated earrings, and found no skin reactions in 40 subjects who showed positive patch tests to palladium.⁵ Another possibility is that palladium permeates through the skin, causing a foreign body reaction. Skin penetration and permeation of palladium nanoparticles in the epidermis and dermis has been shown in damaged skin.⁶ Other cutaneous granulomas have also been reported to be induced by beryllium and zirconium, and in these cases metals have been found in the skin.^{3,4}

To the best of our knowledge, we present the first case in which epithelioid granulomas have been found in the earlobes with a positive patch test reaction to palladium and the confirmed presence of palladium in the skin samples. Although the pathogenesis remains unclear, our findings suggest that palladium might induce granulomas via a foreign body reaction in patients with positive palladium contact allergy.

Acknowledgements

We thank Dr A.P.M Lavrijsen for her support.

Conflict of interest

The authors declare no potential conflict of interests.

ORCID

Nick Marsidi  <http://orcid.org/0000-0002-1637-9962>

REFERENCES

1. Casper C, Groth W, Hunzelmann N. Sarcoidal-type allergic contact granuloma: a rare complication of ear piercing. *Am J Dermatopathol*. 2004;26:59-62.
2. González-Pérez R, Ruiz-Carillo G, Soloeta R. Sarcoid-type allergic contact granuloma caused by earrings in a boy. *Actas Dermosifiliogr*. 2012; 103:73-74.
3. Thijs L, Deraedt K, Goossens A. Granuloma possibly induced by palladium after ear piercing. *Dermatitis*. 2008;19:E26-E29.
4. Goossens A, de Swert A, de Coninck K, Snauwaert JE, Dedeurwaerder M, de Bonte M. Allergic contact granuloma due to palladium following ear piercing. *Contact Dermatitis*. 2006;55:338-341.
5. Tillman C, Engfeldt M, Hindsén M, Bruze M. Usage test with palladium-coated earrings in patients with contact allergy to palladium and nickel. *Contact Dermatitis*. 2013;69:288-295.
6. Larese Filon F, Crosera M, Mauro M, et al. Palladium nanoparticles exposure: evaluation of permeation through damaged and intact human skin. *Environ Pollut*. 2016;214:497-503.



FIGURE 1 Symmetrical nodes on both earlobes

How to cite this article: Marsidi N, Beijnen JH, van Zuuren EJ. Palladium-induced granulomas analysed with inductively coupled plasma mass spectrometry. *Contact Dermatitis*. 2018;79:41-42. <https://doi.org/10.1111/cod.12979>