



Universiteit
Leiden
The Netherlands

Effects of the Namaste Care Family programme on quality of life of nursing home residents with advanced dementia and on family caregiving experiences: study protocol of a cluster-randomised controlled trial

Smaling, H.J.A.; Joling, K.J.; Ven, P.M. van de; Bosmans, J.E.; Simard, J.; Volicer, L.; ... ; Steen, J.T. van der

Citation

Smaling, H. J. A., Joling, K. J., Ven, P. M. van de, Bosmans, J. E., Simard, J., Volicer, L., ... Steen, J. T. van der. (2018). Effects of the Namaste Care Family programme on quality of life of nursing home residents with advanced dementia and on family caregiving experiences: study protocol of a cluster-randomised controlled trial. *Bmj Open*, 8(10), e025411. doi:10.1136/bmjopen-2018-025411

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/76184>

Note: To cite this publication please use the final published version (if applicable).

Correction: *Effects of the Namaste Care Family programme on quality of life of nursing home residents with advanced dementia and on family caregiving experiences: study protocol of a cluster-randomised controlled trial*

Smaling HJA, Joling KJ, van de Ven PM, *et al.* Effects of the Namaste Care Family programme on quality of life of nursing home residents with advanced dementia and on family caregiving experiences: study protocol of a cluster-randomised controlled trial. *BMJ Open* 2018;8:e025411. doi: 10.1136/bmjopen-2018-025411.

This article was previously published with the incorrect sentence about the QUALID below.

We used the mean (M) and SD as calculated in previous work validating the Dutch version of the QUALID in nursing home residents with advanced dementia (M=7.5, SD=4.9).⁴¹For the PES, it was analysed and provided for the subgroup of family caregivers with dementia (M=2.9, SD=1.9).⁹²

Now, this sentence has been removed and the beginning of the next sentence has been revised as given below:

The mean PES was analysed and provided for the subgroup of family caregivers with dementia (M=2.9, SD=1.9).⁹²

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2019;9:e025411corr1. doi:10.1136/bmjopen-2018-025411corr1

