



Universiteit
Leiden
The Netherlands

Recommendations for a nomenclature system for reporting methylation aberrations in imprinted domains

Monk, D.; Morales, J.; Dunnen, J.T. den; Russo, S.; Court, F.; Prawitt, D.; ... ; European Network Human Congenital

Citation


Monk, D., Morales, J., Dunnen, J. T. den, Russo, S., Court, F., Prawitt, D., ... Tumer, Z. (2018). Recommendations for a nomenclature system for reporting methylation aberrations in imprinted domains. *Epigenetics*, 13(2), 117-121. doi:10.1080/15592294.2016.1264561

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/86476>

Note: To cite this publication please use the final published version (if applicable).



Recommendations for a nomenclature system for reporting methylation aberrations in imprinted domains

David Monk, Joannella Morales, Johan T. den Dunnen, Silvia Russo, Franck Court, Dirk Prawitt, Thomas Eggermann, Jasmin Beygo, Karin Buiting, Zeynep Tümer & the Nomenclature group of the European Network for Human Congenital Imprinting Disorders

To cite this article: David Monk, Joannella Morales, Johan T. den Dunnen, Silvia Russo, Franck Court, Dirk Prawitt, Thomas Eggermann, Jasmin Beygo, Karin Buiting, Zeynep Tümer & the Nomenclature group of the European Network for Human Congenital Imprinting Disorders (2018) Recommendations for a nomenclature system for reporting methylation aberrations in imprinted domains, *Epigenetics*, 13:2, 117-121, DOI: [10.1080/15592294.2016.1264561](https://doi.org/10.1080/15592294.2016.1264561)

To link to this article: <https://doi.org/10.1080/15592294.2016.1264561>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



[View supplementary material](#)



Accepted author version posted online: 02 Dec 2016.
Published online: 25 Jan 2018.



[Submit your article to this journal](#)



Article views: 551



[View Crossmark data](#)



Citing articles: 17 [View citing articles](#)

POINT OF VIEW



Recommendations for a nomenclature system for reporting methylation aberrations in imprinted domains

David Monk^a, Joannella Morales^b, Johan T. den Dunnen^c, Silvia Russo^d, Franck Court^e, Dirk Prawitt^f, Thomas Eggermann^g, Jasmin Beygo^h, Karin Buiting^h, Zeynep Tümerⁱ, and the Nomenclature group of the European Network for Human Congenital Imprinting Disorders*

^aImprinting and Cancer group, Cancer Epigenetic and Biology Program, Bellvitge Biomedical Research Institute, Barcelona, Spain; ^bEuropean Molecular Biology Laboratory, European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, UK; ^cHuman Genetics and Clinical Genetics, Leiden University Medical Center, Leiden, the Netherlands; ^dLaboratory of Cytogenetics and Molecular Genetics, Istituto Auxologico Italiano, Milan, Italy; ^eReproduction and Developmental Genetics, Centre National de la Recherche Scientifique, Clermont-Ferrand, France; ^fCenter for Pediatrics and Adolescent Medicine, Johannes Gutenberg University Medical Center, Obere Zahlbacher, Mainz, Germany; ^gInstitute of Human Genetics, Technical University of Aachen, Aachen, Germany; ^hInstitute of Human Genetics, University Hospital Essen, University Duisburg-Essen, Germany; ⁱApplied Human Molecular Genetics, Kennedy Centre, Copenhagen University Hospital, Rigshospitalet, Glostrup, Denmark

ABSTRACT

The analysis of DNA methylation has become routine in the pipeline for diagnosis of imprinting disorders, with many publications reporting aberrant methylation associated with imprinted differentially methylated regions (DMRs). However, comparisons between these studies are routinely hampered by the lack of consistency in reporting sites of methylation evaluated. To avoid confusion surrounding nomenclature, special care is needed to communicate results accurately, especially between scientists and other health care professionals. Within the European Network for Human Congenital Imprinting Disorders we have discussed these issues and designed a nomenclature for naming imprinted DMRs as well as for reporting methylation values. We apply these recommendations for imprinted DMRs that are commonly assayed in clinical laboratories and show how they support standardized database submission. The recommendations are in line with existing recommendations, most importantly the Human Genome Variation Society nomenclature, and should facilitate accurate reporting and data exchange among laboratories and thereby help to avoid future confusion.

ARTICLE HISTORY

Received 7 October 2016
Revised 8 November 2016
Accepted 17 November 2016

KEYWORDS

Imprinting; imprinting disorders; methylation; nomenclature


Mammalian genomic imprinting is an epigenetic regulatory mechanism that results in parent-of-origin specific gene expression in diploid somatic cells (for a review, see ref¹). Several features of the imprinting mechanism have been identified including allelic DNA methylation, histone modifications, and noncoding RNAs.² Clustering and coordinate regulation is a key feature of imprinted domains with much effort invested in understanding how multiple genes are regulated by long-range cis-acting differentially methylated regions (DMRs).

In 1993, two publications reported parent-of-origin specific methylation associated with imprinted domains. Both of these studies were in mouse, the first described the paternally methylated regions associated with the *H19-Igf2* gene cluster,³ the second identified a region of methylation on the maternal allele within the *Igf2r* gene associated with the *T-associated maternal effect (Tme)* deletion.⁴ Since these first pivotal reports, with the advent of genome-wide methylation screening technologies, the number of imprinted DMRs in mammalian species has steadily increased, including those originating from the respective germlines and those that are somatically acquired.

Primary methylation defects of some well characterized imprinted DMRs are directly responsible for developmental

disorders, including Beckwith-Wiedemann syndrome (OMIM 130,650) (BWS), Silver-Russell syndrome (OMIM 180,860) (SRS), Transient Neonatal Diabetes Mellitus (OMIM 601,410) (TNDM), Kagami-Ogata syndrome (OMIM 608,149) (KOS14), Temple syndrome (OMIM 616,222) (TS14), Prader-Willi syndrome (OMIM 176,270) (PWS), Angelman syndrome (OMIM 105,830) (AS), and Pseudohypoparathyroidism Ib (OMIM 103,580) (PHP1b) (a detailed description of human imprinting disorders is available at the webpage of the European Network for Human Congenital Imprinting Disorders, EUCID.net). The aberrant methylation often affects solely the disease-associated locus but, recently, some individuals have been reported to have additional imprinted DMRs affected, leading to a scenario termed multi-locus imprinting disturbance (MLID) (for a review, see ref⁵). Reporting these epigenomic data from molecular tests in laboratory reports or for publication is troubled by the lack of a uniform nomenclature. In this article we recommend unified names for imprinted DMRs and give details of their precise locations and suggest nomenclature for describing the results similar to those routinely used for DNA sequence variants.

CONTACT David Monk  Cancer Epigenetics and Biology Program (PEBC), Bellvitge Biomedical Research Institute (IDIBELL), Av. Gran Via de L'Hospitalet 199–203.

 Supplemental data for this article can be accessed on the publisher's website.

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Consensus for names of imprinted DMR

From the earliest days of the molecular descriptions of imprinting aberrations it became obvious that recording methylation defects would be challenging. This was evident simply because some imprinted genes have historically had multiple names, since many were identified simultaneously by independent groups who have termed the transcripts and DMRs differently. This is exemplified by the maternally methylated region overlapping the promoter of *KCNQ1OT1* within intron 10 of *KCNQ1* on chromosome 11, which has more than five aliases (Table 1). Ultimately, this causes confusion when cross-referencing original literature and modern databases. To improve this situation, the 41 EUCID members from 22 countries, have developed a uniform nomenclature system for reporting methylation aberrations. The final consensus after careful consideration was that the name of an imprinted DMR should be attributed to the nearest transcript with an approved symbol from the Human Genome Organization (HUGO) Gene Nomenclature Committee (HGNC).

Furthermore, we named a DMR in such a way that it gives basic information regarding its localization in relation to the nearest RefSeq transcript with the use of several prefixes outlined in Fig. 1 (e.g., TSS for transcription start site, IG for intergenic, Int for intronic and alt-TSS for alternative transcription start site). The precise location of each imprinted DMR is derived from methyl-seq data from whole blood samples as described by Court et al.,⁶ which has base-pair resolution. To ensure that the same genomic regions are identifiable in subsequent genome builds, all imprinted domains, including their corresponding DMRs, have been submitted to obtain Locus Reference Genomic (LRG) identifiers.⁷ LRGs are manually curated reference sequence records specifically designed for the reporting of variants with clinical implications. The inclusion of stable and unique genomic, transcript, and protein reference sequences ensures that variants are unambiguously and consistently reported over time (www.lrg-sequence.org). The records will contain all relevant DMR annotations. Information regarding the recommended naming, localization, and sizes of each DMR are given in Table 1.

Standardization of reporting exact sites of imprinted methylation

It has previously been discussed that in order to allow correct identification and eventual reproduction of published observations, a universal system for the description of specific sites of DNA methylation tested needs to be employed.⁸ In the case of imprinted DMRs, this is reasonably straightforward if laboratories use commercially available methods to analyze methylation, such as methylation-sensitive multiplex ligation-dependent probe amplifications (MLPA) or high-density methylation arrays. In such cases, the precise location of the probe identifier, restriction site, or the interrogated CpG probes found on commonly used methylation profiling platforms can easily be identified, the genomic nucleotide tested accurately described, and methylation values reported. As an initial step to assist in this standardization, we have provided a resource listing all probes mapping to imprinted DMRs on the popular Infinium HumanMethylation450 BeadChips (Illumina, USA)(Suppl. Table 1), as well as

the CpG dinucleotides interrogated by commonly used methylation-sensitive MLPA kits (MRC Holland, Netherlands)(Suppl. Table 2). For custom technologies, such as in-house pyrosequencing, for example, different CpG positions within the imprinted DMRs may be examined. In such cases, we recommend that the genomic coordinates targeted by the assays be listed and the methylation status described as an average percentage of all CpGs analyzed. However, such a description lacks resolution at the individual CpG level and, for future standardized reporting, it would be advantageous to have this information, not only for methylation at imprinted DMRs but also for all CpG positions in the genome. Such an approach could be based upon the current annotation of genomic locations as recommended by the Human Genome Variation Society (HGVS), allowing methylation values to be paired to each CpG position.^{9,10}

Use of the suggested nomenclature

Following HGVS recommendations, methylation values at a specific region are described with (A) the chromosome number or LRG followed by (B) a colon ":"; (C) prefix "g." for genomic DNA; (D) the position of the cytosine nucleotide or the range of nucleotides tested for the CpG containing interval; (E) the "]" character to indicate that it is a modification of the sequence not a sequence variant; (F) prefix describing the specific modification.

In collaboration with the HGVS' Sequence Variant Description Working Group (SVD-WG), it was decided to use the abbreviation "gom" to report a gain of methylation and "lom" for a loss. For non-specific methylation resistant to bisulphite conversion we suggest "bis" followed by a methylation value in brackets. If the molecular assay differentiates between 5-methylcytosine and its oxidative derivative 5-hydroxymethylcytosine, we propose the use of the "met" and "hmt," respectively. This is consistent with HGVS standards to use three-letter abbreviations that do not include the nucleotide so that the modification can be added to any DNA base. When utilizing this format, it is important to mention the correct imprinted DMR name, the genome build used, and the technique used to measure the methylation status. This is because the EUCID COST action has previously reported that different methods targeting subtly different locations within the same imprinted DMR having different sensitivities.^{11,12} Furthermore, to help characterize variation due to tissue mosaicism, the tissue source from which the DNA is derived should be stated in any report because the methylation levels can be different in different tissues.¹³

For example, the nomenclature for a bisulphite PCR targeting the *KCNQ1OT1*:TSS-DMR negative DNA strand:

GRCh37/hg19 chr11:g.2,722,063_2,722,087, in detail chr11:g.2,722,063_2,722,087|met {0.52} indicates that the CpGs within this 24 bp interval have an average methylation of 52%. Using the LRGs, the nomenclature would be LRG_1,052:g.94506_94,530; LRG_1,052:g.94506_94,530|met {0.52}. For an example with loss of methylation of the same interval, the nomenclature would be GRCh37/hg19 chr11:g.2722063_2,722,087|lom, in detail chr11:g.2,722,063_2,722,087|met {0.22} or LRG_1,052:g.94506_94,530; LRG_1,052:g.94506_94,530|met {0.22}.

Table 1. The extent of imprinted methylation defined by methyl-seq data sets with the commonly used name for each imprinted DMR, those proposed by of EUCID using HGNC approved gene names, previous aliases and LRG identifiers. For completeness, origin of the allelic methylation is given, as are any associated disorders and information whether the methylation is germline or somatically derived. Secondary DMRs are regions of differential methylation, the establishment of which is often somatically acquired and dependent on hierarchical interactions with a neighboring germline DMR. All coordinates are given as GRCh37/hg19. M, maternally derived methylation; P, paternally derived methylation; gDMR, germline DMR; PHP1b, Pseudohypoparathyroidism; SRS, Silver Russell syndrome; BWS, Beckwith Wiedemann syndrome; AS, Angelman syndrome; PWS, Prader Willi syndrome; MLID, Multilocus imprinting disturbance; TS14, Temple syndrome; KOS14, Kagami-Ogata syndrome; TNDM, Transient Neonatal Diabetes Mellitus. All relevant DMR information and aliases can also be found in the “community” section of each LRG record.

Imprinted DMR name	Chr	Extent of DMR (methyl-seq)		Number CpGs	Methylation origin	Germline derived	LRG identifier	Aliases
		Start	Finish					
<i>PPIEL</i> :Ex1-DMR	1	40,024,626	40,025,540	39	M	Oocyte gDMR		
<i>DIRAS3</i> :TSS-DMR	1	68,515,433	68,517,545	88	M	Oocyte gDMR	LRG_1,034	<i>NOEY2, ARH1</i>
<i>DIRAS3</i> :Ex2-DMR	1	68,512,505	68,513,486	39	M	Oocyte gDMR	LRG_1,034	
<i>GPR1-AS</i> :TSS-DMR	2	207,066,967	207,069,445	86	M	Oocyte gDMR		
<i>ZDBF2/GPR1</i> :IG-DMR	2	207,114,583	207,136,544	439	P	Sperm gDMR-secondary DMR	LRG_1,065	
<i>NAP1L5</i> :TSS-DMR	4	89,618,184	89,619,237	57	M	Oocyte gDMR	LRG_1,072	
<i>VTRNA2-1</i> :DMR	5	135,414,802	135,416,645	76	M	Oocyte gDMR		nc886
<i>FAM50B</i> :TSS-DMR	6	3,849,082	3,850,359	90	M	Oocyte gDMR	LRG_1,062	
<i>PLAGL1</i> :alt-TSS-DMR	6	144,328,078	144,329,888	143	M	Oocyte gDMR	LRG_1,101	<i>LOT1, ZAC1</i>
<i>IGF2R</i> :Int2-DMR	6	160,426,558	160,427,561	74	M	Oocyte gDMR	LRG_1,036	
<i>WDR22</i> :Int13-DMR	6	170,054,504	170,055,618	58	M	Oocyte gDMR	LRG_1,071	
<i>GRB10</i> :alt-TSS-DMR	7	50,848,726	50,851,312	171	M	Oocyte gDMR	LRG_1,032	
<i>PEG10</i> :TSS-DMR	7	94,285,537	94,287,960	119	M	Oocyte gDMR	LRG_1,097	
<i>MEST</i> :alt-TSS-DMR	7	130,130,122	130,134,388	226	M	Oocyte gDMR	LRG_1,033	<i>PEG1</i>
<i>SVOP</i> :alt-TSS-DMR	7	138,348,118	138,349,069	31	M	Oocyte gDMR	LRG_1,103	
<i>HTRSA</i> :TSS-DMR	7	154,862,719	154,863,382	55	M	Oocyte gDMR	LRG_1,057	
<i>ERLIN2</i> :Int6-DMR	8	37,604,992	37,606,088	37	M	Oocyte gDMR	LRG_1,040	
<i>PEG13</i> :TSS-DMR	8	141,108,147	141,111,081	193	M	Oocyte gDMR	LRG_1,041	<i>TRAPPC9</i> intronic DMR
<i>FANCC</i> :Int1-DMR	9	98,075,400	98,075,744	26	M	Oocyte gDMR	LRG_497	
<i>INPP5F</i> :Int2-DMR	10	121,578,046	121,578,727	52	M	Oocyte gDMR	LRG_1,061	
<i>H19/IGF2</i> :IG-DMR	11	2,018,812	2,024,740	250	P	Sperm gDMR	LRG_1,030	<i>ICR1, IC1, H19</i> DMR, <i>H19</i> DMD, <i>IGF2-H19</i> DMR
<i>IGF2</i> :Ex9-DMR	11	2,153,991	2,155,112	63	P	No-secondary DMR	LRG_1,031	
<i>IGF2</i> :alt-TSS-DMR	11	2,168,333	2,169,768	33	P	Sperm gDMR	LRG_1,031	
<i>KCNQ1OT1</i> :TSS-DMR	11	2,719,948	2,722,259	192	M	Oocyte gDMR	LRG_1,052	<i>ICR2, IC2, KvDMR1, KvLQT1-A, LIT1</i>
<i>RB1</i> :Int2-DMR	13	48,892,341	48,895,763	195	M	Oocyte gDMR	LRG_517	
<i>MEG3/DLK1</i> :IG-DMR	14	101,275,427	101,278,058	64	P	Sperm gDMR	LRG_1,044	<i>GLT2-DLK1</i>
<i>MEG3</i> :TSS-DMR	14	101,290,524	101,293,978	188	P	No-secondary DMR	LRG_1,098	<i>GTL2</i>
<i>MEG8</i> :Int2-DMR	14	101,370,741	101,371,419	43	M	No-secondary DMR	LRG_1,058	
<i>MKRN3</i> :TSS-DMR	15	23,807,086	23,812,495	109	M	Oocyte gDMR-secondary DMR	LRG_1,045	
<i>MAGEL2</i> :TSS-DMR	15	23,892,425	23,894,029	51	M	No-secondary DMR	LRG_1,046	
<i>NDN</i> :TSS-DMR	15	23,931,451	23,932,759	108	M	No-secondary DMR	LRG_1,047	
<i>SNRPN</i> :alt-TSS-DMR	15	25,068,564	25,069,481	19	M	No-secondary DMR		
<i>SNRPN</i> :Int1-DMR1	15	25,093,008	25,193,829	44	M	No-secondary DMR		
<i>SNRPN</i> :Int1-DMR2	15	25,123,027	25,123,905	45	M	No-secondary DMR		
<i>SNURF</i> :TSS-DMR	15	25,200,004	25,201,976	113	M	Oocyte gDMR		
<i>IGF1R</i> :Int2-DMR	15	99,408,496	99,409,650	55	M	Oocyte gDMR	LRG_1,055	<i>IRAIN</i>
<i>ZNF597</i> :3' DMR	16	3,481,801	3,482,388	29	M	Oocyte gDMR	LRG_1,066	
<i>ZNF597</i> :TSS-DMR	16	3,492,828	3,494,463	76	P	No-secondary DMR	LRG_1,066	<i>NAT15</i>
<i>ZNF331</i> :alt-TSS-DMR1	19	54,040,510	54,042,212	125	M	Oocyte gDMR	LRG_1,068	
<i>ZNF331</i> :alt-TSS-DMR2	19	54,057,086	54,058,425	102	M	Oocyte gDMR	LRG_1,068	
<i>PEG3</i> :TSS-DMR	19	57,348,493	57,353,271	221	M	Oocyte gDMR		<i>ZIM2, ZNF904</i>
<i>MCTS2P</i> :TSS-DMR	20	30,134,663	30,135,933	47	M	Oocyte gDMR		<i>psiMCT-1, MCTS2</i>
<i>NNAT</i> :TSS-DMR	20	36,148,604	36,150,528	135	M	Oocyte gDMR	LRG_1,048	<i>PEG5</i>
<i>L3MBTL1</i> :alt-TSS-DMR	20	42,142,365	42,144,040	84	M	Oocyte gDMR	LRG_1,049	<i>ZC2HC3, KIAA0681</i>
<i>GNAS-NESP</i> :TSS-DMR	20	57,414,039	57,418,612	257	P	No-secondary DMR	LRG_1,051	<i>NESP55</i>
<i>GNAS-AS1</i> :TSS-DMR	20	57,425,649	57,428,033	128	M	Oocyte gDMR	LRG_1,051	<i>NESP-AS</i>
<i>GNAS-XL</i> :Ex1-DMR	20	57,428,905	57,431,463	200	M	Oocyte gDMR	LRG_1,051	secretogranin VI
<i>GNAS A/B</i> :TSS-DMR	20	57,463,265	57,465,201	198	M	No-secondary DMR	LRG_1,051	secretogranin VI
<i>WRB</i> :alt-TSS-DMR	21	40,757,510	40,758,276	43	M	Oocyte gDMR	LRG_1,060	
<i>SNU13</i> :alt-TSS-DMR	22	42,077,774	42,078,873	63	M	Oocyte gDMR	LRG_1,102	<i>NHP2L1</i>

An example of the same region reporting methylation at individual cytosines of CpG dinucleotides would be:

GRCh37/hg19 chr11:g. 2722063C|met {0.50}; 2722072C|met {0.54}; 2722074C|met {0.52}; 2722077C|met {0.53}; 2722083C|met {0.50}; 2722085C|met {0.51}; 2722087C|met

{0.55} or LRG_1,052:g.94506C|met {0.50}; 94515C|met {0.54}; 94517C|met {0.52}; 94520C|met {0.53}; 94526C|met {0.50}; 94528C|met {0.51}; 94530C|met {0.55} in the sample with normal methylation. For the sample with loss of methylation, the nomenclature would be GRCh37/hg19 chr11:g.




Figure 1. Schematic overview of the naming system for imprinted DMRs. The upper panel shows the three imprinted DMRs located within the chromosome 14q32.2 domain depicting (A) an intergenic DMR, (B) a DMR overlapping a transcription start site, and (C) an intronic DMR. The lower panel illustrates a DMR associated with (D) an alternative first exon as found within the *PLAGL1* locus on chromosome 6q24. Black boxes represent methylated regions while open boxes are unmethylated. Blue arrows signify paternally expressed genes, red arrows denote maternally expressed genes, and gray arrows are biallelically expressed transcripts.

Webpage URLs

EUCID.net webpage. www.imprinting-disorders.eu

285 Locus Reference Genomic identifier. www.lrg-sequence.org

HGVS nomenclature webpage. www.HGVS.org/varnomen

2722063C|met {0.22}; 2722072C|met {0.26}; 2722074C|met {0.23}; 2722077C|met {0.17}; 27220783C|met {0.16}; 2722085C|met {0.26}; 2722087C|met {0.26} or LRG_1,052: g.94506C|met {0.22}; 94515C|met {0.26}; 94517C|met {0.23}; 94520C|met {0.17}; 94526C|met {0.16}; 94528C|met {0.26}; 94530C|met {0.26}.

In light of our suggestions, we encourage comments and discussion from clinical geneticists, molecular geneticists, and researchers from the epigenomics community and trust that the recommendations we have made for standardized reporting format will be useful for accurately communicating results. To give the wider epigenetics community the opportunity to be involved in the final discussions the proposed gom/lom nomenclature is open for community consultation on the HGVS webpage (see <http://varnomen.hgvs.org/bg-material/consultation/svd-wg005/>). We hope that by giving precise methylation values as percentages, it will overcome issues of comparing results between laboratories who often describe abnormalities using different methylation indexes.

The next issue that needs a consensus is defining the criteria to allow the description 'lom' or 'gom'. This is complicated as not only statistical cut-offs need to be discussed (i.e., using mean \pm standard deviation), but also the number of controls analyzed to define the normal range. Furthermore, utilizing fixed statistical criteria will be complicated in cases with mosaic epimutations, as methylation variance at different CpGs within a DMR need to be taken into account, as does the reproducibility of the molecular techniques used.¹¹

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Funding

This work was supported by the European COST action under grant number BM1208, the Bundesministerium für Bildung und Forschung (BMBF) under grant number 01GM1513A, B, C and D, the Spanish Ministerio de

Educación y Competitividad (MINECO) under grant number BFU2014–53,093 co-funded with the European Union Regional Development Fund (FEDER), the Wellcome Trust under grant number WT200990/Z/16/Z and by the European Molecular Biology Laboratory.

** Author appendix

Members of the Nomenclature group of the European Network for Human Congenital Imprinting Disorders

Frederic Brioude, Irene Netchine, Deborah Mackay, Pablo Lapunzina, Eammon Maher, Anne Rochtus, Guiomar Perez de Nanclares, Matthias Begemann, Lukas Soellner, Karen Grønskov, Jet Bliëk, Maria Paola Lombardi, Lidia Larizza, Dorota Jurkiewicz, Francesca Elli, Serap Turan, Alessandra Baumer, Tiina Kehre, O'hlcí Bronagh

AP-HP, Hôpitaux Universitaires Paris Est (AP-HP) Hôpital des Enfants Armand Trousseau, Service d'Explorations Fonctionnelles Endocriniennes, 26 avenue du Dr Arnold Netter, 75012 Paris, France.

Human Genetics and Genomic Medicine, Faculty of Medicine University of Southampton, UK.

Instituto de Genética Médica y Molecular (INGEMM)-IdiPAZ, Hospital Universitario La Paz, Madrid, Spain; CIBERER, Centro de Investigación Biomédica en Red de Enfermedades Raras, ISCIII, Madrid, Spain.

Department of Medical Genetics, University of Cambridge and NIHR Cambridge Biomedical Research Center, Cambridge, UK.

Department of Pediatrics, University Hospitals Leuven, 3000 Leuven, Belgium.

Molecular (Epi)Genetics Laboratory, BioAraba National Health Institute, OSI Araba-Txagorritxu, E-01009 Vitoria-Gasteiz, Spain.

Department of Human Genetics, RWTH Aachen, Aachen, Germany.

Clinical Genetic Clinic, Kennedy Center, Rigshospitalet, Copenhagen University Hospital, Glostrup, Denmark.

Academic Medical Center, Department of Clinical Genetics, Laboratory for Genome Diagnostics, Meibergdreef 15, 1105AZ Amsterdam, Netherlands.

Laboratory of Cytogenetics and Molecular Genetics, Istituto Auxologico Italiano, Milan, Italy.

Department of Medical Genetics, Children's Memorial Health Institute, Warsaw, Poland.

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Department of Clinical Sciences and Community Health, University of Milan, Endocrinology and Diabetology Unit, Milan, Italy.

Pediatric Endocrinology and Diabetes, Marmara University Hospital, Istanbul, Turkey.

Institute of Medical Genetics, University of Zürich, Schlieren-Zürich, Switzerland.

Department of Genetics, Tartu University Hospital, Tartu, Estonia; Department of Pediatrics, Institute of Molecular and Cell Biology, University of Tartu, Tartu, Estonia.

National Center for Medical Genetics, Ireland.

ORCID

David Monk  <http://orcid.org/0000-0001-8991-0497>

References

1. Monk D. Germline-derived DNA methylation and early embryo epigenetic reprogramming: The selected survival of imprints. *Int J Biochem Cell Biol* 2015; 67:128-38; PMID:25966912; <https://doi.org/10.1016/j.biocel.2015.04.014>
2. Ferguson-Smith AC. Genomic imprinting: the emergence of an epigenetic paradigm. *Nat Rev Genet* 2011; 12(8):565-75; PMID:21765458; <https://doi.org/10.1038/nrg3032>
3. Ferguson-Smith AC, Sasaki H, Cattanaach BM, Surani MA. Parental-origin-specific epigenetic modification of the mouse H19 gene. *Nature* 1993; 362(6422):751-5; PMID:8469285; <https://doi.org/10.1038/362751a0>
4. Stöger R, Kubicka P, Liu CG, Kafri T, Razin A, Cedar H, Barlow DP. Maternal-specific methylation of the imprinted mouse *Igf2r* locus identifies the expressed locus as carrying the imprinting signal. *Cell* 1993; 73(1):61-71; PMID:8462104; [https://doi.org/10.1016/0092-8674\(93\)90160-R](https://doi.org/10.1016/0092-8674(93)90160-R)
5. Sanchez-Delgado M, Riccio A, Eggermann T, Maher ER, Lapunzina P, Mackay D, Monk D. Causes and Consequences of Multi-Locus Imprinting Disturbances in Humans. *Trends Genet* 2016; 32(7):444-55; PMID:27235113; <https://doi.org/10.1016/j.tig.2016.05.001>
6. Court F, Tayama C, Romanelli V, Martin-Trujillo A, Iglesias-Platas I, Okamura K, Sugahara N, Simón C, Moore H, Harness JV, et al. Genome-wide parent-of-origin DNA methylation analysis reveals the intricacies of human imprinting and suggests a germline methylation-independent mechanism of establishment. *Genome Res* 2014; 24(4):554-69; PMID:24402520; <https://doi.org/10.1101/gr.164913.113>
7. MacArthur JA, Morales J, Tully RE, Astashyn A, Gil L, Bruford EA, Larsson P, Kileck P, Dalgleish R, Maglott DR, Cunningham F. Locus Reference Genomic: reference sequences for the reporting of clinically relevant sequence variants. *Nucl Acid Res* 2014; 42:D873-D878; PMID:24285302; <https://doi.org/10.1093/nar/gkt1198>
8. Saffery R, Gordon L. Time for a standardized system of reporting sites of genomic methylation. *Genome Biol* 2015; 16:85; PMID:25924664; <https://doi.org/10.1186/s13059-015-0654-9>
9. den Dunnen JT, Dalgleish R, Maglott DR, Hart RK, Greenblatt MS, McGowan-Jordan J, Roux AF, Smith T, Antonarakis SE, Taschner PE. HGVS Recommendations for the Description of Sequence Variants: 2016 Update. *Hum.Mutat* 2016; 37(6):564-569; PMID:26931183; <https://doi.org/10.1002/humu.22981>
10. Ogino S, Gulley ML, den Dunnen JT, Wilson RB, Association for Molecular Pathology Training and Education Committee. Standard mutation nomenclature in molecular diagnostics: practical and educational challenges. *J Mol Diagn* 2007; 9(1):1-6; PMID:17251329; <https://doi.org/10.2353/jmoldx.2007.060081>
11. Soellner L, Monk D, Rezwan FI, Begemann M, Mackay D, Eggermann T. Congenital imprinting disorders: Application of multilocus and high throughput methods to decipher new pathomechanisms and improve their management. *Mol Cell Probes* 2015; 29(5):282-90; PMID:26070988; <https://doi.org/10.1016/j.mcp.2015.05.003>
12. Garin I, Mantovani G, Aguirre U, Barlier A, Brix B, Elli FM, Freson K, Grybek V, IZZI B, Linglart A, et al. European guidance for the molecular diagnosis of pseudohypoparathyroidism not caused by point genetic variants at GNAS: an EQA study. *Eur J Hum Genet* 2015; 23(4):560; PMID:25005735; <https://doi.org/10.1038/ejhg.2014.127>
13. Azzi S, Blaise A, Steunou V, Harbison MD, Salem J, Brioude F, Rossignol S, Habib WA, Thibaud N, Neves CD, et al. Complex tissue-specific epigenotypes in Russell-Silver syndrome associated with 11p15 ICR1 hypomethylation. *Hum Mut* 2014; 35(10):1211-20; PMID:25044976; <https://doi.org/10.1002/humu.22623>

Webpage URLs

EUCID.net webpage. www.imprinting-disorders.eu
 Locus Reference Genomic identifier. www.lrg-sequence.org
 HGVS nomenclature webpage. www.HGVS.org/varnomen