



Universiteit  
Leiden  
The Netherlands

## PLASMA HEPCIDIN LEVELS AND ANEMIA IN OLD AGE. THE LEIDEN 85-PLUS STUDY

Elzen, W. den; Craen, A.J. de; Wiegerinck, E.; Westendorp, R.; Swinkels, D.W.; Gussekloo, J.

### Citation

Elzen, W. den, Craen, A. J. de, Wiegerinck, E., Westendorp, R., Swinkels, D. W., & Gussekloo, J. (2012). PLASMA HEPCIDIN LEVELS AND ANEMIA IN OLD AGE. THE LEIDEN 85-PLUS STUDY, 52, 153-153. Retrieved from <https://hdl.handle.net/1887/116996>

Version: Not Applicable (or Unknown)

License: [Leiden University Non-exclusive license](#)

Downloaded from: <https://hdl.handle.net/1887/116996>

**Note:** To cite this publication please use the final published version (if applicable).

[Display Settings:](#)  Abstract

[Send to:](#) 

*Haematologica*. 2013 Mar;98(3):448-54. doi: 10.3324/haematol.2012.068825. Epub 2012 Oct 12.

## Plasma hepcidin levels and anemia in old age. The Leiden 85-Plus Study.

den Elzen WP, de Craen AJ, Wiegerinck ET, Westendorp RG, Swinkels DW, Gussekloo J.

Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands.  
w.p.j.den\_elzen@lumc.nl

### Abstract

Hepcidin, an important regulator of iron homeostasis, is suggested to be causally related to anemia of inflammation. The aim of this study was to explore the role of plasma hepcidin in anemia among older persons from the general population. The Leiden 85-Plus Study is a population-based study of 85-year olds in Leiden, the Netherlands. Eighty-five-year old inhabitants of Leiden were enrolled between September 1997 and September 1999. At the age of 86, plasma hepcidin was determined with time of flight mass spectrometry in 490 participants [160 (32.7%) male, 114 (23.3%) with anemia]. Anemia was defined according to criteria of the World Health Organization (hemoglobin level <13 g/dL for men and hemoglobin <12 g/dL for women). The median plasma hepcidin level was 3.0 nM [interquartile range (IQR) 1.8-4.9]. We found strong correlations between plasma hepcidin and body iron status, C-reactive protein and erythropoietin levels. Significantly higher hepcidin levels were found in participants with anemia of inflammation ( $P<0.01$ ), in participants with anemia of kidney disease ( $P=0.01$ ), and in participants with unexplained anemia ( $P=0.01$ ) than in participants without anemia. Participants with iron-deficiency anemia had significantly lower plasma hepcidin levels than participants without anemia ( $P<0.01$ ). In conclusion, older persons with anemia of inflammation have higher hepcidin levels than their counterparts without anemia. The potential clinical value of hepcidin in future diagnostic algorithms for anemia has to be explored.

PMID: 23065507 [PubMed - in process] [Free full text](#)

[+](#) **Publication Types**

[+](#) **LinkOut - more resources**

### Save items

Add to Favorites

### Related citations in PubMed

Hepcidin and iron status among preg [Asia Pac J Clin Nutr. 2008]

Anemia in heart failure: associatio [Acta Haematol. 2013]

The A736V TMPRSS6 polymorphi [BMC Nephrol. 2013]

[Review](#) Increased hepcidin-25 and erythr [Future Cardiol. 2010]

[Review](#) Hepcidin: clinical utility as a diagnosti [Kidney Int. 2011]

[See reviews...](#)

[See all...](#)

### Recent activity

[Turn Off](#) [Clear](#)

Plasma hepcidin levels and anemia in old age. TI PubMed

High blood pressure, physical and cognitive PubMed

General practitioners' experiences using cogn PMC

E-selectin is elevated in cord blood of South A PubMed

Wegener and Wegener's disease (1752) PubMed

[See more...](#)

---

<a href="#">NCBI Education</a>	<a href="#">Chemicals &amp; Bioassays</a>	<a href="#">PubMed</a>	<a href="#">Genetic Testing Registry</a>	<a href="#">About NCBI</a>
<a href="#">NCBI Help Manual</a>	<a href="#">Data &amp; Software</a>	<a href="#">Nucleotide</a>	<a href="#">PubMed Health</a>	<a href="#">Research at NCBI</a>
<a href="#">NCBI Handbook</a>	<a href="#">DNA &amp; RNA</a>	<a href="#">BLAST</a>	<a href="#">GenBank</a>	<a href="#">NCBI Newsletter</a>
<a href="#">Training &amp; Tutorials</a>	<a href="#">Domains &amp; Structures</a>	<a href="#">PubMed Central</a>	<a href="#">Reference Sequences</a>	<a href="#">NCBI FTP Site</a>
	<a href="#">Genes &amp; Expression</a>	<a href="#">Gene</a>	<a href="#">Map Viewer</a>	<a href="#">NCBI on Facebook</a>
	<a href="#">Genetics &amp; Medicine</a>	<a href="#">Bookshelf</a>	<a href="#">Human Genome</a>	<a href="#">NCBI on Twitter</a>
	<a href="#">Genomes &amp; Maps</a>	<a href="#">Protein</a>	<a href="#">Mouse Genome</a>	<a href="#">NCBI on YouTube</a>
	<a href="#">Homology</a>	<a href="#">OMIM</a>	<a href="#">Influenza Virus</a>	
	<a href="#">Literature</a>	<a href="#">Genome</a>	<a href="#">Primer-BLAST</a>	
	<a href="#">Proteins</a>	<a href="#">SNP</a>	<a href="#">Sequence Read Archive</a>	
	<a href="#">Sequence Analysis</a>	<a href="#">Structure</a>		
	<a href="#">Taxonomy</a>			
	<a href="#">Training &amp; Tutorials</a>			
	<a href="#">Variation</a>			

[Copyright](#) | [Disclaimer](#) | [Privacy](#) | [Browsers](#) | [Accessibility](#) | [Contact](#)

National Center for Biotechnology Information, U.S. National Library of Medicine  
8600 Rockville Pike, Bethesda MD, 20894 USA

