



Universiteit
Leiden
The Netherlands

Novel formaldehyde-induced modifications of lysine residue pairs in peptides and proteins: identification and relevance to vaccine development
Michiels, T.J.M.; Schöneich, C.; Hamzink, M.R.J.; Meiring, H.D.; Kersten, G.F.A.; Jiskoot, W.; Metz, B.

Citation

Michiels, T. J. M., Schöneich, C., Hamzink, M. R. J., Meiring, H. D., Kersten, G. F. A., Jiskoot, W., & Metz, B. (2020). Novel formaldehyde-induced modifications of lysine residue pairs in peptides and proteins: identification and relevance to vaccine development. *Molecular Pharmaceutics*, 17(11), 4375-4385. doi:10.1021/acs.molpharmaceut.0c00851

Version: Publisher's Version
License: [Leiden University Non-exclusive license](#)
Downloaded from: <https://hdl.handle.net/1887/139038>

Note: To cite this publication please use the final published version (if applicable).

Novel Formaldehyde-Induced Modifications of Lysine Residue Pairs in Peptides and Proteins: Identification and Relevance to Vaccine Development

Thomas J.M. Michiels,* Christian Schöneich, Martin R.J. Hamzink, Hugo D. Meiring, Gideon F.A. Kersten, Wim Jiskoot, and Bernard Metz



Cite This: *Mol. Pharmaceutics* 2020, 17, 4375–4385



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: Formaldehyde-inactivated toxoid vaccines have been in use for almost a century. Despite formaldehyde's deceptively simple structure, its reactions with proteins are complex. Treatment of immunogenic proteins with aqueous formaldehyde results in heterogeneous mixtures due to a variety of adducts and cross-links. In this study, we aimed to further elucidate the reaction products of formaldehyde reaction with proteins and report unique modifications in formaldehyde-treated cytochrome c and corresponding synthetic peptides. Synthetic peptides (Ac-GDVEKGAK and Ac-GDVEKGKK) were treated with isotopically labeled formaldehyde ($^{13}\text{CH}_2\text{O}$ or CD_2O) followed by purification of the two main reaction products. This allowed for their structural elucidation by (2D)-nuclear magnetic resonance and nanoscale liquid chromatography-coupled mass spectrometry analysis. We observed modifications resulting from (i) formaldehyde-induced deamination and formation of α,β -unsaturated aldehydes and methylation on two adjacent lysine residues and (ii) formaldehyde-induced methylation and formylation of two adjacent lysine residues. These products react further to form intramolecular cross-links between the two lysine residues. At higher peptide concentrations, these two main reaction products were also found to subsequently cross-link to lysine residues in other peptides, forming dimers and trimers. The accurate identification and quantification of formaldehyde-induced modifications improves our knowledge of formaldehyde-inactivated vaccine products, potentially aiding the development and registration of new vaccines.

KEYWORDS: formaldehyde, vaccines, antigens, NMR, mass spectrometry, structural elucidation, protein modification

INTRODUCTION

Formaldehyde is involved in a wide range of applications and processes. It is an important precursor in the synthesis of many chemicals, such as polymers and resins.¹ Furthermore, the chemical is a potent disinfectant and sterilant. It is either obtained as a 37 wt % aqueous solution (known as formalin, usually stabilized with 10–15 wt % methanol) or vaporized from paraformaldehyde. As a disinfectant, it is effective against a wide variety of bacteria, fungi, and viruses.² In addition to disinfecting equipment, formaldehyde is also used in other medical applications, for instance in dentistry.³ In histology and pathology, it is used as a fixation agent.⁴ In pharmaceutical applications, formaldehyde inactivation of antigens remains an important method for chemical inactivation of pathogens

(reviewed elsewhere⁵) in the production of vaccines almost a century after its discovery.⁶ Marketed vaccines that are inactivated this way range from toxins (e.g., diphtheria toxin and tetanus toxin) to viruses (such as the poliovirus) and bacteria (such as whole-cell *Bordetella pertussis* vaccines).^{6–8} Formaldehyde inactivation also has potential for the develop-

Received: August 18, 2020

Revised: September 23, 2020

Accepted: September 23, 2020

Published: October 5, 2020



ment of new vaccines, such as enteroviruses (i.e., enterovirus 71⁹ and coxsackieviruses¹⁰) or coronaviruses (i.e., SARS-CoV-1¹¹). Nevertheless, some formaldehyde-inactivated vaccine concepts have failed dramatically, such as the formaldehyde-inactivated respiratory syncytial virus (RSV) vaccine, which enhanced the severity of RSV.⁵ Recently, this has been linked with the usage of suboptimal concentrations of formaldehyde, resulting in misfolding of the RSV fusion protein.^{12,13} This failure underlines the importance of product characterization and a general understanding of the mechanisms involved. The use of formaldehyde itself in vaccine production is sometimes a source of criticism because of its toxicity;¹⁴ however, formaldehyde is also an endogenous product in various metabolic processes in vivo and present at higher concentrations in the human body than in vaccines.¹⁵ It has been shown that endogenous formaldehyde induces immunogenic adducts; increased immunogenicity of formaldehyde-treated proteins is also observed in some vaccine products.^{16,17} The mechanism of formaldehyde-mediated inactivation and other formaldehyde reactions with proteins and amino acids has been studied by several groups.^{18–26} As formaldehyde is mixed with a solution containing proteins, imine and hydroxymethyl adducts are formed on amines, amides, and thiols. These then react further with other amino acid residues in the mixture such as tyrosine and arginine residues. This results in intermolecular (with other proteins or amino acids in the solution) and intramolecular cross-links.¹⁹ An overview of the most common formaldehyde-induced modifications is depicted in Table S1. Although a lot of progress has been made in the identification of formaldehyde-induced modifications, the complexity and heterogeneity of the reaction products still hinder a complete understanding of the processes involved. Thorough vaccine product characterization and understanding is key to monitoring batch-to-batch consistency. If the various chemical modifications and the degree of these modifications are known, batch release could be based on in vitro tests in a so-called consistency approach, instead of traditional in vivo release tests, measuring immunogenicity and (absence of) residual toxicity.^{27–30} Moreover, better understanding of formaldehyde-inactivated vaccines can aid the development and registration of new vaccines.

In previous work, we have analyzed the influence of formaldehyde modifications on the kinetics of proteolytic digestion of various model proteins.³¹ To identify formaldehyde-induced modifications in diphtheria toxoid or model proteins, such as cytochrome c, proteins were treated with aqueous solutions of CH₂O or CD₂O.²⁰ After incubation, the resulting mixtures were pooled in a 1:1 ratio. Subsequent protease digestion was used to obtain peptides that were analyzed using nanoscale liquid chromatography-coupled mass spectrometry (LC–MS). Classic formaldehyde modifications yield mass spectral doublet peaks with a 2 Da mass difference or a multiple of 2 Da^{18,20} with equal intensities. The structure of these formaldehyde modifications in cytochrome c has been assigned.³¹ However, several atypical spectral doublet peaks were observed that could not be addressed to these classic formaldehyde modifications.

In this study, we aim to further elucidate the reaction products of formaldehyde with proteins and report new modifications in formaldehyde-treated cytochrome c and corresponding synthetic peptides. Synthetic peptides were treated with isotopically labeled formaldehyde (¹³CH₂O or CD₂O) followed by purification of the two main reaction

products. This allowed their structural elucidation using nuclear magnetic resonance (NMR) and nanoscale LC–MS analysis. These modifications involved (i) formaldehyde-induced deamination and formation of vinylic aldehydes and methylation on two adjacent lysine residues and (ii) formaldehyde-induced methylation and formylation of two adjacent lysine residues. At higher peptide concentrations, these two main reaction products were found to subsequently cross-link to lysine residues in other peptides, forming dimers and trimers.

MATERIALS & METHODS

Synthetic peptides were purchased from Pepscan with >95% purity as trifluoroacetic acid (TFA) salt. In a typical reaction, the peptides were dissolved in water (LC–MS grade; Biosolve) and added to a 100 mM phosphate buffer (pH 7.4, obtained as a 1 M solution from Sigma-Aldrich) containing 120 mM formaldehyde (Sigma-Aldrich). The reaction mixture was then placed at 40 °C, typically for 2 days. To stop the reaction and allow for nanoscale LC–MS analysis, 1 μL aliquots were diluted in 1 mL of 0.1 vol % formic acid (Merck) or 10-μL aliquots added to 90 μL 1 vol % formic acid for conventional LC–MS. To stop the reaction and allow for purification, the pH of the mixture would be adjusted to ~2 by addition of 10 vol % TFA (Sigma-Aldrich).

Reductive Methylation. Solutions of synthetic Ac-GDVEKGKK and Ac-GDVEKGKKIFVQ (1 mM) were treated with 123 mM formaldehyde (CH₂O) in phosphate-buffered saline (Gibco) with a total volume of 100 μL. After 4 days of incubation at 37 °C, an aliquot was taken for nanoscale LC–MS analysis (diluted 1:100 in 0.1 vol % FA). Subsequently, 10 μL of 1.23 M NaBH₃CN and another 10 μL of 1.23 M formaldehyde were added, and the samples were incubated for 30 min at room temperature after which another sample was taken.

Purification. Purification was achieved by reducing the total reaction volume 10-fold by vacuum concentration using an Eppendorf vacuum centrifuge. The reaction products were separated after subsequent injections of 20 μL of the sample onto an XSelect Peptide CSH C18, 130 Å, 5 μm, 4.6 mm × 250 mm semipreparative column (Waters) using an Agilent 1290 Infinity II high-performance LC (HPLC) system with UV detection at 215 nm. Eluent A consisted of water with 0.02 vol % TFA, and eluent B consisted of 80 vol % acetonitrile (LC–MS grade; Biosolve) with 20 vol % water and 0.02 vol % TFA. To separate the products, a 10 to 17% eluent B gradient in 6 min was used followed by 17 to 50% B in 1 min. Fractions were collected manually and subsequently concentrated in the vacuum centrifuge to approximately 50 μL. For NMR analysis, 600 μL of D₂O (Sigma Aldrich) and 10 μL of D₂O containing 0.75 wt % TMSP (Sigma Aldrich) were added. Products were stored at room temperature in NMR tubes at a pH < 2.5 due to residual TFA.

Stability of Product 3a. Purified 3a (90 μL) (prepared from Ac-GDVEKGKK) was taken from the NMR tube (15 μg) and mixed with 10 μL of 1 M phosphate buffer (pH 7.4). The pH was verified using pH indicator paper. Immediately afterward, a 1 μL aliquot was diluted into 1000 μL of 0.1 vol % formic acid to quench the reaction. The remaining mixture was incubated at 40 °C for 18 h and sampled subsequently.

Acetaldehyde Reaction. Acetaldehyde (Sigma-Aldrich) was dissolved in 100 mM phosphate buffer containing 0.1 mg/mL Ac-GDVEKGAK, and the mixture was placed at room




Figure 1. Representative nanoscale LC–MS base peak (m/z 400–500) chromatogram of synthetic Ac-GDVEKGKK treated with aqueous CH_2O for 46 h ($z = 2$ for all peaks). The exact regioisomers of the two peaks corresponding to product 4 could not be assigned.

temperature for 1 day. Subsequently, a $1 \mu\text{L}$ aliquot was diluted into $1000 \mu\text{L}$ of 0.1 vol % formic acid, and $1 \mu\text{L}$ of an LC–MS sample of purified 8a (from Ac-GDVEKGAK treated with $^{13}\text{C}_2\text{O}$) was added as a ^{13}C -labeled reference for nanoscale LC–MS analysis.

Nanoscale LC–MS. The peptides were analyzed by reversed-phase nanoscale LC–MS using a vented column system as described by Meiring et al.³² For this, an Agilent 1290 Infinity HPLC system was used in conjunction with a $100 \mu\text{m}$ inner diameter (I.D.) trapping column packed to a bed length (L) of 20 mm with Reprosil-Pur C18-AQ $5 \mu\text{m}$ particles and a 31 cm, $L \times 50 \mu\text{m}$ I.D. analytical column packed with Reprosil-Pur C18-AQ $3 \mu\text{m}$ particles coupled to a gold-coated nanoelectrospray ionization spray tip with a $3.5 \mu\text{m}$ tip diameter (all prepared in-house). The sample ($10 \mu\text{L}$) was injected onto the trapping column. The trapping column was then washed with 0.1 vol % formic acid in water at a $5 \mu\text{L}/\text{min}$ flow rate for 10 min. The gradient consisted of water with 0.1 vol % formic acid (eluent A) and acetonitrile with 0.1 vol % formic acid (eluent B). The gradient started at 4% B to 34% in 15 min followed by washing steps and re-equilibration. A flow restrictor was used to ensure a flow rate of $\sim 125 \text{ nL}/\text{min}$ through the analytical column. The LC system was coupled to an Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo Fisher). Analysis was done in data-dependent acquisition mode, with MS^1 scans at 120,000 full width at half-maximum (FWHM) resolution from 300 to 1500 m/z . Collision-induced dissociation (CID) fragmentation MS^2 scans were measured in either iontrap or orbitrap mode, where the orbitrap resolution would be decreased to 7500 FWHM resolution for an increased acquisition speed. All MS^1 orbitrap m/z readouts were corrected using fluoranthene as an internal mass calibrant. Typical mass errors expected on MS^1 were <1 and <20 ppm on MS^2 (in orbitrap measurements). *Note:* for ease of reading, the number of digits behind the decimal separator in some figures is lower than those still considered accurate.

Conventional LC–MS. Reaction kinetics of Ac-GDVEKGKK, Ac-GDVEKGAK, Ac-GDVEAGKK, and Ac-GDVEKGKA were compared using conventional LC–MS. To


this end, we used an Agilent Poroshell 120 EC C18 column ($2.1 \text{ mm I.D.} \times 50 \text{ mm L}$) packed with $1.9 \mu\text{m}$ particles with an Agilent 1200SL HPLC system coupled to an LTQ Orbitrap XL mass spectrometer (Thermo Fisher) with electrospray ionization. The same eluent system as with nanoscale LC was used with a 1 to 30% B in a 4-min gradient. The sample ($5 \mu\text{L}$) at a concentration of 0.01 mg/mL peptide was injected. Measurements were performed in the ion trap at a normal scan rate from 300 to 900 m/z .

Nuclear Magnetic Resonance. NMR experiments were carried out using a JEOL JNM-ECZ400S/L1 400 MHz NMR spectrometer in D_2O with trimethylsilylpropanoic acid as internal reference. Default experimental settings as provided by JEOL were used. ^1H -NMR spectra were recorded using Robust-5 water suppression (1064 scans). Standard ^{13}C spectra and DEPT-135 and DEPT-90 spectra were recorded until the signal/noise ratio was sufficient (typically, $>30,000$ scans). For heteronuclear single quantum coherence (HSQC) and heteronuclear multiple bond correlation (HMBC), the HSQC_wet and HMBC_wet experiments were used to suppress the water signal.

RESULTS

Detection of Unusual Formaldehyde Modifications in Cytochrome c. In a previous study, chemical modifications were identified in cytochrome c after formaldehyde treatment.³¹ However, some unexpected modifications were identified using the PEAKS Studio PTM module, which could not be explained. These observations triggered further investigation. In the previous study, proteins were treated with CH_2O and CD_2O separately, and after incubation, the samples were mixed at a 1:1 ratio and digested with the protease cathepsin S. Instead of the usual 1:1 ratio of the light and heavy peptide, the isotope pattern was skewed toward the light peptide (Figure S1). This observation triggered further investigation. Moreover, the mass increase after formaldehyde treatment was $+42.0106 \text{ Da}$, which commonly corresponds to acetylation ($+C_2H_2O$, on top of the standard acetylation of the cytochrome c N-terminus), but acetylation by formaldehyde is not obvious. As the mass increase between the light peptide

Scheme 1. Ac-GDVEKGXX (X = K or A) Reacts with Formaldehyde in Aqueous Buffer to Form 2a-b (a-b Represent Two Regioisomers), 1a-b, and Tautomers 3a-b^a



^aIf X = K, 4-I and 4-II are formed as well. Reaction conditions: 100 mM formaldehyde, 100 mM phosphate buffer (pH 7.4) in water at 40 °C for 24 h.




Figure 2. Comparison of the reaction between synthetic analogues of peptide Ac-GDVEKGKK (where one of the various lysine residues is replaced with an alanine residue) and the use of CH₂O or CD₂O on the formation of products 1 and 2. The area of the regioisomers has been summed.

and the heavy peptide was 4 Da, two formaldehyde molecules (each containing two deuterons) must have been incorporated in the modification. Peptides derived from formaldehyde-treated cytochrome c that included this mass increase (+42.0106 Da) were Ac-GDVEKGKK, Ac-GDVEKGKKIFVQ, KGKKHKKTGPNL, and AYLKK. All these peptides contain at

least two lysine residues, making the involvement of two lysine residues in one modification or cross-link very likely.

Formaldehyde Modifications in Synthetic Ac-GDVEKGKK. To study this modification in detail, a synthetic peptide with the sequence Ac-GDVEKGKK was subjected to 120 mM formaldehyde (CH₂O or CD₂O) in 100 mM




Figure 3. ^1H - ^{13}C HSQC and ^1H - ^{13}C HMBC of purified **1b** obtained from a reaction with Ac-GDVEKGAK with ^{13}C -labeled formaldehyde (adducts containing ^{13}C are marked in red, and ^{13}C NMR shifts are also marked in red). F1: DEPT-135 and F2: Robust5- ^1H (for water suppression) spectra.

phosphate buffer (pH 7.4). Treatment of the native peptide resulted in the same modification with $\Delta M = +42.0106$ Da, as observed for cytochrome c, along with a variety of other modifications. The most important of these other modifications were $\Delta M = +24.9840$ Da ($+[C_2O]-[NH]$) and $\Delta M = +73.0051$ Da ($+[C_3H_3O_3]-[N]$), where the latter is suggested to be an analogue in which an additional molecule has reacted with the former. A representative chromatogram is shown in Figure 1. Because of the complicated nature and the many reaction products formed, we decided to focus on the most important modifications, which form the basis for the other reaction products: $\Delta M = +24.9840$ Da (**1a-b**) and $\Delta M = +42.0106$ Da (**2a-b**). The proposed structures for these modifications are depicted in Scheme 1. Experimental evidence for these structures is described below.

Reaction Kinetics. To investigate which lysine residues were involved in these formaldehyde modifications ($+24.9860$ and $+42.0106$ Da), peptides with the general sequence Ac-GDVEGXGXX were made, where $X = K$ or A in all possible permutations. Treatment of the peptides that contained only one lysine did not result in the formation of significant additional peaks, with the exception of a small amount of the classic hydroxymethyl and imine adducts (data not shown). The other peptides showed different degrees of modification, where the extent of modification decreased in the following order: Ac-GDVEKGKK \equiv Ac-GDVEKGAK $>$ Ac-GDVEKGKA \gg Ac-GDVEAGKK (Figure 2). The formation of **2** was diminished by the use of deuterated formaldehyde (72%

reduction for Ac-GDVEKGKK), while the effect of deuterated formaldehyde on the formation of **1** was less substantial. Ac-GDVEKGKK and Ac-GDVEKGAK form similar amounts of **2**, with similar kinetics, but **1** was formed faster from Ac-GDVEKGAK. The use of D_2O as a solvent instead of H_2O did not affect the structure of the reaction products but slightly decreased the speed of the reaction (Figure S2).

Reductive Methylation of the Reaction Products. To elucidate the structures of reaction products **1** and **2**, the reaction mixture of Ac-GDVEKGKK with formaldehyde was subjected to reductive methylation by the addition of more formaldehyde and NaCNBH_3 . Unreacted Ac-GDVEKGKK was methylated 6 times, twice on each lysine residue. Although a variety of reaction products were formed, some of the main peaks observed were those corresponding to products **1** and **2** with three additional methyl groups. Thus, the original formaldehyde-induced adducts on **1** and **2** were both stable under mild reducing conditions, and three positions on the amine groups could still form imines and the corresponding methyl groups (Scheme S1). These results are consistent with the proposed structures **1** and **2**.

Purification and Stability of Products 1 and 2. To enable NMR analysis of products **1** and **2**, the reaction mixture of $^{13}\text{CH}_2\text{O}$ and Ac-GDVEKGAK was purified by reversed-phase, preparative HPLC. Ac-GDVEKGAK instead of Ac-GDVEKGKK (found in cytochrome c) was used as the reactions were cleaner, which simplifies the purification. The products were reasonably stable in an aqueous solution at a




Figure 4. ^1H - ^{13}C HSQC and ^1H - ^{13}C HMBC of purified **2a** obtained from a reaction with Ac-GDVEKGAK with ^{13}C -labeled formaldehyde (adducts containing ^{13}C are marked in red, and ^{13}C NMR shifts are also marked in red). F1: DEPT-135 and F2: RobustS- ^1H (for water suppression) spectra.




Figure 5. High-resolution MS^2 of **2a**: MS^2 spectrum of Ac-GDVEKGKK treated with (A) CH_2O , (B) $^{13}\text{CH}_2\text{O}$, and (C) CD_2O .




Figure 6. High-resolution MS² of **1a**: MS² spectrum of Ac-GDVEKGKK treated with (A) CH₂O, (B) ¹³CH₂O, and (C) CD₂O.

low pH (<2.5) but not stable when drying, and thus, no dry products **1** and **2** could be obtained. Instead of completely removing the solvent, the solutions were concentrated with almost complete removal of the acetonitrile. The pH was typically <3 due to residual TFA. Typical yields (NMR) were around 10%.

Despite product **2a** being baseline separated from the chromatographically succeeding products **2b**, **1a**, and **1b**, the concentrated solution always contained some **1a** (Figure S3A). Presumably, **1a** was formed during concentration or other steps of the workup, after collection of **2a**. Likewise, the purification of **1a** always resulted in some **3a** being present, despite this chromatographic peak eluting at the same time as the removed product **2a** (Figure S3 B). The presence of these impurities indicates that **1a** can be formed from **2a** and that **3a** can be formed from **1a** without the presence of additional formaldehyde.

To further investigate the stability of the reaction products, purified **2a** (from Ac-GDVEKGKK) was placed in phosphate buffer at pH 7.4 and stored at 37 °C for 1 day. Despite **2a** being stable for weeks at low pH, exposure to pH 7.4 resulted in the formation of **1a** and **1b**, although **1a** was the more abundant degradation product (Figure S4).


NMR of Purified 1b and 2a. By using ¹³CH₂O, sufficient material could be obtained to perform ¹³C NMR experiments on two purified fractions containing **1b** or **2a** derived from Ac-GDVEKGAK. However, only carbons originating from the formaldehyde modifications could be detected this way. To provide further structural information, ¹H (Robust-5 water suppression), DEPT, and HSQC were combined with HMBC (Figure 3 and Figure 4, described in detail in the Supporting

Information) and confirmed the structures of **1b** and **2a**, respectively. Despite purification, nanoscaleLC–MS showed more than one peak, although the major product was **1b**. Indeed, in the NMR analysis, two ¹³C signals were observed other than the peaks assigned to **1b**. These signals correspond to structure **3** (a tautomer of **1**) and free formaldehyde.

MS² Analysis of Products 1, 2, and 3. Subjecting Ac-GDVEKGKK, exposed to various isotopically enriched formaldehyde solutions (CH₂O, ¹³CH₂O, and CD₂O), to MS² analysis supported the NMR findings. Structures **1** and **2** were both found to exist as regioisomers, with mainly lysine-5 and lysine-8 being modified. Either lysine-8 contained the methyl group (structures **1a** and **2a**), with the other modification on lysine-5, or vice versa (structures **1b** and **2b**) (Figure S5).

Fragmentation of **2a** revealed that the y1 ion contains an additional +14.02 Da, consistent with methylation of the C-terminal lysine (Figure 5A). ¹³C-labeled formaldehyde resulted in the y1, y2, and y3 ions to be 1.00 Da heavier than the corresponding y ions resulting from regular formaldehyde treatment (Figure 5B). ¹³C-labeled formaldehyde resulted in a y4 ion with $\Delta M = +2.01$ Da, which is in compliance with the proposed structure and the incorporation of two formaldehyde molecules on the peptide. Fragmentation of CD₂O-treated Ac-GDVEKGKK resulted in the y1 ion being 3.02 Da heavier than that obtained after CH₂O treatment, indicating that a deuteride was transferred from the modification on lysine-5 to the C-terminal lysine. Indeed, the mass difference between the y3 and y4 ions after CD₂O treatment was 1.01 Da ($\Delta M = K + 29.00$) compared with the y3 and y4 ions obtained after CH₂O treatment ($\Delta M = K + 27.99$).

Scheme 2. Proposed Mechanism for the Formation of 1 and 2



Fragmentation of CH_2O -treated **1a** revealed methylation of lysine-8 and a $\Delta M = +10.97$ Da ($-\text{[NH}_3\text{]} + \text{[CO]}$) on lysine-5 (Figure 6A). Contrary to **2a**, CD_2O -treated **1a** showed a methylation of the C-terminal lysine with just 2 deuterium atoms (Figure 6C). The γ_4 ions of **1a** show that the C-terminal lysine contains a modification that is 2 Da heavier when using CD_2O , corresponding to incorporation of both deuterium atoms on this residue.

MS^2 analysis of the chromatographic peaks corresponding to structure **3b** showed specific fragmentation resulting in only the $\gamma_4\text{-H}_2\text{O}$ ion (data not shown). The lack of backbone fragmentation between lysine-5 and lysine-8 supports a cyclic structure. Ionization of **3a** already resulted in significant fragmentation and detection of the b_4 ion in the MS^1 analysis.

MS^2 of Product 4. Product **4** was only observed for the peptide containing three lysine residues. CID (ion trap detection) fragmentation showed identical ions for the two isobaric chromatographic peaks but at different relative intensities (Figure S6). This would fit having both the Markovnikov and anti-Markovnikov product present but with different intensities. From these data, the two isomers could not be assigned to a specific chromatographic peak.

Dimerization and Oligomerization. In order to obtain as much purified reaction product as possible, the reaction of formaldehyde was performed with various concentrations of synthetic Ac-GDVEKGAK. At the lowest peptide concentration (0.25 mg/mL), nanoscale LC-MS analysis revealed a few reaction products, mainly **1** and **2** (Figure S7). When the peptide concentrations were increased to as high as 5.75 mg/mL, reaction products other than **1** and **2** started to form. Some of the most intense peaks may belong to cross-linked structures, such as trimer **7** (Figure S7) and dimers **8** and **9**. These proposed structures are based on their total mass (MS^1 mass error < 1 ppm). Both **1** and **2** seem to be involved in oligomerization, where **1** can form multiple reaction products, such as hydrogen-bond-stabilized enolates (**7**) or hetero-Diels-Alder products (**8**). A variety of reaction products with different numbers of methyl groups were observed.

Reactions with Acetaldehyde. The formation of isobaric reaction products under the influence of acetaldehyde is described in the Supporting Information.

DISCUSSION

Two new groups of formaldehyde-induced modifications on peptide sequences containing (at least) two lysine residues were identified. The first modification involved methylation and formylation of the lysine pair, resulting in $\Delta M = +42.01$ Da (product **2**). In the second modification, one lysine of the lysine pair was converted to an α,β -unsaturated aldehyde, while the other lysine residue was methylated (product **1**). Structure **1** could also form a tautomer where both lysine residues are cross-linked (product **3**). While lysine-lysine cross-links in the form of amins have been suggested,⁵ to our knowledge, no NMR evidence for formaldehyde-induced lysine-lysine cross-links has been reported yet.

The formaldehyde modification (product **2**) ($\Delta M = +42.01$ Da) was not found in our previous studies,²⁰ where we used proteins treated with either CH_2O or CD_2O in a 1:1 mixture. Hydroxymethyl adducts exchange their oxygen atom with the oxygen atom in water through the equilibrium with the corresponding imine. However, deuterons of the common formaldehyde-induced protein modifications were not expected to be exchanged.¹⁹ This assumption implies that formaldehyde-induced modifications are always present at an almost perfect 1:1 ratio. In our current study, we identified a new modification where slower reaction kinetics with deuterated formaldehyde resulted in a skewed isotope pattern. These reaction kinetics point to the transfer of a hydride (or deuteride) from one part of the molecule to another. This transfer would be the rate-limiting step in the formation of **2**. Our suggested mechanism for the formation of **2** (Scheme 2) involves the formation of an imine on the $\epsilon\text{-NH}_2$ group of one of two lysine residues and the corresponding hydroxymethyl on the other lysine residue. We propose that the nitrogen of the imine acts as a base, deprotonating the hydroxymethyl OH; and subsequent transfer of the hydroxymethyl α -hydride to the adjacent imine forms product **2**. This last step would explain the observed kinetic isotope effect, which is further supported by MS^2 analysis of the deuterated **2a**, showing a methylated γ_1 ion 3 Da heavier than its nondeuterated counterpart. NMR evidence for this structure is strong with ^1H , DEPT-90, DEPT-135, and ^1H - ^{13}C HMBC supporting this structure. Formylation in combination with methylation by formaldehyde has been

observed before in solutions of the amino acid ornithine but not on lysine,²¹ indicating that the orientation of the amino groups is pivotal for this reaction. Indeed, the spacing between the lysine residues was important for this reaction to occur; the fastest reaction rates were observed when two spacing amino acids were located between the two lysine residues.

The modification with $\Delta M = +24.98$ Da (**1**) was less affected by the use of deuterated formaldehyde. Our suggested mechanism for the formation of **1** (Scheme 2) again starts with formation of an imine on one of the lysine residues. However, instead of involvement of another formaldehyde molecule, the side chain of the nearby lysine residue is oxidized to an imine. The hydrogen atom of the lysine side chain is transferred to the imine adduct originating from formaldehyde to form intermediate **12**. Subsequent hydrolysis of imine **12** results in aldehyde **13**. Similar to the formation of **2**, this results in methylation of one lysine residue; however, when deuterated formaldehyde was used, not three but two deuterons were present on the methyl group, and the other hydrogen originates from the lysine side chain. Intermediate **13** has been observed by LC–MS. Tautomerization of **13** leads to enolate **14**, which can then perform a nucleophilic attack on a formaldehyde molecule to give **17**. Subsequent β -elimination gives product **1**. Product **1** could tautomerize to form enol **3**. This enol could be stabilized by a hydrogen bond between the lysine nitrogen and the enol proton. Hydrogen-bond-stabilized enols are well described.³³ The limited fragmentation of **3** during MS² analysis further supports a cyclic structure, with a bridge between the two lysine residues, which hinders complete fragmentation of the γ_4 ion. The high amount of reactive intermediates would suggest that multiple reaction products could be formed through various pathways. Indeed, the NMR spectra of purified reaction products always contained some impurities, likely formed due to side reactions after purification. The main impurity in purified **2a** is **3**, a tautomer of **1**, which happens to co-elute with **2a**. Furthermore, formaldehyde is present in the purified samples. While it is possible that this is due to incomplete removal of formaldehyde through the preparative HPLC, it is more likely that formaldehyde is released from the reaction products through equilibria, as formaldehyde is present in a similar amount. Storage at a low pH (<3) seemed to keep both products **1** and **2** stable for weeks probably because protonation of the amines decreases nucleophilicity and prevents them from acting as a Brønsted base, which could catalyze further reactions. Increasing the pH (to 7.4) resulted in relatively fast (days) conversion of **2** to **1**. Due to the reactivity of α,β -unsaturated aldehydes, we expect these reaction products to react further when exposed to suitable nucleophiles in biological matrices. Indeed, the modification with $\Delta M = +24.98$ Da (**1**) was observed only in reactions with synthetic peptides, presumably, either because the modification is not formed in cytochrome *c* due to folding or because of the modified lysine residues reacting further with other amino acids to form other reaction products. As a large number of low-intensity side products were observed by nanoscale LC–MS after formaldehyde treatment of a simple purified peptide, high reactivity of the reaction products combined with a more complicated protein would be a likely explanation for the difference we observed between peptide reactivity and the presence of these modifications in cytochrome *c*.

Product **4** is likely formed from **1b** (Ac-GDVEKGKK), as the modification of lysine-8 involves loss of a nitrogen atom,

similar to product **1b**. The presence of the b_7 ion supports methylation on either lysine-5 or lysine-7 and not lysine-8, while y_1 -related ions confirm modification of lysine-8 by $\Delta M = +58.9895$ Da. Overall, product **4** is thought to be one of many reaction products that are eventually formed from **1** (or **2**). Our suggested mechanism involves the formation of a hydroxymethylated product (**1b**) as an intermediate (Scheme S2). Purification and structural elucidation of all reaction products present in this peptide–formaldehyde reaction mixture are beyond the scope of the present study; thus, no further purification and subsequent NMR analysis of **4** or other side products were attempted.

Several dimers and trimers were also observed. The most probable structures are suggested based on their accurate mass ($\Delta M < 1$ ppm), as MS² fragmentation was not informative and purification and subsequent NMR analysis were deemed unfeasible. Because of the relatively simple reaction components being the peptide and formaldehyde, the possible combinations that lead to the same exact mass are limited. The bulk of the mass has to be made up by a number of peptides, where the residual mass is likely a combination of the main reaction products **1** and **2** along with other formaldehyde adducts. Trimer **7** is most likely a ring of three peptides, one peptide is one of the regioisomers of **1**, and the other two are unmodified peptides. Although a linear structure with one aminal and a hetero-Diels–Alder product similar to dimer **8** is possible, a ring would be more thermodynamically favorable. Hydrogen bonding between the enol hydrogen and the amine could further stabilize this product. Aminals are not very stable in water, especially at a lower pH, but in a ring system held together in a hetero-Diels–Alder product, this reaction could reverse, making the cyclic structure more favorable than a linear one. It is interesting that we managed to observe these aminals, as prior to analysis, aliquots of the reaction mixture were diluted in 0.1 vol % formic acid, which would usually result in rapid hydrolysis of the aminal.³⁴ Two species of dimers were observed, one consisting of reaction product of **2** with an unmodified peptide (**9**) and one consisting of a reaction product of **1** with an unmodified peptide (**8**). To get to the mass of product **9**, two extra formaldehyde molecules and three extra ring double bond equivalents are required. Thus, an aminal formed from the methylated lysine and lysine with an imine adduct, and a new ring involving the formylated lysine and a lysine with an imine adduct is the most logical explanation. Product **8** most likely consists of the product of a hetero-Diels–Alder reaction between an imine adduct on a lysine and the vinyl aldehyde of product **1** and an aminal connecting the other lysine residues. The imine corresponding to the aminal of **8** would have the same total mass, but it would be entropically favorable that the aminal forms first, so that the intramolecular Diels–Alder reaction can happen afterward as these reactions require decent activation energy.^{35,36} In reality, the reaction mixture probably contains a mixture of the cyclic reaction product and the imine, which are in equilibrium.

CONCLUSIONS

Overall, we identified two new groups of formaldehyde-induced modifications of lysine residue pairs in peptides. Both the formylation/methylation modification pair and the α,β -unsaturated aldehyde/methylation modification pair were involved in further cross-linking reactions, which should be addressed in future studies. These reaction products should be taken into consideration when analyzing biological samples

exposed to exogenous or endogenous formaldehyde. Especially, biologicals, such as toxoid vaccines, are notoriously heterogeneous because of their formaldehyde modifications.^{20,37} These newly discovered modifications could help us to further understand the nature of these vaccine products. As formaldehyde-induced modifications are linked to alterations in immunogenicity,^{16,17} their accurate identification and quantification may improve our understanding of formaldehyde-inactivated vaccine products, potentially aiding the development and registration of future formaldehyde-inactivated vaccines.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.molpharmaceut.0c00851>.

Examples of the most common classic formaldehyde modifications; reaction kinetics of synthetic Ac-GDVEKGAK in D₂O; Chromatograms and spectra of structure **1** in cytochrome c; Chromatograms of purified **1a** and **2a**; high-resolution and CID fragmentation MS² spectra comparing **2a** and **2b**; base peak LC–MS chromatogram and representative averaged MS¹ spectra of dimers and trimers; a detailed description of NMR spectra; a description of similar reactions with acetaldehyde instead of formaldehyde; Methylated products **1a** and **2a** after additional formaldehyde and reduction by NaCNBH₃; proposed mechanism for the formation of **4** from **1b**; proposed reaction of peptide Ac-GDVEKGAK with acetaldehyde; chromatogram of acetaldehyde-treated Ac-GDVEKGAK; and MS² comparison between products **1a** and **5a** (PDF).

■ AUTHOR INFORMATION

Corresponding Author

Thomas J.M. Michiels – Division of BioTherapeutics, Leiden Academic Centre for Drug Research (LACDR), Leiden University, 2333 CC Leiden, The Netherlands; Intravacc, Institute for Translational Vaccinology, 3721 MA Bilthoven, The Netherlands; orcid.org/0000-0003-1517-0312; Phone: +31307920584; Email: Thomas.michiels@intravacc.nl

Authors

Christian Schöneich – Department of Pharmaceutical Chemistry, The University of Kansas, Lawrence, Kansas 66047, United States; orcid.org/0000-0001-5082-8672

Martin R.J. Hamzink – Intravacc, Institute for Translational Vaccinology, 3721 MA Bilthoven, The Netherlands

Hugo D. Meiring – Intravacc, Institute for Translational Vaccinology, 3721 MA Bilthoven, The Netherlands

Gideon F.A. Kersten – Division of BioTherapeutics, Leiden Academic Centre for Drug Research (LACDR), Leiden University, 2333 CC Leiden, The Netherlands; Intravacc, Institute for Translational Vaccinology, 3721 MA Bilthoven, The Netherlands

Wim Jiskoot – Division of BioTherapeutics, Leiden Academic Centre for Drug Research (LACDR), Leiden University, 2333 CC Leiden, The Netherlands

Bernard Metz – Intravacc, Institute for Translational Vaccinology, 3721 MA Bilthoven, The Netherlands; orcid.org/0000-0001-6814-7656

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.molpharmaceut.0c00851>

Funding

This work was supported, in part, by the Ministry of Agriculture, Nature, and Food Quality, the Netherlands.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors thank Joost Uittenbogaard and Maarten Danial for their advice, Marjolein Zohlandt for critical review of the manuscript, and Aldolfo Botana of JEOL for assistance with the NMR analysis.

■ REFERENCES

- (1) Christjanson, P.; Pehk, T.; Siimer, K. Hydroxymethylation and polycondensation reactions in urea–formaldehyde resin synthesis. *J. Appl. Polym. Sci.* **2006**, *100*, 1673–1680.
- (2) Rutala, W.; Weber, D. HICPAC Guideline for Disinfection and Sterilization in Healthcare Facilities. <https://www.cdc.gov/infectioncontrol/guidelines/disinfection/> (31-Jul-2020).
- (3) Chandrashekhara, S.; Shashidhar, J. Formocresol, still a controversial material for pulpotomy: A critical literature review. *J. Restor. Dent.* **2014**, *2*, 114.
- (4) Buesa, R. J. Histology without formalin? *Ann. Diagn. Pathol.* **2008**, *12*, 387–396.
- (5) Delrue, I.; Verzele, D.; Madder, A.; Nauwynck, H. J. Inactivated virus vaccines from chemistry to prophylaxis: merits, risks and challenges. *Expert Rev. Vaccines* **2012**, *11*, 695–719.
- (6) Glenny, A. T.; Hopkins, B. E. Diphtheria Toxoid as an Immunising Agent. *Br. J. Exp. Pathol.* **1923**, *4*, 283–288.
- (7) Gupta, R. K.; Sharma, S. B.; Ahuja, S.; Saxena, S. N. The effects of different inactivating agents on the potency, toxicity and stability of pertussis vaccine. *J. Biol. Stand.* **1987**, *15*, 87–98.
- (8) WHO Production and Control of Tetanus Vaccine: Module III: Principles of Tetanus Vaccine Production. (WHO/VSQ/94.4). (26th of March).
- (9) Reed, Z.; Cardoso, M. J. Status of research and development of vaccines for enterovirus 71. *Vaccine* **2016**, *34*, 2967–2970.
- (10) Stone, V. M.; Hankaniemi, M. M.; Laitinen, O. H.; Sioofy-Khojine, A. B.; Lin, A.; Diaz Lozano, I. M.; Mazur, M. A.; Marjomaki, V.; Lore, K.; Hyoty, H.; Hytonen, V. P.; Flodstrom-Tullberg, M. A hexavalent Coxsackievirus B vaccine is highly immunogenic and has a strong protective capacity in mice and nonhuman primates. *Sci. Adv.* **2020**, *6*, eaaz2433.
- (11) Darnell, M. E.; Plant, E. P.; Watanabe, H.; Byrum, R.; St Claire, M.; Ward, J. M.; Taylor, D. R. Severe acute respiratory syndrome coronavirus infection in vaccinated ferrets. *J. Infect. Dis.* **2007**, *196*, 1329–1338.
- (12) Killikelly, A. M.; Kanekiyo, M.; Graham, B. S. Pre-fusion F is absent on the surface of formalin-inactivated respiratory syncytial virus. *Sci. Rep.* **2016**, *6*, 34108.
- (13) Zhang, W.; Zhang, L. J.; Zhan, L. T.; Zhao, M.; Wu, G. H.; Si, J. Y.; Chen, L.; Lin, X.; Sun, Y. P.; Lin, M.; Yu, C.; Fang, M. J.; Wang, Y. B.; Zheng, Z. Z.; Xia, N. S. The Optimal Concentration of Formaldehyde is Key to Stabilizing the Pre-Fusion Conformation of Respiratory Syncytial Virus Fusion Protein. *Viruses* **2019**, *11*, 628.
- (14) Kata, A. A postmodern Pandora's box: Anti-vaccination misinformation on the Internet. *Vaccine* **2010**, *28*, 1709–1716.
- (15) Mitkus, R. J.; Hess, M. A.; Schwartz, S. L. Pharmacokinetic modeling as an approach to assessing the safety of residual formaldehyde in infant vaccines. *Vaccine* **2013**, *31*, 2738–2743.

- (16) Nakamura, J.; Shimomoto, T.; Collins, L. B.; Holley, D. W.; Zhang, Z.; Barbee, J. M.; Sharma, V.; Tian, X.; Kondo, T.; Uchida, K.; Yi, X.; Perkins, D. O.; Willis, M. S.; Gold, A.; Bultman, S. J. Evidence that endogenous formaldehyde produces immunogenic and atherogenic adduct epitopes. *Sci. Rep.* **2017**, *7*, 10787.
- (17) Metz, B.; Jiskoot, W.; Hennink, W. E.; Crommelin, D. J.; Kersten, G. F. Physicochemical and immunochemical techniques predict the quality of diphtheria toxoid vaccines. *Vaccine* **2003**, *22*, 156–167.
- (18) Metz, B.; Kersten, G. F.; Baart, G. J.; de Jong, A.; Meiring, H.; ten Hove, J.; van Steenberghe, M. J.; Hennink, W. E.; Crommelin, D. J.; Jiskoot, W. Identification of formaldehyde-induced modifications in proteins: reactions with insulin. *Bioconjugate Chem.* **2006**, *17*, 815–822.
- (19) Metz, B.; Kersten, G. F.; Hoogerhout, P.; Brugghe, H. F.; Timmermans, H. A.; de Jong, A.; Meiring, H.; ten Hove, J.; Hennink, W. E.; Crommelin, D. J.; Jiskoot, W. Identification of formaldehyde-induced modifications in proteins: reactions with model peptides. *J. Biol. Chem.* **2004**, *279*, 6235–6243.
- (20) Metz, B.; Michiels, T.; Uittenbogaard, J.; Danial, M.; Tilstra, W.; Meiring, H. D.; Hennink, W. E.; Crommelin, D. J. A.; Kersten, G. F. A.; Jiskoot, W. Identification of formaldehyde-induced modifications in diphtheria toxin. *J. Pharm. Sci.* **2019**, *109*, 543–557.
- (21) Kamps, J. J. A. G.; Hopkinson, R. J.; Schofield, C. J.; Claridge, T. D. W. How formaldehyde reacts with amino acids. *Commun. Chem.* **2019**, *2*, 126.
- (22) Trezl, L.; Rusznak, I.; Tyihak, E.; Szarvas, T.; Szende, B. Spontaneous N epsilon-methylation and N epsilon-formylation reactions between L-lysine and formaldehyde inhibited by L-ascorbic acid. *Biochem. J.* **1983**, *214*, 289–292.
- (23) Yamada, M.; Funaki, S.; Miki, S. Formaldehyde interacts with RNA rather than DNA: Accumulation of formaldehyde by the RNA-inorganic hybrid material. *Int. J. Biol. Macromol.* **2019**, *122*, 168–173.
- (24) Gold, T. B.; Smith, S. L.; Digenis, G. A. Studies on the influence of pH and pancreatin on 13C-formaldehyde-induced gelatin cross-links using nuclear magnetic resonance. *Pharm. Dev. Technol.* **1996**, *1*, 21–26.
- (25) Toews, J.; Rogalski, J. C.; Clark, T. J.; Kast, J. Mass spectrometric identification of formaldehyde-induced peptide modifications under in vivo protein cross-linking conditions. *Anal. Chim. Acta* **2008**, *618*, 168–183.
- (26) Toews, J.; Rogalski, J. C.; Kast, J. Accessibility governs the relative reactivity of basic residues in formaldehyde-induced protein modifications. *Anal. Chim. Acta* **2010**, *676*, 60–67.
- (27) Halder, M.; Depraetere, H.; Delannois, F.; Akkermans, A.; Behr-Gross, M. E.; Bruysters, M.; Dierick, J. F.; Jungback, C.; Kross, I.; Metz, B.; Pennings, J.; Rigsby, P.; Riou, P.; Balks, E.; Dobby, A.; Leroy, O.; Stirling, C. Recommendations of the VAC2VAC workshop on the design of multi-centre validation studies. *Biologicals* **2018**, *52*, 78–82.
- (28) De Mattia, F.; Chapsal, J. M.; Descamps, J.; Halder, M.; Jarrett, N.; Kross, I.; Mortiaux, F.; Ponsar, C.; Redhead, K.; McKelvie, J.; Hendriksen, C. The consistency approach for quality control of vaccines - a strategy to improve quality control and implement 3Rs. *Biologicals* **2011**, *39*, 59–65.
- (29) De Mattia, F.; Hendriksen, C.; Buchheit, K. H.; Chapsal, J. M.; Halder, M.; Lambrigts, D.; Redhead, K.; Rommel, E.; Scharton-Kersten, T.; Sesardic, T.; Viviani, L.; Ragan, I. The vaccines consistency approach project: an EPAA initiative. *Pharmeur. Bio. Sci. Notes* **2015**, *2015*, 30–56.
- (30) Metz, B.; Jiskoot, W.; Mekkes, D.; Kingma, R.; Hennink, W. E.; Crommelin, D. J.; Kersten, G. F. Quality control of routine, experimental and real-time aged diphtheria toxoids by in vitro analytical techniques. *Vaccine* **2007**, *25*, 6863–6871.
- (31) Michiels, T. J. M.; Meiring, H. D.; Jiskoot, W.; Kersten, G. F. A.; Metz, B. Formaldehyde treatment of proteins enhances proteolytic degradation by the endo-lysosomal protease cathepsin S. *Sci. Rep.* **2020**, *10*, 11535.
- (32) Meiring, H. D.; van der Heeft, E.; ten Hove, G. J.; de Jong, A. P. J. M. Nanoscale LC–MS(n): technical design and applications to peptide and protein analysis. *J. Sep. Sci.* **2002**, *25*, 557–568.
- (33) Imashiro, F.; Maeda, S.; Takegoshi, K.; Terao, T.; Saika, A. Structure-dependent intermolecular hydrogen-bond effects on carbon-13 NMR chemical shifts in enol forms of 1,3-diketones. *J. Am. Chem. Soc.* **1987**, *109*, 5213–5216.
- (34) Sawatzky, E.; Drakopoulos, A.; Rölz, M.; Sotriffer, C.; Engels, B.; Decker, M. Experimental and theoretical investigations into the stability of cyclic amins. *Beilstein J. Org. Chem.* **2016**, *12*, 2280–2292.
- (35) Carneiro de Oliveira, J.; Laborie, M. P.; Roucoules, V. Thermodynamic and Kinetic Study of Diels-Alder Reaction between Furfuryl Alcohol and N-Hydroxymaleimides-An Assessment for Materials Application. *Molecules* **2020**, *25*, 243.
- (36) Otto, S.; Blokzijl, W.; Engberts, J. B. F. N. Diels-Alder Reactions in Water. Effects of Hydrophobicity and Hydrogen Bonding. *J. Organ. Chem.* **1994**, *59*, 5372–5376.
- (37) Bayart, C.; Peronin, S.; Jean, E.; Paladino, J.; Talaga, P.; Borgne, M. L. The combined use of analytical tools for exploring tetanus toxin and tetanus toxoid structures. *J. Chromatogr. B: Analyt. Technol. Biomed. Life Sci.* **2017**, *1054*, 80–92.