



Universiteit
Leiden
The Netherlands

Safe motherhood : severe maternal morbidity in the Netherlands. The LEMMoN study

Zwart, J.J.

Citation

Zwart, J. J. (2009, September 17). *Safe motherhood : severe maternal morbidity in the Netherlands. The LEMMoN study*. Retrieved from <https://hdl.handle.net/1887/14001>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/14001>

Note: To cite this publication please use the final published version (if applicable).

CHAPTER 3

Severe maternal morbidity during pregnancy, delivery and puerperium in the Netherlands: a nationwide population based study of 371 000 pregnancies

Zwart JJ, Richters JM, Öry F, de Vries JIP, Bloemenkamp KWM, van Roosmalen J.

BJOG 2008;115:842-50



Abstract

Objective: To assess incidence, case fatality rate, risk factors and substandard care in severe maternal morbidity in the Netherlands.

Design: Prospective population based cohort study.

Setting: All 98 Dutch maternity units in the Netherlands.

Methods: Cases of severe maternal morbidity were collected during a two-year period. All pregnant women in the Netherlands in the same period acted as reference cohort (n=371,021). As immigrant women are disproportionately represented in Dutch maternal mortality statistics, special attention was paid to the ethnic background. In a subset of 2.5% of cases substandard care was assessed through clinical audit.

Main outcome measures: Incidence, case fatality rates, possible risk factors, substandard care.

Results: Severe maternal morbidity was reported in 2552 cases, giving an overall incidence of 7.1 per 1000 deliveries. ICU admission was reported in 847 cases (incidence 2.4 per 1000), uterine rupture in 218 cases (incidence 6.1/10,000), eclampsia in 222 cases (incidence 6.2/10,000) and major obstetric haemorrhage in 1606 cases (incidence 4.5 per 1000). Non-Western immigrant women had a 1.3 fold increased risk of severe maternal morbidity (95% CI 1.2-1.5) when compared with Western women. Overall case fatality rate was 1 in 53. Substandard care was found in 39 of a subset of 63 women (62%) through clinical audit.

Conclusions: Severe maternal morbidity complicates at least 0.71% of all pregnancies in the Netherlands, immigrant women experiencing an increased risk. Since substandard care was found in the majority of assessed cases, reduction of severe maternal morbidity seems a mandatory challenge.

Introduction

Severe maternal morbidity gains interest as a new quality indicator of obstetric care.¹⁻⁴ The most important reason is the extremely low maternal mortality rate in Western countries, so that it takes years to collect the numbers needed to be able to draw valid conclusions from analysing cases of maternal mortality. Maternal deaths also tend to be more and more the result of rare complications, whereas regular life-threatening complications like major obstetric haemorrhage (MOH) are relatively underexposed as they less frequently lead to death nowadays.^{2;3} The most important and difficult issue, however, is the definition of severe maternal morbidity. Different research groups have already addressed this issue and the World Health Organisation is in the process of integrating these efforts into internationally accepted criteria for severe maternal morbidity.⁵⁻¹³ Recent studies demonstrate an increase in severe maternal morbidity in Western countries, possibly due to changes in management of obstetric complications and increasing age of pregnant women.^{2;14;15}

A nationwide cohort study of severe maternal morbidity, called LEMMoN, was conducted in the Netherlands to assess incidence, case fatality rates, risk factors and substandard care overall and for different subgroups. As ethnicity appeared to be a significant risk factor for pregnancy related death^{2;16;17} and seemed to be a risk factor for severe maternal morbidity, we are especially interested in the association of ethnicity with severe maternal morbidity.^{18;19}

Methods

Women were included from 1st August 2004 until 1st August 2006. All 98 hospitals (100%) with a maternity unit in the Netherlands participated in the survey: 10 tertiary care centres, 33 non-university teaching hospitals and 55 other general hospitals. The annual number of deliveries per unit in 2005 ranged from 93 to 2655 (average: 1162). Women with high risk pregnancies and those with low risk pregnancies who develop complications deliver in hospital under the guidance of obstetricians (secondary or tertiary care, 59% of all births). Women with low risk pregnancies without complications, deliver under the guidance of midwives and family physicians (primary care), either at home (30% of all births) or in hospital under their responsibility (11% of all births).²⁰

Final inclusion criteria were defined after searching the literature and after agreement with the national Maternal Mortality Committee of the Dutch Society of Obstetrics and Gynaecology. An expert panel of obstetricians advised about the design of the study. The main issues for setting our criteria were easy clinical applicability and univocality. Inclusion criteria are listed in Figure 1.

Figure 1. Inclusion criteria

<p>Group 1: ICU admission</p> <ul style="list-style-type: none"> • Admission to intensive care unit or coronary care unit, other than for standard postoperative recovery <p>Group 2: Uterine rupture</p> <ul style="list-style-type: none"> • Clinical symptoms (pain, fetal distress, acute loss of contractions, haemorrhage) that led to an emergency caesarean section, at which the presumed diagnosis of uterine rupture was confirmed • Peripartum hysterectomy or laparotomy for uterine rupture <p>Group 3: Eclampsia / HELLP syndrome</p> <ul style="list-style-type: none"> • Eclampsia • HELLP-syndrome only when accompanied by liver haematoma or rupture <p>Group 4: Major Obstetric Haemorrhage</p> <ul style="list-style-type: none"> • Transfusion need of ≥ 4 units of packed cells • Embolisation or hysterectomy for major obstetric haemorrhage <p>Group 5: Miscellaneous</p> <ul style="list-style-type: none"> • Other cases of severe maternal morbidity to the opinion of the treating obstetrician, not to be included in group 1-4
--

The last group was meant to include rare conditions of severe maternal morbidity (e.g. acute fatty liver of pregnancy), as well as severe manifestations of generally less severe conditions (e.g. severe early pre-eclampsia which did not require admission to ICU). Proven cases of pulmonary embolism could also be reported in this group. All cases during pregnancy, delivery and puerperium (limited to 6 weeks postpartum) were regarded as cases, including complications in early pregnancy. Cases were classified as 'early pregnancy' if one could not speak of a 'partus' and hence not of 'ante partum' or 'post partum'. This by definition applies to cases in which pregnancy ends before 17 weeks of gestational age in the Netherlands, and it also applies to cases of second trimester instrumental abortion.

Ethnicity was defined by country of origin ('geographical ethnic origin'). We used the definitions of Statistics Netherlands, based on country of birth of the woman. When the woman was born in the Netherlands with at least one of her parents born abroad, she was considered to be from the same origin as her parent(s) from abroad. Women from other Western European countries and from North America, Japan and Indonesia were considered Western immigrants according to Statistics Netherlands because of their socio-economic and cultural position in the Netherlands. All other immigrant women were considered non-Western.

Maternal deaths were reported to the national Maternal Mortality Committee of the Dutch Society of Obstetrics and Gynaecology by the attending obstetrician as usual. These cases were added to our database. Women who had more than one condition were considered only once in the overall incidence figures, only the first group was counted. For example, a woman with MOH (group 4) who was admitted to the ICU (group 1) was only counted as an ICU admission. However, these cases were counted for

each condition in the sub analysis of the different groups.

In each hospital, a local coordinator reported all cases monthly using a standardised web based form. Absence of cases in a particular month was also communicated in order to control for underreporting. Cases were identified in the respective hospitals using multiple strategies, including maternity computer databases, labour ward diaries, staff reports, intensive care admission registers, blood transfusion registers, discharge data and personal communication. At the central office, cases were collected using the national electronic surveillance system of the Netherlands Surveillance Centre for Obstetrics and Gynaecology (NSCOG), a newly established non-profit organisation for scientific data collection, analogous to the NSCK for paediatrics (TNO Quality of Life, Leiden, the Netherlands).²¹ Cases were initially notified by reporting initials and date of birth to LEMMoN, to minimise underreporting. Anonymised data were then obtained, consisting of a case record form with photocopies of relevant parts of the patient file. All cases were entered into an Access database by trained staff and each case was finally checked for correctness by the first author. We recorded maternal characteristics (age, Body Mass Index, parity, ethnicity, income, single household, language skills, smoking), all data on pregnancy and delivery, and data on the specific complication. A minimum of 87 items were entered into the database for each case, depending on the subgroup(s) of severe morbidity in which the case was included. We also recorded characteristics of each hospital (university or teaching hospital, annual number of deliveries). Socio-economic status was ascribed using the validated zip-code/socio-economic status indicator of Statistics Netherlands, based on home price and income, stratified into low, modest and high.²²

Although 30% of women in the Netherlands deliver at home, all women with severe maternal morbidity as defined in our inclusion criteria will eventually have been referred to one of the maternity units. Therefore, this study represents all deliveries in the Netherlands during the study period. As a consequence, nationwide statistics could be used as reference values whenever appropriate. To control for underreporting, we crosschecked our data with different other databases: underreporting of uterine rupture and eclampsia was controlled for using the national perinatal database (LVR-2).²³ Underreporting of MOH was controlled for using data from a large representative sample of local blood transfusion laboratories in the Netherlands during a 20-month period. Cases that were found to be not reported to our study were only counted and were not added to the database.

Seven audit meetings were held throughout the country to assess substandard care in a selection of cases, using the audit criteria developed by the Dutch Maternal Mortality Committee.²⁴ Assessors were members of the LEMMoN expert panel as well as local staff. After individual assessment by each assessor, a plenary meeting was held to discuss all items found. At this meeting, complete patient files were present to optimize assessability. Substandard care was assumed if the majority of assessors judged this to be present.

For each group, incidence was calculated using the total number of births in the Netherlands during

the study period as the denominator. Denominator data for the number of deliveries in the Netherlands were obtained from Statistics Netherlands (CBS).²⁵ They were based on birth registries during the exact study period, corrected for multiple births and stillbirths of 24 weeks or over. Denominator data for the number of women from the different ethnic groups were obtained from Statistics Netherlands. For the four large immigrant groups and for non-Western immigrants overall, numbers of mothers of newborns were available. For the smaller subgroups, we had to rely on numbers of women of fertile age (15-40 years old) to calculate the denominator, thereby disregarding the difference in fertility rate among the different ethnic groups. Relative risks were calculated when reference data were available. National reference values for possible risk factors for severe maternal morbidity were obtained from Statistics Netherlands and from the LVR-2 database. Incidence figures in LVR-2 were multiplied by 59/100 to also represent all deliveries under primary care (41% in 2002).²³ Case fatality rates were calculated by dividing the number of deaths due to a specific condition by the number of severe maternal morbidities due to that condition. Possible risk factors were identified by calculating relative risks and 95% confidence intervals. Significance was assumed when the confidence interval did not cross one. Statistical analysis was performed using the SPSS statistical package 14.0 (SPSS Inc., Chicago, IL, USA).

Results

During the study period, there were 371,021 deliveries in the Netherlands. All 98 hospitals with an obstetric ward in the Netherlands agreed to participate. A maximum of 2352 (98*24) 'hospital-months' could be reported. Mainly due to later enrolment of some hospitals into the study, a total of 2275 'hospital months' were actually returned (97%). Regarding only those maternities occurring during the months each hospital actively participated in the study, the study represents 358,874 deliveries. A total of 2552 cases were reported during the study period. We received detailed data of 2513 of 2552 cases (98.5%). The overall incidence of severe maternal morbidity in The Netherlands was 7.1 per 1000 deliveries. Cases were divided over the five groups as indicated in Table 1.

Table 1. Numbers, incidence and case fatality rate per inclusion group.

	ICU admission	Uterine rupture	Eclampsia/HELLP	MOH	Miscellaneous	Total
patients	847 (33.2%)	191 (7.5%)	135 (5.3%)	1146 (44.9%)	233 (9.1%)	2552 (100%)
complications*	847 (26.9%)	218 (6.9%)	239 (7.6%)	1606 (51.1%)	233 (7.4%)	3143 (100%)
data available	837 (98.8%)	218 (100%)	230 (96.2%)	1590 (99.0%)	228 (97.9%)	3102 (98.7%)
incidence (/1000 deliveries)	2.4	0.6	0.7	4.5	0.7	7.1
case fatality rate	1:29 (3.4%)	- (0%)	1:55 (1.8%)	1:201 (0.5%)	1:14 (7.3%)	1:53 (1.9%)

ICU=intensive care unit admission. MOH=Major obstetric haemorrhage. *one patient can have more than one complication.

Among the 2552 women included, 3143 complications were noted, 21.4% and 1.7% having two and three complications simultaneous, respectively. One woman had all four eligible complications. Two women were included twice during the study period in distinct pregnancies. Forty eight cases of pregnancy related death were reported to the Maternal Mortality Committee during the study period, giving an overall case fatality rate of 1 in 53 (1.9%). Incidence varied largely between hospitals in the Netherlands, ranging from 0 to 39.1 per 1000 deliveries. The mean hospital-incidence (regarding only the secondary and tertiary care deliveries in the respective hospitals) was 10.8 per 1000, 9.3 for non-university hospitals and 26.7 for university hospitals. In 2.8% of cases, the (first) complication occurred in early pregnancy, in 26.5% antepartum and in 70.7% postpartum. Characteristics of women included are shown in Table 2.

Table 2. Characteristics of women in the study.

	<i>n</i>	%
Age (mean 31.6)		
< 20 year	31	1.2%
20-35 year	1770	70.4%
35-40 year	590	23.5%
≥ 40 year	122	4.9%
Socio-economic status indicator		
low	701	31.6%
middle	994	44.9%
high	520	23.5%
unknown	298	
Smoking during pregnancy		
yes	175	12.0%
no	1290	88.0%
unknown	1048	
Body mass index (BMI)		
<18.5	48	2.8%
18.5-24.9	1018	60.2%
25.0-29.9 (overweight)	404	23.9%
30.0-34.9 (obese)	125	7.4%
≥ 35.0 (morbidly obese)	95	5.6%
unknown	823	
Geographical ethnic origin		
Netherlands	1864	74.4%
Morocco	116	4.6%
Turkey	87	3.5%
Surinam/Dutch Antilles	111	4.4%
sub-Saharan Africa	90	3.6%
other non-Western	146	5.8%
other Western	92	3.7%
unknown	7	

Possible risk factors for severe maternal morbidity are shown in Table 3. Overall, 21.1% of women were non-Western immigrants. The relative risk for non-Western women to experience severe maternal morbidity was 1.3 (95%-CI: 1.2-1.5) as compared to Western women. This elevated risk remained significant for each separate inclusion group (Table 4). Of the four largest immigrant groups in the Netherlands (Morocco, Turkey, Surinam and Netherlands Antilles), only Surinam women showed a significantly elevated risk as compared with native Dutch women (RR 1.4; 95% CI 1.1-1.7). Sub-Saharan African women had the highest risk (RR 3.5; 95% CI 2.8-4.3). Overall relative risks of women from the Middle East and South East Asia were 1.5 (95% CI 1.1-2.1) and 2.2 (95% CI 1.7-2.8).

Table 3. Risk factors for severe maternal morbidity.

<i>risk factor</i>	<i>LEMMoN</i>	<i>Netherlands</i>	<i>RR (95% CI)</i>
<i>patient</i>			
age \geq 35	29.3%	24.7% [†]	1.2 (1.1-1.3)
age \geq 40	4.8%	3.4% [†]	1.4 (1.2-1.7)
low income	31.6%	n/a	
single household	3.0%	n/a	
smoking during pregnancy	12.0%	n/a	
BMI \geq 25 (overweight)	36.9%	31.7% [†]	1.3 (1.1-1.4)
BMI \geq 30 (obese)	13.0%	9.1% [†]	1.5 (1.3-1.7)
non-Western immigrants	21.1%	16.8% [†]	1.3 (1.2-1.5)
chronic disease in general history	9.7%	n/a	
<i>pregnancy</i>			
initial antenatal care by obstetrician	35.8%	14.3% [‡]	3.3 (3.1-3.6)
prior caesarean section	19.3%	6.0% ²⁶	3.7 (3.4-4.1)
parity 0	49.9%	45.2% [†]	1.2 (1.1-1.3)
parity \geq 3	5.1%	5.0% [†]	1.0 (0.9-1.2)
parity \geq 6	0.4%	0.4% [‡]	1.2 (0.7-2.2)
multiple pregnancy	8.0%	1.7% [†]	4.9 (4.3-5.7)
artificial reproduction techniques: IVF/ICSI	4.7%	1.9% ²⁷	2.5 (2.1-3.0)
<i>delivery</i>			
home delivery	6.3%	31.6% [†]	0.1 (0.1-0.2)
induction of labour	26.5%	12.5% [‡]	3.1 (2.8-3.4)
caesarean section without labour	22.3%	5.9% [‡]	4.6 (4.2-5.0)
ventouse/forceps	12.7%	8.6% [‡]	1.6 (1.4-1.7)
caesarean section overall	43.6%	13.0% [‡]	5.2 (4.8-5.6)
breech presentation	7.9%	4.9% [‡]	1.7 (1.4-1.9)
preterm birth (<37w)	28.8%	5.8% [‡]	6.6 (6.0-7.2)
post term birth (\geq 42w)	5.3%	4.3% [‡]	1.3 (1.0-1.5)

n/a=data not available. *includes hypertension, diabetes, cardiac disease and coagulation disorders. National reference values from [†]Statistics Netherlands (exact study period) and [‡]The Netherlands Perinatal Registry (LVR-2, 2005).

ICU admission

A total of 847 cases of ICU admission were reported, giving an incidence of 2.4 per 1000 deliveries. Of all cases of pregnancy related death, 29 were admitted to ICU before death, giving a case fatality rate of 1 in 29. The mean duration of ICU stay was 3.6 days (range 1-74). The main reasons for admission were MOH (47%), hypertensive disorders of pregnancy (33%), respiratory complications (8%) and cardiac complications (7%). Assisted ventilation, inotropic support and renal dialysis were necessary in 34.2, 8.6 and 1.9% of cases, respectively.

Table 4. Relative risk of severe maternal morbidity for non-Western immigrants

Geographical Ethnic Origin	non-Western			RR (95% CI)
	immigrants	Western women	unknown	
ICU admission (n=837)	189	645	3	1.5 (1.2-1.7)
Uterine rupture (n=218)	48	170	0	1.4 (1.0-1.9)
Eclampsia/HELLP (n=230)	59	170	1	1.7 (1.3-2.3)
MOH (n=1590)	318	1268	4	1.3 (1.1-1.4)
Total (n=2513)	529	1977	7	1.3 (1.2-1.5)

ICU= Intensive Care Unit; MOH=Major Obstetric Haemorrhage.

Uterine rupture

A total of 218 cases of uterine rupture were reported, giving an incidence of 6.1 per 10,000 deliveries. No cases of pregnancy related death due to uterine rupture occurred. Admission to ICU occurred in 12% of cases and 21% experienced MOH. In 87% of cases, obstetric history revealed at least one caesarean section. Of the other 28 cases, 3 had a history of a known uterine scar due to myomectomy, tubectomy for isthmic pregnancy or dilatation and curettage. In 25 cases, there was no known uterine scar, 12 of whom did not have any possible known risk factor in general or obstetric history. In two cases, rupture occurred as a complication of second trimester dilatation and curettage for unwanted pregnancy. In six cases, uterine rupture complicated medically induced termination of pregnancy after 16 weeks of gestation.

Eclampsia and haemolysis, elevated liver enzymes, low platelets syndrome accompanied by liver haematoma or rupture

A total of 222 cases of eclampsia were reported, giving an incidence in the Netherlands of 6.2 per 10,000 deliveries. In addition, 19 cases of haemolysis, elevated liver enzymes, low platelets (HELLP) syndrome accompanied by liver haematoma (n=12) and/or liver rupture (n=7) were included. Two women had both eclampsia and HELLP syndrome accompanied by liver haematoma. There were four cases of pregnancy related death due to eclampsia/HELLP, case fatality rate being 1 in 55. Admission to ICU occurred in 42% of cases.

Table 5. Major Obstetric Haemorrhage, primary diagnoses

<i>timing</i>	<i>diagnosis[†]</i>	<i>n</i>	<i>(%)</i>
Early pregnancy (n=51)	Ectopic pregnancy	29	(56.9%)
	Spontaneous Abortion	10	(19.6%)
	Termination of pregnancy	10	(19.6%)
	Miscellaneous [‡]	2	(3.9%)
Antepartum (n=135)*	Placental abruption	61	(45.5%)
	Placenta praevia	54	(40.3%)
	Miscellaneous [§]	7	(5.2%)
	Unknown diagnosis	12	(9.0%)
Postpartum (n=1480)*	Retained placenta or placental rests	703	(47.8%)
	Uterine atony	567	(38.5%)
	Haemorrhage following CS	183	(12.4%)
	Perineal tears / episiotomy	148	(10.1%)
	Clotting disorders	116	(7.9%)
	Placenta accreta/increta/percreta	109	(7.4%)
	Rupture of cervix	58	(3.9%)
	Uterine rupture	44	(3.0%)
	Uterine inversion	13	(0.9%)
	Miscellaneous	65	(4.4%)
	Unknown diagnosis	10	(0.7%)

*In 76 cases both antepartum and postpartum diagnoses were coded; †Up to three diagnoses could be coded postpartum; ‡molar pregnancy and placenta percreta; §rupture of uterine/ovarian artery, rupture of ovarian cyst, placenta percreta, vasa praevia, retro placental haematoma, rupture of uterine vein.

Major obstetric Haemorrhage

A total of 1606 cases were reported, giving an incidence of MOH in our study of 4.5 per 1.000 deliveries. There were eight cases of pregnancy related death due to MOH, case fatality rate being 1 in 201. Admission to ICU occurred in 27% of cases. Three percent of cases occurred in early pregnancy, 9% antepartum and 88% postpartum. Primary diagnoses are shown in Table 5. In 107 and 114 cases (6.7 and 7.2%), respectively hysterectomy and arterial embolisation were performed. In 14 cases (13%) hysterectomy was necessary after arterial embolisation failed to stop haemorrhage. Vice versa, in two cases embolisation was performed after hysterectomy. Intrauterine balloon catheters were used in 154 (9.7%) cases, (re)laparotomy was performed in 202 (12.5%) with B-lynch suture in 12 (0.7%) and ligation of arterial vessels in 21 (1.3%). The average amount of estimated blood loss was 3150cc (maximum 20,000cc). An average of 6.6 units of blood were transfused (range 0-50). Fresh frozen plasma and pooled thrombocyte suspension was given in 48.3% and 16.2% of cases, respectively. Recombinant factor seven (Novo-seven[®]) was administered in 64 cases (4.0%). In 178 cases (11.2%), MOH was accompanied by pre-eclampsia. Five cases of Jehovah's witnesses with MOH were included on the basis of hysterectomy or arterial embolisation

without transfusion (n=4), or final acceptance of blood products (n=1). Six others were reported because of ICU admission or as 'other severe maternal morbidity'.

Table 6. Cases reported as 'Other severe maternal morbidity' (n=233)

	n	(%)	examples
<i>Obstetric</i>	100	(43%)	
pre-eclampsia/HELLP	70	(30%)	early severe HELLP necessitating termination of pregnancy or with retinal detachment
Genital tract sepsis	8	(3%)	Group A streptococcal sepsis
AFLP	5	(2%)	
MOH in Jehovah's witnesses	3	(1%)	
Miscellaneous obstetric	14	(6%)	maternal hydrops syndrome, abdominal bleeding after laser therapy for TTTS
<i>Non-obstetric</i>	128	(55%)	
Cardio respiratory	20	(9%)	myocardial infarction after CS, cardiomyopathy, pleural empyema
Cerebral/neurological	19	(8%)	viral meningo-encephalitis, sagittal sinus thrombosis, cerebrovascular accident
Thrombo-embolism	30	(13%)	pulmonary embolism, portal vein thrombosis, vena cava thrombosis
Sepsis	4	(2%)	urosepsis, sepsis from cholangitis
Surgical	21	(9%)	splenectomy after trauma, colectomy due to Crohn's disease, appendectomy
Oncological	12	(5%)	Kaposi sarcoma with AIDS, vulvar cancer, acute lymphatic leukaemia
Anaesthetic complication	2	(1%)	acute life-threatening danger during intubation due to extreme obesity
Miscellaneous non-obstetric	20	(9%)	acute pancreatitis, sickle cell crises, severe immune thrombocytopenic purpura
<i>Unknown</i>	5	(2%)	

HELLP=haemolysis, elevated liver enzymes, low platelets syndrome; AFLP= acute fatty liver of pregnancy; MOH= major obstetric haemorrhage; TTTS= twin-to-twin-transfusion syndrome

Other severe maternal morbidity

A total of 233 cases were reported as other severe maternal morbidity (Table 6). Divided over all inclusion groups, 69 cases of thrombo-embolism were included: pulmonary embolism (n=44), amniotic fluid embolism (n=9), and thrombosis of pelvic, ovarian, mesenteric, portal vein, sagittal sinus, mesenteric vein, portal vein and vena cava (n=16). In addition to the 239 cases of eclampsia/severe HELLP included in group 3, 360 women with pre-eclampsia were included in the other groups. Thus, 23.9% of all cases had pre-eclampsia. Sepsis was reported in 84 cases (estimated incidence 2.3 per 10,000).

Underreporting

During the first five months of the study, we found only one case of uterine rupture and two cases of eclampsia that were not reported to LEMMoN, underreporting being estimated at 2 and 3%, respectively. Underreporting of MOH to the LEMMoN study appeared to be 35% (range 0-83%) in a large representative sample. Cases not reported to our survey appeared to be mainly the

relatively less severe cases of MOH, with 68% of unreported cases being transfused only 4 units of blood. Taking into account this degree of underreporting, the overall incidence of MOH in the Netherlands would be 5.8 per 1000 deliveries. Controlling for underreporting of ICU admission on a national level appeared unfeasible.

Table 7. Substandard care items and their contribution*

	n	%†
<i>Patient</i>	55	6.6
Delay in consulting doctor	30	3.6
Refusal of medical help or advise	15	1.8
Language barrier	10	1.2
<i>GP/midwife</i>	164	19.9
Inadequate antenatal care	44	5.3
Delay in recognition of symptoms/signs	58	7.0
Delay in referral to obstetrician	62	7.5
<i>Obstetrician</i>	441	53.4
Inadequate antenatal care	70	8.5
Delay in recognition of symptoms/signs	146	17.7
Delay in treatment after diagnosis	200	24.2
Delay in referral to tertiary care centre	25	3.0
<i>Other consultant</i>	5	0.6
Delay in consulting obstetrician	5	0.6
<i>Health care system</i>	84	10.2
Home birth influenced outcome	19	2.3
Birth in general hospital influenced outcome	40	4.8
Quality of transport influenced outcome	25	3.0

*after individual assessment of 59/63 cases by on average 14 assessors; †each item could maximally be scored 826 times (59 cases times 14 assessors)

Audit

During seven audit sessions throughout the country, substandard care was judged to be present by the majority of assessors in 39 (62%) of 63 cases (Table 7).

Discussion

This study represents the first nationwide survey of severe maternal morbidity in the Netherlands and to date, by far the largest study in the literature. All aspects of severe maternal morbidity in the Netherlands have been mapped. The incidence in our study was 7.1 per 1000 deliveries, indicating that the average obstetric ward in the Netherlands encounters one case every month. We realize that incidence figures largely depend on the denominator chosen. We deliberately chose to express the incidence as the total pregnancy related risk of severe maternal morbidity per delivery. That means, we included complications of early pregnancy while we did only include deliveries from 24 weeks onward in the denominator. Therefore, the incidence is expressed as a ratio, not a rate. Leaving out all cases in early pregnancy and

before 24 weeks of gestation would have resulted in a slightly lower incidence of 6.8 per 1000.

There have been only two other large surveys addressing the overall incidence of severe maternal morbidity in Western countries.^{7,9} Their reported incidences were 12.0/1000 in South-West England (n=48,865) and 3.8/1000 in Scotland (n=51,165). Our incidence is well within the range of the other two studies. However, the incidence figures are strikingly different, bearing in mind the relatively comparable health care systems and populations. These differences can be mainly explained by the differences in inclusion criteria. The inclusion criteria for MOH, which affect the overall incidence to a great extent as it concerns about half of all inclusions, was more liberal in the English study (blood loss of more than 1500cc) as compared to the Scottish (transfusion of five or more units of blood). Our definition of MOH was in between those two (transfusion of four or more units of blood), and so was our overall incidence. The large differences in reported incidences due to different inclusion criteria clearly reflect the need for internationally accepted definitions of severe maternal morbidity.

The high incidence of eclampsia in the Netherlands is worrying and should be subject of further research, especially seen in the light of the also high maternal mortality due to hypertensive disorders of pregnancy in the Netherlands.³

Case fatality rates for the different types of severe maternal morbidity ranged from 0% for uterine rupture to 3.4% for ICU admission, indicating the severity of cases included. Despite the difference in inclusion criteria, our overall case fatality rate is comparable to that in Scotland.⁹

Several risk factors were identified in this study. Due to the nationwide design of the study, we were able to reliably calculate relative risks based on available national reference data. However, for some of them it is important to realise that the condition could be the cause of severe maternal morbidity, but it could also represent the result of it. This is especially true for caesarean section and induction of labour, which were often performed because of the compromised maternal condition. Preterm birth is also closely related. With respect to artificial reproductive techniques, the trend towards single embryo transfer might well lead to reduction of severe maternal morbidity because women with multiple pregnancies were at higher risk.²⁸ We found Body Mass Index to be an important risk factor for severe maternal morbidity. Since the incidence of overweight and obesity is increasing rapidly in Western countries, we expect severe maternal morbidity to increase in the future. Home delivery appeared to be a strong protective factor for severe maternal morbidity in the Netherlands with a relative risk of 0.1 (95% CI 0.1-0.2). This again demonstrates the proper functioning of the Dutch risk selection system, in which low-risk pregnancies are cared for by midwives in private practices or family physicians outside the scope of the obstetrician.

As was already expected from maternal mortality statistics, non-Western ethnic background appeared to be an independent risk factor for experiencing severe maternal morbidity. Three of the four large immigrant groups in the Netherlands -Moroccan, Turkish, and Dutch Caribbean- however, were not

over-represented in our study. Surinam women were slightly over-represented, especially due to a high incidence of eclampsia. Members of the smaller ethnic minority groups were disproportionately more often represented in our study. Preliminary findings of the qualitative study which complemented the present registration study reveal several issues that may play an important role in their increased risk. Namely, women's relative short stay in the Netherlands, their lack of a social support network, their lack of knowledge of our health care system, communication problems with care providers, and health illiteracy. More qualitative in-depth research into the non-medical backgrounds of these women and the course of events preceding their complication is currently carried out and will shed light on patient-related backgrounds of the increased risk.

The main limitation of this study is that we did not record the individual characteristics of all maternities without severe maternal morbidity during the study period. Therefore, we could not adjust relative risks for confounding variables. Furthermore, despite our efforts, we cannot guarantee the completeness of data. Therefore, our reported incidence figures only represent a minimum level of severe maternal morbidity. Due to the nationwide nature of the study, we depended on the active participation of local coordinating obstetricians for completeness of data. We tried to meet this by keeping coordinators actively involved and providing help in collecting the data. Local coordinators not responding to our monthly request for cases were reminded repeatedly by e-mail and phone. And finally, we thoroughly controlled for underreporting. Underreporting of eclampsia and uterine rupture appeared to be very low during the first five months of the study, and we therefore decided not to control for it in the remainder of the study period. Underreporting of MOH was estimated at 35%, but appeared to be mainly due to relatively less severe complications requiring "only" four units of blood. We expect underreporting to be relatively low for the most severe conditions like acute fatty liver of pregnancy or amniotic fluid embolism, as these are very impressive clinical events that draw the attention of many clinicians involved and will not likely be missed.

The institution of Obstetric High Care (OHC) facilities in university hospitals posed a methodological challenge. Women, who would without any doubt have been treated at an ICU in a non-university hospital, were not admitted to ICU in university hospitals. Inclusion of all admissions to OHC facilities was not merely possible, as most of these admissions were for fetal rather than maternal indication. Coordinators in university hospitals were instructed to report OHC admissions for strict maternal indications as 'other severe maternal morbidity'. More than half of all inclusions in this group came from university hospitals.

The incidence of severe maternal morbidity in the different hospitals varied largely. Although this could be caused by differences in local case management, other possible explanations are bias due to the management based criteria, the degree of underreporting per hospital, and chance. Hospitals, of which we knew that the local coordinator was dedicated and the local system of recognition and reporting of

cases was supported by all clinicians, appeared to have higher incidence figures. Furthermore, incidence was by definition influenced by the management-based criteria 'admission to ICU' and 'MOH requiring four or more units of blood'. We noticed substantial variations in local policy for admission to ICU and transfusion of blood products.

The next step in the process of improving maternal care is to critically assess the course of events that led to the severe condition, as substandard care analysis serves as a basis to improve guidelines and clinical protocols. Substandard care was judged to be present in the majority of assessed cases, mainly at the level of the care providers, indicating that further analysis of cases and improvement of guidelines could reduce severe maternal morbidity.²⁴ Several local initiatives for auditing of severe maternal morbidity came to our knowledge since this study.

In order to confirm the apparent increase in severe maternal morbidity in Western countries, and to evaluate our own clinical practice in the Netherlands, we would need to establish an ongoing registration of severe maternal morbidity in the Netherlands. This should be incorporated in the Dutch Perinatal Database and be very comprehensive to assure cooperation of each maternity unit. Furthermore, the implementation of national registration of rare obstetric conditions, like UKOSS in the United Kingdom³⁰, would be valuable and the infrastructure is already in place in the Netherlands.²¹

This survey appears to be a valuable addition to the maternal mortality statistics which have already been registered for decades in the Netherlands. It gives a clear new insight in the problems encountered in obstetric practice nowadays and can serve as a reference work for severe maternal morbidity in the Netherlands and other Western countries. Improvement of the quality of obstetric care through auditing of cases of severe maternal morbidity seems a mandatory challenge.

Conclusion

Severe maternal morbidity complicates at least 0.71% of all pregnancies in the Netherlands, immigrant women experiencing an increased risk. Severe maternal morbidity should be considered internationally as a new indicator of the quality of obstetric care next to maternal mortality statistics. Since substandard care was found in the majority of assessed cases, reduction of severe maternal morbidity seems a mandatory challenge. Therefore, auditing of severe maternal morbidity at local or regional level should be encouraged in order to improve the quality of obstetric care and decrease the incidence of severe maternal morbidity and maternal mortality. Audit should be incorporated in our national public health policy as an instrument of quality control.

References

- 1 Drife JO. Maternal "near miss" reports? *BMJ* 1993;307:1087-8.
- 2 G. Lewis (ed) 2007. *The Confidential Enquiry into Maternal and Child Health (CEMACH). Saving mother's lives: reviewing maternal deaths to make motherhood safer - 2003-2005. The Seventh Report on Confidential Enquiries into Maternity Deaths in the United Kingdom.* London: CEMACH
- 3 Schutte JM, de Boer K, Briët JW, et al. [Maternal Mortality in the Netherlands: the tip of the iceberg]. *Ned Tijdschr Obstet Gynaecol* 2005;118:89-91.
- 4 Hill K, Thomas K, AbouZahr C, et al. Estimates of maternal mortality worldwide between 1990 and 2005: an assessment of available data. *Lancet* 2007;370:1311-9.
- 5 Stones W, Lim W, Al Azzawi F, Kelly M. An investigation of maternal morbidity with identification of life-threatening 'near miss' episodes. *Health Trends* 1991;23:13-5.
- 6 Mantel GD, Buchmann E, Rees H, Pattinson RC. Severe acute maternal morbidity: a pilot study of a definition for a near-miss. *Br J Obstet Gynaecol* 1998;105:985-90.
- 7 Waterstone M, Bewley S, Wolfe C. Incidence and predictors of severe obstetric morbidity: case-control study. *BMJ* 2001;322:1089-93.
- 8 Say L, Pattinson RC, Gulmezoglu AM. WHO systematic review of maternal morbidity and mortality: the prevalence of severe acute maternal morbidity (near miss). *Reprod Health* 2004;1:3-7.
- 9 Brace V, Penney G, Hall M. Quantifying severe maternal morbidity: a Scottish population study. *BJOG* 2004;111:481-4.
- 10 Geller SE, Rosenberg D, Cox S, Brown M, Simonson L, Kilpatrick S. A scoring system identified near-miss maternal morbidity during pregnancy. *J Clin Epidemiol* 2004;57:716-20.
- 11 Zhang WH, Alexander S, Bouvier-Colle MH, Macfarlane A. Incidence of severe pre-eclampsia, postpartum haemorrhage and sepsis as a surrogate marker for severe maternal morbidity in a European population-based study: the MOMS-B survey. *BJOG* 2005;112:89-96.
- 12 Penney G, Brace V. Near miss audit in obstetrics. *Curr Opin Obstet Gynecol* 2007;19:145-50.
- 13 Baskett TF, O'Connell CM. Severe obstetric maternal morbidity: a 15-year population-based study. *J Obstet Gynec* 2005;25:7-9.
- 14 Wen SW, Huang L, Liston R, et al. Severe maternal morbidity in Canada, 1991-2001. *CMAJ* 2005;173:759-64.
- 15 Joseph KS, Rouleau J, Kramer MS, Young DC, Liston RM, Baskett TF. Investigation of an increase in postpartum haemorrhage in Canada. *BJOG* 2007;114:751-9.
- 16 van Roosmalen J, Schuitemaker NW, Brand R, van Dongen PW, Bennebroek GJ. Substandard care in immigrant versus indigenous maternal deaths in The Netherlands. *BJOG* 2002;109:212-3.
- 17 Schuitemaker N, van Roosmalen J, Dekker G, van Dongen P, van Geijn H, Bennebroek GJ. Confidential enquiry into maternal deaths in The Netherlands 1983-1992. *Eur J Obstet Gynecol Reprod Biol* 1998;79:57-62.
- 18 Bouvier-Colle MH, Varnoux N, Salanave B, Ancel PY, Breart G. Case-control study of risk factors for obstetric patients' admission to intensive care units. *Eur J Obstet Gynecol Reprod Biol* 1997;74:173-7.
- 19 Keizer JL, Zwart JJ, Meerman RH, Harinck BI, Feuth HD, van Roosmalen J. Obstetric intensive care admissions: a 12-year review in a tertiary care centre. *Eur J Obstet Gynecol Reprod Biol* 2006;128:152-6.
- 20 The Netherlands Perinatal Registry. [Report Home Delivery 2001-2002]. [http://www.perinatreg.nl/uploads/153/99/Rapportage_Thuisbevalling_2001-2002.pdf]. Accessed 2 October 2007.
- 21 Hira Sing RA, Rodrigues PR. [The Dutch Pediatric Surveillance System; a quality focused instrument for prevention and research]. *Ned Tijdschr Geneesk* 2002;146:2409-14.
- 22 Duin C van, Keij I. Sociaal-economische status indicator op postcodeniveau. *Statistics Netherlands. Maandstatistiek bevolking* 2002;2:32-5.
- 23 The Netherlands Perinatal Registry. *Perinatal Care in the Netherlands 2005.* Bilthoven, the Netherlands: the Netherlands Perinatal Registry, 2007.

- 24 van Dillen J, Zwart JJ, Lim F, Vredevoogd C, van Roosmalen J. [LEMMoN Audit; a pilot study into incidence and assessment of severe maternal morbidity in The Hague]. *Ned Tijdschr Obstet Gynaecol* 2006;119:29-34.
- 25 Statistics Netherlands (CBS). Statline, Central Bureau of Statistics, 2007. [<http://www.cbs.nl/en-GB/>]. Accessed 2 October 2007.
- 26 Kwee A, Bots ML, Visser GH, Bruinse HW. Obstetric management and outcome of pregnancy in women with a history of caesarean section in the Netherlands. *Eur J Obstet Gynecol Reprod Biol* 2007;132:171-6.
- 27 Dutch Society of Obstetricians and Gynaecologists (NVOG). Nationwide IVF results 2005 [<http://www.nvog.nl/files/landelijkeivfcijfers2005.pdf>]. Accessed 23 October 2007.
- 28 Kjellberg AT, Carlsson P, Bergh C. Randomized single versus double embryo transfer: obstetric and paediatric outcome and a cost-effectiveness analysis. *Hum Reprod* 2006;21:210-6.
- 29 Amelink-Verburg M, Verloove-Vanhorick S, Hakkenberg R, Veldhuijzen I, Bennebroek Gravenhorst J, Buitendijk S. Evaluation of 280 000 cases in Dutch midwifery practices: a descriptive study. *BJOG* 2008;115:570-8.
- 30 Knight M, Kurinczuk JJ, Tuffnell D, Brocklehurst P. The UK Obstetric Surveillance System for rare disorders of pregnancy. *BJOG* 2005;112:263-5.

