



Universiteit
Leiden
The Netherlands

Typical representations for $GL_n(F)$

Nadimpalli, S.

Citation

Nadimpalli, S. (2015, June 16). *Typical representations for $GL_n(F)$* . Retrieved from <https://hdl.handle.net/1887/33218>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/33218>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/33218> holds various files of this Leiden University dissertation

Author: Nadimpalli, Santosh

Title: Typical representations for $\mathrm{GL}_n(F)$

Issue Date: 2015-06-16

Acknowledgements

I express my deepest gratitude to my thesis supervisor Guy Henniart for his guidance during my masters and doctorate years. I learnt a great deal of mathematics, about research and about writing mathematics. I learned the theory of smooth representations of p -adic reductive groups essentially from him. I thank him for the encouragement and support through difficult times in graduate school.

I spent almost a year at Leiden university and had numerous discussions on various aspects of modular forms, Galois representations and geometry among other things with Bas Edixhoven. I am deeply influenced by those discussions we had every week. I would like to thank my thesis co-supervisor Bas Edixhoven for these stimulating discussions and for his encouragement. I also thank him for helping me with the French translation.

I would like to thank Shaun Stevens for his interest in my work. It is indeed his comments on my work on GL_3 encouraged me to think about the certain difficult cases of GL_4 . I would like to thank Corinne Blondel for her detailed comments on the thesis and some simplifications she suggested. I thank David Helm for his report on the thesis.

I would like to thank Jyoti Prakash Saha, Ramon M. Nunes, Shaunik Deo, Vineeth Ch and Sourav Ghosh for the mathematical discussions we had. I am very grateful for the libraries of Orsay and Indian Statistical Institute, Bangalore. I thank all my teachers in Indian statistical Institute, Bangalore, University of Padova and Université Paris sud. Last but not the least I thank my family amma, nanna, satish and tanu for keeping up with me.

Abstract

In this thesis we classify typical representations for certain non-cuspidal Bernstein components of GL_n over a non-Archimedean local field. Following the work of Henniart in the case of GL_2 and Paskunas for the cuspidal Bernstein components, we classify typical representations for Bernstein components of level-zero for GL_n for $n \geq 3$, principal series components, components with Levi subgroup of the form $(n, 1)$ for $n > 1$ and certain components with Levi subgroup of the form $(2, 2)$.

Each of the above component is treated in a separate chapter. The classification uses the theory of types developed by Bushnell and Kutzko in a significant way. We will give the classification in terms of Bushnell-Kutzko types for a given inertial class.

Samenvatting

In deze scriptie classificeren wij typische representaties voor bepaalde niet-cuspidale Bernstein componenten van GL_n over een niet Archimedisch lokaal lichaam. Gebruik makend van het werk van Henniart in het geval van GL_2 voor de cuspidale Bernstein componenten, classificeren wij representaties voor Bernstein componenten van niveau nul voor GL_n voor $n \geq 3$, voor principal series components, voor componenten met Levi ondergroep van de vorm $(n, 1)$ met $n > 1$ en sommige componenten met Levi ondergroep van de vorm $(2, 2)$.

Alle bovenstaande componenten worden in hun eigen hoofdstuk behandeld. De classificatie berust zwaar op de theorie van typen ontwikkeld door Bushnell en Kutzko en deze zal ook gegeven worden in de termen van de Bushnell-Kutzko typen behorend bij een gegeven inertie klasse.

Résumé

Dans cette thèse, nous classifions les représentations typiques pour certaines composantes de Bernstein de GL_n sur un corps localement compact non Archimédien. Suite aux travaux de Henniart dans le cas de GL_2 et de Paskunas pour les composantes de Bernstein cuspidales, nous classifions les représentations typiques pour les composants de Bernstein de niveau zéro pour $n \geq 3$, les composantes de séries principales, les composantes dont le sous-groupe de Levi est de forme $(n, 1)$ pour $n > 1$ et certaines composantes dont le sous-groupe de Levi est de la forme $(2, 2)$.

Chacune des composantes ci-dessus est traitée dans un chapitre distinct. La classification utilise d'une manière significative la théorie des types développée par Bushnell-Kutzko, et elle est établie en termes de tels types.

Curriculum Vitae

Santosh Nadimpalli was born on 4th-January 1990 in Vijayawada, India. He attended Visakha valley school in Visakhapatnam until his 10th class. He completed his intermediate education in Hyderabad aiming to be an engineer. He got admission in Indian Statistical Institute to pursue bachelors of mathematics in 2007.

In 2010 he was offered Algant Masters scholarship to study in University of Padova and Université Paris Sud. He spent the first year of masters in Padova (2010 – 2011) and the second year (2011 – 2012) in Orsay. As a second year masters student he approached Prof. Guy Henniart for Masters thesis. He wrote his masters thesis with Henniart and completed his masters in 2012.

With Algant Doctorate scholarship, in 2012 he began his Phd jointly hosted by University of Leiden and Université de Paris Sud. Following the suggestion of Henniart he is working on the topic of typical representations for the Phd.

Bibliography

- [BH06] Colin J. Bushnell and Guy Henniart, *The local Langlands conjecture for $\mathrm{GL}(2)$* , Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 335, Springer-Verlag, Berlin, 2006. MR 2234120 (2007m:22013)
- [BH13] ———, *Intertwining of simple characters in $\mathrm{GL}(n)$* , Int. Math. Res. Not. IMRN (2013), no. 17, 3977–3987. MR 3096916
- [BK93] Colin J. Bushnell and Philip C. Kutzko, *The admissible dual of $\mathrm{GL}(N)$ via compact open subgroups*, Annals of Mathematics Studies, vol. 129, Princeton University Press, Princeton, NJ, 1993. MR 1204652 (94h:22007)
- [BK98] ———, *Smooth representations of reductive p -adic groups: structure theory via types*, Proc. London Math. Soc. (3) **77** (1998), no. 3, 582–634. MR 1643417 (2000c:22014)
- [BK99] ———, *Semisimple types in GL_n* , Compositio Math. **119** (1999), no. 1, 53–97. MR 1711578 (2000i:20072)
- [Blo06] Corinne Blondel, *Propagation de paires couvrantes dans les groupes symplectiques*, Represent. Theory **10** (2006), 399–434 (electronic). MR 2266698 (2007m:22012)
- [BM02] Christophe Breuil and Ariane Mézard, *Multiplicités modulaires et représentations de $\mathrm{GL}_2(\mathbf{Z}_p)$ et de $\mathrm{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$ en $l = p$* , Duke Math. J. **115** (2002), no. 2, 205–310, With an appendix by Guy Henniart. MR 1944572 (2004i:11052)
- [BZ76] I. N. Bernštejn and A. V. Zelevinskii, *Representations of the group $\mathrm{GL}(n, F)$, where F is a local non-Archimedean field*, Uspehi Mat. Nauk **31** (1976), no. 3(189), 5–70. MR 0425030 (54 #12988)
- [BZ77] I. N. Bernstein and A. V. Zelevinsky, *Induced representations of reductive p -adic groups. I*, Ann. Sci. École Norm. Sup. (4) **10** (1977), no. 4, 441–472. MR 0579172 (58 #28310)
- [Cas73] William Casselman, *The restriction of a representation of $\mathrm{GL}_2(k)$ to $\mathrm{GL}_2(\mathfrak{o})$* , Math. Ann. **206** (1973), 311–318. MR 0338274 (49 #3040)
- [EG14] Matthew Emerton and Toby Gee, *A geometric perspective on the Breuil-Mézard conjecture*, J. Inst. Math. Jussieu **13** (2014), no. 1, 183–223. MR 3134019
- [Hen00] Guy Henniart, *Une preuve simple des conjectures de Langlands pour $\mathrm{GL}(n)$ sur un corps p -adique*, Invent. Math. **139** (2000), no. 2, 439–455. MR 1738446 (2001e:11052)

- [How73] Roger E. Howe, *On the principal series of GL_n over p -adic fields*, Trans. Amer. Math. Soc. **177** (1973), 275–286. MR 0327982 (48 \#6324)
- [HT01] Michael Harris and Richard Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich. MR 1876802 (2002m:11050)
- [LRS93] G. Laumon, M. Rapoport, and U. Stuhler, *D -elliptic sheaves and the langlands correspondence*, Inventiones mathematicae **113** (1993), no. 1, 217–338 (English).
- [Pas05] Vytautas Paskunas, *Unicity of types for supercuspidal representations of GL_N* , Proc. London Math. Soc. (3) **91** (2005), no. 3, 623–654. MR 2180458 (2007b:22018)
- [Ren10] David Renard, *Représentations des groupes réductifs p -adiques*, Cours Spécialisés [Specialized Courses], vol. 17, Société Mathématique de France, Paris, 2010. MR 2567785 (2011d:22019)
- [Roc98] Alan Roche, *Types and Hecke algebras for principal series representations of split reductive p -adic groups*, Ann. Sci. École Norm. Sup. (4) **31** (1998), no. 3, 361–413. MR 1621409 (99d:22028)
- [SZ99] P. Schneider and E.-W. Zink, *K -types for the tempered components of a p -adic general linear group*, J. Reine Angew. Math. **517** (1999), 161–208, With an appendix by Schneider and U. Stuhler. MR 1728541 (2001f:22029)
- [Zel81] Andrey V. Zelevinsky, *Representations of finite classical groups*, Lecture Notes in Mathematics, vol. 869, Springer-Verlag, Berlin-New York, 1981, A Hopf algebra approach. MR 643482 (83k:20017)