



Universiteit
Leiden
The Netherlands

Molecular pathology of mismatch repair deficient tumours with emphasis on immune escape mechanisms

Dierssen, J.W.F.

Citation

Dierssen, J. W. F. (2010, November 17). *Molecular pathology of mismatch repair deficient tumours with emphasis on immune escape mechanisms*. Retrieved from <https://hdl.handle.net/1887/16151>

Version: Corrected Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/16151>

Note: To cite this publication please use the final published version (if applicable).

HNPCC versus sporadic microsatellite-unstable colon cancers follow different routes toward loss of HLA class I expression

Jan Willem F Dierssen, Noel FCC de Miranda, Soldano Ferrone,
Marjo van Puijenbroek, Cees J Cornelisse, Gert Jan Fleuren, Tom
van Wezel and Hans Morreau

BMC Cancer 2007, 7:33

This study was supported by the Dutch Cancer Society (grant number 2000/2135).

1. ABSTRACT

1.1. Background

Abnormalities in Human Leukocyte Antigen (HLA) class I expression are common in colorectal cancer. Since HLA expression is required to activate tumor antigen-specific cytotoxic T-lymphocytes (CTL), HLA class I abnormalities represent a mechanism by which tumors circumvent immune surveillance. Tumors with high microsatellite instability (MSI-H) are believed to face strong selective pressure to evade CTL activity since they produce large amounts of immunogenic peptides. Previous studies identified the prevalence of HLA class I alterations in MSI-H tumors. However, those reports did not compare the frequency of alterations between hereditary and sporadic MSI-H tumors neither the mechanisms that led to HLA class I alterations in each subgroup.

1.2. Methods

To characterize the HLA class I expression among sporadic MSI-H and microsatellite-stable (MSS) tumors, and HNPCC tumors we compared immunohistochemically the expression of HLA class I, β 2-microglobulin (β 2m), and Antigen Processing Machinery (APM) components in 81 right-sided sporadic and 75 HNPCC tumors. Moreover, we investigated the genetic basis for these changes.

1.3. Results

HLA class I loss was seen more frequently in MSI-H tumors than in MSS tumors ($p < 0.0001$). Distinct mechanisms were responsible for HLA class I loss in HNPCC and sporadic MSI-H tumors. Loss of HLA class I expression was associated with β 2m loss in HNPCC tumors, but was correlated with APM component defects in sporadic MSI-H tumors ($p < 0.0001$). In about half of the cases, loss of expression of HLA class I was concordant with the detection of one or more mutations in the β 2m and APM components genes.

1.4. Conclusion

HLA class I aberrations are found at varying frequencies in different colorectal tumor types and are caused by distinct genetic mechanisms. Chiefly, sporadic and hereditary MSI-H tumors follow different routes toward HLA class I loss of expression supporting the idea that these tumors follow different evolutionary pathways in tumorigenesis. The resulting variation in immune escape mechanisms may have repercussions in tumor progression and behavior.

2. BACKGROUND

During cancer development, tumor cells may elicit cytotoxic T-lymphocyte (CTL)-mediated immune responses—partly a consequence of accumulated gene mutations that are translated into altered peptides [1]. Tumor cell expression of HLA class I-antigen complexes is essential for

CTL recognition of aberrant peptides and subsequent activation [2]. Consequently, alteration of HLA class I cell surface expression provides an effective mechanism by which tumors can escape immune detection [3,4]. Multiple mechanisms have been shown to underlie defects in HLA class I expression by tumor cells. They include mutations in the individual HLA class I

genes *HLA-A*, *-B* and *-C*, located on chromosome 6p21.3) [5]; mutations in $\beta 2$ -microglobulin ($\beta 2m$) [6-9], molecule required for cell surface expression of HLA class I antigens; and defects in components of the HLA class I-associated antigen-processing machinery (APM) [9-11]. The APM consists of proteasome components delta, MB1 and Z; the immunoproteasome components LMP2, LMP7 and LMP10; peptide transporters TAP1 and TAP2; and chaperones Calnexin, Calreticulin, ERp57, and Tapasin. The immunoproteasome generates peptides mostly, although not exclusively from endogenous proteins, TAP1 and TAP2 facilitate peptide translocation from the cytosol into the lumen of the endoplasmic reticulum, where the peptides are loaded onto the HLA class I molecules with the aid from the several chaperones [12].

Chromosomal instability (CIN) and microsatellite instability (MIN) are the two major forms of genetic instability in colorectal cancer. Combined with distinct somatic mutation patterns and epigenetic modifications, CIN and MIN lead to the development of sporadic colorectal cancer [13]. MIN sporadic tumors, which constitute approximately 15% of all colorectal cancer cases and up to 40% of the tumors localized on the right side (preceding the splenic flexure) of the colon [14], have a phenotype resulting from the epigenetic inactivation of the mismatch repair gene *hMLH1*. Its inactivation destroys a cell's ability to repair base-base mismatches and small insertions or deletions in repetitive stretches, leading to an accumulation of frameshift mutations that get translated into abnormal peptide sequences. When these mutations are accumulated to large extent in the cell genome the tumors are said to possess high-microsatellite instability (MSI-H) [15]. Hence, it is expected that genes containing microsatellite sequences within their coding regions are more susceptible to somatic mutations, as seen in the *TGF β -RII*

gene. *TGF β -RII*'s third exon contains a microsatellite repeat of 10 adenines that is frequently targeted by frameshift mutations in MSI-H tumors [16]. MSI-H is also the hallmark of hereditary non-polyposis colorectal cancer (HNPCC), in which germline mutations of *hMLH1*, *hMSH2*, *hMSH6* and *PMS2* can be found. HNPCC constitutes approximately 2–4% of all CRC cases [17]. Tumors with MSI-H are thought to be more able to stimulate a CTL-mediated immune response due to their frequent generation of the aberrant frameshift peptides [18]. Therefore, these tumors are subjected to a greater selective pressure which favors the outgrowth of tumor cells with the ability to escape from recognition and destruction by host immune system.

Various studies have identified HLA alterations in colorectal cancer [19-21], including the prevalence of HLA class I alterations in MSI-H tumors [8,22]. However, the latter studies did not compare the frequency of alterations between hereditary and sporadic MSI-H tumors neither the mechanisms that led to HLA class I alterations in each subgroup. It was suggested that MSI-H sporadic and hereditary tumors follow parallel evolutionary pathways during tumorigenesis in terms of both genotype and phenotype [23]. As far as HLA class I defects are concerned it was never investigated whether these different tumors present distinct escape mechanisms from the immune system. In the present study, we compared the frequency of defects in HLA class I expression in right-sided sporadic (MSI-H and microsatellite-stable (MSS) sub-groups) colon tumors and in HNPCC tumors and studied the mechanisms underlying any abnormalities in these subgroups.

3. METHODS

3.1. Patient material and tissue microarrays

Two tissue microarrays were constructed from formalin-fixed, paraffin-embedded tissues as described previously [24]. One array, previously described [25], included colorectal tumor specimens from 129 suspected HNPCC patients with MSI-H colon tumors of which 75 cases were analyzed in the present study after confirmation of their HNPCC status: 73.3% ($n = 55$) of the latter possessed a germline pathogenic mutation in *hMLH1* ($n = 24$), *hMSH2* ($n = 18$), *hMSH6* ($n = 12$) or *PMS2* ($n = 1$), the remaining were MSI-H, without methylation of the *hMLH1* promoter and/or with immunohistochemical loss of the MSH2/MSH6 heterodimer and/or possessed a very young age at diagnosis of colon cancer (<50 yrs old). All cases possessed a positive family history for MSI-H tumors. The second tissue array included 3 tumor tissue cores from 81 sporadic right-sided colon cancer cases resected between 1990 and 2005 at the Leiden University Medical Center (Leiden, The Netherlands) and at the Rijnland Hospital (Leiderdorp, The Netherlands). The 81 patients in the latter array consisted of 47 females and 34 males with a mean age of 71.15 years ($SD = 9.958$). Approximately 60% ($n = 48$) of these cases were classified as MSS while the remaining ($n = 33$) possessed a MSI-H phenotype. The microsatellite instability status of the tumors was determined according to recommendations of the National Cancer Institute/ICG-HNPCC [15]. Moreover all MSI-H sporadic cases have lost the expression of the MLH1/PMS2 heterodimer as assessed by immunohistochemistry. The sporadic status of the MSI-H right-sided tumors (RST) was confirmed by methylation analysis of the *hMLH1* promoter using a methylation-specific MLPA assay as previously described [26]. All MSI-H sporadic cases

presented with hypermethylation at the *hMLH1* promoter.

The present study falls under approval by the Medical Ethical Committee of the LUMC (protocol P01-019). Cases were analyzed following the medical ethnical guidelines described in the Code Proper Secondary Use of Human Tissue established by the Dutch Federation of Medical Sciences [27].

3.2. Immunohistochemistry

Standard three-step, indirect immunohistochemistry was performed on 4- μ m tissue sections transferred to glass slides using a tape-transfer system (Instrumedics, Hackensack, NJ), including citrate antigen retrieval, blockage of endogenous peroxidase and endogenous avidin-binding activity, and di-aminobenzidine development.

The following primary antibodies were used: the mAb HCA2 which recognizes β 2m-free HLA-A (except -A24), -B7301 and -G heavy chains [28,29]; the mAb HC10, which recognizes a determinant expressed on all β 2m-free HLA-B and C heavy chains and on β 2m-free HLA-A10, -A28, -A29, -A30, -A31, -A32 and -A33 heavy chains (supernatant kindly provided by Dr. J. Neefjes, NKI, Amsterdam, The Netherlands and Dr. H. L. Ploegh, MIT, Boston, MA) [28,30]; TAP1 specific mAb NOB1; LMP2-specific mAb SY-1; LMP7-specific mAb HB2; LPM10-specific mAb TO-7; Calnexin-specific mAb TO-5; Calreticulin-specific mAb TO-11; Tapasin-specific mAb TO-3; ERp57-specific mAb TO-2 [31-33]; TAP2-specific mAb (BD Biosciences Pharmingen, San Diego, CA); rabbit anti- β 2m polyclonal Ab (A 072; DAKO Cytomation, Glostrup, Denmark); anti-MLH1 (clone G168-728; BD Biosciences) and anti-PMS2 (clone A16-4; BD Biosciences). Secondary reagents used were biotinylated rabbit anti-mouse IgG antibodies (DAKO Cytomation), goat anti-rabbit IgG antibodies (DAKO Cytomation),

and biotinylated-peroxidase streptavidin complex (SABC; DAKO Cytomation).

Loss of expression was defined by complete lack of staining in membrane and cytoplasm (HCA2, HC10, and anti- β 2m), in the nucleus (anti-MLH1 and anti-PMS2), in the peri-nucleus/endoplasmic reticulum (NOB1, anti-TAP2, TO-2, TO-3, TO-5, TO-7, and TO-11), or in the cytoplasm (SY-1, HB2, and TO-7), but with concurrent staining in normal epithelium, stroma or infiltrating leukocytes. HLA class I expression was considered to be lost when one of the HLA class I antigen-specific antibodies gave a negative result alongside a positive internal control (lymphocytic infiltrate).

3.3. Flow cytometric sorting

The flow cytometric sorting procedure, including tissue preparation, staining and flow cytometry analysis was performed as described previously [34]. Briefly, 2 mm diameter punches from selected areas of formalin-fixed paraffin embedded colorectal carcinomas were digested enzymatically in a mixture of 0.1% collagenase I-A (Sigma-Aldrich, St Louis, MO, USA) and 0.1% dispase (Gibco BRL, Paisley, UK). After determination of cell concentration, one million cells were incubated with 100 μ l of mAb mixture directed against keratin and vimentin containing clones MNF116 (anti-keratin; IgG1; DAKOCytomation, Glostrup, Denmark), AE1/AE3 (anti-keratin; IgG1; Chemicon International Inc, Temecula, CA, USA), and V9-2b (anti-vimentin; IgG2b; Department of Pathology, LUMC [35]). Next day, cells were incubated with 100 μ l of premixed FITC and RPE-labelled goat F(ab')₂ anti-mouse subclass-specific secondary reagents (Southern Biotechnology Associates, Birmingham, AL, USA). After washing, cells were incubated with 10 μ M propidium iodide (PI) and 0.1% DNase-free RNase (Sigma). The next day cells were analyzed by flow cytometry. A standard FACSCalibur (BD

Biosciences) was used for the simultaneous measurement of FITC, RPE, and PI. Tumor and normal cell populations were flow-sorted using a FACSVantage flow-sorter (BD Biosciences) using the FACSCalibur filter settings. Sorting was only performed on samples included in the RST array due to shortage of material from the HNPCC cases. DNA from flow-sorted tumor material was isolated as described by Jordanova et al. [36]. DNA from non-sorted material was isolated using Chelex extraction as described previously [37].

3.4. LOH and fragment analysis

Markers for loss of heterozygosity (LOH) analysis were chosen from the dbMHC database [38] to map the chromosome 6p21.3 region between *HLA-A* and *TAP2*. They were MOGc, D6S510, C125, C141, D6S2444, *TAP1* and M2426. A "linker" sequence of 5'-GTTTCTT was added to the 5' terminus of all reverse primers [39]. LOH was defined as allelic imbalance >2 in the HNPCC cases (non-sorted) and allelic imbalance >5 in the sorted RST [40].

To detect frame-shift mutations in the *HLA-A*, *HLA-B*, β 2m, *LMP2*, *LMP7*, *LMP10*, *TAP1*, *TAP2*, *Calnexin*, *Calreticulin*, *Erp57* and *Tapasin* genes, 28 pairs of primers (Table 1) were constructed surrounding non-polymorphic microsatellite regions within the coding regions.

3.5. Statistics

Significance values were calculated using the software package SPSS 10.0.7 (SPSS Inc., Chicago, IL, USA).

Gene	Forward primer	Reverse primer	Microsatellite Repeat
<i>HLA A 4th ex</i>	CCTGAATTTTCTGACTCTTCCCGT	GTTTCTTCCCGCTGCCAGGTCAGTGT	7(C)
<i>HLA A 5th ex</i>	CCATCGTGGGCATCATTTG	GTTTCTTTCAGTGAGACAAGAAATCTC	3(GGA)
<i>HLA B 2nd ex</i>	GCTTCATCTCAGTGGGCTAC	GTTTCTTCTCGTCTGGTTGTA	3(GA) + 3(CA)
<i>β2m 1st ex</i>	GGCTGGGCACGCGTTTAAT	GTTTCTTAGGGAGAGAAGGACCAGAG	4(CT)
<i>β2m 2nd ex (1)</i>	TACCTCGGCAATATTAATGTG	GTTTCTTGATAGAAAGACCAGTCCTTGC	4(GA) + 5(A)
<i>β2m 2nd ex (2)</i>	CTTACTGAAGAATGGAGAGAG	GTTTCTTGACTACTCATACACAATTTC	5(A)
<i>TAP1 1st ex</i>	TAAATGGCTGAGCTTCTCGC	GTTTCTTAGAGCTAGCCATTGGCA	5(C)
<i>TAP1 3rd ex</i>	ACAGCCACTTGACAGGGAG	GTTTCTTATGAACAGTACATGGCGTAT	5(T)
<i>TAP1 8th ex</i>	CTGCCCTGCTGCAGAATCTG	GTTTCTTCAAGCCACCTGCTTCCAT	5(G)
<i>TAP1 10th ex</i>	CTCGCAGAGGTAGACGAGG	GTTTCTTATTAAGAAGATGACTGCCTCAC	5(G)
<i>TAP1 11th ex</i>	AGCACCTCAGCCTGGTGGA	GTTTCTTGCAAGTCTGAGAAGGCTTTC	6(G) + 5(A)
<i>TAP2 2nd ex</i>	TTCTCAAGGGCTGCCAGGAC	GTTTCTTGCTCCAAGGGCTGAAG	6(C)
<i>TAP2 9th ex</i>	CCTACGCTCTGGTGAGGTGA	GTTTCTTCTGGCTGTGCAGGTAGC	5(G)
<i>Tapasin 2nd ex</i>	TTGGTTCGTGGAGGATGC	GTTTCTCTAGAGACTCACCCTGTAC	5(G)
<i>Tapasin 3rd ex</i>	CTTCTTCTCTACACTCAGACC	GTTTCTTAGGACTGGGCTGGATATGC	5(C)
<i>Tapasin 4th ex</i>	CCTGTCTTCTCAGTGGTAC	GTTTCTTGAGCAGATGTCCCTTACCC	6(C)
<i>Tapasin 5th ex</i>	TGCTCATTTCTGCTCTTTCC	GTTTCTTGTCCCACTCCACCTCCAG	5(G)
<i>Calnexin 7th ex</i>	GAAGGATCAGTTCATGACAAG	GTTTCTTCTGCATCTGGCCTCTTAGC	5(A)
<i>Calnexin 8th ex</i>	TCTGCTCAATGACATGACTCC	GTTTCTTTGAAGACAGTTCCTCAAGAC	5(A)
<i>Calnexin 11th ex</i>	AACCTTTCAGAATGACTCCTTTTAG	GTTTCTTCAAGCAGCAACACGAACC	8(T)
<i>Calreticulin 3rd ex</i>	CTACCGTCCCGTCTCAGG	GTTTCTTCTGTCTGGTCCAACTATTAGG	5(G)
<i>Calreticulin 6th ex</i>	GACAAGCCCCGAGCATATCC	GTTTCTTACCTTGTACTCAGGGTTCTG	5(C)
<i>Erp57 5th ex</i>	CACTTATTGCTTCTTCTTGTG	GTTTCTTAATACTTGGTCAGGAGATTCAAC	6(T)
<i>Erp57 6th ex</i>	CTTCTGCTATGCGCTACTGAG	GTTTCTTTCAAGCAAATAATCCAGACAAG	6(A)
<i>Erp57 13th ex</i>	ACTTTTAAGCTGATCTTCTGTTTT	GTTTCTTTTAGAGATCTCTGTGCCTT	6(C)
<i>LMP2 2nd ex</i>	GAGGGCATCAAGGCTGTTC	GTTTCTTGACAGACTCTGGGAATCAG	5(G)
<i>LMP2 6th ex</i>	CCCTCTCTCCAACCTGAAACC	GTTTCTTTGTAATAGTGACCAGGTAGATGAC	5(G)
<i>LMP7 1st ex</i>	GGCTTTCGCTTCACTTCC	GTTTCTTGAGATCGCATAGAAACTGTAG	6(C)

Table 1. Primers used in fragment analysis

4. RESULTS

4.1. HLA class I, β2m and APM component expression

In order to compare the expression of HLA class I in sporadic MSI-H and MSS right sided tumors (RST) and HNPCC MSI-H cases, we used an antibody panel recognizing monomorphic determinants expressed on HLA class I heavy chains, β2m and APM components (Figure 1).

In total, we identified loss of HLA class I expression in about 34.6% of the RST and 42.7%

of the HNPCC cases. The frequency of alterations differed significantly between the sporadic MSS and MSI-H RST. The lack of HLA class I expression was more frequent in MSI-H sporadic cases than in MSS cases ($p < 0.0001$), as it was 16.7% in the latter group, but 60.6% in the former (Table 2).

Subsequently, we have investigated the frequency of a concomitant loss of HLA class I expression with that of either the β2m molecule or of any APM component. In the sporadic subset, loss of HLA class I expression was more often associated with that of one of the APM




Figure 1. Example of immunohistochemical analysis performed on the RST and HNPCC tumors (Amplification 10×). A, Positive expression of HLA class I antigens detected with the HCA2 antibody. The epithelial (large arrow) membranous expression of HLA class I antigens is identical to the lymphocytic infiltrate (small arrow). B, Loss of expression of HLA class I identified with the HCA 2 antibody. The lymphocytic infiltrate (small arrow) was used as a positive control to determine the loss of expression on the epithelial cells. C, Loss of expression of β 2m in a HNPCC case. D, Loss of expression of one of the APM members (Tapasin) in a RST case.

			HLA [†]	APM loss [‡]	β 2m loss [‡]
RST N=81	MSS N=48 (59,3%)	+	40	0	0
		-	8 (16,7%)	3 (37,5%)*	0
	MSI-H N=33 (41,7%)	+	13	0	0
		-	20 (60,6%)	7 (35,0%)*	1 (5,0%)*
HNPCC N=75	MSI-H	+	43	0	0
		-	32 (42,7%)	2 (6,3%)*	15 (46,9%)*

Table 2. HLA, APM and β 2m expression in RST and HNPCC colon cancer

* – The percentage of cases that lost APM or β 2m expression is relative to the number of HLA negative cases of each subset

† – HLA expression differs significantly between sporadic MSI-H and MSS tumors ($p < 0.0001$) and between HNPCC and sporadic MSS tumors ($p < 0.005$)

‡ – The association of HLA class I loss with that of β 2m and of APM components differs significantly ($p < 0.0001$) between sporadic RST and HNPCC tumors.




Figure 2. Loss of expression of $\beta 2m$ and different APM members was detected by immunohistochemistry in the RST that presented with HLA loss. The shadowing (in black) is indicative for loss of expression of the respective molecules (tpsn – Tapasin, calnx – Calnexin, crtcln – Calreticulin).

components, occurring in about 37% of HLA-negative tumors regardless of their mismatch repair status (Table 2). $\beta 2m$ loss was only found in one HLA class I negative MSI-H sporadic tumor (case 65) that interestingly also presented loss of the APM molecules TAP2, Calreticulin and Tapasin (Figure 2). In contrast, loss of HLA class I expression in HNPCC cases was more frequently associated with that of $\beta 2m$ (Table 2), as it was found in 46.9% of the HLA class I-negative tumors. In contrast loss of any APM component was observed in only 6% of these cases (h38, h49) which also showed loss of $\beta 2m$ expression.

In sporadic RST, the simultaneous loss of more than one APM molecule per case was frequent (Figure 2). Only 3 out of 10 cases lost a single APM component. The TAP2 molecule was most frequently lost (6 cases), followed by TAP1, Tapasin and LMP2 (5 cases), Calreticulin (4 cases), LMP7 (2 cases), and Calnexin and ERp57 (1 case). Loss of the LPM10 protein was detected in neither sporadic RST nor HNPCC tumors. The HNPCC cases h38 and h49 lost the expression of TAP2 and LMP7 respectively.

4.2. LOH and frameshift analysis

Polymorphic markers around the classical *HLA* genes (*A*, *B* and *C*), *TAP1* and *TAP2* genes were used to study LOH and reveal possible chromosomal aberrations that could relate to loss of HLA class I expression (Figure 5A). In HNPCC cases, LOH analysis was only performed around the *HLA* genes since loss of the *TAP1* and *TAP2* proteins was rarely associated with HLA class I loss. LOH was more frequent in the MSS tumors (50%) than in the MSI-H sporadic (20%) and HNPCC (6%) tumors with loss of HLA class I expression ($P < 0.05$) (Figure 3, 4). Furthermore, the patterns of LOH in the MSS cases might indicate loss of the entire 6p21.3 region, in contrast to the MSI-H cases (hereditary and sporadic forms) where LOH seems to be limited.

Frameshift mutation screening of the microsatellite sequences present in the coding regions of the *HLA class I*, APM components and $\beta 2m$ genes was performed on all cases with aberrant HLA class I expression (Figure 5B,C). However specific genes were only analyzed when lack of expression of the encoded proteins was detected by immunohistochemistry. Of the classical HLA class I genes only *HLA-A* and *-B* were analyzed since *HLA-C* does not carry any microsatellite repeat in its coding region. Ten RST cases and 20 HNPCC control cases with normal expression of $\beta 2m$ and APM components were screened for frameshift mutations and none was detected.

Frameshift mutations were mainly found in the MSI-H cases (both sporadic and hereditary forms). At least one mutation in an APM component gene was found in 6 of 7 sporadic MSI-H tumors that lost expression of one or more APM components. The single sporadic MSI-H case that lost $\beta 2m$ expression also presented with a single frameshift mutation in the $\beta 2m$ gene. Of the 13 sporadic MSI-H cases in which loss of HLA class I expression was associated neither with APM component nor with $\beta 2m$ loss as detected by

	LOH							HLA-A		HLA-B		$\beta 2$ -m			TAP1					TAP2		TPSN				CALN3			CRTCLN		ERP57			LMP2		LMP7
	1	2	3	4	5	6	7	1	2	1	2	1	2	3	1	2	3	4	1	2	1	2	3	4	1	2	3	1	2	3	1	2				
MSH																																				
7								0	0	0																										
14								0	0	0																										
18								0	0	0																										
19								0	0	0																										
20								0	0	0																										
39								ins	0	0																										
44								0	0	0																										
45								0	0	0																										
51								0	0	0																										
65								0	0	0		0	0	del					0	0	0	0	del	0			ins	0								
67								0	0	0																										
69								0	0	0					del	0	0	0	0	0													0	0	0	
70								0	0	0																										
76								0	0	0																										
78								0	0	0																										
86								0	0	0																										
90								ins	0	0					0	0	0	0	0						0	0	del									
92								0	0	0																										
94								0	0	0																										
100								0	0	0																										
101								0	0	0																										
MSS																																				
41								0	0	0																										
57								0	0	0					0	0	0	0	0																	
61								0	0	0					0	0	0	0	0																	
62								0	0	0					0	0	0	0	0																	
83								0	0	0					0	0	0	0	0																	
96								0	0	0																										
99								0	0	0																										
102								0	0	0																										
Frameshift mutations - total								2		1				1				1		3				1		1		2		0		0				

Figure 3. LOH and frameshift analysis was performed on sporadic RST that lost HLA class I expression. Only the tumors that presented with loss of one of the APM molecules or $\beta 2m$ were subjected to fragment analysis in their respective genes. The following repeats were analyzed for frameshift mutations: HLA A: 1 – 4th exon 7(C), 2 – 5th exon 3 (GGA); HLA – B: 3(GA) & 3(CA); $\beta 2m$: 1 – 1st exon 4(CT), 2 – 2nd exon 4(GA) & 5(A), 3 – 2nd exon 5(A); TAP1: 1 – 1st exon 5(C), 2 – 3rd exon 5(T), 3 – 8th exon 5(G), 4 – 10th exon 5(G), 5 – 11th exon 6(G) & 5(A); TAP2: 1 – 2nd exon 6(C), 2 – 9th exon 5(G); Tapasin: 1 – 2nd exon 5(G), 2 – 3rd exon 5(C), 3 – 4th exon 6(C), 4 – 5th exon 5(G); Calnexin: 1 – 7th exon 5(A), 2 – 8th exon 5(A), 3 – 11th exon 8(T); Calreticulin: 1 – 3rd exon 5(G), 6th exon 5(C); ERp57: 1 – 5th exon 6(T), 2 – 6th exon 6(A), 3 – 13th exon 6(C); LMP2: 1 – 2nd exon 5(G), 2 – 6th exon 5(G); LMP7: 1st exon 6(C) (key: ins – insertion; del – deletion; 0 – no mutation). LOH analysis of the 6p chromosome was also performed with the following markers: 1 – MOGc, 2 – D6S510, 3 – C125, 4 – C141, 5 – D6S2444, 6 – TAP1, 7 – M2426 (Key: Black – Loss of heterozygosity; Striped – non informative marker; White – Retention of heterozygosity).

	LOH			HLA-A		HLA-B			$\beta 2$ -m			Criteria for HNPCC	
	1	2	3	1	2	1	2	3	1	2	3		
h4				0	0	0			del	del	0	<i>hMLH1</i> mutant	
h8				0	0	0			0	0	0	<i>hMLH1</i> mutant	
h26				0	0	0			0	0	0	MSI-H, no methylation on <i>hMLH1</i> , IHC MSH6 neg	
h38				0	0	0			del	0	0	<i>hMLH1</i> mutant	
h45				0	0	0			0	0	0	MSI-H, no methylation on <i>hMLH1</i>	
h49				0	0	0			0	0	0	<i>hMSH2</i> mutant	
h54				0	0	0			del	0	ins	<i>hMLH1</i> mutant	
h87				0	0	0			del	0	0	<i>hMSH6</i> mutant	
h93				0	0	0			0	0	0	<i>hMSH6</i> mutant	
h97				0	0	0			0	0	0	<i>hMLH1</i> mutant	
h99				0	0		ins		0	0	0	<i>hMLH1</i> mutant	
h103				0	0	0			0	0	ins	MSI-H, no methylation on <i>hMLH1</i> , IHC MSH6 neg	
h118				0	0	0			0	0	ins	<i>PMS2</i> mutant	
h119				0	0	0			0	0	del	<i>hMLH1</i> mutant	
h123				0	0	0			0	del	del	<i>hMSH6</i> mutant	
Total				0		1			11				

Figure 4. LOH and frameshift analysis was performed on HNPCC tumors that simultaneously lost HLA class I and $\beta 2m$ expression. LOH markers: (see legend from Figure 3 for key). Frameshift markers: HLA A: 1 – 4th exon 7(C) 2 – 5th exon 3 (GGA); HLA – B: 3(GA) & 3(CA); $\beta 2m$: 1 – 1st exon 4(CT), 2 – 2nd exon 4(GA), & 5(A), 3 – 2nd exon 5(A) (key: ins – insertion; del – deletion; 0 – no mutation; IHC -immunohistochemistry).




Figure 5. Genetic analysis performed on tumors that have lost HLA class I expression. The different peaks correspond to different sizes from the PCR-amplified products. Peaks corresponding to the normal samples are represented in green whereas tumor samples are represented in blue. A, LOH analysis performed on the RST 41 sample with the polymorphic marker C141. The total loss of a normal allele (on top) illustrates the technical advantage of using flow cytometric sorting to identify LOH events. B, Frameshift mutations identified in different members of the APM machinery. On top a homozygous deletion in the sample RST 65 on the 4th exon of the Tapasin gene is shown. On the bottom, a heterozygous deletion in the sample RST 18 on the second exon of the TAP2 gene is shown. C, Frameshift mutation identified in one HNPCC case (h4) in the 2nd exon of the $\beta 2m$ gene. Because flow sorting was not performed in the HNPCC cases, we cannot determine whether the frameshifts are homo- or heterozygous due to contamination with normal DNA.

immunohistochemical staining, one presented with a frameshift in an *HLA* gene (case 39, Figure 3, Table 3) while 3 other cases showed LOH of the markers adjacent the *HLA* genes (cases 7, 44 and 67). One frameshift mutation was found in the *Tapasin* gene in a MSS case (case 62). From 15 HNPCC tumors that lost $\beta 2m$ expression at least one mutation was found in 8 cases (Figure 4). Three of the latter showed 2 mutations localized in different stretches. In the remaining 17 HNPCC cases that solely lost HLA class I expression, only 4 showed genetic abnormalities. LOH was found in the HLA region in cases h16, h56 and h120 (data not shown). A frameshift mutation in one of the HLA genes (*HLA B*) was found in one case (h99, Figure 4). In neither of the 2 HNPCC cases that immunohistochemically lost the expression of one of the APM components an APM frameshift mutation was found.

5. DISCUSSION

Abnormalities in HLA class I cell surface expression are commonly observed in tumors and are interpreted as a mechanism by which tumor cells evade the host immune system [1]. In colorectal cancer, especially in MSI-H tumors, the high degree of lymphocytic infiltrate in some cases may suggest an active immune response during tumor development [41,42]. Moreover, MSI-H tumors might cause increased immune reactivity as a consequence of the high amounts of aberrant frameshift peptides they generate [8,18]. A selective pressure by CTLs upon these tumors would favor the outgrowth of tumor cells that lost HLA class I expression at the cell surface allowing them to surpass the action of the immune system.

RST	Mutation description
18	<i>TAP2</i> (341–346)del1 – het
39	<i>HLA-A</i> (621–627)ins1 – het
65	$\beta 2m$ (341–345)del1 – hom; <i>Tapasin</i> (1217–1222)del1 – hom; <i>Calreticulin</i> (418–422)ins1 – hom
69	<i>TAP1</i> (362–366)del1 – hom
78	<i>Erp57</i> (675–680)del2 – hom; <i>Erp57</i> (1459–1464)del1 – hom
90	<i>HLA-A</i> (621–627)ins1 – het; <i>Calnexin</i> (1476–1483)del1 – het
94	<i>Tapasin</i> (1217–1222)del1 – het

Table 3. Frameshift mutations description in RST

ins – insertion; del – deletion; hom – homozygous; het – heterozygous; the numbers before and after del/ins indicate the location of the microsatellite repeat containing the frameshift mutation and the number of affected nucleotides respectively.

Applying immunohistochemistry on tissue arrays, we compared HLA class I expression in both sporadic RST (MSI-H and MSS sub-groups) and HNPCC tumors. RST were chosen because of the high percentage of MSI-H cases in this specific tumor type [43]. Indeed, immunohistochemical staining with mAb showed that HLA class I loss was frequent in the MSI-H cases analyzed when compared to their MSS counterpart. This finding supports the hypothesis that MSI-H tumors face greater selective pressure to lose HLA class I expression, as described by Kloor et al[8]. However, we have shown for the first time that distinct molecular mechanisms underlie HLA class I loss in sporadic MSI-H and HNPCC colon cancers. In the latter, HLA class I loss was preferentially associated with that of $\beta 2m$, while in the former HLA class I loss was associated with that of one or more APM components ($p < 0.0001$).

We investigated the genetic abnormalities underlying the HLA class I loss of expression. They included LOH on chromosome region 6p21.3 (encompassing HLA class I and TAP genes), mutations in APM components and mutations in $\beta 2m$.

Loss of heterozygosity at 6p21.3 was most prevalent in MSS tumors. This is consistent with the observation that these tumors frequently possess gross chromosomal aberrations and are often aneuploid [13]. Moreover, since LOH

events in MSS tumors normally comprise large areas of a chromosome, LOH on 6p21 might not be a direct consequence of selective pressure directed to the loss of HLA expression but instead to other genes within the same chromosomal region. The general absence of LOH in MSI-H tumors suggests that this is not the major mechanism by which the cells abrogate HLA class I expression.

The genome's coding regions contain multiple microsatellite repeats, which are considered hotspots for mutations in mismatch repair-deficient tumors [43]. Such repeats are also present within the exons of the APM components, $\beta 2m$, *HLA-A* and *HLA-B* genes. In about half of the MSI-H cases, loss of expression of HLA class I was concordant with the detection of one or more mutations in these genes. We have discovered novel mutations in the antigen presenting machinery genes; *Tapasin*, *Erp57*, *Calreticulin* and *Calnexin* in colorectal cancer. Previous reports associated the loss of HLA class I expression in MSI-H tumors with defects on $\beta 2m$ molecule [7,9]. However, the authors did not distinguish the sporadic/hereditary nature of the tumors that were studied. We cannot exclude that the MSI-H cases included in these studies were mainly HNPCC tumors.

The reason sporadic MSI-H tumors would target APM members for inactivation and HNPCC

would target the $\beta 2m$ chaperon is unclear. One possibility worth further exploration is that the various mutations suggest different immune-escape mechanisms for thwarting distinct anti-tumor responses. HNPCC tumors can have an age of onset before the 5th decade of life while sporadic MSI-H tumors appear generally around the 7th decade of life [43]; one would therefore predict that the alertness and robustness of the immune system would be higher in HNPCC patients leading to a stronger, or at least different selective pressure on the latter. Furthermore it has been recently suggested that the JC polyoma virus plays a role in the oncogenicity of colon tumors with an identical phenotype to sporadic MSI-H tumors [44]. Although speculative, the presence of the JC virus might be implicated in a different immune response between sporadic MSI-H and HNPCC tumors.

The advantages of different escape mechanisms (loss of APM members vs. abrogation of $\beta 2m$) are not understood. The only known function of APM members is facilitating the expression of HLA classical molecules in complex with endogenous peptides. Thus, one would expect that only these HLA molecules would be affected by failure of the antigen processing machinery. On the other hand, it is accepted that cell surface expression of non-classical HLA molecules (e.g. HLA -G, -E) also depends on $\beta 2m$, so the function of these highly specialized molecules would be compromised if $\beta 2m$ were mutated or lost. These molecules might play an important role in regulation of immune cell activity by inhibiting or activating its function. Therefore, MSI-H sporadic tumors that have lost expression of both HLA and an APM component and HNPCC tumors with lost $\beta 2m$ expression might behave differently or present a different kind of interaction with cells from the immune system. For instance, Yamamoto et al. have described a correlation between

$\beta 2m$ mutations and unfavorable prognosis in colorectal cancer [45].

We separately analyzed the presence of the characteristic BRAF V600E somatic mutations in the RST cohort (data not shown). Forty-percent of MSI-H sporadic tumors presented with this mutation which was absent in the MSS tumors. It was previously described that this mutation is also absent in HNPCC tumors [46]. V600E was distributed equally between tumors that lost vs. retained expression of HLA class I in the sporadic MSI-H cases.

6. CONCLUSION

HLA class I aberrations are found at varying frequencies in different colorectal tumor types and are caused by distinct genetic mechanisms. Chiefly, sporadic and hereditary MSI-H tumors follow different routes toward HLA class I loss of expression supporting the idea that these tumors follow different evolutionary pathways in tumorigenesis. The resulting variation in immune escape mechanisms may have repercussions in tumor progression and behavior.

7. ACKNOWLEDGEMENTS

We thank Dr. Hans Vasen and Dr. Carli Tops for crucial information on the HNPCC cohort. We thank Dr. Jan Calame for providing us with RST tumors.

8. REFERENCES

1. Melief CJM: **Cancer immunology: Cat and mouse games.** *Nature* 2005, **437**:41-42.
2. Stevanovic S, Schild H: **Quantitative aspects of T cell activation - peptide generation and editing**

- by MHC class I molecule. *Semin Immunol* 1999, 11:375-384.
3. Algarra I, García-Lora A, Cabrera T, Ruiz-Cabello F, Garrido F: **The selection of tumor variants with altered expression of classical and nonclassical MHC class I molecules: implications for tumor immune escape.** *Cancer Immunol Immunother* 2004, 53:904-910.
 4. Chang CC, Ferrone S: **Immune selective pressure and HLA class I antigen defects in malignant lesions.** *Cancer Immunol Immunother* 2007, 56: 227-36.
 5. Koopman LA, van der Slik AR, Giphart MJ, Fleuren GJ: **Human leukocyte antigen class I gene mutations in cervical cancer.** *J Natl Cancer Inst* 1999, 91:1669-1677.
 6. Jordanova ES, Riemersma SA, Philippo K, Schuur- ing E, Kluin PM: **beta(2)-microglobulin aberrations in diffuse large B-cell lymphoma of the testis and the central nervous system.** *Int J Cancer* 2003, 103:393-398.
 7. Bicknell DC, Kaklamanis L, Hampson R, Bodmer WF, Karran P: **Selection for beta(2)-microglobulin mutation in mismatch repair-defective colorectal carcinomas.** *Curr Biol* 1996, 6:1695-1697.
 8. Kloor M, Becker C, Benner A, Woerner SM, Gebert J, Ferrone S, Knebel Doeberitz M: **Immunoselec- tive Pressure and Human Leukocyte Antigen Class I Antigen Machinery Defects in Microsat- ellite Unstable Colorectal Cancers.** *Cancer Res* 2005, 65:6418-6424.
 9. Cabrera CM, Jimenez P, Cabrera T, Esparza C, Ruiz-Cabello F, Garrido F: **Total loss of MHC class I in colorectal tumors can be explained by two molecular pathways: beta(2)-microglobulin inactivation in MSI-positive tumors and LMP7/ TAP2 downregulation in MSI-negative tumors.** *Tissue Antigens* 2003, 61:211-219.
 10. Ferris RL, Whiteside TL, Ferrone S: **Immune Escape Associated with Functional Defects in Antigen- Processing Machinery in Head and Neck Cancer.** *Clin Cancer Res* 2006, 12:3890-3895.
 11. Seliger B, Atkins D, Bock M, Ritz U, Ferrone S, Huber C, Storkel S: **Characterization of human lymphocyte antigen class I antigen-process- ing machinery defects in renal cell carcinoma lesions with special emphasis on transporter- associated with antigen-processing down-reg- ulation.** *Clin Cancer Res* 2003, 9:1721-1727.
 12. Klein J, Sato A: **The HLA System- First of Two Parts.** *N Engl J Med* 2000, 343:702-709.
 13. Takayama T, Miyanishi K, Hayashi T, Sato Y, Nirrsu Y: **Colorectal cancer: genetics of development and metastasis.** *J Gastroenterol* 2006, 41:185-192.
 14. Gervaz P, Bucher P, Morel P: **Two colons-two can- cers: Paradigm shift and clinical implications.** *J Surg Oncol* 2004, 88:261-266.
 15. Boland CR, Thibodeau SN, Hamilton SR, Sidransky D, Eshleman JR, Burt RW, Meltzer SJ, Rodriguez- Bigas MA, Fodde R, Ranzani GN, Srivastava S: **A National Cancer Institute Workshop on Mic- rosatellite Instability for cancer detection and familial predisposition: development of inter- national criteria for the determination of micro- satellite instability in colorectal cancer.** *Cancer Res* 1998, 58:5248-5257.
 16. Shin KH, Park YJ, Park JG: **Mutational Analysis of the Transforming Growth Factor {beta} Recep- tor Type II Gene in Hereditary Nonpolyposis Colorectal Cancer and Early-onset Colorectal Cancer Patients.** *Clin Cancer Res* 2000, 6:536-540.
 17. Umar A, Risinger JI, Hawk ET, Barrett JC: **Guide- lines - Testing guidelines for hereditary non- polyposis colorectal cancer.** *Nat Rev Cancer* 2004, 4:153-158.
 18. Linnebacher M, Gebert J, Rudy W, Woerner S, Yuan YP, Bork P, Doeberitz MV: **Frameshift pep- tide-derived T-cell epitopes: a source of novel tumor-specific antigens.** *Int J Cancer* 2001, 93: 6-11.
 19. Maleno I, Cabrera CM, Cabrera T, Paco L, Lopez- Nevot MA, Collado A, Ferrer A, Garrido F: **Dis- tribution of HLA class I altered phenotypes in colorectal carcinomas: high frequency of HLA**

- haplotype loss associated with loss of heterozygosity in chromosome region 6p21.** *Immunogenetics* 2004, **56**:244-253.
20. Watson NFS, Ramage JM, Madjd Z, Spendlove I, Ellis IO, Scholefield JH, Durrant LG: **Immunosurveillance is active in colorectal cancer as down-regulation but not complete loss of MHC class I expression correlates with a poor prognosis.** *Int J Cancer* 2006, **118**:6-10.
 21. Cabrera T, Collado A, Fernandez MA, Ferron A, Sancho J, Ruiz-Cabello F, Garrido F: **High frequency of altered HLA class I phenotypes in invasive colorectal carcinomas.** *Tissue Antigens* 1998, **52**:114-123.
 22. Dierssen JW, de Miranda N, Mulder A, van Puijenbroek M, Verduyn W, Claas F, van de Velde C, Jan Fleuren G, Cornelisse C, Corver W, Morreau H: **High-resolution analysis of HLA class I alterations in colorectal cancer.** *BMC Cancer* 2006, **6**: 233.
 23. Young J, Simms LA, Biden KG, Wynter C, Whitehall V, Karamatic R, George J, Goldblatt J, Walpole I, Robin SA, Borten MM, Stitz R, Searle J, McKeone D, Fraser L, Purdie DR, Podger K, Price R, Botten-shaw R, Walsh MD, Barker M, Leggett BA, Jass JR: **Features of colorectal cancers with high-level microsatellite instability occurring in familial and sporadic settings - Parallel pathways of tumorigenesis.** *Am J Pathol* 2001, **159**:2107-2116.
 24. van Puijenbroek M, van Asperen CJ, van Mil A, Devilee P, van Wezel T, Morreau H: **Homozygosity for a CHEK2* 1100delC mutation identified in familial colorectal cancer does not lead to a severe clinical phenotype.** *J Pathol* 2005, **206**: 198-204.
 25. Hendriks Y, Franken P, Dierssen JW, de Leeuw W, Wijnen J, Dreef E, Tops C, Breuning M, Brocker-Vriends A, Vasen H, Fodde R, Morreau H: **Conventional and Tissue Microarray Immunohistochemical Expression Analysis of Mismatch Repair in Hereditary Colorectal Tumors.** *Am J Pathol* 2003, **162**:469-477.
 26. Nygren AOH, Ameziane N, Duarte HMB, Vijzelaar RNCP, Waisfisz Q, Hess CJ, Schouten JP, Errami A: **Methylation-specific MLPA (MS-MLPA): simultaneous detection of CpG methylation and copy number changes of up to 40 sequences.** *Nucleic Acids Res* 2005, **33**: e128.
 27. *Code Proper Secondary Use of Human Tissue* [<http://www.federa.org/?s=1&m=78&p=&v=4>]
 28. Stam NJ, Spits H, Ploegh HL: **Monoclonal antibodies raised against denatured HLA-B locus heavy chains permit biochemical characterization of certain HLA-C locus products.** *J Immunol* 1986, **137**:2299-2306.
 29. Sernee MF, Ploegh HL, Schust DJ: **Why certain antibodies cross-react with HLA- A and HLA-G: Epitope mapping of two common MHC class I reagents.** *Mol Immunol* 1998, **35**:177-188.
 30. Perosa F, Luccarelli G, Prete M, Favoino E, Ferrone S, Dammacco F: **beta2-Microglobulin-Free HLA Class I Heavy Chain Epitope Mimicry by Monoclonal Antibody HC-10-Specific Peptide.** *J Immunol* 2003, **171**:1918-1926.
 31. Ogino T, Wang X, Kato S, Miyokawa N, Harabuchi Y, Ferrone S: **Endoplasmic reticulum chaperone-specific monoclonal antibodies for flow cytometry and immunohistochemical staining.** *Tissue Antigens* 2003, **62**:385-393.
 32. Bando N, Ogino T, Cho HS, Hur SY, Shen J, Wang X, Kato S, Miyokawa N, Harabuchi Y, Ferrone S: **Development and characterization of human constitutive proteasome and immunoproteasome subunit-specific monoclonal antibodies.** *Tissue Antigens* 2005, **66**:185-194.
 33. Wang X, Campoli M, Cho HS, Ogino T, Bando N, Shen J, Hur SY, Kageshita T, Ferrone S: **A method to generate antigen-specific mAb capable of staining formalin-fixed, paraffin-embedded tissue sections.** *J Immunol Methods* 2005, **299**:139-151.
 34. Corver WE, ter Haar NT, Dreef EJ, Miranda NFCC, Prins FA, Jordanova ES, Cornelisse CJ, Fleuren GJ: **High-resolution multi-parameter DNA flow**

- cytometry enables detection of tumour and stromal cell subpopulations in paraffin-embedded tissues. *J Pathol* 2005, **206**:233-241.
35. Vanmuijen GNP, Ruiter DJ, Warnaar SO: **Coexpression of Intermediate Filament Polypeptides in Human-Fetal and Adult Tissues.** *Lab Invest* 1987, **57**:359-369.
 36. Jordanova ES, Corver WE, Vonk MJ, Leers MPG, Riemersma SA, Schuurung E, Kluin PM: **Flow cytometric sorting of paraffin-embedded tumor tissues considerably improves molecular genetic analysis.** *Am J Clin Pathol* 2003, **120**:327-334.
 37. de Jong AE, van Puijenbroek M, Hendriks Y, Tops C, Wijnen J, Ausems MGEM, Meijers-Heijboer H, Wagner A, van Os TAM, Bocker-Vriend AHJT, Vasen HFA, Morreau H: **Microsatellite Instability, Immunohistochemistry, and Additional PMS2 Staining in Suspected Hereditary Nonpolyposis Colorectal Cancer.** *Clin Cancer Res* 2004, **10**:972-980.
 38. dbMHC [<http://www.ncbi.nlm.nih.gov/mhc/>]
 39. de Leeuw WJF, van Puijenbroek M, Merx R, Wijnen JT, Bocker-Vriend AHJT, Tops C, Vasen H, Cornelisse CJ, Morreau H: **Bias in detection of instability of the (C)8 mononucleotide repeat of MSH6 in tumours from HNPCC patients.** *Oncogene* 2001, **20**:6241-6244.
 40. Devilee P, Cleton-Jansen AM, Cornelisse CJ: **Ever since Knudson.** *Trends Genet* 2001, **17**:569-573.
 41. Phillips AB: **Tumour-infiltrating lymphocytes in colorectal cancer with microsatellite instability are activated and cytotoxic.** *Br J Surg* 2004, **91**:469-475.
 42. Dolcetti R, Viel A, Doglioni C, Russo A, Guidoboni M, Capozzi E, Vecchiato N, Macri E, Fornasari M, Boiocchi M: **High prevalence of activated intraepithelial cytotoxic T lymphocytes and increased neoplastic cell apoptosis in colorectal carcinomas with microsatellite instability.** *Am J Pathol* 1999, **154**:1805-1813.
 43. Ward R, Meagher A, Tomlinson I, O'Connor T, Norrie M, Wu R, Hawkins N: **Microsatellite instability and the clinicopathological features of sporadic colorectal cancer.** *Gut* 2001, **48**:821-829.
 44. Goel M, Li MS, Nagasaka T, Shin SK, Fuerst F, Ricciardiello L, Wasserman L, Boland CR: **Association of JC virus T-antigen expression with the methylator phenotype in sporadic colorectal cancers.** *Gastroenterology* 2006, **130**:1950-1961.
 45. Yamamoto H, Yamashita K, Perucho M: **Somatic mutation of the beta(2)-microglobulin gene associates with unfavorable prognosis in gastrointestinal cancer of the microsatellite mutator phenotype.** *Gastroenterology* 2001, **120**:1565-1567.
 46. Domingo E, Laiho P, Ollikainen M, Pinto M, Wang L, French AJ, Westra J, Frebourg T, Espin E, Armengol M, Hamelin R, Yamamoto H, Hofstra RMW, Seruca R, Lindblom A, Peltomaki P, Thibodeau SN, Aaltonen LA, Schwartz S: **BRAF screening as a low-cost effective strategy for simplifying HNPCC genetic testing.** *J Med Genet* 2004, **41**:664-668.

