



Universiteit
Leiden
The Netherlands

The MUSE 3D view of the Hubble Deep Field South

Bacon, R.; Brinchmann, J.; Richard, J.; Contini, T.; Drake, A.; Franx, M.; ... ; Zeeuw, T.

Citation

Bacon, R., Brinchmann, J., Richard, J., Contini, T., Drake, A., Franx, M., ... Zeeuw, T. (2015). The MUSE 3D view of the Hubble Deep Field South. *Astronomy & Astrophysics*, 575, A75. Retrieved from <https://hdl.handle.net/1887/48405>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/48405>

Note: To cite this publication please use the final published version (if applicable).

The MUSE 3D view of the *Hubble* Deep Field South^{★,★★}

R. Bacon¹, J. Brinchmann², J. Richard¹, T. Contini^{3,4}, A. Drake¹, M. Franx², S. Tacchella⁵, J. Vernet⁶, L. Wisotzki⁷, J. Blaizot¹, N. Bouché^{3,4}, R. Bouwens², S. Cantalupo⁵, C. M. Carollo⁵, D. Carton², J. Caruana⁷, B. Clément¹, S. Dreizler⁸, B. Epinat^{3,4,9}, B. Guiderdoni¹, C. Herenz⁷, T.-O. Husser⁸, S. Kamann⁸, J. Kerutt⁷, W. Kollatschny⁸, D. Krajnovic⁷, S. Lilly⁵, T. Martinsson², L. Michel-Dansac¹, V. Patricio¹, J. Schaye², M. Shirazi⁵, K. Soto⁵, G. Soucail^{3,4}, M. Steinmetz⁷, T. Urrutia⁷, P. Weilbacher⁷, and T. de Zeeuw^{6,2}

¹ CRAL, Observatoire de Lyon, CNRS, Université Lyon 1, 9 avenue Ch. André, 69561 Saint Genis-Laval Cedex, France
e-mail: roland.bacon@univ-lyon1.fr

² Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

³ IRAP, Institut de Recherche en Astrophysique et Planétologie, CNRS, 14 avenue Édouard Belin, 31400 Toulouse, France

⁴ Université de Toulouse, UPS-OMP, 31400 Toulouse, France

⁵ ETH Zurich, Institute of Astronomy, Wolfgang-Pauli-Str. 27, 8093 Zurich, Switzerland

⁶ ESO, European Southern Observatory, Karl-Schwarzschild Str. 2, 85748 Garching bei Muenchen, Germany

⁷ AIP, Leibniz-Institut für Astrophysik Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

⁸ AIG, Institut für Astrophysik, Universität Göttingen, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany

⁹ Aix-Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, 13388 Marseille, France

Received 27 November 2014 / Accepted 20 January 2015

ABSTRACT

We observed *Hubble* Deep Field South with the new panoramic integral-field spectrograph MUSE that we built and have just commissioned at the VLT. The data cube resulting from 27 h of integration covers one arcmin² field of view at an unprecedented depth with a 1σ emission-line surface brightness limit of 1×10^{-19} erg s⁻¹ cm⁻² arcsec⁻², and contains ~90 000 spectra. We present the combined and calibrated data cube, and we performed a first-pass analysis of the sources detected in the *Hubble* Deep Field South imaging. We measured the redshifts of 189 sources up to a magnitude $I_{814} = 29.5$, increasing the number of known spectroscopic redshifts in this field by more than an order of magnitude. We also discovered 26 Ly α emitting galaxies that are not detected in the HST WFPC2 deep broad-band images. The intermediate spectral resolution of 2.3 Å allows us to separate resolved asymmetric Ly α emitters, [O II]3727 emitters, and C III]1908 emitters, and the broad instantaneous wavelength range of 4500 Å helps to identify single emission lines, such as [O III]5007, H β , and H α , over a very wide redshift range. We also show how the three-dimensional information of MUSE helps to resolve sources that are confused at ground-based image quality. Overall, secure identifications are provided for 83% of the 227 emission line sources detected in the MUSE data cube and for 32% of the 586 sources identified in the HST catalogue. The overall redshift distribution is fairly flat to $z = 6.3$, with a reduction between $z = 1.5$ to 2.9, in the well-known redshift desert. The field of view of MUSE also allowed us to detect 17 groups within the field. We checked that the number counts of [O II]3727 and Ly α emitters are roughly consistent with predictions from the literature. Using two examples, we demonstrate that MUSE is able to provide exquisite spatially resolved spectroscopic information on the intermediate-redshift galaxies present in the field. This unique data set can be used for a wide range of follow-up studies. We release the data cube, the associated products, and the source catalogue with redshifts, spectra, and emission-line fluxes.

Key words. cosmology: observations – galaxies: evolution – galaxies: high-redshift – techniques: imaging spectroscopy – galaxies: formation

1. Introduction

The *Hubble* Deep Fields North and South (see e.g. Williams et al. 1996; Ferguson et al. 2000; Beckwith et al. 2006) are still among the deepest images ever obtained in the optical/infrared, providing broad band photometry for sources up to $V \sim 30$. Coupled with extensive multi-wavelength follow-up campaigns they and the subsequent *Hubble* Ultra Deep Field have been instrumental in improving our understanding of galaxy formation and evolution in the distant Universe.

Deep, broad-band, photometric surveys provide a wealth of information on galaxy populations, such as galaxy morphology, stellar masses, and photometric redshifts. Taken together, this can be used to study the formation and evolution of the *Hubble* sequence (e.g. Mortlock et al. 2013; Lee et al. 2013), the change in galaxy sizes with time (e.g. van der Wel et al. 2014; Carollo et al. 2013), and the evolution of the stellar mass function with redshift (e.g. Muzzin et al. 2013; Ilbert et al. 2013). But photometric information alone gives only a limited view of the Universe: essential physical information, such as the kinematic state of the galaxies and their heavy element content, require spectroscopic observations. Furthermore, while photometric redshifts work well on average for many bright galaxies (e.g. Ilbert et al. 2009), they have insufficient precision for environmental studies and are occasionally completely wrong, and their performance on very faint galaxies is not well known.

* Advanced data products are available at <http://muse-vlt.eu/science>. Based on observations made with ESO telescopes at the La Silla Paranal Observatory under program ID 60.A-9100(C).

** Appendices are available in electronic form at <http://www.aanda.org>

Ideally, one would like to obtain spectroscopy for all sources at the same depth as the broad-band photometry. However current technology does not match these requirements. For example, the VIMOS ultra-deep survey (Le Fevre et al. 2015), which is the largest spectroscopic deep survey today with 10 000 observed galaxies in 1 deg^2 , is in general limited to $R \sim 25$, and only 10% of the galaxies detected in the *Hubble* Deep Fields North and South are that bright.

Another fundamental limitation, when using multi-object spectrographs, is the need to pre-select a sample based on broad band imaging. Even if it was feasible to target all objects found in the *Hubble* deep fields WFPC2 deep images (i.e. ~ 6000 objects), the sample would still not include all galaxies with high equivalent-width emission lines, even though determining redshifts for these would be relatively easy. For example, faint low-mass galaxies with high star formation rate at high redshifts may not have an optical counterpart even in very deep HST broad-band imaging, although their emission lines arising in their star-forming interstellar medium might be detectable spectroscopically.

Long-slit observations are not a good alternative because of the limited field of view and other technical limitations due to slits such as unknown slit light losses, loss of positional information perpendicular to the slit and possible velocity errors. For example, Rauch et al. (2008) performed a long slit integration of 92 h with FORS2 at the VLT. They targeted the redshift range of 2.67–3.75, and their observations went very deep, with a emission-line surface-brightness limit (1σ) depth of $8.1 \times 10^{-20} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ arcsec}^{-2}$. However, this performance was obtained with a field of view of 0.25 arcmin^2 and only one spatial dimension, limiting the usefulness of this technique for any follow-up surveys of *Hubble* deep fields.

To overcome some of these intrinsic limitations, a large, sensitive integral-field spectrograph is required. It must be sensitive and stable enough to be able to reach a depth commensurate to that of the *Hubble* deep fields, while at the same time having a high spatial resolution, large multiplex, spectral coverage, and good spectral resolution. This was at the origin of the Multi Unit Spectroscopic Explorer (MUSE) project to build a panoramic integral field spectrograph for the VLT (Bacon et al. 2010; and in prep.). The commissioning of MUSE on the VLT was completed in August 2014 after development by a consortium of seven European institutes: CRAL (lead), AIG, AIP, ETH-Zurich, IRAP, NOVA/Leiden Observatory and ESO. The instrument has a field of view of $1 \times 1 \text{ arcmin}^2$ sampled at 0.2 arcsec, an excellent image quality (limited by the 0.2 arcsec sampling), a wide simultaneous spectral range (4650–9300 Å), a medium spectral resolution ($R \approx 3000$), and a very high throughput (35% end-to-end including telescope at 7000 Å).

Although MUSE is a general purpose instrument and has a wide range of applications (see Bacon et al. 2014, for a few illustrations), it has been, from the very start of the project in 2001, designed and optimised for performing deep field observations. Some preliminary measurements during the first commissioning runs had convinced us that MUSE was able to reach its combination of high throughput and excellent image quality. However it is only by performing a very long integration on a deep field that one can assess the ultimate performance of the instrument. This was therefore one key goal of the final commissioning run of ten nights in dark time late July 2014. The *Hubble* Deep Field South (HDFS) which was observable during the second half of the nights for a few hours, although at relatively high airmass, was selected as the ideal target to validate the performance of

MUSE, the observing strategy required for deep fields to limit systematic uncertainties and to test the data reduction software.

The HDFS was observed with the *Hubble* Space Telescope in 1998 (Williams et al. 2000). The WFPC2 observations (Casertano et al. 2000) reach a 10σ limiting AB magnitude in the *F606W* filter (hereafter V_{606}) at 28.3 and 27.7 in the *F814W* filter (hereafter I_{814}). The field was one of the first to obtain very deep Near-IR multi-wavelength observations (e.g., Labbé et al. 2003). But, contrarily to the *Hubble* Ultra Deep Field (Beckwith et al. 2006) which had very extensive spectroscopic follow-up observations, the main follow-up efforts on the HDFS have been imaging surveys (e.g. Labbé et al. 2003, 2005).

In the present paper we show that the use of a wide-field, highly sensitive IFU provides a very powerful technique to target such deep HST fields, allowing a measurement of ~ 200 redshifts per arcmin^2 , ~ 30 of which are not detected in the HDF continuum images. We present the first deep observations taken with the MUSE spectrograph, and we determine a first list of secure redshifts based on emission-line and absorption-line features. We show that the high spectral resolution, broad wavelength range, and three dimensional nature of the data help to disentangle confused galaxies and to identify emission lines securely.

The paper is organised as follows: the observations, data reduction and a first assessment of instrument performance are described in the next two sections. In Sects. 4 and 5, we proceed with the source identifications and perform a first census of the field content. The resulting redshift distribution and global properties of the detected objects are presented in Sect. 6. Examples of kinematics analysis performed on two spatially resolved galaxies is given in Sect. 7. Comparison with the current generation of deep spectroscopic surveys and conclusions are given in the last section. The data release is described in the Appendices.

2. Observations

The HDFS was observed during six nights in July 25–29, 31 and August 2, 3 2014 of the last commissioning run of MUSE. The $1 \times 1 \text{ arcmin}^2$ MUSE field was centred at $\alpha = 22\text{h}32^{\text{m}}55.64^{\text{s}}$, $\delta = -60^{\circ}33'47''$. This location was selected in order to have one bright star in the Slow Guiding System (SGS) area and another bright star in the field of view (Fig. 1). We used the nominal wavelength range (4750–9300 Å) and performed a series of exposures of 30 min each. The spectrograph was rotated by 90° after each integration, and the observations were dithered using random offsets within a 3 arcsec box. This scheme ensures that most objects will move from one channel¹ to a completely different one while at the same time minimizing the field loss. This is, however, not true for the objects that fall near the rotation centre.

In addition to the standard set of calibrations, we obtained a flat field each hour during the night. These single flat field exposures, referred to as attached flats in the following, are used to correct for the small illumination variations caused by temperature variations during the night. The Slow Guiding System was activated for all exposures using a bright $R = 19.2$ star located in the SGS field. The SGS also gives an accurate real-time estimate of the seeing which was good for most of the nights (0.5–0.9 arcsec). Note that the values given by the SGS are much closer to the seeing achieved in the science exposure than the values given by the DIMM seeing monitor. An astrometric solution was derived using an off-centre field of a globular cluster

¹ The field of view of MUSE is first split in 24 “channels”, each channel is then split again in 48 “slices” by the corresponding image slicer.




Fig. 1. Location of the MUSE field of view within the HDF-S *F814W* image. The star used in the slow guiding system is indicated in red and the brightest star in the field ($R = 19.6$) in blue.

with HST data. A set of spectrophotometric standard stars was also observed when the conditions were photometric.

In total, 60 exposures of 30 min integration time were obtained. A few exposures were obtained in cloudy conditions and were discarded. One exposure was lost due to an unexpected VLT guide star change in the middle of the exposure. The remaining number of exposures was 54, with a total integration time of 27 h. One of these exposures was offset by approximately half the field of view to test the performance of the SGS guiding on a faint galaxy.

3. Data reduction and performance analysis

3.1. Data reduction process

The data were reduced with version 0.90 of the MUSE standard pipeline. The pipeline will be described in detail in Weilbacher et al. (in prep.)². We summarize the main steps to produce the fully reduced data cube:

1. Bias, arcs and flat-field master calibration solutions were created using a set of standard calibration exposures obtained each night.
2. Bias images were subtracted from each science frame. Given its low value, the dark current ($\sim 1 \text{ e}^- \text{ h}^{-1}$, that is 0.5 e^- per exposure) was neglected. Next, the science frames were flat-fielded using the master flat field and renormalized using the attached flat field as an illumination correction. An additional flat-field correction was performed using the twilight sky exposures to correct for the difference between sky and calibration unit illumination. The result of this process is a large table (hereafter called a pixel-table) for each science frame. This table contains all pixel values corrected for bias and flat-field and their location on the detector.

A geometrical calibration and the wavelength calibration solution were used to transform the detector coordinate positions to wavelengths in Ångström and focal plane spatial coordinates.

3. The astrometric solution was then applied. The flux calibration was obtained from observations of the spectrophotometric standard star Feige 110 obtained on August 3, 2014. We verified that the system response curve was stable between the photometric nights with a measured scatter below 0.2% rms. The response curve was smoothed with spline functions to remove high frequency fluctuations left by the reduction. Bright sky lines were used to make small corrections to the wavelength solution obtained from the master arc. All these operations have been done at the pixel-table level to avoid unnecessary interpolation. The formal noise was also calculated at each step.
4. To correct for the small shifts introduced by the derotator wobble between exposures, we fitted a Gaussian function to the brightest star in the reconstructed white-light image of the field. The astrometric solution of the pixel-tables of all exposures was normalized to the HST catalogue coordinate of the star ($\alpha = 22\text{h}37'57.0''$, $\delta = -60^\circ34'06''$). The fit to the star also provides an accurate measurement of the seeing of each exposure. The average Gaussian white-light full width at half maximum (FWHM) value for the 54 exposures is 0.77 ± 0.15 arcsec. We also derived the total flux of the reference star by simple aperture photometry, the maximum variation among all retained exposures is 2.4%.
5. To reduce systematic mean zero-flux level offsets between slices, we implemented a non-standard self-calibration process. From a first reconstructed white-light image produced by the merging of all exposures, we derived a mask to mask out all bright continuum objects present in the field of view. For each exposure, we first computed the median flux over all wavelengths and the non-masked spatial coordinates. Next we calculated the median value for all slices, and we applied an additive correction to each slice to bring all slices to the same median value. This process very effectively removed residual offsets between slices.
6. A data cube was produced from each pixel-table using a 3D drizzle interpolation process which include sigma-clipping to reject outliers such as cosmic rays. All data cubes were sampled to a common grid in view of the final combination ($0'.2 \times 0'.2 \times 1.25 \text{ \AA}$).
7. We used the software ZAP (Soto et al., in prep.) to subtract the sky signal from each of the individual exposures. ZAP operates by first subtracting a baseline sky level, found by calculating the median per spectral plane, leaving any residuals due to variations in the line spread function (LSF) and system response. The code then uses principal component analysis on the data cube to calculate the eigenspectra and eigenvalues that characterize these residuals, and determines the minimal number of eigenspectra that can reconstruct the residual emission features in the data cube.
8. The 54 data cubes were then merged in a single data cube using 5σ sigma-clipped mean. The variance for each combined volume pixel or “voxel” was computed as the variance derived from the comparison of the N individual exposures divided by $N-1$, where N is the number of voxels left after the sigma-clipping. This variance data cube is saved as an additional extension of the combined data cube. In addition an exposure map data cube which counts the number of

² A short description is also given in Weilbacher et al. (2012).




Fig. 2. Derived FWHM as function of wavelength from the Moffat fit to the brightest star in the field.

exposures used for the combination of each voxel was also saved.

9. Telluric absorption from H₂O and O₂ molecules was fitted to the spectrum of a white dwarf found in the field ($\alpha = 22\text{h}32'58.77''$, $\delta = -60^\circ33'23.52''$) using the `molecfits` software described in [Smette et al. \(2015\)](#) and [Kausch et al. \(2015\)](#). In the fitting process, the LSF was adjusted using a wavelength dependent Gaussian kernel. The resulting transmission correction was then globally applied to the final data cube and variance estimation.

The result of this process is a fully calibrated data cube of 3 Gb size with spectra in the first extension and the variance estimate in the second extension, as well as an exposure cube of 1.5 Gb size giving the number of exposures used for each voxel.

3.2. Reconstructed white-light image and point-spread functions

The image quality was assessed using a Moffat fit of the reference star as a function of wavelength. The PSF shape is circular with a fitted Moffat β parameter of 2.6 and a FWHM of 0.66 arcsec at 7000 Å. While β is almost constant with wavelength, the FWHM shows the expected trend with wavelength decreasing from 0.76 arcsec in the blue to 0.61 arcsec in the red (Fig. 2). Note that the FWHM derived from the MOFFAT model is systematically 20% lower than the Gaussian approximation.

The spectral LSF was measured on arc calibration frames. We obtain an average value of 2.1 ± 0.2 pixels which translates into a spectral resolution of $R\ 3000 \pm 100$ at 7000 Å. A precise measurement of the LSF shape is difficult because it is partially under sampled. In the present case this uncertainty is not problematic because the spectral features of the identified objects are generally broader than the LSF.

A simple average over all wavelengths gave the reconstructed white light image (Fig. 3). Inspection of this image reveals numerous objects, mostly galaxies. The astrometric accuracy, derived by comparison with the [Casertano et al. \(2000\)](#) catalogue, is ~ 0.1 arcsec. At lower flux levels, $F \sim 2 \times 10^{-21}$ erg s⁻¹ cm⁻² Å⁻¹ pixel⁻¹, some residuals of the instrument channel splitting can be seen in the reconstructed white-light image in the form of a series of vertical and horizontal stripes.




Fig. 3. Reconstructed white-light image of the combined exposures. The flux scale shown on the right is in 10^{-21} erg s⁻¹ cm⁻² Å⁻¹ pixel⁻¹. Orientation is north(up)-east(left).

3.3. Signal to noise ratios

Characterising the noise in a MUSE data cube is not trivial, as each voxel of the cube is interpolated not only spatially, but also in the spectral domain, and each may inherit flux from just one up to ~ 30 original CCD pixels. While readout and photon noise are formally propagated by the MUSE pipeline as pixel variances, correlations between two neighbouring spatial pixels cause these predicted variances to be systematically too low. Furthermore, the degree of correlation between two neighbouring spatial pixels varies substantially across the field (and with wavelength), on spatial scales comparable to or larger than real astronomical objects in a deep field such as the HDFs. A correct error propagation also accounting for the covariances between pixels is theoretically possible, but given the size of a MUSE dataset this is unfortunately prohibitive with current computing resources. We therefore have to find other ways to estimate the “true” noise in the data.

For the purpose of this paper we focus on faint and relatively small sources, and we therefore neglect the contribution of individual objects to the photon noise. In addition to readout and sky photon noise, unresolved low-level systematics can produce noise-like modulations of the data, especially when varying rapidly with position and wavelength. Such effects are certainly still present in MUSE data, e.g. due to the residual channel and slice splitting pattern already mentioned in Sect. 3. Another issue are sky subtraction residuals of bright night sky emission lines which are highly nonuniform across the MUSE field of view. Future versions of the data reduction pipeline will improve on these features, but for the moment we simply absorb them into an “effective noise” budget.

In the HST image of the HDFs we selected visually a set of 100 “blank sky” locations free from any continuum or known emission line sources. These locations were distributed widely over the MUSE field of view, avoiding the outskirts of the brightest stars and galaxies. We extracted spectra through circular apertures from the sky-subtracted cube which consequently should have an expectation value of zero at all wavelengths. Estimates of the effective noise were obtained in two different ways: (A) By measuring the standard deviation inside of

spectral windows selected such that no significant sky lines are contained; (B) by measuring the standard deviation of aperture-integrated fluxes between the 100 locations, as a function of wavelengths. Method (A) directly reproduces the “noisiness” of extracted faint-source spectra, but cannot provide an estimate of the noise for all wavelengths. Method (B) captures the residual systematics also of sky emission line subtraction, but may somewhat overestimate the “noisiness” of actual spectra. We nevertheless used the latter method as a conservative approach to construct new “effective” pixel variances that are spatially constant and vary only in wavelength. Overall, the effective noise is higher by a factor of ~ 1.4 than the local pixel-to-pixel standard deviations, and by a factor of ~ 1.6 higher than the average propagated readout and photon noise. Close to the wavelengths of night sky emission lines, these factors may get considerably higher, mainly because of the increased residuals.

The median effective noise per spatial and spectral pixel for the HDFs data cube is then $9 \times 10^{-21} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ \AA}^{-1}$, outside of sky lines. For an emission line extending over 5 \AA (i.e. 4 spectral pixels), we derive a 1σ emission line surface brightness limit of $1 \times 10^{-19} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ arcsec}^{-2}$.

An interesting comparison can be made with the Rauch et al. (2008) deep long slit integration. In 92 h, they reached a 1σ depth of $8.1 \times 10^{-20} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ arcsec}^{-2}$, again summed over 5 \AA . Rauch et al. covered the wavelength range $4457\text{--}5776 \text{ \AA}$, which is $3.4\times$ times smaller than the MUSE wavelength range, and they cover an area (0.25 arcmin^2) which is four times smaller. Folding in the ratio of exposure times and the differences in achieved flux limits, the MUSE HDFs data cube is then in total over 32 times more effective for a blind search of emission line galaxies than the FORS2 observation. This is not a surprise given that FORS2 was not designed for this specific application but for imaging and slit spectroscopy over a $\sim 20 \text{ arcmin}^2$ field of view.

From the limiting flux surface brightness one can also derive the limiting flux for a point source. This is, however, more complex because it depends of the seeing and the extraction method. A simple approximation is to use fix aperture. For a 1 arcsec diameter aperture, we measured a light loss of 40% at 7000 \AA for the brightest star in the MUSE field. Using this value, we derive an emission line limiting flux at 5σ of $3 \times 10^{-19} \text{ erg s}^{-1} \text{ cm}^{-2}$ for a point source within a 1 arcsec aperture.

For a better understanding of the contribution of systematics to the noise budget, we also investigated the scaling of the noise with exposure time. Taking the effective noise in a single 30 min exposure as unit reference, we also measured the noise in coadded cubes of 4, 12, and the full set of 54 exposures. While systematic residuals are also expected to decrease because of the rotational and spatial dithering, they would probably not scale with $1/\sqrt{n}$ as perfect random noise. The result of this exercise is shown in Fig. 4, which demonstrates that a significant deviation from $1/\sqrt{n}$ is detected, but that it is quite moderate (factor ~ 1.2 for the full HDFs cube with $n = 54$). Even without reducing the systematics, adding more exposures would make the existing HDFs dataset even deeper.

4. Source identification and redshift determination

The three-dimensional nature of the MUSE observations presents unique challenges, while at the same time offering multiple ways to extract spectra and to determine and confirm redshifts. We have found that constructing 1D and 2D projections of the sources is essential to ascertain redshifts for the fainter




Fig. 4. Overall scaling of pixel noise measured in the cubes as a function of the number of exposures combined, relative to the noise in a single exposure. The solid line represents the ideal $1/\sqrt{n}$ behaviour.

sources. In particular we have constructed 1D spectra and for each tentative emission line we construct a continuum subtracted narrow-band image over this line, typically with a width of 10 \AA , and only if this produces a coherent image of the source we consider the emission line real. Likewise, 2D spectra can be useful additional tools for understanding the spatial distribution of emission.

The extraction of spectra from a very deep data cube can be challenging since the ground based seeing acts to blend sources. Here too the construction of 2D images can help disentangling sources that otherwise would be blended. For this step the existence of deep HST images is very helpful to interpret the results.

This method does not lend itself to absorption line redshifts. In this case we examine the spatial variation of possible absorption line features by extracting spectra at various spatial positions. A real absorption line should be seen in multiple spectra across the galaxy.

4.1. Redshift determination of continuum detected objects

We extracted subcubes around each object in the Casertano et al. (2000) catalogue that fell within the FoV sampled by the observations. We defined our spectrum extraction aperture by running SExtractor (Bertin & Arnouts 1996), version 2.19.5 on the reconstructed white-light images. In the case when no object was clearly detected in the white-light image, a simple circular extraction aperture with diameter $1.4''$ was used. When a redshift was determined, we constructed narrow-band images around Ly- α , C III]1909, [O II]3727, H β , [O III]5007, and H α , whenever that line fell within the wavelength range of MUSE, and ran SExtractor on these as well. The union of the emission-line and the white-light segmentation maps define our object mask. The SExtractor segmentation map was also used to provide a sky mask. The object and sky masks were inspected and manually adjusted when necessary to mask out nearby sources and to avoid edges.

The local sky residual spectrum is constructed by averaging the spectra within the sky mask. The object spectrum was constructed by summing the spectra within the object mask, subtracting off the average sky spectrum in each spatial pixel (spaxel). We postpone the optimal extraction of spectra (e.g. Horne 1986) to future work as this is not essential for the present paper. Note also that we do not account for the wavelength variation of the PSF in our extraction – this is a significant concern for optimal extraction but for the straight summation we found




Fig. 5. Example of the extraction process for objects 5 and 144, at $z = 0.58$ and $z = 4.02$ respectively. *Top row the left panel:* MUSE reconstructed white-light image, while the *left panel in the bottom row* shows the $\text{Ly}\alpha$ narrow-band image. *Middle panels:* object and sky masks, with the object aperture shown in black and the sky aperture in blue. Part of the final extracted spectrum is shown in the *right-hand panel* on each row.

by testing on stars within the data cube that the effect is minor for extraction apertures as large as ours.

An example of the process can be seen in Fig. 5. The top row shows the process for a $z \approx 0.58$ galaxy with the white-light image shown in the left-most column, and the bottom row the same for a $\text{Ly}\alpha$ -emitter at $z = 4.02$ with the $\text{Ly}\alpha$ -narrow-band image shown in the left-most column. The region used for the local sky subtraction is shown in blue in the middle column, while the object mask is shown in black.

The resulting spectra were inspected manually and emission lines and absorption features were identified by comparison to template spectra when necessary. In general an emission line redshift was considered acceptable if a feature consistent with an emission line was seen in the 1D spectrum and coherent spatial feature was seen in several wavelength planes over this emission line. In some cases mild smoothing of the spectrum and/or the cube was used to verify the reality of the emission line. In the case of absorption line spectra several absorption features were required to determine a redshift.

In many cases this process gives highly secure redshifts, with multiple lines detected in 72 galaxies and 8 stars. We assign them a Confidence = 3. In general the identification of single line redshifts is considerably less challenging than in surveys carried out with low spectral resolution. The [O II]3726, 3729 and C III]1907, 1909 doublets are in most cases easily resolved and the characteristic asymmetric shape of $\text{Ly}\alpha$ is easily identified. In these cases we assign a Confidence = 2 for single-line redshifts with high signal-to-noise (S/N). In a number of cases we do see unresolved [O II]3726, 3729 – in these cases we still have a secure redshift from Balmer absorption lines and/or [Ne III]3869 – but these lines are unresolved due to velocity broadening and we do not expect to see this behaviour in spectra of very faint galaxies. The other likely cases of single line redshifts with a *symmetric* line profile are: $\text{H}\alpha$ -emitters with undetected [N II]6584 and no accompanying strong [O III]5007; [O III]5007 emitters with undetectable [O III]4959 and $\text{H}\beta$; and $\text{Ly}\alpha$ -emitters with symmetric line profiles.

To distinguish between these alternatives, we make use of two methods. The first is to examine the continuum shape of the spectra. For the brighter objects breaks in the spectra can be used to separate between the various redshift solutions. The second method is to check the spectrum at the location of any other possible line, and to extract narrow-band images for all possible strong lines – this is very useful for [O III]5007 and $\text{H}\alpha$ emitters,

and can exclude or confirm low redshift solutions. If this process does not lead to a secure redshift, we assign a confidence 0 to these sources. The redshift confidence assignments is summarized below:

- 0: No secure or unique redshift determination possible;
- 1: Redshift likely to be correct but generally based on only one feature;
- 2: Redshift secure, but based on one feature;
- 3: Redshift secure, based on several features.

In the case of overlapping sources we do not attempt to optimally extract the spectrum of each source, leaving this to future papers. In at least four cases we see two sets of emission lines in the extracted spectrum and we are unable to extract each spectrum separately. Despite this we are still generally able to associate a redshift to a particular object in the HST catalogue by looking at the distribution of light in narrow-band images. We also use these narrow band images to identify cases where a strong emission line in a nearby object contaminates the spectrum of a galaxy.

4.2. Identification of line emitters without continuum

In parallel to the extraction of continuum-selected objects, we also searched for sources detected only by their line emission. Two approaches were used: a visual inspection of the MUSE data cube, and a systematic search using automatic detection tools.

Two of the authors (JR, TC) visually explored the data cube over its full wavelength range in search of sources appearing only in a narrow wavelength range, typically 4–5 wavelength planes ($\sim 6\text{--}7 \text{ \AA}$), and seemingly extended over a number of pixels at least the size of the seeing disk. We then carefully inspected the extracted line profile around this region to assess the reality of the line.

Any visual inspection has obvious limits and we also employed more automatic tools for identifying sources dominated by emission lines. One such tool is based on SExtractor (Bertin & Arnouts 1996) which was run on narrow-band images produced by averaging each wavelength plane of the cube with the 4 closest wavelength planes, adopting a weighting scheme which follows the profile on an emission line with a velocity $\sigma = 100 \text{ km s}^{-1}$. This procedure was performed across the full

wavelength range of the cube to enhance the detection of single emission lines (see also [Richard et al. 2015](#)). All SExtractor catalogues obtained from each narrow-band image were merged and compared with the continuum estimation from the white light image to select emission lines.

A different approach was used in the LSDCat software (Herenz, in prep.), which was specifically designed to search for line emitters not associated with continuum sources in the MUSE data cube. The algorithm is based on matched filtering (e.g. [Das 1991](#)): by cross-correlating the data cube with a template that resembles the expected 3D-signal of an emission line, the S/N of a faint emission line is maximized. The optimal template for the search of compact emission line objects is a function that resembles the seeing PSF in the spatial domain and a general emission line shape in the spectral domain. In practice we use a 3D template that is a combination of a 2D Moffat profile with a 1D Gaussian spectral line. The 2D Moffat parameters is taken from the bright star fit (see Fig. 2 in Sect. 3) and the FWHM of the Gaussian is fixed to 300 km s^{-1} in velocity space. To remove continuum signal, we median-filter the data cube in the spectral direction and subtract this cube from the original cube. In the following cross-correlation operation the variances are propagated accordingly, and the final result is a data cube that contains a formal detection significance for the template in each cube element (voxel). Thresholding is performed on this data cube, where regions with neighbouring voxels above the detection threshold are counted as one object and a catalogue of positions (x, y, λ) of those detections is created. To limit the number of false detections due to unaccounted systematics from sky-subtraction residuals in the redder part of the data cube, a detection threshold of 10σ was used. The candidate sources were then visually inspected by 3 authors (CH, JK & JB). This process results in the addition of 6 new identifications that had escaped the previous inspections.

All the catalogues of MUSE line emitters described above were cross-correlated with the [Casertano et al. \(2000\)](#) catalogue of continuum sources presented in Sect. 4.1. A few of the emission lines were associated with continuum sources based on their projected distance in the plane of the sky. Isolated emission lines not associated with HST continuum sources were treated as separate entries in the final catalogue and their spectra were extracted blindly at the locations of the line emission. The spectral extraction procedure was identical to the one described in Sect. 4.1.

The very large majority of line emitters not associated with HST continuum detections show a clear isolated line with an asymmetric profile, which we associate with $\text{Ly}\alpha$ emission. In most case this is corroborated by the absence of other strong lines (except possibly C III] emission at $2.9 < z < 4$) and the absence of a resolved doublet (which would be expected in case of $[\text{O II}]$ emission).

4.3. Line flux measurements

We measure emission line fluxes in the spectra using the platefit code described by [Tremonti et al. \(2004\)](#) and [Brinchmann et al. \(2004\)](#) and used for the MPA-JHU catalogue of galaxy parameters from the SDSS³. This fits the stellar spectrum using a non-negative least-squares combination of theoretical spectra broadened to match the convolution of the velocity dispersion and the instrumental resolution. It then fits Gaussian profiles to emission lines in the residual spectrum. Because of

its asymmetric shape, this is clearly not optimal for $\text{Ly}\alpha$ and we describe a more rigorous line flux measurement for this line in Sect. 6.3 below.

The S/N in most spectra is insufficient for a good determination of the stellar velocity dispersion, so for the majority of galaxies we have assumed a fixed intrinsic velocity dispersion of 80 km s^{-1} . This resulted in good fits to the continuum spectrum for most galaxies and changing this to 250 km s^{-1} changes forbidden line fluxes by less than 2%, while for Balmer lines the effect is $< 5\%$ for those galaxies for which we cannot measure a velocity dispersion. These changes are always smaller than the formal flux uncertainty and we do not consider these further here.

The emission lines are fit jointly with a single width in velocity space and a single velocity offset relative to the continuum redshift. Both the $[\text{O II}]3726, 3729$ and C III]1906, 1908 doublets are fit separately so the line ratios can be used to determine electron density. We postpone this calculation for future work.

4.4. Comparison between MUSE and published spectroscopic redshifts

Several studies have provided spectroscopic redshifts for sources in the HDFs and its flanking fields:

- [Sawicki & Mallén-Ornelas \(2003\)](#) presented spectroscopic redshifts for 97 $z < 1$ galaxies with FORS2 at the VLT. Their initial galaxy sample was selected based on photometric redshifts, $z_{\text{phot}} \lesssim 0.9$ and their resulting catalogue is biased towards $z \sim 0.5$ galaxies. The spectral resolution ($R \sim 2500\text{--}3500$) was sufficient to resolve the $[\text{O II}]3727$ doublet, enabling a secure spectroscopic redshift determination. The typical accuracy that they quote for their spectroscopic redshifts is $\delta z = 0.0003$.
- [Rigopoulou et al. \(2005\)](#) followed up 100 galaxies with FORS1 on the VLT, and measured accurate redshifts for 50 objects. Redshifts were determined based on emission lines (usually $[\text{O II}]3727$) or, in a few cases, absorption features such as the Ca II H, K lines. The redshift range of the spectroscopically-detected sample is $0.6\text{--}1.2$, with a median redshift of 1.13. These redshifts agree well with the [Sawicki & Mallén-Ornelas \(2003\)](#) estimates for sources in common between the two samples.
- [Iwata et al. \(2005\)](#) presented VLT/FORS2 spectroscopic observations of galaxies at $z \sim 3$; these were selected to have $2.5 < z_{\text{phot}} < 4$ based on HST/WFPC2 photometry combined with deep near-infrared images obtained with ISAAC at the VLT by [Labbé et al. \(2003\)](#). They firmly identified five new redshifts as well as two additional tentative redshifts of $z \sim 3$ galaxies.
- [Glazebrook et al. \(2006\)](#) produced 53 additional extragalactic redshifts in the range $0 < z < 1.4$ with the AAT Low Dispersion Survey Spectrograph by targeting 200 objects with $R > 23$.
- Finally [Wuyts et al. \(2009\)](#) used a variety of optical spectrographs on 8–10 m class telescopes (LRIS and DEIMOS at Keck Telescope, FORS2 at the VLT and GMOS at Gemini South) to measure redshifts for 64 optically faint distant red galaxies.

In total these long term efforts have provided a few hundred spectroscopic redshifts. They are, however, distributed over a much larger area than the proper HDFs deep imaging WFPC2 field, which has only ~ 88 confirmed spectroscopic redshifts. In the MUSE field itself, which covers 20% of the WFPC2 area,

³ <http://www.mpa-garching.mpg.de/SDSS>




Fig. 6. Comparison between the MUSE (red) and FORS1 spectra of Rigopoulou et al. (2005; blue) for the galaxy ID#13, a strong [O II] emitter at $z_{\text{MUSE}} = 1.2902$. The strongest spectral features are indicated in black. The grey lines show the position of the sky lines. *Upper panel:* entire spectra; *lower panel:* zoom on the rest-frame 2300–2800 Å region, which contains strong MgII and FeII absorption features.

we found 18 sources in common (see Table A.3). As shown in Fig. 16, most of the 18 sources cover the bright part ($I_{814} < 24$) of the MUSE redshift-magnitude distribution.

Generally speaking there is an excellent agreement between the different redshift estimates over the entire redshift range covered by the 18 sources. Only one major discrepancy is detected: ID#2 (HDFS J223258.30-603351.7), which Glazebrook et al. (2006) estimated to be at redshift 0.7063 while MUSE reveals it to be a star. These authors gave however a low confidence grade of $\sim 50\%$ to their identification. After excluding this object, the agreement is indeed excellent with a normalized median difference of $\langle \Delta z / (1+z) \rangle = 0.00007$ between MUSE and the literature estimates.

For two of the galaxies in common with other studies, ID#13 (HDFS J223252.16-603323.9) and ID#43 (HDFS J223252.03-603342.6), the actual spectra have been published together with the estimated redshifts. This enables us to perform a detailed comparison between these published spectra and our MUSE spectra (integrated over the entire galaxy). This comparison is shown in Figs. 6 and 7 for ID#13 (Rigopoulou et al. 2005) and ID#43 (Iwata et al. 2005), respectively.

The galaxy ID#13 is a strong [O II]3727 emitter at $z_{\text{MUSE}} = 1.2902$. In the top panel of Fig. 6 its full FORS1 and MUSE spectra are shown in blue and red, respectively. The FORS1 spectrum covers only the rest-frame $\lambda < 3500$ Å wavelength region, which does not include the [O II] feature. The lower panel of the figure shows a zoom on the rest-frame ~ 2300 –2800 Å window, which includes several strong Fe and Mg absorption features. The MUSE spectrum resolves the FeII and MgII doublets well; it is also clear that some strong features in the FORS1 spectrum, e.g., at ~ 2465 Å, 2510 Å, 2750 Å and 2780 Å are not seen in the high S/N MUSE spectrum.

The galaxy ID#43 is a Ly α emitter at $z_{\text{MUSE}} = 3.2925$. Figure 7 shows the MUSE spectrum in red and the FORS2 spectrum of Iwata et al. (2005) in blue. The middle and bottom panels show zooms on the Ly α emission line and the rest-frame ~ 1380 –1550 Å Si absorption features, respectively. The higher S/N and resolution of the MUSE spectrum opens the




Fig. 7. Comparison between the MUSE (red) and FORS2 spectra of Iwata et al. (2005; blue) for the galaxy ID#43, a strong Ly α emitter at $z_{\text{MUSE}} = 3.2925$. The strongest spectral features are indicated in black. The grey lines show the position of the sky lines; grey areas show wavelength regions for which no FORS2 spectrum was published. *Upper panel:* entire spectra; *middle and lower panels:* zoom on the Ly α and 1380–1550 Å region, respectively; the latter contains strong SiIV absorption features.

way to quantitative astrophysical studies of this galaxy, and generally of the early phases of galaxy evolution that precede the $z \sim 2$ peak in the cosmic star-formation history of the Universe.

4.5. Comparison between MUSE and published photometric redshifts

Our analysis of the HDFS allows for a quantitative comparison with photometric redshifts from the literature. We make a first comparison to the photometric redshift catalogue of Labbé et al. (2003) who used the FIRES survey to complement existing HST imaging with J , H , and K_s band data reaching $K_{s,AB}^{\text{tot}} \leq 26$. We find 89 objects in common between the two catalogues, including 8 stars. The comparison is given in Fig. 8. Considering the 81 non-stellar objects, we quantify the agreement between the MUSE spectroscopic redshifts and the 7-band photometric redshifts of Labbé et al. (2003) by calculating σ_{NMAD} (Eq. (1) of Brammer et al. 2008). This gives the median absolute deviation of Δz , and quantifies the number of “catastrophic outliers”, defined as those objects with $|\Delta z| > 5\sigma_{\text{NMAD}}$.

$$\sigma_{\text{NMAD}} = 1.48 \times \text{median} \left(\left| \frac{\Delta z - \text{median}(\Delta z)}{1 + z_{\text{sp}}} \right| \right) \quad (1)$$

where $\Delta z = (z_{\text{sp}} - z_{\text{ph}})$.

We find $\sigma_{\text{NMAD}} = 0.072$ with 6 catastrophic outliers, equating to 7.4 percent of the sample. Excluding outliers and




Fig. 8. Comparison of MUSE spectroscopic redshifts with the photometric redshifts of Labbé et al. (2003). *Upper panel:* distribution of Δz as a function of MUSE z_{sp} with outliers highlighted in red. The error bars show the uncertainties reported by Labbé et al. The grey shaded area depicts the region outside of which objects are considered outliers. *Lower panel:* direct comparison of MUSE z_{sp} and Labbé et al. (2003) z_{ph} with outliers again highlighted in red.

recomputing results in $\sigma_{\text{NMAD}} = 0.064$ reduces the number of catastrophic outliers to 2 objects, 2.7 percent of the remaining 75 sources.

Of the 6 outlying objects, 5 are robustly identified as [O II] emitters in our catalogue, but the photometric redshifts put all 5 of these objects at very low redshift, most likely due to template mismatch in the SED fitting. The final object's spectroscopic redshift is $z = 0.83$, identified through absorption features, but the photometric redshift places this object at $z = 4.82$. This is not a concern, as the object exhibits a large asymmetric error on the photometric redshift bringing it into agreement with z_{sp} . As noted in Labbé et al. (2003), this often indicates a secondary solution to the SED-fitting with comparable probability to the primary solution, at a very different redshift.

The advantage of a blind spectroscopic survey such as ours is highlighted when considering the reliability of photometric redshifts for the faint emission-line objects we detect in abundance here. Figure 9 shows values of $\Delta z/(1+z)$ for objects in our catalogue with an HST detection in the *F814W* filter. For galaxies with magnitude below $I_{814} = 24$, the measured scatter (rms) 3.7%, is comparable to what is usually measured (e.g. Saracco et al. 2006; Chen et al. 1998). However, at fainter magnitudes we see an increase with a measured scatter of 11% (rms) for galaxies in the 24–27 I_{814} magnitude range, making the photometric redshift less reliable and demonstrating the importance of getting spectroscopic redshifts for faint sources.

4.6. Spectrophotometric accuracy

The spectral range of MUSE coincides almost perfectly with the union of the two HST/WFPC2 filters *F606W* and *F814W*. It is therefore possible to synthesize broad-band magnitudes in these two bands directly from the extracted spectra, without any extrapolation or colour terms. In principle, a comparison between synthetic MUSE magnitudes with those measured in the HST images, as provided by Casertano et al. (2000), should give a straightforward check of the overall fidelity of the spectrophotometric calibration. In practice such a comparison is




Fig. 9. Relation between HST I_{814} magnitude and the scatter in $\Delta z = (z_{\text{phot}} - z_{\text{spec}})/(1 + z_{\text{spec}})$. An increase in scatter is seen towards fainter I_{814} magnitudes, highlighting the importance of spectroscopic redshifts for emission-line objects with very faint continuum magnitudes.




Fig. 10. Differences between broad-band magnitudes synthesized from extracted MUSE spectra and filter magnitudes measured by HST; *top:* *F606W*, *bottom:* *F814W*. The different symbols represent different object types: blue filled circle denote stars and red open circle stand for galaxies with $FWHM < 0\prime.4$ (in the HST images).

complicated by the non-negligible degree of blending and other aperture effects, especially for extended sources but also for the several cases of multiple HST objects falling into one MUSE seeing disk. We have therefore restricted the comparison to stars and compact galaxies with spatial $FWHM < 0\prime.4$ (as listed by Casertano et al. 2000). We also remeasured the photometry in the HST data after convolving the images to MUSE resolution, to consistently account for object crowding.

The outcome of this comparison is shown in Fig. 10, for both filter bands. Considering only the 8 spectroscopically confirmed stars in the field (blue dots in Fig. 10), all of which are relatively bright and isolated, Star ID#0 is partly saturated in the HST images and consequently appears 1 mag brighter in MUSE than in HST. For the remaining 7 stars, the mean magnitude differences (MUSE – HST) are +0.05 mag in both bands, with a formal statistical uncertainty of ± 0.04 mag. The compact galaxies (red dots in Fig. 10) are much fainter on average, and the MUSE measurement error at magnitudes around 28 or fainter




Fig. 11. Location of sources with secure redshifts in the HDFs MUSE field. In grey the WFPC2 *F814W* image. The object categories are identified with the following colours and symbols: blue: stars, cyan: nearby objects with $z < 0.3$, green: [O II] emitters, yellow: objects identified solely with absorption lines, magenta: C III] emitters, orange: AGN, red circles: Ly α emitters with HST counterpart, red triangles: Ly α emitters without HST counterpart. Objects which are spatially extended in MUSE are represented by a symbol with a size proportional to the number of spatially resolved elements.

is probably dominated by flat fielding and background subtraction uncertainties. Nevertheless, the overall flux scales are again consistent. We conclude that the spectrophotometric calibration provides a flux scale for the MUSE data cube that is fully consistent with external space based photometry, at least for sources with V_{606} brighter than 28 mag.

5. Census of the MUSE HDFs field

Given the data volume, its 3D information content and the number of objects found, it would be prohibitive to show all sources in this paper. Instead, detailed information content for all objects will be made public as described in Appendix B. In this section we carry out a first census of the MUSE data cube with a few illustrations on a limited number of representative objects.

A total of 189 objects in the data cube have a securely determined redshift. It is a rich content with 8 stars and 181 galaxies of various categories. Table 1 and Fig. 11 give a global view of the sources in the field. The various categories of objects are described in the following subsections.

Table 1. Census of the objects in the MUSE HDFs field sorted by categories.

Category	Count	z range	I_{814} range
Stars	8	0	18.6–23.9
Nearby galaxies	7	0.12–0.28	21.2–25.9
[O II] emitters	61	0.29–1.48	21.5–28.5
Absorption lines galaxies	10	0.83–3.90	24.9–26.2
AGN	2	1.28	22.6–23.6
C III] emitters	12	1.57–2.67	24.6–27.2
Ly α emitters	89	2.96–6.28	24.5–30+

5.1. Stars

We obtained spectra for 8 stars in our field. Seven were previously identified by Kilic et al. (2005) from proper motion measurements of point sources in the HDFs field. Among these stars we confirm that HDFs 1444 (ID#18) is a white dwarf. We also identify one additional M star (ID#31) that was not identified by Kilic et al. (2005).




Fig. 12. ID#160 is a $z = 1.28$ [O II] emitter with a faint ($I_{814} \sim 26.7$) HST counterpart. The HST images in the *F606W* and *F814W* filters are shown at the *top left*, the MUSE reconstructed white-light and [O II] continuum subtracted narrow band images at the *top right*. The one arcsec radius red circles show the object location derived from the HST image. At the *bottom left*, the full spectrum (in blue), smoothed with a 4 \AA boxcar, and its 3σ error (in grey) are displayed. A zoom of the unsmoothed spectrum, centred around the [O II]3726, 3729 emission lines, is also shown at the bottom right.

5.2. Nearby galaxies

In the following we refer to objects whose [O II] emission line is redshifted below the 4800 \AA blue cut-off of MUSE as nearby galaxies; that is all galaxies with $z < 0.29$. Only 7 galaxies fall into this category. Except for a bright (ID#1 $V_{606} = 21.7$) and a fainter (ID#26 $V_{606} = 24.1$) edge-on disk galaxy, the 5 remaining objects are faint compact dwarfs ($V_{606} \sim 25\text{--}26$).

5.3. [O II] emitters

A large fraction of identified galaxies have [O II]3726, 3729 in emission and we will refer to these as [O II]-emitters in the following even if this is not the strongest line in the spectrum. In Fig. 12 we show an example of a faint [O II] emitter at $z = 1.28$ (ID#160). In the HST image it is a compact source with a $26.7 V_{606}$ and I_{814} magnitude.

The average equivalent width of [O II]3727 is 40 \AA in galaxies spanning a wide range in luminosity from dwarfs with $M_B \approx -14$ to the brightest galaxy at $M_B \approx -21.4$, and sizes from marginally resolved to the largest (ID#4) with an extent of $0.9''$ in the HST image.

It is also noticeable that the [O II]-emitters often show significant Balmer absorption. In the $D4000_N\text{--}H\delta_A$ diagram they fall in the region of star forming and post-starburst galaxies. This frequent strong Balmer absorption does fit with previous results from the GDDS (Le Borgne et al. 2007) and VVDS (Wild et al. 2009) surveys.

5.4. Absorption line galaxies

For 10 galaxies, ranging from $z = 0.83$ to $z = 3.9$, the redshift determination has been done only on the basis of absorption lines. This can be rather challenging for faint sources because establishing the reality of an absorption feature is more difficult than for an emission line. For that reason the faintest source with a secure absorption line redshift has $I_{814} = 26.2$ and

$z = 3.9$, while the faintest source (ID#83) with absorption redshifts in the $1.5 < z < 2.9$ so-called MUSE “redshift desert” (Steidel et al. 2004) has I_{814} of 25.6.

A notable pair of objects is ID#50 and ID#55 which is a merger at $z = 2.67$ with a possible third companion based on the HST image, which can not be separated in the MUSE data. And while not a pure absorption line galaxy, as it does have [O II]3727 in emission, object ID#13 shows very strong Mg II and Fe II absorption lines.

5.5. C III] emitters

At $1.5 < z < 3$, well into the “redshift desert”, the main emission line identified is C III]1907, 1909, which is typically resolved as a doublet of emission lines at the resolution achieved with MUSE. Among the clear C III] emitters identified, the most interesting one (ID#97) is displayed in Fig. 13. It is a $z = 1.57$ galaxy with strong C III]1907, 1909 and Mg II 2796, 2803 emission lines. It appears as a compact source in the HST images with $V_{606} = 26.6$ and $I_{814} = 25.8$. The object is unusually bright in C III] with a total flux of $2.7 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$ and a rest-frame equivalent width of 16 \AA . These are relatively rare objects, with only 17 found in our field of view, but such C III] emission is expected to appear for younger and lower mass galaxies, typically showing a high ionization parameter (Stark et al. 2014).

5.6. Ly α emitters

The large majority of sources at $z > 3$ are identified through their strong Lyman- α emission line. Interestingly, 26 of the discovered Ly α emitters are below the HST detection limit, i.e. $V_{606} > 29.6$ and $I_{814} > 29$ (3σ depth in a 0.2 arcsec^2 aperture, Casertano et al. 2000). Approximately 60% of them have a high S/N and exhibit the typical asymmetric Ly α profile. We produced a stacked image in the *WFPC2-F814W* filter of these 26 Ly α emitters not individually detected in HST and measured an average continuum at the level of $I_{814} = 29.8 \pm 0.2 \text{ AB}$




Fig. 13. ID#97 is a $z = 1.57$ strong C III] emitter. The HST images in *F606W* and *F814W* filters are shown at *the top left*, the MUSE reconstructed white-light, the C III] and Mg II continuum subtracted narrow band images at *the top right*. The one arcsec radius red circles show the object location derived from the HST image. At *the bottom left*, the full spectrum (in blue), smoothed with a 4 \AA boxcar, and its 3σ error (in grey) are displayed. A zoom of the unsmoothed spectrum, centred around the C III] 1907, 1909 \AA and Mg II 2796, 2803 \AA emission lines, are also shown at *the bottom right*.




Fig. 14. ID#553 is a $z = 5.08$ Ly α emitter without HST counterpart. The HST images in *F606W* and *F814W* filters are shown at *the top left*, the MUSE reconstructed white-light and Ly α narrow band images at *the top right*. The one arcsec radius red circles show the emission line location. The spectrum is displayed on the *bottom figures*; including a zoom at the emission line. At *the bottom left*, the full spectrum (in blue), smoothed with a 4 \AA boxcar, and its 3σ error (in grey) are displayed. A zoom of the unsmoothed spectrum, centred around the Ly α emission line, is also shown at *the bottom right*.

(Drake et al., in prep.). We present in Fig. 14 one such example, ID#553 in the catalogue. With a total Ly α flux of $4.2 \times 10^{-18} \text{ erg s}^{-1} \text{ cm}^{-2}$ the object is one of the brightest of its category. It is also unambiguously detected in the reconstructed Ly α narrow band image. With such a low continuum flux the emission corresponds to a rest-frame equivalent width higher than 130 \AA .

Note that we have found several even fainter line emitters that have no HST counterpart. However, because of their low S/N, it is difficult to firmly identify the emission line and they have therefore been discarded from the final catalogue.

5.7. Active galactic nuclei

Among the [O II] emitting galaxies we identify two objects (ID#10 and ID#25) that show significant [Ne V] 3426 emission, a strong signature of nuclear activity. Both galaxies show pronounced Balmer breaks and post-starburst characteristics, and their forbidden emission lines are relatively broad with a FWHM $\sim 230 \text{ km s}^{-1}$. There is, however, no clear evidence for broader permitted lines such as Mg II 2798, thus both objects are probably type 2 AGN. Both objects belong to the same group of galaxies at $z \approx 1.284$ (Sect. 6.2).




Fig. 15. Example of spatially overlapping objects: a $z = 3.09$ Ly α emitter (ID#71) with a $z = 0.83$ [O II] emitter (ID#72). The HST images in *F606W* and *F814W* filters are shown at the top left, the MUSE Ly α and [O II] images at the top right. The one arcsec radius red circles show the object location derived from the HST image. At the bottom left, the full spectrum (in blue), smoothed with a 4 Å boxcar, and its 3σ error (in grey) are displayed. A zoom of the unsmoothed spectrum, centred around the Ly α and [O II]3727 emission line, is also shown at the bottom right.

Object ID#144 was classified by [Kilic et al. \(2005\)](#) as a probable QSO at $z = 4.0$ on the basis of its stellar appearance and its *UBVI* broad band colours. The very strong Ly α line of this objects confirms the redshift ($z = 4.017$), but as the line is relatively narrow (~ 100 km s $^{-1}$) and no other typical QSO emission lines are detected, a definite spectroscopic classification as an AGN is not possible.

5.8. Spatially resolved galaxies

Twenty spatially resolved galaxies up to $z \sim 1.3$ are identified in the MUSE data cube (see Fig. 11). We consider a galaxy as resolved if it extends over a minimum area of twice the PSF. To compute this area we performed emission line fitting (see Sect. 7) on a list of 33 galaxies that had previously been identified to be extended in the HST images. Flux maps were built for each fitted emission line and we computed the galaxy size (FWHM of a 2D fitted Gaussian) using the brightest one (usually [O II]). Among these 20 resolved galaxies, 3 are at low redshift ($z \leq 0.3$) and 4 are above $z \sim 1$. Note that 5 of the resolved galaxies are in the group identified at ~ 0.56 (see Sect. 6.2) including ID#3 which extends over ~ 5 times the MUSE PSF.

5.9. Overlapping objects

While searching for sources in the MUSE data cube, we encountered a number of spatially overlapping objects. In many cases a combination of high spatial resolution HST images and MUSE narrow-band images has been sufficient to assign spectral features and redshifts to a specific galaxy in the HST image. But in some cases, the sources cannot be disentangled, even at the HST resolution. This is illustrated in Fig. 15, where the HST image shows only one object but MUSE reveals it to be the result of two galaxies that are almost perfectly aligned along the line of sight: an [O II] emitter at $z = 0.83$ and a $z = 3.09$ Ly α emitter. There are other cases of objects that potentially could be identified as mergers on the basis of the HST images but are

in fact just two galaxies at different redshifts. The power of the 3D information provided by MUSE is nicely demonstrated by these examples.

6. Redshift distribution and global properties

6.1. Redshift distribution

We have been able to measure a redshift at confidence ≥ 1 for 28% of the 586 sources reported in HST catalogue of [Casertano et al. \(2000\)](#) in the MUSE field. The redshift distribution is presented in Fig. 16. We reach 50% completeness with respect to the HST catalogue at $I_{814} = 26$. At fainter magnitudes the completeness decreases, but it is still around 20% at $I_{814} = 28$. In addition to the sources identified in the HST images, we found 26 Ly α emitters, i.e. 30% of the entire Ly α emitter sample, that have no HST counterparts and thus have $I_{814} > 29.5$.

Redshifts are distributed over the full $z = 0-6.3$ range. Note the decrease in the $z = 1.5-2.8$ window – the well known redshift desert – corresponding to the wavelengths where [O II] is beyond the 9300 Å red limit of MUSE and Ly α is bluer than the 4800 Å blue cut-off of MUSE.

Although the MUSE HDFS field is only a single pointing, and thus prone to cosmic variance, one can compare the measured redshift distribution with those from other deep spectroscopic surveys (e.g. *z*COSMOS-Deep – [Lilly et al. 2007](#); [Lilly et al. 2009](#), VVDS-Deep – [Le Fèvre et al. 2013](#), VUDS – [Le Fèvre et al. 2015](#)). The latter, with 10 000 galaxies in the $z \sim 2-6$ range, is the most complete.

We show in Fig. 17 the MUSE-HDFS and VUDS normalized redshift distributions. They look quite different. This was expected given the very different observational strategy: the VUDS redshift distribution is the result of a photometric redshift selection $z_{\text{phot}} > 2.3 \pm 1\sigma$ (with first and second peaks of PDF) combined with continuum selection $I_{\text{AB}} < 25$, while MUSE does not make any pre-selection. With 22% of galaxies at $z > 4$ in contrast to 6% for the VUDS, MUSE demonstrates a higher efficiency for finding high redshift galaxies.




Fig. 16. Redshift distribution of sources in the HDF5 MUSE field. *Left:* redshift versus HST I_{814AB} magnitude. The symbol size is proportional to the object's HST size. The green circles are for the published spectroscopic redshifts and the brown symbols for the new MUSE measurements. The red arrows show objects without HST counterparts. The dashed line shows the 3σ detection limit for the HST $F814W$ images. *Right:* the grey histogram shows the distribution of all magnitudes in the HST catalogue while the light blue histogram shows the magnitude distribution of those galaxies with confirmed redshifts and HST counterparts. The last histogram bar centred at $I_{814} = 29.75$ is for the $\text{Ly}\alpha$ emitters not detected in the HST image. The blue curve gives the completeness of the redshift identification with respect to the identified sources in the HST images.




Fig. 17. Comparison of the HDF5-MUSE normalized redshift distribution (*bottom*) with the one from the VUDS (*top*).

The number density of observed galaxies is also very different between the two types of observations. With 6000 galaxies per square degree, the VUDS is the survey which achieved the

highest density with respect to the other spectroscopic surveys (see Fig. 30 of [Le Fevre et al. 2015](#)). This translates, however, to only 1.7 galaxies per arcmin², which is two orders of magnitude smaller than the number density achieved by MUSE in the HDF5.

6.2. Galaxy groups

The high number density of accurate spectroscopic redshifts allows us to search for galaxy groups in the field. We applied a classical friend-of-friend algorithm to identify galaxy groups, considering only secure redshifts in our catalogue, i.e. those with confidence index ≥ 1 . We adopted a maximum linking length Δr of 500 kpc in projected distance and $\Delta v = 700 \text{ km s}^{-1}$ in velocity following the z Cosmos high redshift group search by [Diener et al. \(2013\)](#). The projected distance criterium has very little influence since our field-of-view of $1'$ on a side corresponds to ~ 500 kpc at most. This results in the detection of 17 groups with more than three members. These are listed in Table 2 together with their richness, redshift, and nominal rms size and velocity dispersion as defined in [Diener et al. \(2013\)](#).

Among the 181 galaxies with a secure redshift in our catalogue, 43% reside in a group, 29% in a pair and 28% are isolated. The densest structure we find lies at $z = 1.284$ and has nine members, including 2 AGN and an interacting system showing a tidal tail, all concentrated to the north west of the field. The structure at $z \sim 0.578$ first spotted by [Vanzella et al. \(2002\)](#) and further identified as a rich cluster by [Glazebrook et al. \(2006\)](#) shows up here as a 5-member group. The richer group (6 members) at slightly lower redshift ($z = 0.565$) is also within the redshift range 0.56–0.60 considered by [Glazebrook et al. \(2006\)](#)

Table 2. Galaxy groups detected in the HDFs ordered by redshift.

z	v_{rms} km s $^{-1}$	r_{rms} kpc	N_{m}	Member IDs
0.172	65	43	3	1, 63, 70
0.421	262	54	4	6, 57, 101, 569
0.564	52	142	7	3, 4, 9, 23, 32, 135
0.578	424	150	5	5, 8, 11, 17, 122
0.972	56	201	3	24, 68, 129
1.284	354	92	9	10, 13, 15, 25, 27, 35, 64, 114, 160
2.672	101	87	4	50, 51, 55, 87
3.013	350	115	3	40, 56, 155
3.124	329	92	4	422, 437, 452, 558
3.278	36	144	4	162, 202, 449, 513
3.349	35	90	3	139, 200, 503
3.471	324	139	4	433, 469, 478, 520
3.823	161	93	4	238, 514, 563, 581
4.017	113	181	4	89, 144, 216, 308
4.699	430	109	6	325, 441, 453, 474, 499, 548
4.909	370	164	6	186, 218, 334, 338, 484, 583
5.710	26	101	3	546, 547, 574

and could be part of the same large scale structure. The highest redshift group identified is also the one with the lowest v_{rms} of all groups: three Ly- α emitters at $z = 5.71$ within 26 km s $^{-1}$. We also find two relatively rich groups both including six members at redshift ~ 4.7 and ~ 4.9 .

6.3. Number counts of emission line galaxies

With more than 100 emission line galaxies identified by MUSE in this field, a meaningful comparison of the observed number counts with expectations from the literature is possible. We consider Ly α and [O II]3727 emitters.

While integrated emission line fluxes can, in principle, be measured easily from the fits to the extracted spectra (see Appendix A), the significant level of crowding among faint sources resulted in extraction masks that were often too small for capturing the total fluxes. This is especially relevant for Ly α emitters which in many cases show evidence for extended line emission. In order to derive robust total fluxes of [O II] and Ly α emitters we therefore adopted a more manual approach. We first produced a ‘‘pure emission line cube’’ by median-filtering the data in spectral direction and subtracted the filtered cube. We then constructed narrow-band images centred on each line, which were now empty of sources except for the emitters of interest. Finally, total line fluxes were determined by a growth curve analysis in circular apertures around each source. We adjusted the local sky background for each line such that the growth curve became flat within $\sim 2''$ from the centroid, and integrated the emission line flux inside that aperture. Avoiding the edges of the field, this procedure yielded subsamples of 74 Ly α and 41 [O II] emitters, with well-measured emission line fluxes.

The resulting number counts of the two object classes, expressed as the cumulative number of objects per arcmin 2 brighter than a given flux, are depicted by the blue step functions in Fig. 18. Both panels show a relatively steep increase at bright fluxes which then flattens and eventually approaches a constant number density for line fluxes of $f \lesssim 2 \times 10^{-18}$ erg s $^{-1}$ cm $^{-2}$, as no fainter objects are added to the sample. The detailed shapes of these curves depend on the emission line luminosity function and its evolution with redshift, but also on the selection functions which are different for the present samples of Ly α and




Fig. 18. Cumulative number counts of emission line galaxies in the HDFs, as a function of line flux. *Top panel:* [O II] emitters; *bottom panel:* Ly α emitters. The green lines show the predictions for the relevant redshift ranges ($0.288 < z < 1.495$ and $2.95 < z < 6.65$, respectively), using a compilation of published luminosity functions as described in the text.

[O II] emitters. A detailed discussion of this topic is beyond the scope of this paper. We note that at 5×10^{-18} erg s $^{-1}$ cm $^{-2}$, the numbers of detected Ly α and [O II] emitters are equal; at fainter flux levels, Ly α emitters are more numerous than [O II] emitters.

Contrasting the MUSE results for the HDFs to expectations from the literature is not straightforward, as the relevant surveys have very diverse redshift coverages. In the course of MUSE science preparations, we combined several published results into analytic luminosity functions for both Ly α and [O II] emitters, which we now compare with the actual measurements. For the statistics of [O II] emitters we used data by Ly et al. (2007), Takahashi et al. (2007), and Rauch et al. (2008), plus some guidance from the predictions for H α emitters by Geach et al. (2010). For the Ly α emitters we adopted a luminosity function with non-evolving parameters following Ouchi et al. (2008), with luminosity function parameters estimated by combining the Ouchi et al. (2008) results with those of Gronwall et al. (2007) and again Rauch et al. (2008). Note that these luminosity functions are just intended to be an overall summary of existing observations, and no attempt was made to account for possible tensions or inconsistencies between the different data sets.

Figure 18 shows the predicted cumulative number counts based on these prescriptions as solid green curves (dotted where we consider these predictions as extrapolations). While the overall match appears highly satisfactory, cosmic variance is of course a strong effect in a field of this size, particularly at the bright end. We reiterate that any more detailed comparison with published luminosity functions would be premature. Nevertheless, it is reassuring to see that the present MUSE source catalogue for the HDFs gives number counts consistent with previous work.




Fig. 19. Morpho-kinematics of HDF5140 (*left*) and HDF54070 (*right*). For each galaxy and from left to right. *Top*: HST/WFPC2 $F814W$ image in log scale (the yellow rectangle shows the GIRAFFE FoV), the MUSE [O III] $\lambda 5007$ flux map, and the deconvolved [O III] $\lambda 5007$ flux map from GalPaK3D (for HDF5140 only). *Middle*: MUSE observed velocity field from H β and [O III], velocity field of the 2D rotating disk model, residual velocity field, and deconvolved [O III] $\lambda 5007$ velocity field from GalPaK3D (for HDF5140 only). *Bottom*: MUSE observed velocity dispersion map from H β and [O III], velocity dispersion map deduced from the 2D velocity field model (beam-smearing effect and spectral PSF), deconvolved velocity dispersion map, and deconvolved [O III] $\lambda 5007$ velocity dispersion from GalPaK3D (for HDF5140 only). In each map, north is up and east is left. The centre used for kinematical modelling is indicated as a white cross, the position angle is indicated by the black line which ends at the effective radius. For comparison, the published velocity field and dispersion map obtained with GIRAFFE for HDF5140 (Puech et al. 2006) and HDF54070 (Flores et al. 2006) are shown in the *bottom* row.

7. Spatially resolved kinematics

To illustrate the power of MUSE for spatially-resolved studies of individual galaxies, we derive the kinematics for two galaxies with published spatially-resolved spectroscopy. These galaxies, namely ID#6 (hereafter HDF54070) at $z = 0.423$ and ID#9 (hereafter HDF5140) at $z = 0.5645$, were observed earlier with the GIRAFFE multi-IFU at the VLT as part of the IMAGES survey (Flores et al. 2006; Puech et al. 2006). These observations have a spectral resolution of $22\text{--}30\text{ km s}^{-1}$, were taken with a seeing in the $0.35\text{--}0.8''$ range and used an integration time of 8 h. These data have mainly been used to derive the ionized gas kinematics from the [O II] emission lines, which was published in Flores et al. (2006) and Puech et al. (2006) for HDF54070 and HDF5140, respectively. As shown in the lower panels of Fig. 19, the velocity fields appear very perturbed. This led those authors to conclude that these galaxies show complex gas kinematics.

With the MUSE data in hand, we have the opportunity to revisit the ionized gas kinematics for these two galaxies and to compare the derived parameters with those obtained with GIRAFFE. To probe the gas kinematics, we make use of the three brightest emission lines available in the MUSE spectral range: the [O II] doublet, H β , and [O III].

The flux and velocity maps are presented in Fig. 19. Rotating disk models, both in 2D (see Epinat et al. 2012 for details) and with GalPaK3D (Bouché et al. 2013, 2014) are also shown. Compared to the GIRAFFE maps, the kinematics revealed by MUSE give a very different picture for these galaxies.

The barred structure of HDF54070 is clearly detected in the MUSE velocity field with its typical S-shape, whereas the velocity field of HDF5140 is much more regular and typical of an early-type spiral with a prominent bulge. The maximum rotation

velocity derived for HDF5140 is consistent between the 2D and 3D models ($\sim 140\text{ km s}^{-1}$), but is significantly lower than the value ($\sim 220\text{ km s}^{-1}$) obtained from the GIRAFFE data. In HDF54070, the velocity dispersion is quite uniform over the disk whereas it is clearly peaked at the centre of HDF5140. Note, however, that there are some structures in the velocity dispersion residual map of this galaxy, with velocity dispersion of the order of $\sim 80\text{--}100\text{ km s}^{-1}$. Such a broad component, aligned along the south-west side of the minor axis of HDF5140, could well be produced by superwind-driven shocks as revealed in a significant population of star-forming galaxies at $z \sim 2$ (see e.g. Newman et al. 2012). Note that outflows were already suspected in this galaxy from the GIRAFFE data, but not for the same reasons. Puech et al. (2006) argued that the velocity gradient of HDF5140 is nearly perpendicular to its main optical axis, which is clearly not the case.

The quality of the 2D MUSE maps uniquely enables reliable modelling and interpretation of the internal physical properties of distant galaxies. A complete and extensive analysis of galaxy morpho-kinematics probed with MUSE in this HDFS field will be the scope of a follow-up paper (Contini, in prep.).

8. Summary and conclusion

The HDFS observations obtained during the last commissioning run of MUSE demonstrate its capability to perform deep field spectroscopy at a depth comparable to the HST deep photometry imaging.

In 27 h, or the equivalent of 4 nights of observations with overhead included, we have been able to get high quality spectra and to measure precise redshifts for 189 sources (8 stars and 181 galaxies) in the 1 arcmin^2 field of view of MUSE. This is to be compared with the 18 spectroscopic redshifts that had been




Fig. 20. Relative counts of MUSE HDFs Ly α emitters (in %) versus UV continuum magnitude. Note that the last bin of the histogram contains the 12 Ly α emitters with magnitude $29.25 < I_{814} < 29.75$ (taken from Casertano et al. 2000, catalogue) and the 26 MUSE Ly α emitters not detected by HST which have been arbitrarily assigned the average magnitude estimated from the stacked image $I_{814} = 29.5$ (see Sect. 5.6).

obtained before for (relatively bright) sources in the same area. Among these 181 galaxies, we found 26 Ly α emitters which were not even detected in the deep broad band WFPC2 images. The redshift distribution is different from the ones derived from deep multi-object spectroscopic surveys (e.g. Le Fevre et al. 2015) and extends to higher redshift. The galaxy redshifts revealed by MUSE extend to very low luminosities, as also shown in Fig. 20. Previous deep optical surveys like those of Stark et al. (2010) in the GOODS field did not push to such faint magnitudes. Stark et al. obtained emission line redshifts to $AB = 28.3$ and it can be seen that MUSE can be used to derive Ly α fluxes and redshifts to fainter limits. We achieved a completeness of 50% of secure redshift identification at $I_{814} = 26$ and about 20% at $I_{814} = 27-28$.

In the MUSE field of view, we have detected 17 groups with more than three members. The densest group lies at $z = 1.284$ and has nine members, including 2 AGN and an interacting system showing a tidal tail. We also find three groups of Ly α emitters at high redshifts ($z \sim 4.7, 4.9$ and 5.7).

We have also investigated the capability of MUSE for spatially resolved spectroscopy of intermediate redshift galaxies. Thanks to its excellent spatial resolution, MUSE enables reliable modelling and interpretation of the internal physical properties of distant galaxies. Although the number of objects with relevant spatial information is limited to twenty in the MUSE HDFs field, this is still a large multiplex factor compared to single object pointing observations used e.g. for the MASSIV survey (Contini et al. 2012) with SINFONI at the VLT.

The MUSE observations of the HDFs are already among the deepest spectroscopic survey ever performed in optical astronomy. It is a unique data set and can be used for a wide range of follow-up studies. We release the reduced data world-wide, and make the associated products public (see Appendix B).

The unique performance in ultra-deep spectroscopic fields achieved by MUSE is very complementary to wide field multi-object spectroscopic surveys (e.g. Le Fevre et al. 2015). The latter cover a much larger field of view and is much more efficient to probe the bright end of the continuum luminosity function while MUSE can probe the faint end of emission line luminosity function where the blind selection is a great helper. In addition MUSE is able to obtain several hundred redshifts per arcmin² in a moderate amount of time, a spatial density two orders of

magnitude higher than what is achieved by current multi-object spectroscopic surveys.

With its capability to obtain high quality spectroscopic information of the population of faint galaxies at high-redshift without having to perform any pre-selection, MUSE is also highly complementary to the present (*Hubble*, ALMAs) and future (James Webb) ultra-deep surveys. As we have demonstrated using the HDFs observations, MUSE has in addition its own large potential for discoveries. MUSE is not limited to follow-up observations, but is also able to find large number of objects that are not (or barely) detected by broad band deep imaging.

These observations performed during the commissioning mark the end of the realization of the instrument and the start of the science exploitation. This is, however, only the first step. Further improvements of the data reduction process – such as super flat-fields or optimal combination of exposures – are still possible and will be implemented in the near future. We are also developing new methods for 3D source detections, optimised for the large size of the MUSE data cubes (Paris et al. 2013; Herenz, in prep.; Meillier, in prep.; Cantalupo, in prep.). With these new tools and an improved data reduction, we expect to further extend the number of secure redshift identifications in the field. This will be the subject of a future public release for the HDFs field.

Another major milestone is expected in a few years, when the Adaptive Optics Facility will come in operations at UT4 (Arsenault et al. 2012). The four laser guide stars, the deformable VLT secondary mirror and GALACSI, the module dedicated to MUSE (Ströbele et al. 2012), will improve the spatial resolution of MUSE up to 50% without impacting its superb throughput and efficiency.

Acknowledgements. We thank the referee Michele Cirasuolo and Olivier Le Fevre for proving useful comments on the submitted version and Laure Piqueras and Gerard Zins for their support with analysis and observation software. We also warmly thank Fernando Selman and all Paranal staff for their enthusiastic support of MUSE during the commissioning runs. R.B. acknowledges support from the ERC advanced grant 339659-MUSICOS. J.R., B.C., V.P. acknowledge support from the ERC starting grant CALENDs. T.C., B.G., A.D. acknowledge support from the ANR FOGHAR. L.M. acknowledges support from the Lyon Institute of Origins under grant ANR-10-LABX-66. S.J.L. and K.S. acknowledge support from the Swiss National Science Foundation.

References

- Arsenault, R., Madec, P.-Y., Paufigue, J., et al. 2012, SPIE Conf. Ser., 8447
 Bacon, R., Accardo, M., Adjali, L., et al. 2010, SPIE Conf. Ser., 7735, 8
 Bacon, R., Vernet, J., Borosiva, E., et al. 2014, *The Messenger*, 157, 21
 Beckwith, S. V. W., Stiavelli, M., Koekemoer, A. M., et al. 2006, *AJ*, 132, 1729
 Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
 Bouché, N., Murphy, M. T., Kacprzak, G. G., et al. 2013, *Science*, 341, 50
 Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, *ApJ*, 686, 1503
 Brinchmann, J., Charlot, S., White, S. D. M., et al. 2004, *MNRAS*, 351, 1151
 Carollo, C. M., Bschorr, T. J., Renzini, A., et al. 2013, *ApJ*, 773, 112
 Casertano, S., de Mello, D., Dickinson, M., et al. 2000, *AJ*, 120, 2747
 Chen, H.-W., Fernandez-Soto, A., Lanzetta, K. M., et al. 1998
 [arXiv:astro-ph/9812339]
 Contini, T., Garilli, B., Le Fèvre, O., et al. 2012, *A&A*, 539, A91
 Das, P. K. 1991, *Optical Signal Processing – Fundamentals* (Berlin, Heidelberg, New York: Springer)
 Diener, C., Lilly, S. J., Knobel, C., et al. 2013, *ApJ*, 765, 109
 Epinat, B., Tasca, L., Amram, P., et al. 2012, *A&A*, 539, A92
 Ferguson, H. C., Dickinson, M., & Williams, R. 2000, *ARA&A*, 38, 667
 Flores, H., Hammer, F., Puech, M., Amram, P., & Balkowski, C. 2006, *A&A*, 455, 107
 Geach, J. E., Cimatti, A., Percival, W., et al. 2010, *MNRAS*, 402, 1330
 Glazebrook, K., Verma, A., Boyle, B., et al. 2006, *AJ*, 131, 2383
 Gronwall, C., Ciardullo, R., Hickey, T., et al. 2007, *ApJ*, 667, 79
 Horne, K. 1986, *PASP*, 98, 609
 Ilbert, O., Capak, P., Salvato, M., et al. 2009, *ApJ*, 690, 1236
 Ilbert, O., McCracken, H. J., Le Fèvre, O., et al. 2013, *A&A*, 556, A55

- Iwata, I., Inoue, A. K., & Burgarella, D. 2005, *A&A*, **440**, 881
- Kilic, M., Mendez, R. A., von Hippel, T., & Winget, D. E. 2005, *ApJ*, **633**, 1126
- Kausch, W., Noll, S., Smette, A., et al. 2015, *A&A*, in press
DOI: [10.1051/0004-6361/201423909](https://doi.org/10.1051/0004-6361/201423909)
- Labbé, I., Franx, M., Rudnick, G., et al. 2003, *AJ*, **125**, 1107
- Labbé, I., Huang, J., Franx, M., et al. 2005, *ApJ*, **624**, L81
- Le Borgne, J. F., Paschke, A., Vandenbroere, J., et al. 2007, *A&A*, **476**, 307
- Le Fèvre, O., Cassata, P., Cucciati, O., et al. 2013, *A&A*, **559**, A14
- Le Fevre, O., Tasca, L. A. M., Cassata, P., et al. 2015, *A&A*, in press,
DOI: [10.1051/0004-6361/201423829](https://doi.org/10.1051/0004-6361/201423829)
- Lee, B., Giavalisco, M., Williams, C. C., et al. 2013, *ApJ*, **774**, 47
- Lilly, S. J., Le Fèvre, O., Renzini, A., et al. 2007, *ApJS*, **172**, 70
- Lilly, S. J., Le Brun, V., Maier, C., et al. 2009, *ApJS*, **184**, 218
- Ly, C., Malkan, M. A., Kashikawa, N., et al. 2007, *ApJ*, **657**, 738
- Mortlock, A., Conselice, C. J., Hartley, W. G., et al. 2013, *MNRAS*, **433**, 1185
- Muzzin, A., Marchesini, D., Stefanon, M., et al. 2013, *ApJ*, **777**, 18
- Newman, S. F., Genzel, R., Förster-Schreiber, N. M., et al. 2012, *ApJ*, **761**, 43
- O'Donnell, J. E. 1994, *ApJ*, **422**, 158
- Ouchi, M., Shimasaku, K., Akiyama, M., et al. 2008, *ApJS*, **176**, 301
- Paris, S., Suleiman, R. F. R., Mary, D., & Ferrari, A. 2013, in *Acoustics, Speech and Signal Processing (ICASSP)*, IEEE Int. Conf., 3947
- Puech, M., Hammer, F., Flores, H., Östlin, G., & Marquart, T. 2006, *A&A*, **455**, 119
- Rauch, M., Haehnelt, M., Bunker, A., et al. 2008, *ApJ*, **681**, 856
- Richard, J., Patricio, V., Martinez, J., et al. 2015, *MNRAS*, **446**, L16
- Rigopoulou, D., Vacca, W. D., Berta, S., Franceschini, A., & Aussel, H. 2005, *A&A*, **440**, 61
- Saracco, P., Fiano, A., Chincarini, G., et al. 2006, *MNRAS* **367**, 349
- Sawicki, M., & Mallén-Ornelas, G. 2003, *AJ*, **126**, 1208
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, **500**, 525
- Smette, A., Sana, H., Noll, S., et al. 2015, *A&A*, in press,
DOI: [10.1051/0004-6361/201423932](https://doi.org/10.1051/0004-6361/201423932)
- Stark, D. P., Ellis, R. S., Chiu, K., Ouchi, M., & Bunker, A. 2010, *MNRAS*, **408**, 1628
- Stark, D. P., Richard, J., Charlot, S., et al. 2014, *MNRAS*, submitted
[[arXiv:1408.3649](https://arxiv.org/abs/1408.3649)]
- Steidel, C. C., Shapley, A. E., Pettini, M., et al. 2004, *ApJ*, **604**, 534
- Ströbele, S., La Penna, P., Arsenault, R., et al. 2012, *SPIE Conf. Ser.*, 8447
- Takahashi, M. I., Shioya, Y., Taniguchi, Y., et al. 2007, *ApJS*, **172**, 456
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, **613**, 898
- van der Wel, A., Franx, M., van Dokkum, P. G., et al. 2014, *ApJ*, **788**, 28
- Vanzella, E., Cristiani, S., Arnouts, S., et al. 2002, *A&A*, **396**, 847
- Weilbacher, P. M., Streicher, O., Urrutia, T., et al. 2012, *SPIE Conf. Ser.*, 8451
- Wild, V., Walcher, C. J., Johansson, P. H., et al. 2009, *MNRAS*, **395**, 144
- Williams, R. E., Blacker, B., Dickinson, M., et al. 1996, *AJ*, **112**, 1335
- Williams, R. E., Baum, S., Bergeron, L. E., et al. 2000, *AJ*, **120**, 2735
- Wuyts, S., van Dokkum, P. G., Franx, M., et al. 2009, *ApJ*, **706**, 885

Appendix A: Catalogue

Table A.1 lists the basic properties of the galaxies studied in the paper. All the sources from the Casertano et al. (2000) catalogue that fall within the field of view of the final MUSE data cube are included as well as sources without an entry in the Casertano et al. catalogue. Note that *F814W* magnitudes have been manually computed for a few objects that were missing in the Casertano et al. (2000) catalogue but were clearly present in the HST image. The table is sorted by increasing apparent *F814W* magnitude from the catalogue with no particular ordering of the sources without a catalogue magnitude.

The first column gives a running number which is the one used for ID#XX entries in the text. The subsequent two columns give the right ascension and declination from the MUSE observations, then follows the *F814* SExtractor BEST magnitude from the Casertano et al. catalogue and the *F606W*–*F814W* colour from the BEST magnitudes. The redshift and its confidence follows thereafter, with the subsequent column indicating features identified in the spectrum. The final column gives the running number of the object in the Casertano et al. catalogue.

The second-to-last column, N_{exp} , gives the median number of exposures going into the reduction of the region where the spectrum was extracted. Recall that the exposures are 30 min in

duration, so a value of 40 corresponds to an exposure time of 20 h. Our redshift catalogue is naturally less complete where $N_{\text{exp}} < 30$.

Table A.2 gives the emission line fluxes measured off the 1D spectra produced by straight summation. In this table only the 181 sources with redshift >0 with redshift confidence ≥ 1 are included. The procedure adopted for emission line flux measurements is given in Sect. 4.3. Note that the fluxes have been corrected for Galactic reddening using the Schlegel et al. (1998) dust maps using the reddening curve from O’Donnell (1994) for consistency with that work.

We emphasise that there has been no attempt to correct these fluxes to true total fluxes. In particular these fluxes are generally lower than those used in Sect. 6.3 which are aperture corrected. However the technique used in Sect. 6.3 is not suitable for all galaxies so we use the simpler approach here. Note also that the Ly α fluxes have been measured by fitting a Gaussian to an asymmetric line so are likely to give suboptimal flux measurements. Finally observe also that flux measurements are provided for all lines within the wavelength range of the spectrum, regardless of whether they were reliably detected or not.

Table A.3 gives the list of the redshift comparison between MUSE and published spectroscopic redshifts discussed in Sect. 4.4.

Table A.1. Basic data and redshifts on sources with spectra in the HDF-S.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
0	22:32:57.00	-60:34:05.72	... ^a	...	0.0000	3	A3-A4	54	10 692
1	22:32:54.68	-60:33:33.16	21.22 ± 0.002	0.49 ± 0.003	0.1723	3	H α , [O III], H β	54	1635
2	22:32:58.30	-60:33:51.66	21.36 ± 0.001	2.20 ± 0.004	0.0000	3	M5	54	10 617
3	22:32:53.74	-60:33:37.57	21.52 ± 0.004	0.58 ± 0.005	0.5637	3	[O III], H β , H γ , H δ , [Ne III], [O II]	53	1703
4	22:32:52.14	-60:33:59.56	21.78 ± 0.003	0.80 ± 0.005	0.5638	3	[O III], H β , H γ , [Ne III], [O II]	53	2141
5	22:32:52.24	-60:34:02.75	21.97 ± 0.004	0.87 ± 0.006	0.5798	3	[O III], H β , H γ , [Ne III], [O II]	53	2142
6	22:32:58.21	-60:33:31.64	21.98 ± 0.004	0.50 ± 0.006	0.4223	3	[O III], H β , H γ , [Ne III], [O II]	54	1603
7	22:32:59.43	-60:33:39.82	21.99 ± 0.004	0.56 ± 0.006	0.4637	3	[O III], H β , H γ , H δ , [Ne III], [O II]	51	1753
8	22:32:51.51	-60:33:37.57	22.08 ± 0.003	0.86 ± 0.005	0.5772	3	[O III], [O II]	35	1733
9	22:32:56.08	-60:34:14.17	22.08 ± 0.004	0.61 ± 0.006	0.5638	3	[O III], H β , H γ , H δ , [Ne III], [O II]	53	2467
10	22:32:53.03	-60:33:28.53	22.56 ± 0.004	1.13 ± 0.008	1.2840	3	[O II], Mg II	53	1539
11	22:32:52.35	-60:33:33.05	22.72 ± 0.004	0.64 ± 0.005	0.5779	3	[O III], H β , H γ , H δ , [Ne III], [O II]	53	1657
12	22:32:57.06	-60:33:23.03	22.79 ± 0.005	0.57 ± 0.007	0.6701	3	[O III], H β , H γ , H δ , [Ne III], [O II]	54	1429
13	22:32:52.16	-60:33:23.92	22.83 ± 0.005	0.60 ± 0.008	1.2900	3	[O II], Mg II . Fe II	53	1465
14	22:32:53.24	-60:34:13.36	22.83 ± 0.004	0.38 ± 0.006	0.0000	3	K1	53	2469
15	22:32:52.88	-60:33:17.12	22.86 ± 0.008	1.11 ± 0.014	1.2838	3	[O II], Mg II	12	1334
16	22:32:55.25	-60:34:07.50	22.87 ± 0.013	0.37 ± 0.017	0.4647	3	[O III], H δ , [Ne III], [O II]	54	2305
17	22:32:55.87	-60:33:17.75	22.93 ± 0.010	0.63 ± 0.014	0.5810	3	[O III], H β , H γ , [O II]	21	1348
18	22:32:58.77	-60:33:23.52	23.02 ± 0.004	-0.22 ± 0.005	0.0000	3	White Dwarf B8-B9	54	1444
19	22:32:57.72	-60:34:08.65	23.04 ± 0.004	2.28 ± 0.014	0.0000	3	M5	54	2364
20	22:32:57.90	-60:33:49.12	23.06 ± 0.009	0.55 ± 0.012	0.4275	3	[O III], H β , H γ , [O II]	54	1942
21	22:32:58.11	-60:33:37.37	23.12 ± 0.004	0.48 ± 0.006	0.0000	3	K3	54	1724
22	22:32:55.36	-60:33:54.97	23.31 ± 0.007	2.14 ± 0.021	0.0000	3	M5	54	2072
23	22:32:55.74	-60:33:33.75	23.42 ± 0.019	0.48 ± 0.024	0.5641	3	[O III], H β , H γ , H δ , [Ne III], [O II]	54	1655
24	22:32:55.22	-60:34:10.15	23.42 ± 0.011	0.88 ± 0.019	0.9723	3	H γ , H δ , [O II], Mg II	54	2389
25	22:32:54.98	-60:33:28.91	23.64 ± 0.011	1.39 ± 0.023	1.2826	3	[Ne III], [O II]	54	1557
26	22:32:55.83	-60:33:30.20	23.67 ± 0.012	0.45 ± 0.016	0.2249	3	H α , [O III], H β , H γ	54	1577
27	22:32:53.41	-60:33:33.37	23.75 ± 0.012	0.97 ± 0.020	1.2853	3	[O II], Mg II	53	1650
28	22:32:52.25	-60:33:19.48	23.79 ± 0.012	0.13 ± 0.014	0.3179	3	[O III], H β , [Ne III], [O II]	50	1391
29	22:32:57.09	-60:33:28.91	23.90 ± 0.013	1.64 ± 0.034	0.8309	3	[O II], Balmer abs.	54	1554
30	22:32:59.85	-60:34:05.48	23.92 ± 0.014	0.68 ± 0.020	...	0		5	2281
31	22:32:55.86	-60:33:51.37	23.92 ± 0.008	1.25 ± 0.016	0.0000	3	M-star	54	2003
32	22:32:54.33	-60:33:20.70	23.94 ± 0.012	0.53 ± 0.016	0.5644	3	[O III], H β , [O II]	53	1400
33	22:32:59.56	-60:34:20.60	23.94 ± 0.018	1.48 ± 0.050	...	0		2	2602
34	22:32:54.26	-60:33:20.07	24.11 ± 0.013	0.48 ± 0.018	0.3649	1	[O II], contamination 32	53	1399
35	22:32:54.00	-60:33:41.48	24.15 ± 0.014	1.03 ± 0.025	1.2806	3	[O II], Mg II	54	1800
36	22:32:53.86	-60:34:20.18	24.19 ± 0.025	0.21 ± 0.031	...	0		1	2584
37	22:32:59.46	-60:33:28.94	24.19 ± 0.010	0.66 ± 0.015	1.0978	3	H δ , [Ne III], [O II], Mg II	48	1560
38	22:32:59.44	-60:33:56.10	24.41 ± 0.019	0.60 ± 0.026	1.0217	3	H γ , [O II]	48	2071
39	22:32:52.56	-60:33:57.72	24.50 ± 0.021	0.61 ± 0.028	1.2658	3	[O II], Balmer abs.	53	2205
40	22:32:54.24	-60:33:50.58	24.50 ± 0.017	0.39 ± 0.022	3.0121	3	Ly α , C III], Si II, CII	54	1983
41	22:32:54.18	-60:34:08.94	24.57 ± 0.019	0.18 ± 0.023	2.4061	3	C III], C IV, Si II, CII	54	2376
42	22:32:59.06	-60:34:14.70	24.58 ± 0.015	0.61 ± 0.022	...	0		53	2493
43	22:32:52.03	-60:33:42.59	24.62 ± 0.018	0.23 ± 0.022	3.2903	3	Ly α , Si II, C IV, Si IV	53	1825
44	22:32:59.20	-60:33:57.53	24.63 ± 0.043	0.66 ± 0.061	...	0		54	2109
45	22:32:54.97	-60:33:36.72	24.69 ± 0.048	0.94 ± 0.078	1.1552	3	[Ne III], [O II], Balmer abs.	54	1719
46	22:32:51.31	-60:34:01.64	24.70 ± 0.016	0.31 ± 0.020	...	0		15	2196
47	22:32:58.30	-60:33:29.02	24.82 ± 0.029	0.40 ± 0.037	...	0		54	1567
48	22:32:58.32	-60:33:46.92	24.84 ± 0.022	0.40 ± 0.029	1.2154	3	[Ne III], [O II], Balmer abs.	54	1921
49	22:32:53.89	-60:34:02.01	24.84 ± 0.022	0.74 ± 0.033	0.9991	3	[O II], Balmer abs.	53	2228
50	22:32:53.22	-60:33:20.28	24.86 ± 0.024	0.38 ± 0.030	2.6721	3	C III], OI/SiII, CII, CIV, FeII, AlIII	52	1396
51	22:32:52.89	-60:33:21.74	24.87 ± 0.020	0.27 ± 0.025	2.6728	3	C II, SiII, C IV	53	1422
52	22:32:59.57	-60:33:25.42	24.90 ± 0.031	0.21 ± 0.038	...	0		41	1491
53	22:32:56.54	-60:33:36.64	24.92 ± 0.019	0.31 ± 0.023	0.2297	3	H α , [O III], H β	54	1717

Notes. ^(a) Saturated in the HST image. ^(b) Not in the Casertano et al. (2000) catalogue but clearly detected in HST, thus photometry was done manually on the HST images using the same aperture in each band.

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
54	22:32:55.03	-60:33:25.44	24.93 ± 0.024	0.22 ± 0.029	...	0		54	1496
55	22:32:53.11	-60:33:20.25	24.94 ± 0.027	0.49 ± 0.034	2.6735	3	C III], OI/SiIII, CII, SiIV, SiII, CIV, FeII, AlIII	52	1512
56	22:32:54.32	-60:33:52.49	25.03 ± 0.027	0.45 ± 0.034	3.0082	3	Si II, C IV, C II	54	2012
57	22:32:57.60	-60:33:18.32	25.07 ± 0.027	0.47 ± 0.035	0.4200	3	[O III], H β , [O II]	30	1359
58	22:32:56.82	-60:33:31.54	25.09 ± 0.028	1.61 ± 0.073	0.8317	1	Ca H+K, Mg II	54	1631
59	22:32:58.60	-60:33:46.58	25.12 ± 0.040	1.93 ± 0.136	...	0		54	1912
60	22:32:58.79	-60:33:33.04	25.13 ± 0.030	0.23 ± 0.037	2.2230	3	Si II, C IV, Al II	54	1652
61	22:32:53.91	-60:33:46.32	25.14 ± 0.024	0.28 ± 0.030	0.3210	3	H α , [O III], H β , H γ , [O II]	53	1928
62	22:32:54.52	-60:33:24.47	25.14 ± 0.020	0.45 ± 0.026	0.2836	3	[O III], H β	54	1485
63	22:32:57.34	-60:33:26.07	25.14 ± 0.027	0.29 ± 0.034	0.1718	3	H β	54	1505
64	22:32:54.00	-60:33:32.87	25.21 ± 0.029	0.46 ± 0.038	1.2813	2	[Ne III], [O II]	54	1661
65	22:32:53.67	-60:34:03.14	25.25 ± 0.025	0.25 ± 0.030	2.0200	3	[Ne III], [O II]	53	2253
66	22:32:57.32	-60:33:28.75	25.27 ± 0.023	2.45 ± 0.066	...	0		54	1573
67	22:32:56.73	-60:33:37.42	25.29 ± 0.025	0.43 ± 0.033	0.2946	3	[O III], H β , [O II]	54	1735
68	22:32:53.74	-60:33:50.70	25.29 ± 0.029	0.60 ± 0.041	0.9715	3	H γ , H δ , [Ne III], [O II]	53	1994
69	22:32:59.77	-60:34:10.35	25.30 ± 0.035	0.10 ± 0.042	...	0		9	2387
70	22:32:59.08	-60:33:41.77	25.31 ± 0.031	0.41 ± 0.040	0.1722	3	H α , [O III], H β	54	1803
71	22:32:53.64	-60:33:55.03	25.34 ± 0.032	0.52 ± 0.043	3.0856	2	Ly α	53	2078
72	22:32:53.64	-60:33:55.03	25.34 ± 0.032	0.52 ± 0.043	0.8391	3	[O III], H β , [O II]	53	2078
73	22:32:59.11	-60:34:05.39	25.36 ± 0.040	0.27 ± 0.049	2.8673	1	OI, SiII, C II, CIV, Fe I, AlIII, C III]?	54	2289
74	22:32:53.31	-60:34:04.50	25.41 ± 0.033	0.07 ± 0.040	1.9830	1	C III], Fe II, Al II	53	2280
75	22:32:56.97	-60:33:49.97	25.43 ± 0.058	0.94 ± 0.096	0.8414	2	[O III], [O II]	54	1975
76	22:32:58.45	-60:33:51.69	25.46 ± 0.030	0.38 ± 0.038	...	0		54	10623
77	22:32:53.20	-60:34:20.17	25.48 ± 0.088	0.54 ± 0.124	...	0		5	2609
78	22:32:52.46	-60:34:12.58	25.51 ± 0.037	-0.02 ± 0.044	2.3730	2	C III], C IV, Si II	53	2454
79	22:32:55.43	-60:34:07.74	25.52 ± 0.025	0.45 ± 0.034	1.1459	2	[Ne III], [O II]	54	2306
80	22:32:52.56	-60:33:16.28	25.52 ± 0.030	0.17 ± 0.036	...	0		4	1333
81	22:32:55.89	-60:33:55.75	25.56 ± 0.042	0.47 ± 0.056	...	0		54	2093
82	22:32:53.80	-60:34:18.98	25.57 ± 0.113	-0.15 ± 0.130	...	0		19	2583
83	22:32:59.39	-60:33:48.13	25.60 ± 0.036	0.09 ± 0.042	2.8158	2	Si II, Fe III.	52	1940
84	22:32:59.42	-60:33:55.05	25.61 ± 0.025	0.23 ± 0.030	...	0		49	2070
85	22:32:56.64	-60:34:16.60	25.67 ± 0.050	0.16 ± 0.063	...	0		52	2521
86	22:32:58.50	-60:33:42.12	25.72 ± 0.051	0.25 ± 0.062	...	0		54	1808
87	22:32:54.86	-60:33:42.12	25.74 ± 0.031	-0.02 ± 0.035	2.6707	3	C III], C IV, Si IV abs	54	1827
88	22:32:56.68	-60:33:27.54	25.76 ± 0.051	0.28 ± 0.063	1.3600	1	[O II]	54	1540
89	22:32:51.53	-60:34:17.12	25.78 ± 0.088	0.54 ± 0.133	4.0140	2	Ly α	39	2556
90	22:32:56.00	-60:33:59.80	25.80 ± 0.046	0.64 ± 0.065	0.6304	3	[O III], H β , [O II]	54	2169
91	22:32:52.74	-60:34:00.02	25.83 ± 0.047	-0.02 ± 0.054	...	0		53	2167
92	22:32:54.82	-60:34:14.12	25.83 ± 0.077	1.25 ± 0.169	4.5804	3	Ly α	53	2487
93	22:32:52.77	-60:34:13.67	25.83 ± 0.050	0.73 ± 0.079	...	0		53	2480
94	22:32:58.71	-60:34:16.56	25.84 ± 0.052	1.30 ± 0.118	...	0		53	2524
95	22:32:58.60	-60:34:09.01	25.84 ± 0.049	0.80 ± 0.077	4.2248	3	Ly α	54	2374
96	22:32:56.42	-60:34:05.41	25.85 ± 0.041	0.71 ± 0.062	1.1516	2	[Ne III], [O II]	54	2294
97	22:32:55.50	-60:33:57.42	25.87 ± 0.036	0.04 ± 0.042	1.5707	3	C III], Mg II	54	2129
98	22:32:59.04	-60:34:06.31	25.88 ± 0.037	0.46 ± 0.047	0.1188	2	H α , [O III]	54	2290
99	22:32:51.42	-60:34:00.84	25.89 ± 0.042	0.60 ± 0.057	...	0		29	2195
100	22:32:55.13	-60:33:35.07	25.97 ± 0.068	0.77 ± 0.104	...	0		54	1691
101	22:32:58.68	-60:33:27.09	25.98 ± 0.039	0.45 ± 0.051	0.4223	3	[O III], H β , [O II]	54	1530
102	22:32:58.47	-60:33:52.69	25.98 ± 0.047	0.17 ± 0.057	...	0		54	10629
103	22:32:54.05	-60:34:19.36	26.00 ± 0.064	0.89 ± 0.102	...	0		14	2596
104	22:32:52.03	-60:34:05.75	26.00 ± 0.044	0.63 ± 0.063	1.1387	1	[O II]	53	2307
105	22:32:54.44	-60:34:19.14	26.02 ± 0.070	0.30 ± 0.084	...	0		19	2579
106	22:32:56.46	-60:33:31.95	26.03 ± 0.073	0.71 ± 0.107	...	0		54	1632
107	22:32:54.84	-60:33:23.72	26.03 ± 0.052	0.49 ± 0.069	1.1464	1	[O II]	54	1469
108	22:32:56.19	-60:34:00.34	26.03 ± 0.042	0.13 ± 0.049	...	0		54	2201
109	22:32:56.16	-60:34:11.74	26.04 ± 0.066	-0.24 ± 0.073	2.1933	3	C III], Si II, Al II	54	2422
110	22:32:54.52	-60:34:18.74	26.04 ± 0.053	0.38 ± 0.072	...	0		23	2578
111	22:32:54.97	-60:33:47.93	26.07 ± 0.049	0.13 ± 0.058	...	0		54	1944
112	22:32:57.51	-60:33:48.47	26.08 ± 0.055	0.35 ± 0.070	3.9076	3	Ly α	54	1946
113	22:32:51.54	-60:33:58.40	26.08 ± 0.048	1.20 ± 0.093	...	0		41	2153
114	22:32:54.24	-60:33:47.27	26.09 ± 0.064	0.48 ± 0.083	1.2844	1	[O II]	54	1931
115	22:32:56.99	-60:33:35.02	26.10 ± 0.043	0.57 ± 0.059	0.6708	3	[O III], H β , [O II]	54	1689
116	22:32:59.49	-60:34:06.35	26.11 ± 0.041	0.07 ± 0.048	1.9202	1	C III], Mg I, Fe III	39	2314
117	22:32:59.67	-60:33:33.35	26.12 ± 0.084	0.29 ± 0.104	0.6454	3	[O III], H β , [O II]	24	1651

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
118	22:32:51.14	-60:34:11.89	26.13 ± 0.078	0.24 ± 0.102	...	0		3	2441
119	22:32:52.86	-60:33:47.29	26.14 ± 0.052	0.15 ± 0.062	...	0		53	1930
120	22:32:58.27	-60:33:56.48	26.16 ± 0.064	0.54 ± 0.087	0.8479	3	[O III], H β , [O II]	54	2102
121	22:32:51.83	-60:33:51.74	26.16 ± 0.066	0.33 ± 0.082	...	0		53	2010
122	22:32:57.43	-60:33:28.16	26.17 ± 0.054	0.05 ± 0.063	0.5752	1	[O II], H β	54	1550
123	22:32:54.70	-60:33:36.14	26.18 ± 0.040	1.11 ± 0.074	3.9014	1	Si IV, Si II, C II, Al II	54	1718
124	22:32:53.29	-60:33:34.88	26.18 ± 0.040	1.13 ± 0.094	...	0		53	1688
125	22:32:53.69	-60:33:57.72	26.19 ± 0.056	0.07 ± 0.065	2.0188	1	C III], Al II, Fe II, Mg II	53	2133
126	22:32:59.06	-60:33:53.72	26.19 ± 0.041	-0.15 ± 0.046	2.3714	1	C III]	54	2042
127	22:32:54.37	-60:34:08.47	26.25 ± 0.059	0.00 ± 0.069	...	0		54	2372
128	22:32:57.36	-60:34:10.13	26.27 ± 0.045	0.34 ± 0.055	1.5656	1	C III], Fe II, Al II	54	2390
129	22:32:59.03	-60:33:23.40	26.28 ± 0.045	0.19 ± 0.054	0.9719	3	H δ , [Ne III], [O II]	54	1461
130	22:32:57.35	-60:34:05.65	26.28 ± 0.069	-0.12 ± 0.078	...	0		54	10710
131	22:32:52.26	-60:33:45.37	26.30 ± 0.075	0.73 ± 0.112	...	0		53	1901
132	22:32:51.34	-60:34:18.43	26.33 ± 0.206	0.59 ± 0.279	...	0		11	2576
133	22:32:52.38	-60:33:53.78	26.33 ± 0.078	0.27 ± 0.096	...	0		53	2048
134	22:32:54.69	-60:33:35.51	26.35 ± 0.064	1.56 ± 0.124	...	0		54	1700
135	22:32:55.11	-60:33:22.72	26.37 ± 0.071	0.60 ± 0.097	0.5637	3	[O III], [O II]	54	1452
136	22:32:55.80	-60:33:41.90	26.38 ± 0.076	0.46 ± 0.101	...	0		54	1815
137	22:32:52.31	-60:33:38.63	26.39 ± 0.070	0.09 ± 0.083	...	0		53	1760
138	22:32:54.15	-60:33:31.61	26.39 ± 0.063	0.07 ± 0.074	...	0		54	1636
139	22:32:55.45	-60:33:40.23	26.40 ± 0.098	0.32 ± 0.121	3.3490	3	Ly α , C III]	54	1788
140	22:32:57.34	-60:34:17.88	26.41 ± 0.114	-0.04 ± 0.132	...	0		41	2560
141	22:32:57.95	-60:34:08.65	26.42 ± 0.062	0.75 ± 0.093	0.8263	3	[O III], H β , [O II]	54	2370
142	22:32:54.33	-60:34:12.94	26.43 ± 0.076	0.48 ± 0.105	...	0		54	2473
143	22:32:54.58	-60:33:39.66	26.46 ± 0.080	0.30 ± 0.099	...	0		54	1772
144	22:32:58.94	-60:34:00.07	26.47 ± 0.046	0.24 ± 0.056	4.0160	2	Ly α	54	2178
145	22:32:57.71	-60:33:34.73	26.48 ± 0.077	0.21 ± 0.094	...	0		54	1681
146	22:32:54.07	-60:33:42.71	26.48 ± 0.056	0.73 ± 0.078	3.7990	1	Ly α , Si II, Si IV, C IV	54	1834
147	22:32:57.54	-60:34:17.07	26.49 ± 0.089	0.26 ± 0.115	...	0		50	2547
148	22:32:53.14	-60:33:31.36	26.49 ± 0.176	1.19 ± 0.339	...	0		53	1633
149	22:32:53.88	-60:34:00.26	26.51 ± 0.065	0.16 ± 0.078	...	0		53	2185
150	22:32:58.98	-60:34:09.35	26.51 ± 0.080	0.30 ± 0.101	...	0		54	2382
151	22:32:58.02	-60:34:15.43	26.56 ± 0.075	0.14 ± 0.091	...	0		53	2505
152	22:32:55.86	-60:34:16.75	26.56 ± 0.057	0.25 ± 0.073	...	0		51	2538
153	22:32:54.21	-60:33:22.54	26.58 ± 0.128	1.03 ± 0.223	...	0		54	1451
154	22:32:57.69	-60:33:16.81	26.60 ± 0.056	0.64 ± 0.080	...	0		5	1341
155	22:32:51.61	-60:33:26.81	26.61 ± 0.117	0.07 ± 0.135	3.0176	1	Ly α ?, C II, Si II, Si IV?	45	1535
156	22:32:51.98	-60:33:41.68	26.62 ± 0.047	2.29 ± 0.123	...	0		53	1822
157	22:32:58.64	-60:33:33.34	26.65 ± 0.048	0.73 ± 0.069	0.7691	2	H β , [O II]	54	1663
158	22:32:55.24	-60:33:29.74	26.65 ± 0.086	0.59 ± 0.119	...	0		54	1598
159	22:32:59.24	-60:33:46.94	26.65 ± 0.082	0.51 ± 0.109	3.7450	2	Ly α	54	1932
160	22:32:52.65	-60:33:20.52	26.66 ± 0.063	0.06 ± 0.073	1.2836	3	[Ne III], [O II]	52	1409
161	22:32:56.38	-60:33:30.31	26.66 ± 0.054	0.36 ± 0.067	...	0		54	1607
162	22:32:59.03	-60:33:22.08	26.67 ± 0.078	0.65 ± 0.111	3.2776	1	Ly α	54	1427
163	22:32:51.53	-60:33:53.45	26.69 ± 0.072	0.64 ± 0.102	...	0		38	2050
164	22:32:56.21	-60:34:17.75	26.70 ± 0.093	0.35 ± 0.127	...	0		42	2562
165	22:32:51.28	-60:34:07.57	26.73 ± 0.070	0.95 ± 0.121	...	0		14	2355
166	22:32:54.69	-60:33:44.79	26.74 ± 0.086	0.02 ± 0.098	...	0		54	1886
167	22:32:57.22	-60:33:29.96	26.74 ± 0.055	0.54 ± 0.073	...	0		54	1602
168	22:32:54.07	-60:33:53.39	26.74 ± 0.106	0.18 ± 0.126	...	0		54	2039
169	22:32:58.83	-60:34:11.70	26.75 ± 0.072	0.38 ± 0.092	...	0		54	2424
170	22:32:58.24	-60:33:36.14	26.78 ± 0.075	0.89 ± 0.108	...	0		54	1706
171	22:32:59.74	-60:34:07.45	26.79 ± 0.087	0.09 ± 0.104	0.7725	1	[O II]	10	2336
172	22:32:57.46	-60:34:21.82	26.79 ± 0.219	1.32 ± 0.488	...	0		1	2617
173	22:32:59.77	-60:33:47.69	26.80 ± 0.110	0.69 ± 0.161	...	0		7	1935
174	22:32:57.42	-60:33:24.47	26.83 ± 0.085	1.27 ± 0.174	...	0		54	1490
175	22:32:52.26	-60:34:11.79	26.84 ± 0.130	1.62 ± 0.381	...	0		53	2444
176	22:32:54.34	-60:33:40.35	26.84 ± 0.106	0.03 ± 0.122	...	0		54	1787
177	22:32:53.75	-60:33:26.93	26.84 ± 0.108	0.21 ± 0.131	...	0		53	1529
178	22:32:52.85	-60:33:52.23	26.88 ± 0.070	0.06 ± 0.082	...	0		53	2024
179	22:32:59.52	-60:34:18.23	26.89 ± 0.121	0.51 ± 0.173	...	0		25	2567
180	22:32:54.74	-60:33:49.21	26.89 ± 0.143	-0.06 ± 0.161	...	0		54	1968
181	22:32:59.05	-60:33:25.45	26.90 ± 0.096	0.07 ± 0.112	3.3368	3	Ly α	54	1498
182	22:32:51.45	-60:34:17.16	26.91 ± 0.089	0.72 ± 0.138	...	0		33	2557
183	22:32:52.66	-60:34:16.63	26.91 ± 0.155	0.30 ± 0.213	3.3745	2	Ly α	52	2543
184	22:32:54.02	-60:33:27.06	26.92 ± 0.071	-0.04 ± 0.081	...	0		54	1538

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
185	22:32:54.91	-60:33:54.82	26.93 ± 0.195	0.26 ± 0.237	...	0		54	2075
186	22:32:56.19	-60:33:55.42	26.94 ± 0.072	2.68 ± 0.484	4.9017	2	Ly α	54	2092
187	22:32:59.33	-60:34:09.23	26.95 ± 0.108	0.22 ± 0.133	...	0		54	2379
188	22:32:52.77	-60:33:59.23	26.96 ± 0.064	0.80 ± 0.094	...	0		53	2244
189	22:32:56.66	-60:34:13.88	26.99 ± 0.124	0.11 ± 0.150	...	0		53	2483
190	+22:32:51.38	-60:34:08.02	27.01 ± 0.073	1.01 ± 0.134	...	0		24	2369
191	+22:32:55.88	-60:34:16.76	27.04 ± 0.086	0.47 ± 0.113	...	0		51	2533
192	+22:32:53.50	-60:34:02.40	27.05 ± 0.141	0.26 ± 0.172	...	0		53	2316
193	+22:32:57.79	-60:34:17.52	27.05 ± 0.107	0.84 ± 0.179	...	0		50	2551
194	+22:32:59.01	-60:34:19.62	27.07 ± 0.096	0.60 ± 0.140	...	0		1	2614
195	+22:32:58.58	-60:34:19.44	27.08 ± 0.102	0.26 ± 0.133	...	0		24	2585
196	+22:32:53.08	-60:33:18.98	27.09 ± 0.081	0.76 ± 0.112	...	0		41	1387
197	+22:32:53.19	-60:33:36.42	27.09 ± 0.092	0.43 ± 0.119	...	0		53	1726
198	+22:32:59.01	-60:34:08.09	27.10 ± 0.161	0.68 ± 0.239	...	0		54	2358
199	+22:32:52.05	-60:33:47.40	27.10 ± 0.120	1.02 ± 0.208	...	0		53	1939
200	+22:32:56.37	-60:33:22.15	27.11 ± 0.126	0.05 ± 0.146	3.3494	3	Ly α	54	1430
201	22:32:53.21	-60:33:43.60	27.11 ± 0.112	0.84 ± 0.174	...	0		53	1866
202	+22:32:59.33	-60:34:06.56	27.11 ± 0.122	-0.04 ± 0.140	3.2775	3	Ly α	54	2322
203	22:32:55.60	-60:33:18.90	27.12 ± 0.092	0.85 ± 0.134	...	0		37	1383
204	22:32:58.42	-60:34:03.88	27.15 ± 0.123	0.27 ± 0.153	...	0		54	2261
205	22:32:55.63	-60:34:06.79	27.17 ± 0.103	1.09 ± 0.192	...	0		54	2329
206	22:32:58.39	-60:33:29.67	27.19 ± 0.077	2.11 ± 0.188	...	0		54	1593
207	22:32:59.39	-60:33:45.05	27.20 ± 0.089	-0.03 ± 0.103	...	0		52	1929
208	+22:32:54.30	-60:33:34.33	27.21 ± 0.113	0.41 ± 0.145	...	0		54	1679
209	22:32:53.37	-60:33:45.12	27.22 ± 0.167	0.60 ± 0.232	...	0		53	1896
210	22:32:55.16	-60:33:21.03	27.22 ± 0.201	0.48 ± 0.263	...	0		53	1417
211	22:32:52.15	-60:34:12.54	27.23 ± 0.127	1.33 ± 0.302	...	0		53	2462
212	+22:32:54.00	-60:33:59.67	27.24 ± 0.120	-0.19 ± 0.133	1.9740	1	C III]	54	2174
213	22:32:54.60	-60:33:29.83	27.24 ± 0.091	0.40 ± 0.107	...	0		54	1600
214	22:32:59.35	-60:33:16.51	27.25 ± 0.074	1.07 ± 0.134	...	0		4	1332
215	22:32:52.98	-60:33:49.00	27.26 ± 0.129	0.34 ± 0.161	...	0		53	1969
216	+22:32:56.73	-60:33:38.55	27.26 ± 0.117	0.74 ± 0.173	4.0175	2	Ly α	54	1758
217	22:32:51.76	-60:34:18.66	27.27 ± 0.360	0.37 ± 0.458	...	0		22	2587
218	+22:32:57.95	-60:33:20.23	27.28 ± 0.102	0.88 ± 0.167	4.9014	2	Ly α	52	1404
219	22:32:57.69	-60:34:15.41	27.32 ± 0.168	0.84 ± 0.278	...	0		53	2507
220	22:32:55.46	-60:34:05.88	27.33 ± 0.133	0.25 ± 0.166	...	0		54	2311
221	22:32:51.25	-60:33:27.86	27.34 ± 0.123	0.38 ± 0.158	...	0		5	1556
222	22:32:57.52	-60:34:19.58	27.34 ± 0.170	0.05 ± 0.205	...	0		14	2597
223	22:32:51.98	-60:33:35.11	27.34 ± 0.130	1.36 ± 0.280	...	0		53	1697
224	22:32:58.02	-60:33:58.99	27.35 ± 0.200	-0.18 ± 0.221	...	0		54	2157
225	+22:32:55.60	-60:33:19.32	27.35 ± 0.107	0.79 ± 0.150	4.3118	2	Ly α	47	1395
226	22:32:56.42	-60:34:16.97	27.35 ± 0.125	-0.09 ± 0.147	...	0		50	2545
227	22:32:53.29	-60:34:20.67	27.36 ± 0.155	0.60 ± 0.213	...	0		1	2613
228	22:32:57.01	-60:33:25.70	27.37 ± 0.091	0.78 ± 0.139	...	0		54	1508
229	+22:32:54.26	-60:33:37.78	27.37 ± 0.132	-0.16 ± 0.147	1.4756	1	[O II]	54	1746
230	22:32:57.36	-60:33:36.57	27.39 ± 0.129	-0.01 ± 0.147	...	0		54	1721
231	22:32:58.78	-60:34:12.74	27.40 ± 0.097	0.23 ± 0.121	...	0		54	2455
232	+22:32:52.57	-60:33:39.92	27.41 ± 0.203	0.39 ± 0.258	5.2147	2	Ly α	53	1779
233	22:32:55.02	-60:33:54.79	27.41 ± 0.114	0.51 ± 0.137	...	0		54	2076
234	22:32:59.78	-60:33:56.97	27.44 ± 0.191	0.19 ± 0.228	...	0		7	2118
235	22:32:55.91	-60:33:59.44	27.44 ± 0.098	0.50 ± 0.121	...	0		54	2168
236	22:32:52.56	-60:33:50.74	27.44 ± 0.118	0.76 ± 0.173	...	0		53	1999
237	22:32:54.50	-60:34:00.02	27.44 ± 0.165	0.51 ± 0.221	...	0		54	2181
238	+22:32:55.41	-60:33:56.16	27.44 ± 0.130	1.01 ± 0.180	3.8195	1	Ly α	54	2096
239	22:32:53.86	-60:33:23.56	27.45 ± 0.158	0.03 ± 0.183	...	0		53	1471
240	22:32:55.82	-60:33:39.95	27.45 ± 0.105	0.61 ± 0.147	...	0		54	1781
241	22:32:55.82	-60:33:54.96	27.46 ± 0.116	1.14 ± 0.178	...	0		54	2079
242	22:32:59.46	-60:33:35.46	27.46 ± 0.146	0.33 ± 0.181	...	0		46	1694
243	22:32:59.16	-60:33:54.52	27.46 ± 0.178	0.86 ± 0.280	...	0		54	2063
244	22:32:57.12	-60:34:18.14	27.49 ± 0.152	0.31 ± 0.192	...	0		37	2570
245	22:32:57.08	-60:33:40.01	27.50 ± 0.162	0.37 ± 0.207	...	0		54	1775
246	+22:32:56.42	-60:33:30.47	27.51 ± 0.081	0.89 ± 0.160	5.6802	2	Ly α	54	1615
247	+22:32:57.22	-60:33:25.03	27.52 ± 0.109	0.23 ± 0.134	...	0		54	1517
248	22:32:54.62	-60:33:45.05	27.52 ± 0.102	0.37 ± 0.122	...	0		54	1887
249	22:32:59.51	-60:33:16.81	27.52 ± 0.158	1.46 ± 0.379	...	0		8	1340
250	22:32:54.95	-60:34:05.70	27.52 ± 0.213	-0.37 ± 0.229	...	0		54	2308
251	22:32:54.92	-60:34:07.99	27.53 ± 0.263	1.24 ± 0.520	...	0		54	2362

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
252	22:32:51.72	-60:33:52.60	27.54 ± 0.186	0.79 ± 0.283	...	0		53	2033
253	22:32:56.81	-60:33:48.39	27.54 ± 0.163	0.38 ± 0.208	...	0		54	1953
254	+22:32:59.02	-60:33:58.40	27.54 ± 0.218	0.78 ± 0.329	...	0		54	2243
255	+22:32:53.46	-60:33:44.23	27.55 ± 0.188	0.35 ± 0.236	0.9844	1	[O II]	53	1885
256	22:32:58.46	-60:33:25.76	27.56 ± 0.140	0.91 ± 0.226	...	0		54	1506
257	+22:32:52.31	-60:33:50.31	27.56 ± 0.141	0.29 ± 0.173	4.5247	1	Ly α	53	1982
258	22:32:54.82	-60:33:44.92	27.56 ± 0.197	0.64 ± 0.277	...	0		54	1895
259	22:32:53.01	-60:33:57.04	27.56 ± 0.138	1.05 ± 0.244	...	0		53	2124
260	+22:32:51.30	-60:34:16.37	27.58 ± 0.252	1.19 ± 0.646	...	0		11	2534
261	+22:32:52.62	-60:34:16.82	27.58 ± 0.143	0.87 ± 0.210	3.3747	1	Ly α , contamination 183	49	2544
262	22:32:51.17	-60:33:41.51	27.59 ± 0.153	0.25 ± 0.187	...	0		0	1817
263	22:32:57.80	-60:33:26.98	27.59 ± 0.121	0.11 ± 0.144	...	0		54	1532
264	22:32:59.04	-60:33:26.76	27.60 ± 0.114	0.66 ± 0.162	...	0		54	1526
265	22:32:56.33	-60:33:42.39	27.60 ± 0.207	1.10 ± 0.381	...	0		54	1831
266	22:32:59.88	-60:34:19.99	27.62 ± 0.181	0.51 ± 0.264	...	0		1	2601
267	22:32:59.25	-60:33:34.77	27.62 ± 0.156	1.43 ± 0.352	...	0		54	1686
268	22:32:53.92	-60:34:16.68	27.62 ± 0.208	0.00 ± 0.256	...	0		51	2542
269	22:32:52.83	-60:33:33.55	27.63 ± 0.130	0.40 ± 0.165	...	0		53	1672
270	22:32:52.46	-60:34:19.27	27.64 ± 0.378	0.26 ± 0.486	...	0		16	2599
271	+22:32:57.73	-60:33:20.75	27.66 ± 0.140	0.94 ± 0.237	4.1909	2	Ly α	53	1410
272	22:32:51.65	-60:34:02.35	27.67 ± 0.193	-0.11 ± 0.216	...	0		51	2378
273	22:32:52.74	-60:33:29.48	27.67 ± 0.206	1.40 ± 0.448	...	0		53	1594
274	22:32:56.31	-60:33:47.39	27.69 ± 0.203	2.26 ± 0.937	...	0		54	1934
275	22:32:52.19	-60:34:15.14	27.69 ± 0.168	0.40 ± 0.228	...	0		53	2511
276	22:32:53.67	-60:33:59.76	27.70 ± 0.177	-0.05 ± 0.202	...	0		53	2176
277	22:32:51.35	-60:33:54.05	27.70 ± 0.189	0.28 ± 0.234	...	0		17	2065
278	22:32:52.21	-60:33:34.07	27.70 ± 0.117	0.89 ± 0.169	...	0		53	1680
279	22:32:58.74	-60:33:58.00	27.73 ± 0.170	0.70 ± 0.247	...	0		54	2137
280	22:32:54.65	-60:33:57.59	27.73 ± 0.233	0.65 ± 0.329	...	0		54	2134
281	22:32:52.98	-60:33:42.98	27.73 ± 0.156	0.31 ± 0.194	...	0		53	1856
282	22:32:54.68	-60:33:52.10	27.73 ± 0.135	1.09 ± 0.244	...	0		54	2019
283	22:32:58.62	-60:34:03.83	27.73 ± 0.149	0.36 ± 0.190	...	0		54	2262
284	22:32:51.83	-60:33:22.23	27.74 ± 0.153	-0.14 ± 0.171	...	0		53	1448
285	22:32:57.49	-60:33:31.91	27.75 ± 0.121	0.05 ± 0.141	...	0		54	1643
286	22:32:57.25	-60:33:22.21	27.77 ± 0.199	0		54	1438
287	22:32:52.63	-60:34:02.62	27.78 ± 0.113	0.57 ± 0.145	...	0		53	2238
288	22:32:57.81	-60:33:53.01	27.78 ± 0.137	0.28 ± 0.169	...	0		54	2031
289	22:32:57.48	-60:33:32.54	27.79 ± 0.123	-0.11 ± 0.139	...	0		54	1653
290	+22:32:58.18	-60:33:40.10	27.81 ± 0.130	1.21 ± 0.247	6.0856	2	Ly α	54	1784
291	22:32:51.77	-60:34:02.60	27.82 ± 0.167	0.44 ± 0.218	...	0		53	2248
292	22:32:58.58	-60:34:07.68	27.82 ± 0.208	0.64 ± 0.304	...	0		54	2345
293	22:32:52.65	-60:34:02.22	27.83 ± 0.185	-0.14 ± 0.207	...	0		53	2237
294	+22:32:52.70	-60:33:34.85	27.84 ± 0.222	0.94 ± 0.361	3.9924	2	Ly α	53	1693
295	22:32:54.47	-60:33:42.02	27.85 ± 0.215	0.19 ± 0.257	...	0		54	1826
296	22:32:59.78	-60:33:54.18	27.85 ± 0.254	0.15 ± 0.301	...	0		8	2054
297	22:32:53.34	-60:33:56.14	27.85 ± 0.215	0.31 ± 0.267	...	0		53	2106
298	22:32:57.44	-60:34:10.75	27.86 ± 0.149	0.53 ± 0.205	...	0		54	2407
299	22:32:53.36	-60:34:16.78	27.86 ± 0.169	0.65 ± 0.228	...	0		51	2549
300	22:32:52.68	-60:33:45.39	27.87 ± 0.229	1.00 ± 0.386	...	0		53	1905
301	22:32:53.94	-60:34:03.02	27.88 ± 0.254	0.10 ± 0.300	...	0		53	2251
302	22:32:52.50	-60:33:38.05	27.88 ± 0.136	0.93 ± 0.186	...	0		53	1888
303	22:32:58.79	-60:33:46.02	27.89 ± 0.227	2.77 ± 1.626	...	0		54	1910
304	22:32:55.47	-60:33:23.99	27.90 ± 0.148	1.34 ± 0.316	...	0		54	1478
305	22:32:58.87	-60:34:17.85	27.90 ± 0.251	0.32 ± 0.335	...	0		43	2572
306	22:32:52.65	-60:34:10.78	27.90 ± 0.320	-0.12 ± 0.363	...	0		53	2412
307	22:32:53.96	-60:34:15.32	27.90 ± 0.247	-0.01 ± 0.294	...	0		53	2519
308	+22:32:57.99	-60:33:42.25	27.92 ± 0.155	0.50 ± 0.206	4.0183	2	Ly α	54	1829
309	22:32:56.89	-60:34:01.86	27.92 ± 0.226	1.44 ± 0.538	...	0		54	2225
310	22:32:57.79	-60:33:18.01	27.93 ± 0.291	0.36 ± 0.373	...	0		24	1362
311	+22:32:57.20	-60:33:51.67	27.93 ± 0.230	0.59 ± 0.318	3.8879	2	Ly α	54	2009
312	22:32:53.83	-60:34:04.93	27.94 ± 0.204	0.39 ± 0.264	...	0		53	2292
313	22:32:57.41	-60:33:21.72	27.95 ± 0.193	1.21 ± 0.381	...	0		54	1424
314	22:32:51.91	-60:33:18.08	27.95 ± 0.159	-0.14 ± 0.178	...	0		28	1373
315	22:32:56.28	-60:34:07.80	27.95 ± 0.264	1.21 ± 0.543	...	0		54	2359
316	22:32:58.14	-60:34:11.85	27.95 ± 0.161	0.59 ± 0.229	...	0		54	2433
317	+22:32:54.06	-60:33:52.86	27.96 ± 0.263	0.30 ± 0.324	...	0		54	2020
318	22:32:52.07	-60:34:15.35	27.97 ± 0.303	1.03 ± 0.579	...	0		53	2517

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
319	22:32:58.49	-60:33:48.89	27.98 ± 0.241	0.29 ± 0.300	...	0		54	1961
320	22:32:54.98	-60:34:17.05	27.98 ± 0.393	0.17 ± 0.483	...	0		49	2552
321	22:32:54.33	-60:34:17.81	28.00 ± 0.229	1.10 ± 0.525	...	0		39	2569
322	22:32:57.74	-60:33:42.46	28.00 ± 0.142	0.49 ± 0.190	...	0		54	1864
323	22:32:55.74	-60:33:55.25	28.00 ± 0.311	0.22 ± 0.374	...	0		54	2080
324	+22:32:58.63	-60:33:32.97	28.00 ± 0.131	0.22 ± 0.148	3.5010	1	Ly α	54	1662
325	+22:32:52.30	-60:33:46.73	28.00 ± 0.216	0.93 ± 0.354	4.7006	2	Ly α	53	1927
326	22:32:59.89	-60:33:53.46	28.01 ± 0.166	0.47 ± 0.221	...	0		2	2040
327	22:32:58.67	-60:33:25.78	28.01 ± 0.177	0.75 ± 0.266	...	0		54	1507
328	22:32:53.12	-60:34:02.15	28.01 ± 0.225	0.78 ± 0.350	...	0		53	2236
329	22:32:52.20	-60:33:36.91	28.02 ± 0.202	1.22 ± 0.398	...	0		53	1737
330	22:32:53.45	-60:34:02.66	28.02 ± 0.268	0.15 ± 0.317	...	0		53	2247
331	22:32:56.93	-60:34:12.27	28.03 ± 0.290	0.70 ± 0.436	...	0		54	2447
332	22:32:51.31	-60:33:43.46	28.03 ± 0.153	0.24 ± 0.187	...	0		12	1867
333	22:32:52.54	-60:33:27.77	28.03 ± 0.146	1.30 ± 0.305	...	0		53	1552
334	+22:32:52.84	-60:33:35.57	28.03 ± 0.125	2.70 ± 0.827	4.9152	2	Ly α	53	1711
335	22:32:58.70	-60:33:16.52	28.04 ± 0.318	0.37 ± 0.404	...	0		4	1335
336	22:32:58.11	-60:33:52.83	28.04 ± 0.207	2.03 ± 0.775	...	0		54	2030
337	22:32:58.25	-60:33:25.98	28.04 ± 0.215	1.24 ± 0.429	...	0		54	1513
338	+22:32:52.11	-60:33:21.86	28.04 ± 0.262	...	4.9007	2	Ly α	53	1433
339	22:32:52.88	-60:33:30.10	28.05 ± 0.370	0.25 ± 0.448	...	0		53	1621
340	22:32:52.31	-60:33:41.66	28.06 ± 0.210	0.35 ± 0.267	...	0		53	1821
341	22:32:53.39	-60:34:17.08	28.07 ± 0.163	0.85 ± 0.276	...	0		47	2550
342	22:32:58.64	-60:34:00.33	28.08 ± 0.239	0.48 ± 0.318	...	0		54	2183
343	22:32:56.95	-60:33:39.24	28.10 ± 0.180	0.21 ± 0.219	...	0		54	1765
344	22:32:54.00	-60:33:18.14	28.10 ± 0.162	0.62 ± 0.228	...	0		26	1372
345	22:32:54.63	-60:33:30.13	28.10 ± 0.145	0.53 ± 0.174	...	0		54	1601
346	22:32:57.40	-60:33:22.37	28.11 ± 0.188	0.35 ± 0.238	...	0		54	1439
347	22:32:53.12	-60:34:12.56	28.11 ± 0.403	-0.14 ± 0.462	...	0		53	2465
348	22:32:57.95	-60:34:17.25	28.12 ± 0.195	0.56 ± 0.275	...	0		50	2553
349	22:32:52.55	-60:33:38.15	28.13 ± 0.155	1.09 ± 0.231	...	0		53	1755
350	22:32:54.67	-60:33:42.72	28.15 ± 0.153	1.34 ± 0.259	...	0		54	1847
351	22:32:54.50	-60:33:22.90	28.16 ± 0.246	0.11 ± 0.289	...	0		54	1450
352	22:32:53.04	-60:33:59.06	28.17 ± 0.287	-0.39 ± 0.308	...	0		53	2165
353	22:32:55.97	-60:33:41.32	28.18 ± 0.294	0.51 ± 0.389	...	0		54	1801
354	22:32:59.52	-60:33:59.26	28.18 ± 0.300	0.77 ± 0.457	...	0		37	2163
355	22:32:57.48	-60:33:49.44	28.19 ± 0.186	0.01 ± 0.215	...	0		54	1971
356	22:32:59.52	-60:34:01.70	28.20 ± 0.275	0.27 ± 0.340	...	0		37	2218
357	22:32:53.72	-60:33:32.44	28.21 ± 0.175	0.87 ± 0.276	...	0		53	1656
358	22:32:53.06	-60:34:05.23	28.21 ± 0.197	1.04 ± 0.350	...	0		53	2300
359	22:32:59.28	-60:34:12.08	28.25 ± 0.246	0.07 ± 0.289	...	0		54	2434
360	22:32:51.42	-60:33:57.36	28.27 ± 0.176	0.64 ± 0.253	...	0		28	2136
361	22:32:53.02	-60:33:59.78	28.28 ± 0.241	0.00 ± 0.278	...	0		53	2179
362	22:32:57.48	-60:33:56.60	28.29 ± 0.144	0.71 ± 0.192	...	0		54	2112
363	+22:32:52.46	-60:33:46.51	28.29 ± 0.222	2.60 ± 1.317	4.5272	1	Ly α , contamination 325	53	1925
364	22:32:54.46	-60:33:53.65	28.29 ± 0.424	0.14 ± 0.500	...	0		54	2052
365	22:32:58.50	-60:34:00.90	28.29 ± 0.250	1.08 ± 0.444	...	0		54	2199
366	22:32:58.67	-60:33:41.81	28.29 ± 0.199	0.23 ± 0.241	...	0		54	1816
367	22:32:53.78	-60:34:02.75	28.29 ± 0.334	0		53	2249
368	22:32:51.85	-60:33:37.31	28.31 ± 0.311	0.88 ± 0.498	...	0		53	1745
369	22:32:59.46	-60:33:37.22	28.31 ± 0.192	0.46 ± 0.253	...	0		46	1734
370	22:32:52.61	-60:33:38.35	28.31 ± 0.156	0.96 ± 0.240	...	0		53	1761
371	22:32:58.33	-60:34:14.17	28.31 ± 0.314	0.40 ± 0.416	...	0		53	2491
372	22:32:53.86	-60:33:56.87	28.32 ± 0.290	0.52 ± 0.390	...	0		53	2366
373	22:32:53.40	-60:33:36.11	28.32 ± 0.261	0.33 ± 0.327	...	0		53	1720
374	22:32:58.81	-60:33:46.44	28.32 ± 0.298	2.21 ± 1.296	...	0		54	1918
375	22:32:56.41	-60:33:43.02	28.33 ± 0.355	0.02 ± 0.410	...	0		54	1851
376	22:32:51.96	-60:34:17.00	28.33 ± 0.244	-0.09 ± 0.295	...	0		49	2558
377	22:32:58.17	-60:33:19.04	28.34 ± 0.267	0.80 ± 0.411	...	0		36	1384
378	22:32:54.50	-60:33:22.36	28.34 ± 0.157	0.71 ± 0.204	...	0		54	1449
379	22:32:59.78	-60:33:35.51	28.34 ± 0.178	0.36 ± 0.228	...	0		10	1696
380	22:32:52.97	-60:34:06.56	28.35 ± 0.281	0.10 ± 0.332	...	0		53	2328
381	22:32:55.83	-60:34:10.98	28.36 ± 0.176	1.13 ± 0.342	...	0		54	2414
382	22:32:51.14	-60:33:26.03	28.36 ± 0.300	-0.21 ± 0.329	...	0		0	1520
383	22:32:51.55	-60:33:33.90	28.37 ± 0.365	0.10 ± 0.425	...	0		38	1678
384	22:32:52.19	-60:33:42.18	28.38 ± 0.372	0.78 ± 0.570	...	0		53	1833
385	22:32:54.02	-60:33:16.90	28.38 ± 0.214	0.34 ± 0.268	...	0		8	1346

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
386	+22:32:59.52	-60:33:50.34	28.38 ± 0.357	0.30 ± 0.445	3.7419	1	$\text{Ly}\alpha$	37	1985
387	22:32:58.88	-60:33:48.20	28.38 ± 0.347	-0.27 ± 0.379	...	0		54	1947
388	22:32:53.00	-60:33:23.87	28.39 ± 0.309	-0.17 ± 0.343	...	0		53	1477
389	22:32:54.45	-60:33:16.90	28.40 ± 0.390	0.59 ± 0.537	...	0		7	1347
390	22:32:57.56	-60:34:01.13	28.40 ± 0.287	0.79 ± 0.440	...	0		54	2208
391	22:32:52.76	-60:33:20.76	28.41 ± 0.264	0.25 ± 0.321	...	0		52	1414
392	22:32:55.99	-60:33:57.27	28.42 ± 0.454	-0.07 ± 0.513	...	0		54	2130
393	+22:32:57.94	-60:33:29.10	28.42 ± 0.377	0.29 ± 0.468	4.1886	1	$\text{Ly}\alpha$	54	1584
394	22:32:51.80	-60:34:10.70	28.42 ± 0.381	0.58 ± 0.544	...	0		53	2411
395	22:32:56.04	-60:33:47.72	28.43 ± 0.189	1.43 ± 0.426	...	0		54	1941
396	22:32:52.35	-60:34:16.18	28.43 ± 0.315	-0.22 ± 0.365	...	0		53	2531
397	22:32:54.49	-60:33:20.20	28.43 ± 0.175	1.29 ± 0.290	...	0		53	1428
398	22:32:58.51	-60:33:46.41	28.44 ± 0.172	1.22 ± 0.280	...	0		54	1920
399	22:32:53.25	-60:34:16.08	28.44 ± 0.335	0.23 ± 0.437	...	0		53	2528
400	22:32:56.66	-60:33:17.92	28.45 ± 0.230	0.01 ± 0.263	...	0		24	1364
401	22:32:52.81	-60:34:05.72	28.45 ± 0.351	-0.28 ± 0.384	...	0		53	2315
402	22:32:51.94	-60:33:38.46	28.45 ± 0.187	1.14 ± 0.357	...	0		53	1842
403	22:32:59.93	-60:34:19.05	28.47 ± 0.278	0.40 ± 0.388	...	0		3	2582
404	+22:32:59.60	-60:34:12.55	28.47 ± 0.218	0.40 ± 0.282	0.8533	2	[O III], H β	29	2450
405	22:32:52.37	-60:33:28.69	28.48 ± 0.201	0.79 ± 0.303	...	0		53	1580
406	22:32:53.40	-60:34:01.37	28.48 ± 0.344	-0.39 ± 0.369	...	0		53	2220
407	22:32:57.55	-60:33:36.15	28.48 ± 0.272	1.45 ± 0.637	...	0		54	1713
408	22:32:58.51	-60:33:35.47	28.48 ± 0.233	0.83 ± 0.365	...	0		54	1698
409	22:32:59.81	-60:33:37.91	28.49 ± 0.219	1.96 ± 0.803	...	0		6	1749
410	22:32:53.97	-60:33:53.34	28.50 ± 0.209	1.08 ± 0.372	...	0		53	2045
411	22:32:51.16	-60:34:02.29	28.50 ± 0.589	-0.08 ± 0.667	...	0		0	2240
412	22:32:56.84	-60:33:46.03	28.52 ± 0.246	0.04 ± 0.286	...	0		54	1915
413	22:32:51.75	-60:33:30.25	28.53 ± 0.373	-0.03 ± 0.423	...	0		53	1616
414	22:32:56.23	-60:33:59.73	28.53 ± 0.146	0.88 ± 0.224	...	0		54	2194
415	22:32:57.47	-60:33:56.16	28.53 ± 0.239	0.14 ± 0.284	...	0		54	2101
416	22:32:51.50	-60:33:49.67	28.53 ± 0.338	0.10 ± 0.394	...	0		36	1986
417	22:32:56.61	-60:33:20.66	28.54 ± 0.257	-0.16 ± 0.286	...	0		53	1408
418	22:32:53.03	-60:34:03.55	28.55 ± 0.284	0.85 ± 0.454	...	0		53	2269
419	22:32:56.84	-60:33:51.32	28.55 ± 0.211	0.33 ± 0.264	...	0		54	2006
420	22:32:58.25	-60:34:16.09	28.55 ± 0.305	-0.30 ± 0.336	...	0		53	2520
421	22:32:57.60	-60:33:20.99	28.55 ± 0.227	0.18 ± 0.274	...	0		53	1413
422	+22:32:52.16	-60:34:10.94	28.56 ± 0.315	-0.15 ± 0.356	3.1293	2	$\text{Ly}\alpha$	53	2416
423	22:32:57.98	-60:34:00.46	28.57 ± 0.473	-0.22 ± 0.519	...	0		54	2189
424	22:32:55.63	-60:34:05.34	28.58 ± 0.358	0.42 ± 0.472	...	0		54	2299
425	22:32:51.98	-60:33:28.86	28.59 ± 0.301	1.11 ± 0.555	...	0		53	1582
426	22:32:59.27	-60:33:28.43	28.60 ± 0.345	0.79 ± 0.520	...	0		54	1559
427	22:32:54.26	-60:34:13.81	28.60 ± 0.593	-0.19 ± 0.668	...	0		53	2486
428	22:32:53.04	-60:33:42.66	28.62 ± 0.190	0.88 ± 0.251	...	0		53	1848
429	22:32:54.73	-60:33:42.46	28.62 ± 0.680	-0.60 ± 0.713	...	0		54	1835
430	+22:32:55.98	-60:33:52.07	28.62 ± 0.260	0.68 ± 0.375	6.2797	2	$\text{Ly}\alpha$	54	2021
431	22:32:58.78	-60:33:31.13	28.64 ± 0.268	0.76 ± 0.411	...	0		54	1630
432	22:32:53.14	-60:33:37.33	28.64 ± 0.224	0.11 ± 0.264	...	0		53	1743
433	+22:32:51.65	-60:33:30.24	28.67 ± 0.299	-0.25 ± 0.326	3.4696	2	$\text{Ly}\alpha$	48	1612
434	22:32:52.24	-60:33:18.41	28.67 ± 0.356	0.13 ± 0.423	...	0		36	1380
435	22:32:56.52	-60:34:01.84	28.68 ± 0.336	0.92 ± 0.563	...	0		54	2224
436	22:32:53.78	-60:34:16.38	28.68 ± 0.236	0.95 ± 0.458	...	0		53	2535
437	+22:32:55.85	-60:34:06.59	28.69 ± 0.270	-0.39 ± 0.291	3.1201	2	$\text{Ly}\alpha$	54	2326
438	22:32:58.49	-60:34:14.79	28.69 ± 0.357	0.28 ± 0.456	...	0		53	2497
439	22:32:51.48	-60:33:24.85	28.70 ± 0.355	0.72 ± 0.525	...	0		31	1497
440	22:32:51.74	-60:33:18.44	28.70 ± 0.266	-0.03 ± 0.304	...	0		39	1379
441	+22:32:54.69	-60:33:54.54	28.70 ± 0.241	1.49 ± 0.572	4.6946	2	$\text{Ly}\alpha$	54	2086
442	22:32:51.15	-60:33:37.33	28.70 ± 0.299	0.73 ± 0.443	...	0		0	1748
443	22:32:51.65	-60:33:52.07	28.70 ± 0.378	0.22 ± 0.457	...	0		49	2026
444	22:32:54.11	-60:33:51.14	28.71 ± 0.438	0.33 ± 0.545	...	0		54	2005
445	22:32:53.56	-60:33:24.40	28.72 ± 0.449	-0.15 ± 0.498	...	0		53	1489
446	22:32:57.10	-60:33:39.66	28.72 ± 0.195	0.95 ± 0.267	...	0		54	1774
447	22:32:59.03	-60:33:18.41	28.74 ± 0.300	0.00 ± 0.347	...	0		33	1371
448	22:32:53.22	-60:33:26.28	28.74 ± 0.490	-0.07 ± 0.555	...	0		53	1523
449	+22:32:58.49	-60:34:01.96	28.75 ± 0.243	0.61 ± 0.342	3.2768	2	$\text{Ly}\alpha$	54	2229
450	22:32:51.82	-60:33:48.61	28.75 ± 0.304	0.93 ± 0.494	...	0		53	1964
451	22:32:56.21	-60:33:48.82	28.75 ± 0.370	-0.08 ± 0.418	...	0		54	1963
452	+22:32:52.97	-60:34:19.35	28.75 ± 0.589	-0.16 ± 0.680	3.1201	1	$\text{Ly}\alpha$	18	2591

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
453	+22:32:55.14	-60:33:52.75	28.76 ± 0.368	0.62 ± 0.511	4.7013	1	<i>Lyα</i>	54	2025
454	22:32:56.20	-60:33:30.11	28.76 ± 0.315	-0.07 ± 0.356	...	0		54	1608
455	22:32:58.68	-60:34:06.66	28.76 ± 0.447	0.65 ± 0.658	...	0		54	2324
456	22:32:54.44	-60:34:09.28	28.77 ± 0.355	-0.04 ± 0.407	...	0		54	2400
457	22:32:53.33	-60:34:00.96	28.78 ± 0.215	1.13 ± 0.402	...	0		53	2212
458	22:32:53.32	-60:33:42.66	28.79 ± 0.340	0.01 ± 0.390	...	0		53	1846
459	22:32:55.19	-60:33:40.41	28.79 ± 0.262	0		54	1793
460	22:32:56.53	-60:33:24.05	28.81 ± 0.467	0.15 ± 0.555	...	0		54	1479
461	22:32:59.26	-60:33:52.25	28.83 ± 0.524	-0.30 ± 0.571	...	0		54	2015
462	+22:32:57.42	-60:33:38.70	28.85 ± 0.247	1.28 ± 0.511	5.5961	1	<i>Lyα</i>	54	1876
463	+22:32:52.38	-60:33:24.10	28.85 ± 0.579	-0.75 ± 0.601	...	0		53	1484
464	22:32:52.76	-60:33:26.78	28.85 ± 0.523	0.02 ± 0.594	...	0		53	1536
465	22:32:58.71	-60:33:31.68	28.85 ± 0.307	1.63 ± 0.846	...	0		54	1638
466	22:32:52.99	-60:33:34.56	28.86 ± 0.316	0.17 ± 0.375	...	0		53	1690
467	22:32:58.92	-60:34:11.55	28.86 ± 0.259	0.50 ± 0.348	...	0		54	2423
468	22:32:58.64	-60:33:54.09	28.86 ± 0.282	0.85 ± 0.444	...	0		54	2058
469	+22:32:56.19	-60:33:51.55	28.86 ± 0.395	0.23 ± 0.479	3.4663	1	<i>Lyα</i>	54	2008
470	22:32:54.31	-60:34:11.01	28.86 ± 0.513	-0.44 ± 0.553	...	0		54	2417
471	22:32:59.05	-60:34:16.75	28.91 ± 0.294	-0.01 ± 0.348	...	0		53	2536
472	22:32:52.60	-60:33:22.50	28.91 ± 0.436	0.66 ± 0.630	...	0		53	1456
473	22:32:54.50	-60:33:44.04	28.91 ± 0.276	0.77 ± 0.410	...	0		54	1873
474	+22:32:56.49	-60:33:28.42	28.92 ± 0.294	0.74 ± 0.436	4.7127	1	<i>Lyα</i>	54	1566
475	22:32:55.92	-60:33:57.33	28.92 ± 0.441	0.76 ± 0.665	...	0		54	2131
476	22:32:57.25	-60:33:17.79	28.92 ± 0.327	0.05 ± 0.378	...	0		17	1360
477	22:32:57.28	-60:34:16.37	28.93 ± 0.278	0.44 ± 0.376	...	0		53	2530
478	+22:32:51.86	-60:33:27.82	28.93 ± 0.512	0.04 ± 0.593	3.4691	1	<i>Lyα</i>	53	1558
479	22:32:59.22	-60:33:28.48	28.94 ± 0.419	0.43 ± 0.544	...	0		54	1561
480	22:32:59.85	-60:34:02.23	28.96 ± 0.255	0.86 ± 0.407	...	0		3	2233
481	22:32:53.96	-60:34:08.03	28.98 ± 0.425	-0.35 ± 0.462	...	0		53	2367
482	22:32:53.96	-60:33:58.06	28.99 ± 0.293	0.10 ± 0.341	...	0		53	2148
483	22:32:54.48	-60:34:13.12	28.99 ± 0.479	0.40 ± 0.642	...	0		54	2472
484	+22:32:58.61	-60:34:12.73	29.01 ± 0.293	0.03 ± 0.347	4.9133	2	<i>Lyα</i>	54	2452
485	22:32:52.12	-60:34:11.75	29.01 ± 0.313	0.35 ± 0.414	...	0		53	2442
486	22:32:57.92	-60:34:01.07	29.02 ± 0.498	0.17 ± 0.593	...	0		54	2207
487	22:32:52.68	-60:33:24.69	29.04 ± 0.259	0.64 ± 0.366	...	0		53	1493
488	22:32:59.41	-60:34:02.96	29.05 ± 0.324	0.61 ± 0.457	...	0		50	2275
489	+22:32:57.03	-60:33:44.45	29.06 ± 0.277	0.49 ± 0.366	2.9565	2	<i>Lyα</i>	54	1881
490	+22:32:57.38	-60:33:17.75	29.06 ± 0.392	-0.28 ± 0.429	...	0		4	1342
491	22:32:53.04	-60:33:52.67	29.06 ± 0.384	0.51 ± 0.513	...	0		53	2034
492	+22:32:57.88	-60:33:58.29	29.06 ± 0.405	0.09 ± 0.473	5.7604	2	<i>Lyα</i>	54	2144
493	22:32:58.98	-60:33:56.84	29.07 ± 0.599	0.73 ± 0.877	...	0		54	2119
494	22:32:53.76	-60:34:00.97	29.09 ± 0.477	0.31 ± 0.599	...	0		53	2211
495	22:32:56.28	-60:33:34.45	29.09 ± 0.559	-0.59 ± 0.589	...	0		54	1683
496	22:32:57.29	-60:33:57.20	29.10 ± 0.305	0		54	2122
497	22:32:57.50	-60:33:27.42	29.10 ± 0.358	-0.55 ± 0.379	...	0		54	1545
498	+22:32:59.50	-60:34:09.46	29.10 ± 0.599	0.58 ± 0.836	4.2052	1	<i>Lyα</i>	36	2384
499	+22:32:54.70	-60:33:54.52	29.12 ± 0.271	0.80 ± 0.414	4.6953	2	<i>Lyα</i>	54	2067
500	+22:32:56.72	-60:33:56.29	29.13 ± 0.284	0.49 ± 0.379	3.5009	1	<i>Lyα</i>	54	2107
501	+22:32:54.41	-60:34:02.47	29.15 ± 0.501	-0.40 ± 0.540	...	0		54	2256
502	22:32:54.43	-60:33:38.09	29.16 ± 0.301	0.20 ± 0.360	...	0		54	1857
503	+22:32:56.42	-60:33:50.45	29.17 ± 0.523	-0.37 ± 0.564	3.3484	2	<i>Lyα</i>	54	1990
504	22:32:54.35	-60:33:22.90	29.17 ± 0.577	0.38 ± 0.732	...	0		54	1460
505	22:32:57.53	-60:34:07.66	29.20 ± 0.303	0.57 ± 0.420	...	0		54	2352
506	22:32:53.35	-60:33:55.66	29.20 ± 0.578	-0.50 ± 0.613	...	0		53	2100
507	22:32:54.67	-60:34:19.69	29.20 ± 0.629	-0.02 ± 0.706	...	0		9	2604
508	22:32:58.16	-60:33:26.07	29.21 ± 0.426	0.14 ± 0.505	...	0		54	1515
509	22:32:57.88	-60:33:52.21	29.21 ± 0.350	-0.01 ± 0.399	...	0		54	2017
510	22:32:53.09	-60:33:47.20	29.22 ± 0.263	0.97 ± 0.439	...	0		53	1938
511	22:32:59.67	-60:33:31.01	29.22 ± 0.421	0.33 ± 0.531	...	0		25	1627
512	22:32:59.42	-60:34:07.68	29.22 ± 0.714	-0.02 ± 0.820	...	0		47	2346
513	+22:32:59.49	-60:33:32.05	29.26 ± 0.402	-0.10 ± 0.448	3.2780	2	<i>Lyα</i>	48	1645
514	+22:32:56.28	-60:33:56.54	29.29 ± 0.442	0.54 ± 0.601	3.8227	2	<i>Lyα</i>	54	2115
515	22:32:58.84	-60:34:03.22	29.30 ± 0.362	-0.20 ± 0.399	...	0		54	2252
516	22:32:54.54	-60:33:48.03	29.30 ± 0.545	-0.31 ± 0.590	...	0		54	1950
517	22:32:57.79	-60:34:19.58	29.33 ± 0.734	-0.82 ± 0.768	...	0		14	2600
518	22:32:59.28	-60:33:21.38	29.33 ± 0.277	0.76 ± 0.415	...	0		53	1419
519	22:32:54.20	-60:34:07.45	29.35 ± 0.371	-0.07 ± 0.423	...	0		54	2351

Table A.1. continued.

ID	α_{J2000}	δ_{J2000}	I_{814}	$V_{606} - I_{814}$	Z	Confidence	Comments	N_{exp}	HDFS-ID
520	+22:32:56.19	-60:33:30.72	29.36 ± 0.629	-0.38 ± 0.676	3.4776	2	$\text{Ly}\alpha$	54	1623
521	22:32:58.07	-60:33:59.86	29.38 ± 0.430	-0.54 ± 0.456	...	0		54	2177
522	22:32:58.48	-60:34:06.34	29.38 ± 0.292	0.42 ± 0.389	...	0		54	2320
523	22:32:58.73	-60:33:18.08	29.40 ± 0.433	-0.70 ± 0.452	...	0		28	1365
524	22:32:57.77	-60:33:37.89	29.40 ± 0.523	0.03 ± 0.605	...	0		54	1750
525	22:32:56.04	-60:33:31.24	29.40 ± 0.412	0.11 ± 0.484	...	0		54	1634
526	22:32:52.79	-60:33:56.66	29.41 ± 0.372	-0.01 ± 0.425	...	0		53	2350
527	22:32:56.10	-60:33:42.72	29.41 ± 0.543	0.47 ± 0.720	...	0		54	1843
528	22:32:56.02	-60:33:55.48	29.48 ± 0.550	0.30 ± 0.685	...	0		54	2094
529	22:32:59.48	-60:33:26.75	29.50 ± 0.366	0.20 ± 0.437	...	0		49	1527
530	22:32:54.54	-60:33:28.40	29.52 ± 0.393	-0.21 ± 0.434	...	0		54	1569
531	22:32:58.29	-60:33:20.12	29.53 ± 0.581	-0.31 ± 0.631	...	0		51	1403
532	22:32:51.61	-60:34:09.42	29.58 ± 0.439	0.02 ± 0.514	...	0		51	2392
533	22:32:51.63	-60:34:12.14	29.59 ± 0.895	-1.33 ± 0.910	...	0		53	2453
534	+22:32:59.66	-60:34:06.97	29.61 ± 0.431	-0.24 ± 0.475	4.4323	2	$\text{Ly}\alpha$	22	2332
535	22:32:55.81	-60:34:06.97	29.62 ± 0.489	-0.03 ± 0.562	...	0		54	2335
536	22:32:52.81	-60:33:32.54	29.65 ± 0.763	-0.35 ± 0.821	...	0		53	1664
537	22:32:52.63	-60:34:00.70	29.79 ± 0.890	-0.96 ± 0.914	...	0		53	2203
538	22:32:56.18	-60:33:45.41	29.91 ± 0.667	-0.43 ± 0.714	...	0		54	1904
539	22:32:52.56	-60:34:00.86	30.01 ± 0.880	-0.19 ± 0.970	...	0		53	2210
540	22:32:57.14	-60:33:30.18	30.58 ± 2.068	0		54	1609
541	22:32:56.31	-60:33:18.48	0		32	1376
542	22:32:57.25	-60:33:23.33	0		54	-1
543	+22:32:57.47	-60:33:48.56	3.6326	2	$\text{Ly}\alpha$	54	-1
544 ^b	+22:32:56.82	-60:34:03.46	25.99 ± 0.023	0.39 ± 0.035	1.2369	3	$\text{H}\delta$, [Ne III], [O II]	54	-1
545 ^b	+22:32:52.02	-60:34:03.99	25.83 ± 0.013	0.87 ± 0.022	0.9643	2	$\text{H}\gamma$?, [O II]	53	-1
546	+22:32:53.69	-60:33:41.10	5.7102	2	$\text{Ly}\alpha$	53	-1
547	+22:32:53.94	-60:34:06.19	5.7099	2	$\text{Ly}\alpha$	53	-1
548	+22:32:55.56	-60:34:15.67	4.6886	1	$\text{Ly}\alpha$	53	-1
549	+22:32:55.63	-60:33:40.70	4.6716	2	$\text{Ly}\alpha$	54	-1
550	+22:32:52.02	-60:33:26.13	0		53	-1
551	+22:32:55.11	-60:34:01.09	3.1796	2	$\text{Ly}\alpha$	54	-1
552	+22:32:53.24	-60:33:46.96	5.0805	1	$\text{Ly}\alpha$	53	-1
553	+22:32:52.64	-60:33:55.89	5.0789	2	$\text{Ly}\alpha$	53	-1
554	+22:32:56.15	-60:33:41.53	4.5641	2	$\text{Ly}\alpha$	54	-1
555	+22:32:57.53	-60:33:54.46	4.5090	2	$\text{Ly}\alpha$	54	-1
556	22:32:55.84	-60:34:06.50	0		54	-1
557	+22:32:53.77	-60:33:48.13	5.2024	2	$\text{Ly}\alpha$	53	-1
558	+22:32:54.41	-60:34:02.23	3.1258	1	$\text{Ly}\alpha$	54	-1
559	+22:32:53.56	-60:33:50.93	4.8849	1	$\text{Ly}\alpha$	53	-1
560	+22:32:59.14	-60:33:24.36	5.8769	2	$\text{Ly}\alpha$	54	-1
561	+22:32:51.72	-60:33:39.74	4.8095	2	$\text{Ly}\alpha$	53	-1
562	+22:32:53.45	-60:33:19.13	4.8434	2	$\text{Ly}\alpha$	48	-1
563	+22:32:52.38	-60:34:01.18	3.8258	2	$\text{Ly}\alpha$	53	-1
564	+22:32:58.53	-60:33:18.89	6.0847	1	$\text{Ly}\alpha$	36	-1
565	+22:32:51.31	-60:33:27.98	0		5	-1
566 ^b	+22:32:51.64	-60:33:36.73	26.72 ± 0.031	0.14 ± 0.035	0.3178	3	[O II], contamination 8	46	-1
567 ^b	+22:32:52.03	-60:34:05.76	26.15 ± 0.023	0.56 ± 0.33	1.1389	1	[O II]	53	-1
568	+22:32:53.76	-60:33:35.55	4.6638	2	$\text{Ly}\alpha$	53	-1
569 ^b	+22:32:55.84	-60:33:32.45	27.49 ± 0.043	0.42 ± 0.059	0.4203	3	[O III], $\text{H}\beta$, $\text{H}\gamma$, [O II]	54	-1
570	22:32:52.11	-60:33:21.67	0		53	-1
571	22:32:54.78	-60:34:05.48	0		54	-1
572	22:32:54.51	-60:33:25.28	0		54	-1
573	+22:32:59.48	-60:34:12.96	6.2712	1	$\text{Ly}\alpha$	43	-1
574	+22:32:53.17	-60:33:24.74	5.7110	1	$\text{Ly}\alpha$	53	-1
575	+22:32:57.19	-60:33:35.29	0		54	-1
576	+22:32:58.21	-60:33:23.63	2.4192	1	C III]	54	-1
577	+22:32:58.07	-60:34:01.15	5.7641	1	$\text{Ly}\alpha$	54	-1
578	+22:32:55.09	-60:34:01.58	3.1800	1	$\text{Ly}\alpha$	54	-1
579	+22:32:57.96	-60:33:26.42	0		54	-1
580	+22:32:55.23	-60:33:23.41	3.6339	2	$\text{Ly}\alpha$	54	-1
581	+22:32:55.80	-60:33:43.12	3.8230	1	$\text{Ly}\alpha$	54	-1
582	+22:32:56.41	-60:33:54.53	0		54	-1
583	+22:32:54.34	-60:33:36.21	4.9151	1	$\text{Ly}\alpha$	54	-1
584	+22:32:51.83	-60:34:16.27	5.8956	2	$\text{Ly}\alpha$	53	-1
585	+22:32:57.35	-60:33:47.36	28.95 ± 0.321	0.30 ± 0.399	3.3376	1	$\text{Ly}\alpha$	54	1937

Table A.2. Line fluxes for galaxies with secure redshifts in units of 10^{-18} erg s $^{-1}$ cm $^{-2}$ Å $^{-1}$.

ID	Ly α	C III]1906, 1909	[O II]3727, 3729	[Ne III]3869	H β	[O III]5007	H α	[N II]6584
1	26.24 ± 0.29	35.63 ± 0.32	101.72 ± 0.42	14.07 ± 0.33
3	175.39 ± 0.46	6.36 ± 0.29	76.40 ± 0.90	73.61 ± 0.32
4	84.09 ± 0.92	2.60 ± 0.36	35.95 ± 1.07	31.69 ± 0.42
5	33.59 ± 0.70	1.12 ± 0.21	22.45 ± 0.28	7.29 ± 0.86
6	96.35 ± 0.43	3.60 ± 0.27	29.01 ± 0.25	44.62 ± 0.26
7	111.80 ± 0.39	4.98 ± 0.24	37.96 ± 0.22	72.52 ± 0.34
8	106.59 ± 0.96	4.07 ± 0.38	65.16 ± 0.44	34.72 ± 0.42
9	125.29 ± 0.59	3.98 ± 0.30	60.29 ± 0.89	47.57 ± 0.34
10	54.58 ± 0.65	14.51 ± 0.72
11	68.09 ± 0.39	3.06 ± 0.20	25.93 ± 0.22	37.61 ± 0.27
12	73.85 ± 0.34	4.73 ± 0.27	25.12 ± 0.26	61.73 ± 0.51
13	134.85 ± 0.57	6.70 ± 0.69
15	19.90 ± 0.81	0.33 ± 1.44
16	69.18 ± 0.33	4.87 ± 0.22	27.17 ± 0.19	92.72 ± 0.33
17	41.97 ± 1.35	2.34 ± 0.40	13.71 ± 0.81	31.86 ± 2.22
20	27.78 ± 0.34	0.54 ± 0.23	10.20 ± 0.22	23.53 ± 0.23
23	32.46 ± 0.27	1.83 ± 0.18	12.87 ± 0.50	26.88 ± 0.22
24	28.90 ± 0.36	0.80 ± 0.39
25	8.60 ± 0.48	2.47 ± 0.78
26	3.48 ± 0.16	4.03 ± 0.16	11.79 ± 0.19	1.28 ± 0.16
27	28.22 ± 0.44	2.77 ± 0.57
28	34.62 ± 0.35	2.78 ± 0.22	14.60 ± 0.25	35.99 ± 0.36	50.67 ± 1.65	1.12 ± 0.79
29	9.18 ± 0.35	0.47 ± 0.21	...	1.20 ± 0.39
32	20.88 ± 0.25	0.86 ± 0.16	8.04 ± 0.42	14.16 ± 0.22
34	2.03 ± 0.23	...	0.75 ± 0.21	0.10 ± 0.28	1.46 ± 0.49	0.60 ± 0.67
35	35.59 ± 0.46	1.94 ± 0.79
37	37.21 ± 0.66	1.23 ± 0.29
38	18.56 ± 0.44
39	15.59 ± 0.47
40	3.43 ± 0.40	1.72 ± 0.72
41	...	1.37 ± 0.54
43	19.90 ± 0.35	0.15 ± 0.46
45	9.95 ± 0.66	0.29 ± 0.40
48	22.70 ± 0.28	1.05 ± 0.25
49	15.86 ± 0.21	0.80 ± 0.21
50	...	0.58 ± 0.22
51	...	0.50 ± 0.12
53	1.22 ± 0.12	2.26 ± 0.11	4.93 ± 0.15	0.22 ± 0.13
55
56	0.63 ± 0.21	0.15 ± 0.32
57	4.53 ± 0.34	0.51 ± 0.14	1.79 ± 0.22	3.89 ± 0.25	...	2.63 ± 9.53
58	0.32 ± 0.14
60	...	0.13 ± 0.14
61	7.06 ± 0.18	0.81 ± 0.12	3.40 ± 0.11	8.44 ± 0.11	12.91 ± 0.28	0.62 ± 0.18
62	1.87 ± 0.18	0.07 ± 0.11	0.62 ± 0.11	2.96 ± 0.10	2.32 ± 0.42	...
63	0.60 ± 0.11	0.56 ± 0.11	2.76 ± 0.13	0.06 ± 0.17
64	15.14 ± 0.37
65	...	0.21 ± 0.73
67	5.72 ± 0.20	0.09 ± 0.12	2.77 ± 0.44	3.62 ± 0.10	7.48 ± 0.17	0.54 ± 0.12
68	12.21 ± 0.24	1.38 ± 0.32
70	0.69 ± 0.12	0.73 ± 0.12	3.84 ± 0.16	0.39 ± 0.13
71	23.01 ± 0.24	0.85 ± 0.33
72	7.60 ± 0.17	0.58 ± 0.10	2.69 ± 0.40	8.61 ± 0.23
73	...	0.54 ± 0.13
74	...	2.20 ± 0.16
75	2.24 ± 0.20	0.04 ± 0.07	2.88 ± 0.43	2.56 ± 0.20
78	...	0.83 ± 0.14
79	12.09 ± 0.26	2.27 ± 0.21
83
87	...	2.20 ± 0.20
88	8.50 ± 0.37	0.79 ± 0.31
89	15.00 ± 0.86
90	1.71 ± 0.09	0.14 ± 0.08	1.33 ± 0.17	5.86 ± 0.12
92	10.45 ± 0.23
95	3.71 ± 0.21
96	5.43 ± 0.27	0.60 ± 0.07

Table A.2. continued.

ID	$\text{Ly}\alpha$	$\text{C III}]1906, 1909$	$[\text{O II}]3727, 3729$	$[\text{Ne III}]3869$	$\text{H}\beta$	$[\text{O III}]5007$	$\text{H}\alpha$	$[\text{N II}]6584$
97	...	5.88 ± 0.21
98	0.51 ± 0.09	0.83 ± 0.10	2.02 ± 0.10	0.14 ± 0.20
101	2.91 ± 0.16	0.69 ± 0.11	1.45 ± 0.09	8.71 ± 0.12
104	3.25 ± 0.17	0.17 ± 0.12
107	4.43 ± 0.15	0.57 ± 0.17
109	...	1.02 ± 0.16
112	14.43 ± 0.20
114	1.83 ± 0.18	0.12 ± 0.21
115	2.82 ± 0.11	0.09 ± 0.09	1.34 ± 0.08	4.40 ± 0.11
116	...	96.23 ± 208.22
117	1.52 ± 0.31	0.01 ± 0.34	2.27 ± 0.30	4.91 ± 0.32
120	3.74 ± 0.15	0.52 ± 0.08	2.58 ± 0.34	8.06 ± 0.19
122	1.22 ± 0.16	...	0.26 ± 0.17	0.04 ± 0.21
123	0.14 ± 0.09
125	...	0.81 ± 0.55
126	...	1.72 ± 0.13
128	...	1.47 ± 0.74
129	4.78 ± 0.13	1.07 ± 0.14
135	0.77 ± 0.11	0.07 ± 0.08	0.66 ± 0.24	0.88 ± 0.09
139	6.05 ± 0.17	0.42 ± 0.38
141	1.93 ± 0.09	0.20 ± 0.07	1.38 ± 0.18	3.03 ± 0.15
144	31.14 ± 0.19
146	1.57 ± 0.27	0.28 ± 1.02
155	1.28 ± 0.21
157	1.44 ± 0.12	0.05 ± 0.07	0.53 ± 0.11	1.99 ± 0.18
159	4.31 ± 0.25
160	7.18 ± 0.35	1.30 ± 0.39
162	1.29 ± 0.16	0.15 ± 0.18
171	2.96 ± 0.29	0.12 ± 0.40	0.52 ± 0.98
181	18.15 ± 0.22
183	1.72 ± 0.21	2.23 ± 0.61
186	1.15 ± 0.12
200	1.37 ± 0.14	0.18 ± 0.38
202	10.94 ± 0.26	0.65 ± 0.32
212	...	1.58 ± 0.15
216	6.55 ± 0.16
218	2.69 ± 0.24
225	8.72 ± 0.25
229	2.10 ± 0.26
232	2.04 ± 0.15
238	0.79 ± 0.18
246	5.68 ± 0.20
255	3.47 ± 0.31	0.05 ± 0.12
257	1.45 ± 0.22
261	2.11 ± 0.25	5.09 ± 0.73
271	6.16 ± 0.38
290	4.08 ± 0.28
294	3.26 ± 0.15
308	2.26 ± 0.11
311	3.24 ± 0.15
324	1.71 ± 0.13
325	6.17 ± 0.12
334	2.77 ± 0.14
338	5.74 ± 0.25
363	0.62 ± 0.11
386	1.29 ± 0.28
393	3.22 ± 0.22
404	0.22 ± 0.26	0.32 ± 0.18	0.56 ± 0.22	1.78 ± 0.51
422	2.82 ± 0.20
430	4.77 ± 0.31
433	7.51 ± 0.19
437	7.38 ± 0.22
441	6.54 ± 0.24
449	3.59 ± 0.21
452	8.79 ± 0.67	2.37 ± 1.58
453	1.46 ± 0.16
462	3.68 ± 0.33

Table A.2. continued.

ID	Ly α	C III]1906, 1909	[O II]3727, 3729	[Ne III]3869	H β	[O III]5007	H α	[N II]6584
469	2.64 \pm 0.22
474	1.88 \pm 0.18
478	2.41 \pm 0.21
484	2.94 \pm 0.14
489	2.65 \pm 0.20	0.11 \pm 0.18
492	2.88 \pm 0.15
498	3.61 \pm 0.39
499	6.40 \pm 0.22
500	1.85 \pm 0.15	0.19 \pm 0.26
503	3.43 \pm 0.21	0.36 \pm 0.45
513	3.09 \pm 0.24
514	1.99 \pm 0.15	0.21 \pm 0.24
520	1.35 \pm 0.13
534	5.75 \pm 0.60
543	3.38 \pm 0.12
544	11.01 \pm 0.60	1.70 \pm 0.27
545	5.27 \pm 0.21
546	3.80 \pm 0.15
547	3.56 \pm 0.12
548	1.21 \pm 0.16
549	2.38 \pm 0.12
551	4.04 \pm 0.19	0.05 \pm 0.29
552	1.76 \pm 0.15
553	4.69 \pm 0.14
554	1.36 \pm 0.13
555	1.05 \pm 0.11
557	1.90 \pm 0.15
558	2.85 \pm 0.20	0.29 \pm 0.45
559	0.57 \pm 0.08
560	5.38 \pm 0.23
561	1.78 \pm 0.16
562	2.05 \pm 0.25
563	3.68 \pm 0.17	0.40 \pm 0.31
564	3.46 \pm 1.11
566	1.04 \pm 0.17	0.71 \pm 0.12	2.69 \pm 0.10	9.30 \pm 0.14	9.70 \pm 0.44	0.25 \pm 0.26
567	4.06 \pm 0.29	0.73 \pm 0.36
568	3.42 \pm 0.18
569	1.00 \pm 0.12	0.16 \pm 0.08	0.98 \pm 0.07	1.42 \pm 0.08
573	2.64 \pm 0.61
574	1.79 \pm 0.19
576	...	1.94 \pm 0.16
577	6.55 \pm 0.15
578	2.75 \pm 0.17	0.32 \pm 0.28
580	1.78 \pm 0.18
581	1.59 \pm 0.17	0.58 \pm 0.31
583	1.05 \pm 0.13
584	7.44 \pm 0.30
585	2.55 \pm 0.15

Table A.3. Redshift comparison.

id	Object	ra	Dec	m_{F814W}	z_{MUSE}	z_{lit}	Literature	Comments
2	HDFS J223258.30-603351.7	22:32:58.30	-60:33:51.66	21.3571	0.0	0.7063	G06	star
3	HDFS J223253.74-603337.6	22:32:53.74	-60:33:37.57	21.521	0.564497	0.5645	S03	–
4	HDFS J223252.14-603359.6	22:32:52.14	-60:33:59.56	21.778	0.564616	0.5646	S03	–
5	HDFS J223252.24-603402.7	22:32:52.24	-60:34:02.75	21.9726	0.580445	0.5804	S03	–
6	HDFS J223258.22-603331.6	22:32:58.21	-60:33:31.64	21.9835	0.423033	0.4229	S03	Figure 19
7	HDFS J223259.43-603339.8	22:32:59.43	-60:33:39.82	21.9948	0.464372	0.4644	S03; G06	–
9	HDFS J223256.08-603414.2	22:32:56.08	-60:34:14.17	22.0849	0.56453	0.5645	S03	–
10	HDFS J223253.03-603328.5	22:32:53.03	-60:33:28.53	22.5629	1.284517	1.27	R05	–
13	HDFS J223252.16-603323.9	22:32:52.16	-60:33:23.92	22.8323	1.290184	1.293	R05	Figure 6
15	HDFS J223252.88-603317.1	22:32:52.88	-60:33:17.12	22.8565	1.284184	1.284	W09	–
16	HDFS J223255.24-603407.5	22:32:55.25	-60:34:07.50	22.8682	0.465384	0.4656	S03	–
17	HDFS J223255.87-603317.8	22:32:55.87	-60:33:17.75	22.926	0.581722	0.5817	S03	–
20	HDFS J223257.90-603349.1	22:32:57.90	-60:33:49.12	23.0644	0.428094	0.428	S03	–
23	HDFS J223255.75-603333.8	22:32:55.74	-60:33:33.75	23.4153	0.564872	0.5649	S03; I05	–
41	HDFS J223254.17-603409.1	22:32:54.18	-60:34:08.94	24.571	2.4061	2.412	I05; W09	–
43	HDFS J223252.03-603342.6	22:32:52.03	-60:33:42.59	24.6219	3.29254	3.295	I05; W09	Figure 7
55	HDFS J223253.12-603320.3	22:32:53.11	-60:33:20.25	24.9427	2.67	2.67	I05; W09	–
87	HDFS J223254.87-603342.2	22:32:54.86	-60:33:42.12	25.7363	2.67	2.676	W09	–

References. Sawicki & Mallén-Ornelas (2003, S03), Rigopoulou et al. (2005, R05), Iwata et al. (2005, I05), Glazebrook et al. (2006, G06), and Wuyts et al. (2009, W09).

Appendix B: Public data release

In addition to the source catalogue, we release the reduced data cube and associated files. We also deliver spectra and reconstructed images in the main emission lines for all the catalogue sources⁴.

⁴ All this material is available at <http://muse-vlt.eu/science>