

Growth Until Puberty After In Utero Exposure to Coumarins

D. Van Driel,¹ J. Wesseling,¹ F.R. Rosendaal,² R.J. Odink,¹ E. Van der Veer,^{3*} W.J. Gerver,⁴ L.M. Geven-Boere,⁵ and P.J.J. Sauer¹

¹Department of Pediatrics, Beatrix Children's Hospital, University Hospital Groningen, Groningen, The Netherlands

²Departments of Hematology and Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

³Pathology and Laboratory Medicine, University Hospital Groningen, Groningen, The Netherlands

⁴Department of Pediatrics, University Hospital Maastricht, Maastricht, The Netherlands

⁵Dutch Federation of Thrombosis Services, The Hague, The Netherlands

Anticoagulation with coumarins is an effective therapy during pregnancy. Fetal exposure to coumarin derivatives during the first trimester, however, is associated with skeletal anomalies (warfarin or coumarin embryopathy). Information about long-term effects of prenatal coumarin exposure on the skeletal development is not available. We investigated growth and body proportions at school age of children exposed to coumarins in utero. A blind population-based cohort study was conducted on 307 exposed children and 267 non-exposed controls ages 8–15 years. The exposed cohort was based on a prospective registry of coumarin-treated pregnant women. Anthropometric data included height, weight, head circumference, and measurements to evaluate body proportions. The mean height of exposed children did not differ from that of the non-exposed children (mean difference 0.01 SD). In addition, no differences were found for the proportional measures. As a group, children exposed in the first trimester showed no evidence of growth impairment. Two children in this group, however, were born with signs of coumarin embryopathy and one of these displayed a deficit in height at school age. Long-term growth was not affected by a high cumulative dosage or exposure after the first trimester. We conclude that, when exposure during the first trimester is avoided, coumarin therapy

during pregnancy has no demonstrable risk for the child's skeletal development. *Am. J. Med. Genet.* 95:438–443, 2000.

© 2000 Wiley-Liss, Inc.

KEY WORDS: coumarin embryopathy; long-term growth; anticoagulants; pregnancy

INTRODUCTION

Anticoagulant therapy during pregnancy is indicated for the treatment and prophylaxis of thrombo-embolic disease and prevention of systemic embolism in patients with valvular heart disease or prosthetic heart valves [Ginsberg, 1989]. The prescription of anticoagulants during pregnancy must be carefully considered because of adverse effects in both mother and child. Heparins have the potential for causing adverse effects in the mother when phenprocoumon and warfarin are used. Coumarin therapy during pregnancy is associated with birth defects [Bates and Ginsberg, 1997; Nelson-Piercy, 1997]. Centers in Europe often use the protocol that prescribes heparin during the first trimester of gestation and from the 36th week until birth, whereas coumarins are prescribed during the second and third trimester of pregnancy [Lecuru et al., 1996; Harenberg, 1998]. In the United States, heparin is preferred throughout pregnancy because there is great concern about the teratogenic effect of coumarins [Ginsberg and Barron, 1994; Togliola and Weg, 1996].

Case reports describe central nervous system [Kaplan et al., 1982; Lapiedra et al., 1986] and skeletal abnormalities [Tamburrini et al., 1987; Howe et al., 1997; Wellesley et al., 1998] after in utero exposure to coumarins. The bone defects constitute the so-called 'coumarin (or warfarin) embryopathy,' whereas the term 'coumarin syndrome' refers to the central nervous system anomalies. Reported skeletal anomalies observed shortly after birth include various degrees of nasal hypoplasia, rhizomelia, hypoplastic distal phalanges and stippling in epiphyses. There is one report

Grant sponsor: Dutch Heart Association; Grant number: 94.148; Contract grant sponsor: Praeventiefonds; Grant number: 002824340.

*Correspondence to: E. van der Veer, Pathology and Laboratory Medicine, University Hospital Groningen, PO Box 30.001, 9700 RB Groningen, The Netherlands.
E-mail: e.van.der.veer@med.rug.nl

Received 1 November 1999; Accepted 15 August 2000

of a prenatally exposed child who showed disturbed growth during childhood resulting in a disproportionately short stature at adult age [Hosenfeld and Wiedeman, 1989]. In a review of published cases, it was assumed that exposure during the first trimester of pregnancy, especially from the 6th to the 9th week, confers the greatest risk for developing the coumarin embryopathy [Hall et al., 1980].

Animal studies confirm an effect of warfarin exposure on growth. Rats showed a decreased body length after 12 weeks of postnatal warfarin treatment that was correlated with calcium bridges in epiphyses [Howe and Webster, 1992]. Chronic exposure of developing rats caused growth plate closure resulting in cessation of growth [Price et al., 1982]. It is not clear whether these findings resemble the skeletal abnormalities seen in humans.

So far, growth at school age has not been examined in children who were prenatally exposed to coumarins. Knowledge of long-term effects for the child's development are essential for a careful consideration of anticoagulant therapy in pregnant women. We performed a large cohort study to assess late effects in children exposed to coumarins in utero. The aim of the present study is to determine whether prenatal exposure to coumarin derivatives causes a (disproportionally) short stature or other growth impairment at school age.

SUBJECTS AND METHODS

Subjects

Two groups are included in the study: a cohort of children exposed in utero to coumarin derivatives and a cohort of non-exposed controls. The coumarin-exposed children were offspring of mothers included during pregnancy in a registry by Dutch anticoagulation clinics, charged with monitoring oral anticoagulant therapy in out-patients. Inclusion criteria for the exposed study group were: consent for registration, prescription of coumarin derivatives during pregnancy and childbirth between January 1st, 1982 and December 31st, 1990. Eligible registered women were approached by the anticoagulation clinics, either directly or after consultation with the family's general practitioner. Non-exposed control children were approached by regional vaccination centres, keeping records of the inoculation of all children in The Netherlands. Matching was done for age ($\pm 1/2$ year) and gender to obtain an equal distribution in both study cohorts, and for demographic region (postal code) as a measure of socio-economic status. Exclusion criteria were diseases interfering with growth that are not related to coumarin exposure, including Down syndrome.

The study protocol was approved by the Medical Ethics Committee of the University Hospital Groningen and the parents of all children gave written informed consent.

Data Collection

After enrollment in the study, all children received an appointment for physical examination. Puberty assessment included examination of secondary sexual characteristics according to Tanner [1962]. We consid-

ered breast development Stage 2 in girls and mean testicular volume of 5 ml in boys as minimum criteria for the onset of puberty.

Information about pregnancy, delivery, and the medical history of the child were obtained by questionnaire. We inquired about parental height and weight as references for target height and Body Mass Index of the child. Maternal education and paternal occupation, the latter classified according to Sixma and Ultee [1983], were regarded as measures of socio-economic status.

For the exposed cohort, information about indication, period, coumarin derivative, and prescribed dosage during pregnancy was collected from the initial registration form together with medical records from anticoagulation clinics and gynecologists.

Anthropometric Measurements

The anthropometric data were collected by a trained observer who was not aware of the exposure status of the child. Measurements were performed on subjects wearing light clothing (underwear) and repeated three times, measuring to the nearest 0.1 cm. Height and sitting height were measured by two observers; for sitting height the child was placed on a standard stool with his feet on a footrest. Head circumference was measured with a non-stretchable tape at the most prominent points of occiput and forehead. Arm span was measured (with a non-elastic measuring rod) as the distance between the tips of the stretched middle fingers, the child standing with his arms fully extended. The different skeletal parts were measured on the left side of the body using bony prominences as anatomical landmarks. Upper arm length was measured from the lateral border of the acromion to the head of the radius, lower arm length from the head to the bottom of the radius, and hand length from the bottom of the radius to the tip of the longest finger. Biacromial diameter was measured between the lateral borders of the acromion processes and bi-iliac diameter at the widest point of the iliac crests, using the anterior superior spina iliaca as a landmark. The distance from the proximal medial border of the tibia to the distal border of the medial malleolus was used for tibia length; the distance between the most posterior part of the heel and the tip of the longest toe was used for the foot length. Weight was measured in kilograms, using a digital scale with an accuracy of 100 g.

Statistical Analysis

To adjust for gender and age, we calculated SD-scores using Dutch references [Gerver and De Bruin, 1996]. Values were expressed in means \pm SD. For the comparison of weight, Body Mass Index (kg/m^2) was calculated. Statistical differences between exposed and non-exposed children were assessed by linear regression analysis, controlling for the confounding variables gender, age, target height, puberty, and socio-economic status.

Exposure to coumarins was classified according to trimester (exposure at least during the first trimester of pregnancy or exposure only during the second or

third trimester), derivative (acenocoumarol or phenprocoumon), and daily dosage. In addition, cumulative dosage (duration (days) calculated from start and stop date of therapy multiplied by mean daily dosage) was used to investigate the combined effect of dosage and duration of exposure. To disentangle the effect of coumarin exposure from the effect of maternal disease, we performed a stratified analysis by indication for coumarin treatment during pregnancy. The significance level was set at $P < 0.05$. Statistical analysis was performed with Statistical Package for the Social Sciences (SPSS Inc., Chicago, IL).

RESULTS

Study Population

A total of 574 children were enrolled in the study, 307 in the exposed and 267 in the non-exposed cohort, respectively. For the exposed cohort, 451 pregnancies registered by Dutch anticoagulation clinics were considered for inclusion. Fifty-five of these women could not be traced because of incomplete personal data. Of 14 families, the general practitioner discouraged contact because of death or severe illness of the mother ($n = 6$) or because of other familial circumstances ($n = 5$). In three cases, the general practitioner did not give an explication, but the child concerned was healthy. The remaining 382 registered pregnancies proved eligible for inclusion; 307 of these children actually participated in the study (response rate 80%). In 75 cases, participation was refused because the child was not willing to cooperate ($n = 15$), the mother had no contact with the child because of divorce ($n = 6$), 'taking part costed too much time' ($n = 8$), or participation in the planned investigation period was not possible due to familial circumstances ($n = 4$). Forty-three parents declined participation without explanation.

Characteristics of the exposed and non-exposed cohort are presented in Table I. Gestational age was lower in the exposed study cohort probably related to a higher percentage of labour induction in this high risk group (31% vs. 12%, RR = 2.5 CI₉₅ 1.7 to 3.5). Birth weight and length, when adjusted for gestational age, were not significantly different. Age at study was similar in both groups ranging from 7.6 to 15.3 years. Mean age at the various stages of puberty, both in boys and in girls, were not significantly different for the exposed and non-exposed cohort. For four children (three exposed, one non-exposed) the level of maternal education could not be identified.

Exposure

Indications for coumarin derivatives during pregnancy were treatment ($n = 64$) and prophylaxis ($n = 207$) of thrombo-embolic events, hereditary thrombophilia ($n = 11$), artificial heart valve ($n = 10$), and a group of incidental causes like antiphospholipid antibody syndrome, trauma, and surgery ($n = 9$); for six cases the indication was unknown.

Coumarin derivative and exposure period during pregnancy could be identified for 281 children (92%). Most children ($n = 240$) were exposed during the second or third trimester of pregnancy. The first trimester

TABLE I. Characteristics of Study Population

	Non-exposed	Exposed
Number of patients	267	307
Gender: male	53%	50%
Caucasian origin	93%	93%
Gestational age (wks)	39.4 (± 1.8) [†]	38.9 (± 2.0)*
Birthweight (grams)	3380 (± 598)	3229 (± 577)
Length at birth (cm)	50.2 (± 2.8)	49.7 (± 3.0)
Mean age in yrs (sd)	10.5 (± 2.0)	10.7 (± 2.0)
Puberty:		
prepubertal	161 (60%)	165 (54%)
pubertal	85 (32%)	120 (39%)
unknown	21 (8%)	22 (7%)
Socio-economic status:		
Education mother		
untrained	3 (1%)	4 (1%)
low	118 (44%)	137 (45%)
middle	92 (34%)	108 (35%)
high	53 (20%)	55 (18%)
Occupation father		
unknown/single parent	18 (7%)	14 (5%)
none	10 (4%)	16 (5%)
low	69 (26%)	78 (25%)
middle	108 (40%)	129 (42%)
high	62 (23%)	70 (23%)

* $P = 0.01$, Mann Whitney U-test

study group was comprised of 41 children, of which 20 were exposed during (part of) the teratogenic window (6th to 9th gestational week). The duration of exposure ranged from one to 36 weeks (mean: 16 weeks). In most cases ($n = 239$) the short-acting coumarin derivative acenocoumarol was prescribed, whereas 33 mothers used phenprocoumon. Seven women changed coumarin derivative during pregnancy and in two cases preparation could not be retrieved.

The prescribed dosage during pregnancy was available for 166 subjects (54%) of the exposed cohort. Mean daily dosage was 3.3 mg (± 1.14) for the group exposed to acenocoumarol and 3.3 mg (± 1.20) for the group exposed to phenprocoumon, respectively. Mean cumulative dosage including both derivatives was 408.7 \pm 231.2 mg (range: 34.1–1,501.0 mg).

Growth

The mean and standard deviation of the calculated SD-scores of height, sitting height, the proportional measure sitting height/height and the other growth parameters of exposed children and non-exposed controls are shown in Table II. Using linear regression analysis we found no differences for these parameters between both groups. Body Mass Index was not different between the exposed and non-exposed cohort (mean difference = -0.01 , CI₉₅ -0.42 to 0.41). Children exposed during the first trimester of pregnancy showed no difference in height compared with children exposed during the second or third trimester and compared with non-exposed controls (Fig. 1). Two children exposed during part of the teratogenic window (6th to 9th gestational week) were born with signs of coumarin embryopathy. One of these children was born at 32 weeks manifesting severe growth retardation (SD-score of birth length -3.78 and SD-score of birthweight -1.72 both corrected for gestational age). At the time of

Height-for-age

5.12
4.80

TABLE II Mean SD-Score (Standard Deviation) of Growth Parameters for Coumarin-Exposed and Non-Exposed Children

	Non-exposed (n = 267)	Exposed (n = 307)	Difference ¹	CI ₉₅
Height	0.09 (0.98)	0.20 (1.0)	0.01	-0.14 to 0.14
Sitting height	-0.01 (0.99)	0.07 (1.1)	-0.01	-0.16 to 0.14
Sitting height/height	-0.48 (0.98)	-0.44 (1.1)	0.05	-0.14 to 0.23
Head circumference	-0.07 (0.98)	0.05 (1.1)	0.03	-0.14 to 0.20
Arm span	0.06 (1.0)	0.17 (1.1)	0.03	-0.13 to 0.18
Upper arm length	-0.34 (1.1)	-0.26 (1.1)	<-0.01	-0.17 to 0.16
Lower arm length	1.04 (1.3)	1.17 (1.3)	0.1	-0.11 to 0.30
Hand length ²	-0.55 (0.96)	-0.37 (1.1)	0.05	-0.12 to 0.23
Biacromial diameter	0.33 (1.2)	0.43 (1.2)	0.08	-0.12 to 0.28
Buliacal diameter	0.67 (1.4)	0.73 (1.5)	-0.04	-0.27 to 0.20
Tibia length	0.14 (1.1)	0.20 (1.2)	<0.01	-0.18 to 0.18
Foot length	0.08 (1.0)	0.20 (1.1)	0.05	-0.11 to 0.22

¹Mean difference derived from linear regression analysis comparing exposed versus non exposed children

study, this 11-year-old girl displayed a short stature (SD-score of height-for-age: -1.96, SD-score corrected for target height: -2.79) with relatively short legs (SD-score of sitting height/height: 1.43). The other child was born at 41 weeks with a normal birth length (SD-score corrected for gestational age: 1.56). At 13 years of age, he displayed a normal stature (SD-score of height-for-age: 2.65; SD-score corrected for target height: 1.52), with relatively long legs (SD-score of sitting height/height: -1.15). Figure 2 shows that there is no relationship between cumulative dosage of in utero exposure to coumarin derivatives and height at school age. Using linear regression analysis, we found that prescribed preparation of coumarin therapy did not have an influence on long-term height (mean difference (d) <-0.001 SD for each mg of acenocoumarol and d < 0.001 SD for each mg phenprocoumon). Stratified analysis showed no differences in height between children whose mothers were treated for acute thromboembolism (d = 0.17, CIs -0.08 to 0.43), prophylaxis of thromboembolism (d = -0.06, CIs -0.22 to 0.11), arti-

ficial heart valve (d = 0.45, CIs -0.15 to 1.04), or hereditary thrombophilia (d = 0.22, CIs -0.34 to 0.78) compared with non-exposed control children.

DISCUSSION

Our study demonstrates that children exposed to coumarins during pregnancy who are born without signs of coumarin embryopathy do not differ in growth with age-matched non-exposed controls. At school age, we found no differences in various growth parameters and body proportions between children exposed in utero to the coumarins acenocoumarol or phenprocoumon and non-exposed controls. In addition, time of onset of puberty did not differ between the exposed and non-exposed children. On the other hand, we found two children who were exposed to coumarins in the first trimester of pregnancy and were born with manifestations of coumarin embryopathy. One of these children was growth retarded at school age.

One of the hypotheses about the pathogenesis of cou-




Fig 1 Boxplot of the SD score height for age for non-exposed and coumarin-exposed children in the different exposure periods during pregnancy




Fig 2 Scatterplot of the SD score height for age vs cumulative dosage of coumarins during pregnancy comparing non-exposed (cumulative dosage 0 mg) and coumarin-exposed (cumulative dosage 34.1-1,501.0 mg) children

marin embryopathy proposes a decreased production of Vitamin K-dependent mineralization inhibitors in cartilage resulting in ectopic calcium deposits in epiphyses and nasal septum. Experiments in rats showed maxillofacial hypoplasia together with ectopic calcium deposits in the septal cartilage of the nasal septum and in epiphyseal growth plates after warfarin treatment [Howe and Webster, 1992]. Another theory states that warfarin inhibits the activity of a novel gene, arylsulfatase E, that is assumed to play an important role in the metabolic pathway of some types of chondrodysplasia punctata [Franco et al., 1995].

Exposure to coumarin derivatives during the first trimester of gestation, especially from the 6th to the 9th week, carries the greatest risk for inducing skeletal anomalies. Signs of coumarin embryopathy are described in association with exposure to warfarin as well as acenocoumarol and phenprocoumon [Harrod and Sherrod, 1981; Struwe et al., 1984; De Vries et al., 1993]; reported prevalence rates vary considerably from 0–68% [Iturbe-Alessio et al., 1986; Wong et al., 1993; Sbarouni and Oakley, 1994]. In one case report, an adult with a disproportionate short stature was described after exposure to phenprocoumon during the first trimester [Hosenfeld and Wiedeman, 1989]. In our study, 41 children were exposed during the first trimester of gestation; 20 of these were exposed during (part of) the proposed teratogenic window. Two of the children who were exposed during the critical period, were born with signs of coumarin embryopathy. This suggests a prevalence rate of 10%. Most children ($n = 38$) were exposed to acenocoumarol, whereas three were exposed to phenprocoumon. One child exposed in the critical period was born with signs of classic coumarin embryopathy (nasal hypoplasia and stippled epiphyses) that was diagnosed neonatally. Another child who was also exposed during the teratogenic window showed manifestations (nasal hypoplasia with breathing problems) after birth that retrospectively suggested an embryopathy. As a group, children exposed in the first trimester showed no evidence of long-term growth impairment. The child with classic coumarin embryopathy, however, showed a short stature with relatively short legs at the time of study. This child, born to a mother suffering from systemic lupus erythematosus, displayed severe growth retardation (-3.78 SDS) at birth. Although it is difficult to rule out other influences, the coumarin exposure may have caused the intrauterine growth retardation, as well as the subsequent growth impairment. The other child with signs of embryopathy at birth showed a normal proportional stature at the time of study. The children in our study who were exposed in the first trimester and were born without overt manifestations of embryopathy had a normal growth at school age.

The vast majority of our study population was exposed during the second and third trimester of pregnancy. We found no evidence that a high cumulative dosage of prenatal exposure to coumarin derivatives has an effect on growth at school age. The mothers of these exposed children used either acenocoumarol or phenprocoumon during pregnancy, with a mean daily dosage of 3.3 mg. Warfarin-treated postnatal rats

showed a decreased growth that was related to the exposure duration [Price et al., 1982; Howe and Webster, 1992]. In these experiments, the rats were treated with warfarin in high dosages ranging from 77–100 mg/kg body weight. Differences in dosage and pharmacokinetics may explain the different outcome of our study compared with those in the rat experiments.

Chong et al. [1984] performed a follow-up study to assess the growth and development of coumarin-exposed children compared with non-exposed controls at 4 years of age. They found no differences in centile distribution of height between the two study groups. This study was small ($n = 22$) and no correction for target height was made.

The exposed cohort of our study was based upon a prospective registry of women treated with coumarins during pregnancy. The response rate of parents and children that were approached was high (80%). Of the 75 parents who refused to participate in the study, in 32 cases the reason not to participate was not related to the condition of the child. In 43 cases, parents refused to cooperate without an explication. Due to privacy reasons, it was not possible to inquire about the reasons to decline. Taking part in the study was time consuming; the investigation was carried out during office hours and lasted for one hour and a half. Because refusal without explication regarded only 11% of the approached parents and children ($n = 382$), the possibility of selection bias is unlikely.

The present study is the first large cohort study to date that investigates growth at school age in children who were prenatally exposed to coumarins. We found that exposure to the coumarin derivatives acenocoumarol and phenprocoumon during the second or third trimester of pregnancy has no effect on growth at school age. This implies that coumarin therapy during pregnancy that adheres to a protocol that avoids treatment during the first trimester has no demonstrable risk for the skeletal growth of the exposed child.

ACKNOWLEDGMENTS

We are grateful to the families and children who participated in the study and to the Dutch Federation of Thrombosis Services, the various anticoagulation clinics and regional vaccination administrations for their help in approaching eligible participants. Furthermore, we would like to thank Mrs. Siekmans for her administrative assistance and the various hospitals for providing study accommodation. Last but not least, we would like to thank Prof. B.C.L. Touwen, M Smrkovsky and Prof. H.S.A. Heymans for their support and critical comments.

REFERENCES

- Bates SM, Ginsberg JS 1997 Anticoagulants in pregnancy fetal effects. *Baillieres Clin Obstet Gynecol* 11 479–488
- Chong MKB, Harvey D, De Swiet M 1984 Follow-up study of children whose mothers were treated with warfarin during pregnancy. *Br J Obstet Gynecol* 91 1070–1073
- De Vries TW, Van der Veer E, Heymans HSA 1993 Warfarin embryopathy patient, possibility, pathogenesis and prognosis. *Br J Obstet Gynecol* 100 869–871

- Franco B, Meroni G, Parenti G, Levilliers J, Bernard L, Gebbia M, Cox L, Maroteaux P, Sheffield L, Rappold G, Andria G, Petit C, Ballabio A 1995 A cluster of sulfatase genes on Xp22.3 mutations in *Chondrodysplasia Punctata* (CDPX) and implications for warfarin embryopathy *Cell* 81 15-25
- Gerver WJM, De Bruin R 1996 Pediatric morphometrics Utrecht, The Netherlands Bunge
- Ginsberg JS 1989 Anticoagulants during pregnancy *Ann Rev Med* 40 79-86
- Ginsberg J, Barron W 1994 Pregnancy and prosthetic heart valves *Lancet* 344 1170-1172
- Hall JG, Pauli RM, Wilson KM 1980 Maternal and fetal sequelae of anticoagulation during pregnancy *Am J Med* 68 122-140
- Harenberg J 1998 Antikoagulation bei Patienten mit Herzklappenersatz in der Schwangerschaft *Z Kardiol* 87 63-67
- Harrod JE, Sherrod PS 1981 Warfarin embryopathy in siblings *Obstet Gynecol* 57 673-676
- Hosenfeld D, Wiedeman HR 1989 Chondrodysplasia punctata in an adult recognized as vitamin K antagonist embryopathy *Clin Genet* 35 376-381
- Howe AM, Lipson AH, De Silva M, Ouvrier R, Webster WS 1997 Severe cervical dysplasia and nasal cartilage calcification following prenatal warfarin exposure *Am J Med Genet* 71 391-396
- Howe AM, Webster WS 1992 The warfarin embryopathy a rat model showing maxillofacial hypoplasia and other skeletal disturbances *Teratology* 46 379-390
- Iturbe-Alessio I, Del Carmen Fonseca M, Mutchnik O, Angel Santos M, Zajaras A, Salazar E 1986 Risks of anticoagulant therapy in pregnant women with artificial heart valves *N Engl J Med* 315 1390-1393
- Kaplan LC, Anderson GG, Ring BA 1982 Congenital hydrocephalus and Dandy-Walker malformation associated with warfarin use during pregnancy *Birth Defect Orig Art Ser* 18 79-83
- Lapiedra OJ, Bernal JM, Ninot S, Gonzalez I, Pastor E, Miralles PJ 1986 Open heart surgery for thrombosis of a prosthetic mitral valve during pregnancy Fetal hydrocephalus *J Cardiovasc Surg* 27 217-220
- Lecuru F, Desnos M, Taurelle R 1996 Anticoagulant therapy in pregnancy Report of 54 cases *Acta Obstet Gynecol Scand* 75 217-221
- Nelson-Piercy C 1997 Hazards of heparin allergy, heparin-induced thrombocytopenia and osteoporosis *Baillieres Clin Obstet Gynecol* 11 489-509
- Price PA, Wilhamson MK, Haba T, Dell RB, Jee WSS 1982 Excessive mineralization with growth plate closure in rats on chronic warfarin treatment *Proc Natl Acad Sci USA* 79 7734-7738
- Sbarouni E, Oakley CM 1994 Outcome of pregnancy in women with valve prostheses *Br Heart J* 71 196-201
- Sixma H, Ultee W 1983 Een beroeps-prestigeschaal voor Nederland in de jaren tachtig *Mens en Maatschappij* 4 360-381
- Struwe FE, Reinwein H, Stier R 1984 Coumarin embryopathie *Radiologe* 24 68-71
- Tamburrini O, Bartolomeo De Iuri A, Di Guglielmo GL 1987 Chondrodysplasia punctata after warfarin case report with 18 month follow up *Pediatr Radiol* 17 323-324
- Tanner JM 1962 Growth at adolescence Oxford Blackwell Scientific Publications
- Togha M, Weg J 1996 Venous thromboembolism during pregnancy *N Engl J Med* 335 108-114
- Wellesley D, Moore I, Heard M, Keeton B 1998 Two cases of warfarin embryopathy a re-emergence of this condition? *Br J Obstet Gynecol* 105 805-806
- Wong V, Cheng CH, Chan KC 1993 Fetal and neonatal outcome of exposure to anticoagulants during pregnancy *Am J Med Genet* 45 17-21