



Universiteit  
Leiden  
The Netherlands

## Competition between CO and N<sub>2</sub> desorption from interstellar ices

Oberg, K.; Broekhuizen, F.A. van; Bisschop, S.E.; Dishoeck, E.F. van; Schlemmer, S.; Fraser, H.J.

### Citation

Oberg, K., Broekhuizen, F. A. van, Bisschop, S. E., Dishoeck, E. F. van, Schlemmer, S., & Fraser, H. J. (2005). Competition between CO and N<sub>2</sub> desorption from interstellar ices. *Astrophys. J.*, 621, L33-L36. Retrieved from <https://hdl.handle.net/1887/2218>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/2218>

**Note:** To cite this publication please use the final published version (if applicable).

## COMPETITION BETWEEN CO AND N<sub>2</sub> DESORPTION FROM INTERSTELLAR ICES

K. I. ÖBERG,<sup>1,2</sup> F. VAN BROEKHUIZEN,<sup>1</sup> H. J. FRASER,<sup>1</sup> S. E. BISSCHOP,<sup>1</sup> E. F. VAN DISHOCK,<sup>1</sup> AND S. SCHLEMMER<sup>1</sup>

Received 2004 November 8; accepted 2005 January 18; published 2005 February 2

### ABSTRACT

Millimeter observations of pre- and protostellar cores show that the abundances of the gas-phase tracer molecules, C<sup>18</sup>O and N<sub>2</sub>H<sup>+</sup>, anticorrelate with each other and often exhibit “holes” where the density is greatest. These results are reasonably reproduced by astrochemical models, provided that the ratio between the binding energies of N<sub>2</sub> and CO,  $R_{BE}$ , is taken to be between 0.5 and 0.75. This Letter is the first experimental report of the desorption of CO and N<sub>2</sub> from layered and mixed ices at temperatures relevant to dense cores, studied under ultrahigh vacuum laboratory conditions using temperature programmed desorption. From control experiments with pure ices,  $R_{BE} = 0.923 \pm 0.003$ , given  $E_b(\text{N}_2\text{-N}_2) = 790 \pm 25$  K and  $E_b(\text{CO-CO}) = 855 \pm 25$  K. In mixed (CO:N<sub>2</sub> = 1:1) and layered (CO above or below N<sub>2</sub>) ice systems, both molecules become mobile within the ice matrix at temperatures as low as 20 K and appear miscible. Consequently, although a fraction of the deposited N<sub>2</sub> desorbs at lower temperatures than CO, up to 50% of the N<sub>2</sub> molecules leave the surface as the CO itself desorbs, a process not included in existing gas-grain models. This codesorption suggests that for a fraction of the frozen-out molecules,  $R_{BE}$  is unity. The relative difference between the CO and N<sub>2</sub> binding energies as derived from these experiments is therefore significantly less than that currently adopted in astrochemical models.

*Subject headings:* astrochemistry — infrared: ISM — ISM: molecules — methods: laboratory

*Online material:* color figure

### 1. INTRODUCTION

Dense molecular clouds are the starting point in our understanding of star formation. These regions evolve toward characteristically cold ( $\leq 10$  K), centrally concentrated, prestellar cores, which then collapse to form protostars. The fundamental properties of these cores are determined almost exclusively from observations of optically thin gas-phase tracers. However, as cores evolve toward the collapse phase, densities increase and the timescales for volatile gas-phase molecules to deplete onto grains become shorter than the age of the core (e.g., Caselli et al. 1999; Walmsley et al. 2004). Of the critical processes governing chemical differentiation in starless cores that are not yet well understood, a key issue is the relative freezeout and desorption rates of CO and N<sub>2</sub> (Bergin & Langer 1997, hereafter BL97).

C<sup>18</sup>O accurately traces the distribution of material in cold, dense cores, except in regions of high extinction ( $A_v \geq 10$  mag; e.g., Kramer et al. 1999). When CO becomes frozen out onto the surfaces of cold dust grains, a corresponding drop is observed in the gas-phase C<sup>18</sup>O abundance (e.g., Bergin et al. 2001; Tafalla et al. 2002; Pagani et al. 2005), and solid <sup>12</sup>CO is detected (e.g., Chiar et al. 1998; Whittet et al. 1998 and references therein). Unfortunately, N<sub>2</sub> is not directly observable in the gas phase, and the vibrational band strength of solid <sup>14</sup>N<sub>2</sub> is around 5 orders of magnitude lower than the corresponding band strength of <sup>12</sup>CO, making its detection in interstellar ices impossible (Sandford et al. 2001). The N<sub>2</sub> gas abundance is therefore inferred from observations of its chemical “daughters,” such as N<sub>2</sub>H<sup>+</sup>.

The detailed chemical network of N<sub>2</sub>H<sup>+</sup> formation and destruction is described in detail by Jørgensen et al. (2004), showing that the chemistries of CO, N<sub>2</sub>, N<sub>2</sub>H<sup>+</sup>, HCO<sup>+</sup>, and H<sub>3</sub><sup>+</sup> are inti-

mately entwined. N<sub>2</sub>H<sup>+</sup> is considered a more suitable gas-phase tracer than C<sup>18</sup>O because its abundance can remain constant when CO is depleted (Bergin et al. 2001; Tafalla et al. 2002; Di Francesco et al. 2004; Pagani et al. 2005; Jørgensen 2004). However, depletion of N<sub>2</sub>H<sup>+</sup> (and thus N<sub>2</sub>) has been detected within the densest regions of the prestellar B68 dark cloud (Bergin et al. 2002), around the Class 0 source IRAM 04191 +1522 (Belloche & André 2004), within starless cores in L183 (Pagani et al. 2005), and in a range of sources in Oph A (Di Francesco et al. 2004).

In gas-grain models, the relative abundances of gas-phase CO and N<sub>2</sub> are governed by the balance of the freezeout timescales—which depend mostly on density—and the desorption timescales—which depend mostly on the grain temperature, binding energies, and desorption kinetics. Models reasonably assume that for both molecules, and at grain temperatures lower than 10–15 K, all sticking probabilities are very close to 1, so the freezeout timescales are identical for a given grain density. By using first-order desorption kinetics, with preexponential factors,  $\nu_0 \approx 10^{12}$  s<sup>-1</sup>, models can reproduce the N<sub>2</sub>H<sup>+</sup> and CO observations by assuming that the difference between the desorption rates of CO and N<sub>2</sub> is a direct function of the ratios between their binding energies,  $R_{BE} = E_b(\text{N}_2\text{-N}_2)/E_b(\text{CO-CO})$ . For example, BL97 set  $R_{BE}$  at 0.65, based on the ab initio calculations of Sadlej et al. (1995) and assuming  $E_b(\text{CO-CO}) = 960$  K. Subsequently, Bergin et al. (2002) used an extension of the BL97 chemical model to reproduce their observations of N<sub>2</sub>H<sup>+</sup> in B68, with initial values for the binding energies of CO-CO and N<sub>2</sub>-N<sub>2</sub> of 1210 and 750 K, respectively ( $R_{BE} = 0.62$ ), although best fits to the emission spectra were actually obtained with  $E_b(\text{N}_2\text{-N}_2) = 900$  K ( $R_{BE} = 0.75$ ). Conversely, Aikawa et al. (2003) assume that the value  $R_{BE}$  is even lower than that used in BL97. The range of binding energies applied in each case arises owing to a lack of relevant laboratory data in the literature.

To accurately model the entire CO-N<sub>2</sub> gas-grain system, a minimum of five binding energies need to be known, namely, those for N<sub>2</sub>-N<sub>2</sub>, CO-N<sub>2</sub>, H<sub>2</sub>O-N<sub>2</sub>, CO-CO, and H<sub>2</sub>O-CO, together with a knowledge of the order of reaction and pre-exponential factor in the rate equations describing the desorp-

<sup>1</sup> Raymond and Beverly Sackler Laboratory for Astrophysics at Leiden Observatory, Postbus 9513, 2300 RA, Leiden, Netherlands; fraser@strw.leidenuniv.nl.

<sup>2</sup> Current address: Division of Geological and Planetary Sciences, California Institute of Technology, Mail Stop 150-21, Pasadena, CA 91125.




FIG. 1.—TPD spectra of (a) pure CO (40 L) and N<sub>2</sub> (40 L), (b) an intimately mixed equimolar CO-N<sub>2</sub> ice (80 L total exposure), (c) an N<sub>2</sub> layer (40 L) over a CO layer (40 L), and (d) a CO layer (40 L) over an N<sub>2</sub> layer (40 L). The <sup>13</sup>CO ( $m/e = 29$ ) traces are in black; <sup>15</sup>N<sub>2</sub> ( $m/e = 30$ ) traces are in gray. The heating rate is 0.1 K minute<sup>-1</sup>. The two dashed lines are superposed on the plot to show the positions of the desorption peaks of the 40 L pure samples of N<sub>2</sub> (gray) and CO (black).

tion processes. In light of these needs, the best laboratory analog mimicking the interstellar case would be the tertiary ice system, N<sub>2</sub>-CO-H<sub>2</sub>O, where CO and N<sub>2</sub> are sequentially or coadsorbed on a preexisting porous H<sub>2</sub>O-ice layer. One such qualitative study has been made under high-vacuum conditions showing that, after codeposition of all three molecules (producing a mixed ice), more CO than N<sub>2</sub> becomes trapped in H<sub>2</sub>O ices (Notesco & Bar-Nun 1996). However, no mention is made of the kinetics or binding energies in the paper, and as its focus was CO and N<sub>2</sub> in cometary systems, the ice surface was held at temperatures in excess of 40–50 K, not relevant to cold, prestellar cores.

It is known from observations that in many pre- and protostellar cores, up to 90% of the CO is frozen out in a distinct layer separated from the H<sub>2</sub>O ice (Tielens et al. 1991; Pontoppidan et al. 2003). If N<sub>2</sub> freezes out subsequent to CO, it is likely to form an ice layer *above* the CO ice, creating segregated N<sub>2</sub>/CO ices, or if CO continues to accrete to the grain, a mixed CO-N<sub>2</sub> ice. Consequently, the interplay between CO and N<sub>2</sub> during their desorption from binary CO-N<sub>2</sub> ice systems is equally relevant.

## 2. EXPERIMENTAL PROCEDURE

The apparatus used here, CRYOPAD (F. van Broekhuizen et al. 2005, in preparation), is very similar to the other Leiden instrument, SURFRESIDE, described in detail elsewhere (Fraser & van Dishoeck 2004). Briefly, the experiments were performed in an ultrahigh vacuum chamber, reaching base pressures of better than  $1 \times 10^{-10}$  mbar. At the center of the chamber is a gold-coated copper substrate, mounted in close thermal contact with a closed-cycle He cryostat, which cools the whole substrate to 12 K. The system temperature is monitored with two thermocouples, one mounted on the substrate, the second by the heater element. Ices are grown in situ, by exposing the cold substrate to a steady flow of gas, directed along the surface normal. Desorption is induced by linear heating of the substrate (and ice sample): during film growth or temperature programmed desorption (TPD), gases liberated from the surface are monitored using a quadrupole mass spec-

trometer (Pfeiffer Prisma). TPD is a well-established method in surface science for determining surface-adsorbate binding energies (e.g., Attard & Barnes 1998, p. 72; Woodruff & Delchar 1994).

Mixed (CO:N<sub>2</sub>) and layered (N<sub>2</sub>/CO) ice morphologies were studied. For completeness, control experiments were conducted on pure CO and pure N<sub>2</sub> ice films. Previous spectroscopic studies have shown that almost equal amounts of N<sub>2</sub> and CO can be frozen out onto interstellar grains without detectably affecting the line profile of the solid-CO vibrational band (Elsila et al. 1997). Therefore, the three ice morphologies were studied in equimolar ratios. To enable the two molecules to be discriminated from each other (and the background signal) with mass spectrometry, isotopes of both molecules were used, i.e., <sup>13</sup>CO (Cambridge Isotopes, Inc., 99%),  $m/e = 29$ , and <sup>15</sup>N<sub>2</sub> (Cambridge Isotopes, Inc., 98%),  $m/e = 30$ .

In the pure and layered ice morphologies, the gases were used as supplied; to form the mixed ices, a 1:1 gas mixture of <sup>13</sup>CO:<sup>15</sup>N<sub>2</sub> was pre-prepared then mounted on the chamber. The dosing rate for ice-film growth was set prior to cooling the sample, by sequentially backfilling the chamber with the gas(es) of interest, to a pressure of around  $1 \times 10^{-8}$  mbar, equivalent to an ion reading on the mass spectrometer of  $7.5 \times 10^{-10}$  A for both <sup>15</sup>N<sub>2</sub> and <sup>13</sup>CO. After cooling the substrate to 12 K, the ice films were grown by opening the preset flow valve for exposure times equivalent to a dose of 40 langmuirs (L) per sample gas (where  $1 \text{ L} \approx 1 \times 10^{-6}$  torr s), i.e., the number of molecules of each gas deposited on the surface is equivalent to the number of molecules in 40 monolayers of a nonporous condensed solid, assuming  $1 \times 10^{15}$  molecules cm<sup>-2</sup>. The value of 40 L was chosen for two reasons: First, it is the upper limit to the number of equivalent monolayers of pure CO ice condensed on grains on lines of sight to low-mass stars, as determined by Pontoppidan et al. (2003). Second, additional experiments (not shown here) indicate that at this ice thickness the behavior of the sample is substrate-independent (S. E. Bisschop et al. 2005, in preparation). Finally, the ice sample was warmed from 12 to 80 K at a rate of 0.1 K minute<sup>-1</sup> while the desorbing gases were monitored at the mass spectrometer.

## 3. RESULTS

The TPD spectra from each of the ice systems are shown in Figure 1. Equal amounts of either N<sub>2</sub> (gray traces) or CO (black traces) contribute to the TPD signal such that the integrated area under each TPD curve is constant to within  $\pm 5\%$  and the desorption characteristics of the different ice morphologies can be compared directly.

*Pure ices.*—In Figure 1a, the data from 40 L exposure pure CO and N<sub>2</sub> ices are overlaid. It is clear that N<sub>2</sub> desorbs at lower temperatures than CO. TPD data from layers of higher and lower exposures (not shown) confirm that, at 40 L exposure, the desorption kinetics of N<sub>2</sub> and CO are zeroth order, like those of H<sub>2</sub>O, i.e., the preexponential factor will typically have a value between  $10^{28}$  and  $10^{32}$  molecules cm<sup>-2</sup> s<sup>-1</sup> (Fraser et al. 2001; Collings et al. 2003a), and the desorption rate is independent of the amount of CO or N<sub>2</sub> adsorbed at the surface. If we assume the enthalpy of adsorption and desorption in these systems are equal, the mean desorption energies (in units of kelvins) are equivalent to the mean binding energy and can be approximately estimated from the TPD peak position (Attard & Barnes 1998, p. 72),

$$E_b = T_{\text{peak}} \times 30.068, \quad (1)$$




FIG. 2.—Fraction of N<sub>2</sub> desorbing from pure and mixed ice environments in each of the ice systems studied as a percentage of the total N<sub>2</sub> desorbed at the end of each experiment. See text for details.

giving  $E_b(\text{N}_2\text{-N}_2) = 790 \pm 25$  K and  $E_b(\text{CO-CO}) = 855 \pm 25$  K, respectively ( $R_{\text{BE}} = 0.923 \pm 0.003$ ). The error bars on  $E_b$  arise from a conservative estimate of the range in which  $T_{\text{peak}}$  actually lies, given that in zeroth-order desorption the TPD peak position shifts to higher temperatures and higher intensities as the surface coverage increases. It should be stressed that the mean desorption energies quoted throughout this Letter are *estimates* and the absolute values of  $E_b$  can only be accurately determined from kinetic models of the system, the subject of a future publication. Consequently, the error bars are larger than those obtained from the TPD peak positions alone, which can be read more precisely and are reproducible to  $\pm 0.05$  K for identical surface coverages. Within experimental error, our  $E_b(\text{CO-CO})$  is consistent with that reported by Collings et al. (2003a) of  $830 \pm 20$  K. Our values are also close to the sublimation enthalpies of N<sub>2</sub> and CO determined from IUPAC accredited data (Lide 2002), i.e.,  $756 \pm 5$  K and  $826 \pm 5$  K, respectively, giving a ratio  $R_{\text{BE}} = 0.915$  (upper limit = 0.923, lower limit = 0.903).

**Layered ices.**—In Figures 1c and 1d, TPD data from layered CO-N<sub>2</sub> ices are shown. As with the pure CO-ice samples, the onset of CO desorption occurs at around 26 K, and over 95% of the CO has desorbed by 31–32 K. It seems that CO only desorbs from a CO-dominated environment, although in comparison to the pure CO sample the presence of N<sub>2</sub> may result in a small increase in the range of binding sites from which desorption occurs, accounting for the slight broadening of the TPD profile. The mean desorption energy of CO molecules from the ice,  $E_b(\text{CO-CO})$ , is  $855 \pm 25$  K, and like the pure CO ice, the desorption kinetics of CO in layered CO-N<sub>2</sub> ice systems follow zeroth-order kinetics.

Although a significant fraction of the N<sub>2</sub> desorbs at lower temperatures than CO, i.e., 24–27.5 K, a second peak is clearly visible in the desorption profile of the layered ices, corresponding to codesorbing N<sub>2</sub>. This is interpreted as some fraction of the N<sub>2</sub> molecules becoming mobile and diffusing into the CO ice, where they become mixed. To establish the relative fractions of N<sub>2</sub> desorbing from each environment, a simple analysis was undertaken. The pure N<sub>2</sub> TPD trace (Fig. 1a) was scaled to fit the leading edge of each of the other N<sub>2</sub> TPD traces (Figs. 1b–1d), then subtracted. The integrated area under the scaled, pure-N<sub>2</sub> trace was then attributed to N<sub>2</sub> molecules desorbing from a pure ice environment, and the integrated area under the remaining curve to a mixed ice environment, under the assumption that




FIG. 3.—Summary of the desorption behavior of CO and N<sub>2</sub> within astroscopically relevant ice systems. Pure CO and N<sub>2</sub> are represented by light gray and dark gray shading, respectively; mixed ice regions are checked. Desorption is indicated by the wiggly arrows. (a) Mixed ice: The ice components begin segregating as soon as the ice is heated, forming layers of CO and N<sub>2</sub>, below and above the remaining ice mixture. Further heating leads to the onset of N<sub>2</sub> desorption from the segregated N<sub>2</sub> layer and then codesorption of CO and N<sub>2</sub>, until all the N<sub>2</sub> has desorbed. Finally, only CO is desorbing from the surface. (b) Layered ice: As the N<sub>2</sub>/CO layered ice is heated, the majority of N<sub>2</sub> molecules desorb directly from the upper surface of the N<sub>2</sub> multilayer. From  $\approx 20$  K, the diffusion of CO and/or N<sub>2</sub> across the layer interface leads to mixing between the two ice layers, and a fraction of the N<sub>2</sub> molecules subsequently codesorb with the CO molecules. [See the electronic edition of the Journal for a color version of this figure.]

each TPD trace is a linear combination of these two contributions. The results of this analysis are shown in Figure 2, where the error bars arise from the uncertainty in the initial scaling of the pure N<sub>2</sub> trace. The fraction of N<sub>2</sub> desorbing from the pure and mixed ice environments depends intrinsically on the initial morphology of the ice system. If N<sub>2</sub> is deposited above an existing CO ice layer, the majority of N<sub>2</sub> molecules do desorb prior to CO, whereas when N<sub>2</sub> is deposited below a CO layer about 50% of the N<sub>2</sub> molecules codesorb with the CO. This implies that the desorption kinetics are unchanged between layered and pure ice systems (i.e., still zeroth order) and  $R_{\text{BE}}$  is unaltered, i.e.,  $R_{\text{BE}} = 0.923 \pm 0.003$ . The value  $E_b(\text{CO-N}_2)$  is determined from the codesorbing fraction of N<sub>2</sub>, i.e.,  $\approx 855 \pm 25$  K, i.e.,  $R_{\text{BE}} = 1$ , although further differential N<sub>2</sub> exposure experiments are required to confirm this value.

**CO-N<sub>2</sub> ice mixture.**—In Figure 1b, the mixed ice CO TPD data are broadened and shifted to higher temperatures in comparison to the pure CO ice or layered ice systems. From the leading edge of the CO TPD trace, plus evidence from thickness-dependent studies (not shown), it appears that the CO desorption is hindered by the presence of N<sub>2</sub>. This results in a small shift in the TPD peak position, commensurate with a rise in the mean desorption energy of the CO molecules, i.e.,  $E_b(\text{CO-CO}) = 885 \pm 25$  K. This implies that the majority of the CO molecules in a mixed ice desorb from slightly different binding environ-

ments to those from which desorption occurs in pure and layered ice systems. Figure 2 shows that at least 50% of the  $N_2$  originally deposited in the mixture actually codesorbs with the CO. The remaining  $N_2$  desorbs from an environment resembling a pure  $N_2$  ice layer: it is therefore reasonable to assume that some  $N_2$  becomes segregated in a layer separated from the remaining ice mixture upon heating. For this segregated fraction of  $N_2$ ,  $R_{BE} = 0.893 \pm 0.003$ , due to the rise in the mean desorption energy of CO; for the remainder of the  $N_2$ ,  $R_{BE} = 1$ .

#### 4. DISCUSSION

The results from these experiments, summarized in Figure 3, have a direct impact on the modeling of CO and  $N_2$  desorption in prestellar and star-forming cores. First, since the desorption behavior of layered and mixed ices is not identical, a decision has to be made on which of these systems most suitably emulates the interstellar case. Second, the desorption kinetics of CO and  $N_2$  should be modeled as zeroth- and not first-order processes, with adopted rates independent of the surface concentration of the molecules, with preexponential factors in the rate constant of  $10^{28}$ – $10^{32}$  molecules  $cm^{-2} s^{-1}$  (Fraser et al. 2001; Collings et al. 2003a). Depending on the initial ice morphology and assuming that depleted molecules play no part in any grain-surface chemistry, some fraction of the  $N_2$  originally frozen-out on interstellar grains will only desorb as the CO multilayers themselves desorb.

Irrespective of the ice morphology, this work gives  $E_b(N_2-N_2) = 790 \pm 25$  K and  $E_b(N_2-CO) = 855 \pm 25$  K, except in the mixed ice, where  $E_b(CO-CO) = 885 \pm 25$  K.

Within experimental error, these values are consistent with those obtained by Collings et al. (2003a) but significantly smaller than the  $E_b(CO-CO)$  value of 960 K (Sandford & Allamandola 1990) used in BL97 or  $E_b(CO-SiO_2)$  of 1210 K (Bergin et al. 2002). The latter value is closer to the experimentally determined value of  $E_b(CO-H_2O) = 1180 \pm 20$  K (Collings et al. 2003b; Fraser et al. 2004). It is clear that the most suitable ratio between the binding energies of  $N_2$  and CO,  $R_{BE}$ , is  $0.923 \pm 0.003$  for all cases where  $N_2$  desorbs from “pure” ice environments. However,  $R_{BE} = 1$  for the fraction of  $N_2$  that codesorbs with CO. The lowest empirical value of  $R_{BE}$ ,  $0.893 \pm 0.003$ , is found in mixed ices, where the lower ratio reflects the rise in the mean value of  $E_b(CO-CO)$ . Nevertheless, all these ratios are significantly greater than any of the values adopted by BL97, Bergin et al. (2002), or Aikawa et al. (2003). Such a situation mimics that explored at the end of BL97, where  $R_{BE}$  was set to 1, to study the effects of the  $N_2$  binding energy on  $N_2H^+$  depletion, which was then enhanced by 2 orders of magnitude. These new laboratory data, when applied to models of dense cores, will certainly decrease the anticorrelation between gas-phase CO and  $N_2H^+$  abundances. The next step is clearly to address detailed modeling of the CO- $N_2$  desorption kinetics followed by empirical studies of the tertiary CO- $N_2$ - $H_2O$  ice system.

We are grateful for comments by J. Jørgensen and T. Bergin and funding from NOVA, the Dutch Research School for Astronomy, and an NWO Spinoza grant to E. F. v. D.; K. I. Ö. is grateful to the summer undergraduate research fellowship program at Caltech for sponsoring her visit to Leiden.

#### REFERENCES

- Aikawa, Y., Ohashi, N., & Herbst, E. 2003, *ApJ*, 593, 906  
 Attard, G., & Barnes, C. 1998, *Surfaces* (Oxford: Oxford Univ. Press)  
 Belloche, A., & André, P. 2004, *A&A*, 419, L35  
 Bergin, E. A., Alves, J., Huard, T., & Lada, C. J. 2002, *ApJ*, 570, L101  
 Bergin, E. A., Ciardi, D. R., Lada, C. J., Alves, J., & Lada, E. A. 2001, *ApJ*, 557, 209  
 Bergin, E. A., & Langer, W. D. 1997, *ApJ*, 486, 316 (BL97)  
 Caselli, P., Walmsley, C. M., Tafalla, M., Dore, L., & Myers, P. C. 1999, *ApJ*, 523, L165  
 Chiar, J. E., Gerakines, P. A., Whittet, D. C. B., Pendleton, Y. J., Tielens, A. G. G. M., Adamson, A. J., & Boogert, A. G. A. 1998, *ApJ*, 498, 716  
 Collings, M. P., Dever, J. W., Fraser, H. J., & McCoustra, M. R. S. 2003a, *Ap&SS*, 285, 633  
 Collings, M. P., Dever, J. W., Fraser, H. J., McCoustra, M. R. S., & Williams, D. A. 2003b, *ApJ*, 583, 1058  
 Di Francesco, J., André, P., & Myers, P. C. 2004, *ApJ*, 617, 425  
 Elsila, J., Allamandola, L. J., & Sandford, S. A. 1997, *ApJ*, 479, 818  
 Fraser, H. J., Collings, M. P., Dever, J. W., & McCoustra, M. R. S. 2004, *MNRAS*, 353, 59  
 Fraser, H. J., Collings, M. P., McCoustra, M. R. S., & Williams, D. A. 2001, *MNRAS*, 327, 1165  
 Fraser, H. J., & van Dishoeck, E. F. 2004, *Adv. Space Res.*, 33, 14  
 Jørgensen, J. K. 2004, *A&A*, 424, 589  
 Jørgensen, J. K., Schöier, F. L., & van Dishoeck, E. F. 2004, *A&A*, 416, 603  
 Kramer, C., Alves, J., Lada, C. J., Lada, E. A., Sievers, A., Ungerechts, H., & Walmsley, C. M. 1999, *A&A*, 342, 257  
 Lide, D. R. 2002, *CRC Handbook of Chemistry and Physics* (82nd ed.; Boca Raton: CRC Press)  
 Notesco, G., & Bar-Nun, A. 1996, *Icarus*, 122, 118  
 Pagani, L., Pardo, J.-R., Apponi, A. J., Bacmann, A., & Cabrit, S. 2005, *A&A*, 429, 181  
 Pontoppidan, K. M., et al. 2003, *A&A*, 408, 981  
 Sadlej, J., Rowland, B., Devlin, J. P., & Buch, V. 1995, *J. Chem. Phys.*, 102, 4804  
 Sandford, S. A., & Allamandola, L. J. 1990, *Icarus*, 87, 188  
 Sandford, S. A., Bernstein, M. P., Allamandola, L. J., Goorvitch, D., & Teixeira, T. C. V. S. 2001, *ApJ*, 548, 836  
 Tafalla, M., Myers, P. C., Caselli, P., Walmsley, C. M., & Comito, C. 2002, *ApJ*, 569, 815  
 Tielens, A. G. G. M., Tokunaga, A. T., Geballe, T., & Baas, F. 1991, *ApJ*, 381, 181  
 Walmsley, C. M., Flower, D. R., & Pineau des Forêts, G. 2004, *A&A*, 418, 1035  
 Woodruff, D. P., & Delchar, T. A. 1994, *Modern Techniques of Surface Science* (2nd ed.; Cambridge: Cambridge Univ. Press), chap. 9  
 Whittet, D. C. B., et al. 1998, *ApJ*, 498, L159