



Universiteit
Leiden
The Netherlands

Two challenges in bone marrow transplantation. Graft-versus-Host-Disease and the unrelated donor

Goulmy, E.A.J.M.; Rood, J.J. van; Claas, F.H.J.; Hendriks, G.F.J.; Schreuder, G.M.Th.; Azogui, O.; Gluckman, E.

Citation

Goulmy, E. A. J. M., Rood, J. J. van, Claas, F. H. J., Hendriks, G. F. J., Schreuder, G. M. T., Azogui, O., & Gluckman, E. (1983). Two challenges in bone marrow transplantation. Graft-versus-Host-Disease and the unrelated donor. *Exp. Hemat.*, 11(suppl. 13), 61-66. Retrieved from <https://hdl.handle.net/1887/2918>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/2918>

Note: To cite this publication please use the final published version (if applicable).

TYPE WITHIN IN- NOT TYPE A

W. J. (RJP)

A. J. A. I

TWO CHALLENGES IN BONE MARROW TRANSPLANTATION

Graft versus host disease and the unrelated donor.

(Report of the Working Party on Immunology)

J.J. van Rood, E. Goulmy, F.H.J. Claas, G.F.J. Hendriks, G.M.Th. Schreuder, O. Azogui^x and E. Gluckman^xDepartment of Immunohaematology & Blood Bank, University Hospital, Leiden, the Netherlands, ^xl'Hôpital St. Louis, Paris, France.

GRAFT VERSUS HOST DISEASE (GVHD)

Immunogenetics

Goulmy has reported during the previous meeting the occurrence of cytotoxic lymphocytes (CTL's) in a patient suffering from chronic GVHD after allogeneic bone marrow transplantation for acute myeloid leukaemia (1,2). These CTL's were directed against a non-HLA determinant (HA) and were restricted by the class I antigens of the donor. They allow thus the recognition of genetic differences between HLA-identical siblings, for instance the CTL's reacted with the patient's own pretransplant lymphocytes, but not with the donor's lymphocytes. Similar CTL's have been described by Elkens et al. in a patient, who had become sensitized after multiple blood transfusions (3).

Goulmy et al. suggested that the anti-HA CTL's could and should be used to study their relevance in the occurrence of GVHD. In a collaborative effort with the bone marrow transplant group of the l'Hôpital St. Louis, in Paris (Head: Prof. E. Gluckman) and of the University Hospital Leiden (Head: Prof. Dr. J.J. Veltkamp) preliminary evidence has been collected which indicate that incompatibility for HA (or other non-HLA CTL determinants) can lead to GVHD.

Table I. Relation between mismatches for non-HLA determinants between donor and recipient and GVHD (Leiden-Paris)

	non-HLA mismatches		
	not detected	HA	other
Acute GVHD	19	0	0
Chronic GVHD	2	3	4

Table II. CML reactivity between HLA-identical donor-recipient combinations (Leiden-Paris)


	CML	chron. acute	
		GVHD	GVHD
pos.	6	0	
neg.	2	4	
p = 0.02			

Table I shows that in chronic GVHD but not in acute GVHD differences for HA (or its allele) could be found. Likewise a positive CML was found between post- and pretransplant lymphocytes from patients suffering from chronic GVHD but not from such patients suffering from acute GVHD (Table II). Our preliminary conclusion is that such CTL's can recognize non-HLA but MHC restricted determinants and that incompatibility for these determinants could lead to GVHD.

Immune modulation

We have discussed in previous reports the possibility that the finding that pretransplant blood transfusion can facilitate renal allograft survival might also be relevant in bone marrow transplantation. Recently it has been shown that in primates and mice not only whole blood but also pure platelets can induce this so-called pretransplant blood transfusion effect (4,5). In mice it was shown that heat pretreated leucocytes behaved as platelets as far as the graft protecting effect was concerned. On the basis of these findings it was investigated in a mouse model whether platelets obtained from the future recipient and given to the bone marrow donor would mitigate GVHD. Figure 1 shows that this is indeed the case. Note that if the donor receives recipient's leucocytes GVHD is more severe, while survival is improved (and GVHD is lessened) if pure platelets are given.

Figure 1. Survival of (B10.T(6R) x B10.A(2R))F1 mice following a bone marrow transplantation with B10.AQR spleen cells. Donor pretreatment: ■ = pretreated with recipient leucocytes; ● = not pretreated; + = pretreatment with heat-treated leucocytes of the recipient. (Claas et al. 1982)



Of course these data cannot be extrapolated to man without extensive preclinical testing (dogs, monkeys). Even if such experiments would corroborate the findings in the mouse, ethical considerations would

THE UNRELATED DONOR

Leaving matching for the class II antigens aside, we have analysed the feasibility of finding class I identical bone marrow donors in a file of 10.000. For 28 patients who had no HLA identical sibling donor, we would have been able to provide for 18 of them 5 to over a 100 unrelated class I identical donors. Although the logistics offer thus for about 2/3 of our patients no unsurmountable problems the implementation of such an operation will take considerable time. Such protocols should be reviewed by the medical ethics committee, informed consent should be obtained as well as adequate reimbursement of the costs.

They showed that some AB and DR mismatched renal grafts did extremely well. It was assumed that this was due to a low responder characteristic of the recipient, at least as far as HLA allo-antigens were concerned. The first significant evidence that immune response genes might be involved in developing immunity against non-HLA antigens was unearthed by Baldwin et al. when they showed that only HLA-DRw6 positive individuals formed antibodies against antigens present on some endothelial cells and monocytes (8). HLA-DRw6 appeared to act as an immune response (Ir) gene for these EM antigens.

This study was undermore criticized because it is notoriously difficult to type for HLA-DRw6. Recently however the definition of these class II antigens has been much improved (9) and it seemed to be a good moment to assess the importance of HLA-DRw6 as Ir gene for HLA alloantigens.

Figure 2 shows that overall renal graft survival is poorer in HLA-DRw6 positive than in negative ones. (10) A finding which in the meantime has been confirmed by others (P.J. Morris personal communication). In contrast matching for HLA-DR seems to be very effective in DRw6 positive and only of borderline importance in DRw6 negative recipients. That this was not a red herring is shown in figure 3, which shows the influence of matching for HLA-DR on renal graft survival in Eurotransplant (G.F.J. Hendriks personal communication).

Figure 2.

EFFECT OF DRW6 ON KIDNEY GRAFT SURVIVAL





Figure 3.

RENAL ALLOGRAFT SURVIVAL WITH RESPECT TO THE YEAR OF TRANSPLANTATION.




In the years before 1978 typing for HLA-DR was possible in Leiden only, thus donors were typed but not matched with recipients for HLA-DR. In 1980 most centers had implemented HLA-DR typing and matching became feasible. Over these years graft survival at one year post-transplant in the HLA-DRw6 negative group remained unchanged but in the HLA-DRw6 positive group rose from about 35% to over 70%. This is thus a confirmation on an independent set of data of the findings shown in figure 2. Finally it could be shown that in HLA-DRw6 negative individuals pretransplant blood transfusion improved graft survival quite substantially while in DRw6 positive individuals it appeared to have no effect. These are thus 3 different observations which all suggest that DRw6 is a strong Ir gene for incompatible HLA antigens, and that matching for DR (but not pretransplant blood transfusion) turns the DRw6 positive recipient from a high in a low responder. The situation appears to be inversed if DRw6 is present in the donor. In that situation it appears to act as an activator of suppressor cells. In figure 4 it is shown that if a HLA-DRw6 negative recipient is transplanted with kidney from a donor which is mismatched for one HLA-DR antigen, graft survival is excellent ($> 90\%$), if that antigen is DRw6 but only average ($\pm 70\%$) if it is any other DR antigen ($p =$

0.01). Similar findings were observed when the outcome of grafts which were mismatched for two DR antigens were analysed (Hendriks et al. to be submitted).

In conclusion, it appears that DRw6 in the recipient of a renal allograft acts as a strong Ir gene both for non-HLA and HLA antigens, but when it is present in the donor it acts as an activator of suppressor cells. To which extent these data can be extrapolated to the situation in bone marrow transplantation remains to be seen. The partly mismatched family transplants might be better material to study and this the working party on immunology plans to do.

Figure 4



Even if such studies would not reveal a relation between HLA-DRw6 and the prognosis of partly mismatched bone marrow grafts, the results in renal transplantation remain relevant for haematologists. This is so because they provide for the first time an indication that it is possible to activate the suppressor circuit in man. This is obviously not only important for the control of the homograft reaction, but also of the graft versus host reaction and possibly even of autoimmune disease.

ACKNOWLEDGEMENTS

In part supported by the Dutch Foundation for Medical Research (FUNGO) which is subsidized by the Dutch Organization for the Advancement of Pure Research (ZWO) and J.A. Cohen Institute for Radiopathology and Radiation Protection (IRS).

REFERENCES

1. Goulmy, E, Gratama, JW, Blokland, E, Zwaan, FE & van Rood, JJ. Recognition of an - as yet unknown - minor transplantation antigen by post-transplant lymphocytes from an A.M.L. patient. *Exp Hemat* 1982; 10, suppl. 10: 127-9.
2. Goulmy, E, Gratama, JW, Blokland, E, Zwaan, FE & van Rood, JJ. A minor transplantation antigen detected by MHC restricted CTLs during graft versus host disease, *Nature* 1983; in press.
3. Elkins, WL, Pierson, G, Negendank, W & Zier KS. Recognition of human minor alloantigen(s) by cytotoxic lymphocytes in vitro. *Immunogenetics* 1982; 15: 489-99.
4. Borleffs, JCC, Neuhaus, P, van Rood, JJ & Balner, H. Platelet transfusions improve kidney allograft survival in Rhesus monkeys without inducing cytotoxic antibodies. *The Lancet* 1982; May 15: 1117-8.
5. Claas, FHJ, Blankert, JJ, Ruigrok, R & Moerel, L. Platelet transfusions can induce transplantation tolerance. *Immunology Letters* 1982; 5: 35-9.
6. Wagemaker, G, Heidt, PJ, Merchav, S & van Bekkum, DW. Abrogation of histocompatibility barriers to bone marrow transplantation in rhesus monkeys. In: *Experimental Hematology Today 1982* (JS Baum et al. Eds.). Basel, Karger, 1982, pp. 111-8.
7. Termijtelen, A, Vossen, JMJJ, Nauta, J, van 't Veer-Korthof, EF & van Rood, JJ. Selection of unrelated bone marrow donors for two SCID and one Wiskott Aldrich patient. *Blut* 1980; 41: 241-4.
8. Baldwin, WM, Claas, FHJ, van Es, LA, van Rood, JJ, Paul, LC & Persijn, GG. Renal graft rejection and the antigenic anatomy of human kidneys. In: *Transplantation and Clinical Immunology*, Vol. XIII (JL Touraine et al. Eds.). Amsterdam, Excerpta Med., 1981, pp. 140-6.
9. Schreuder, GMTh, Parlevliet, J, Termijtelen, A & van Rood, JJ. A reanalysis of the HLA-DRw6 complex. *Tissue Antigens* 1982; in press.
10. Hendriks, G.F.J., Schreuder, GMTh, Persijn, GG, Cohen, B & van Rood, JJ. HLA-DRw6 and renal allograft rejection. *Brit.Med.J.* 1983; 286: 85-7.