Universiteit
Leiden
The Netherlands

The Marburg I polymorphism of factor VII-activating protease is not associated with venous thrombosis

Minkelen, R. van; Visser, M.C.H. de; Vos, H.L.; Bertina, R.M.; Rosendaal, F.R.

Citation

Minkelen, R. van, Visser, M. C. H. de, Vos, H. L., Bertina, R. M., & Rosendaal, F. R. (2005). The Marburg I polymorphism of factor VII-activating protease is not associated with venous thrombosis. *Blood*, 105(12), 4898-4898. Retrieved from <https://hdl.handle.net/1887/5046>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/5046>

Note: To cite this publication please use the final published version (if applicable).

To the editor:

Does the FLIPI apply to grade 3 follicular lymphoma?

We welcome the publication of a score (Follicular Lymphoma International Prognostic Index [FLIPI]) assisting in the choice of treatment for patients with newly diagnosed follicular lymphoma (FL).¹ As the authors state, treatment options for this disease range from “watch and wait” to allogeneic bone marrow transplantation, and the selection of the best available option will be favored by a realistic estimation of the expected survival for each particular patient.

The 5 independent prognostic factors retained in the FLIPI reflect important characteristics of the patient (age), of the disease extension and clinical aggressiveness (stage and number of nodal sites involved, lactose dehydrogenase [LDH]), and of the tumor-host interaction (hemoglobin level). Surprisingly enough, histology grade, a disease characteristic considered by many clinicians to be of paramount importance for prognosis and choice of treatment, is missing from the index and was not even significant in the univariate analysis.

The absence of this parameter from the index and the omission of any discussion about this fact in the paper could suggest that histologic grade is indeed not relevant to patient survival and should therefore not be taken as an important information.

We believe that this misunderstanding results from a bias that is not sufficiently covered in the article and that should be clarified.

It was long recognized that an important proportion of large cells (centroblasts) at histologic examination (an observation referred to as grade 3 in the current World Health Organization [WHO] classification) does confer to FL a worse prognosis, despite apparently favorable clinical prognostic features.^{2,3} Nevertheless, the outcome of these cases can be similar to the forms with less centroblasts when they are treated with an anthracycline-containing regimen.^{4,5} This information was known in 1985⁶ (the year of the start of data collection for the FLIPI) and, although details on treatment are not given in the paper, it is probable that the majority of grade 3 FLs of this data set were treated with anthracyclines, thus influencing the prognosis.

Response:

FLIPI in grade 3 follicular lymphoma

We agree with Prof Ghielmini that grade 3 follicular lymphoma is in many cases a more aggressive disease than other follicular lymphomas, although this is an area of ongoing controversy.¹ Grade 3b follicular lymphoma especially shares the prognosis of and requires treatment approaches similar to that of diffuse large B-cell lymphoma.² We welcome the opportunity to comment further on the Follicular Lymphoma International Prognostic Index (FLIPI) observations about grade 3 follicular lymphoma.

First, a central pathology review was not realistic for the 5000 FLIPI cases. Since there are known interobserver variations in the assignment of histologic grade,^{3,4} we felt we would introduce a bias if we excluded these cases arbitrarily. Moreover, they represented only 9% of the entire FLIPI population.

Even though the article states that “none of the treatments given during the period of inclusion has significantly changed the natural history of the disease,”^{1(p1264)} this only applies to cases with histologic grades 1 and 2, while for WHO grade 3 we have reason to think that treatment did indeed influence survival, and therefore interacted with other prognostic factors.

The FLIPI could have been a more reliable index if grade 3 FLs were not included in the analysis. We believe that for grades 1 and 2 FL, treatment should be selected based on a number of factors, including the FLIPI score, but an anthracycline-containing regimen should still be favored for patients with grade 3 FL, independent of their FLIPI index.

Michele Ghielmini and Oreste Mora

Correspondence: Michele Ghielmini and Oreste Mora, Oncology Institute of Southern Switzerland, Ospedale San Giovanni, 6500 Bellinzona, Switzerland; e-mail: mghiellini@ticino.com

References

1. Solal-Célgny P, Roy P, Colombat P, White, et al. Follicular Lymphoma International Prognostic Index. *Blood*. 2004;104:1258-1265.
2. Martin AR, Weisenburger DD, Chan WC, et al. Prognostic value of cellular proliferation and histologic grade in follicular lymphoma. *Blood*. 1995;85:3671-3678.
3. Rodriguez J, McLaughlin P, Hagemeister FB, et al. Follicular large cell lymphoma: an aggressive lymphoma that often presents with favorable prognostic features. *Blood*. 1999;93:2202-2207.
4. Nathwani BN, Harris NL, Weisenburger D. Follicular lymphoma. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. *World Health Organization Classification of Tumours. Tumours of haematopoietic and lymphoid tissues*. Albany, NY: WHO Publication Center; 2001:162-167.
5. Bartlett NL, Rizeq M, Dorfman RF, Halpern J, Horning SJ. Follicular large-cell lymphoma: intermediate or low grade? *J Clin Oncol*. 1994;12:1349-1357.
6. Glick JH, McFadden E, Costello W, Ezdinli E, Berard CW, Bennett JM. Nodular histiocytic lymphoma: factors influencing prognosis and implications for aggressive chemotherapy. *Cancer*. 1982;49:840-845.

Second, the IPI appears to have one of the same shortcomings for grade 3 follicular lymphoma as it does for all follicular lymphomas, namely the identification of only a small (15%-20%) fraction of cases with high risk.^{5,6} Thus the inclusion of grade 3 cases in the FLIPI, and the applicability of the FLIPI to grade 3 follicular lymphoma, represents a potential benefit to patient management.

But we completely agree with Prof Ghielmini's observation that an anthracycline-containing regimen is favored for patients with grade 3 follicular lymphoma. It was beyond the scope of the initial FLIPI project to analyze in any detail the various treatment modalities used. Moreover, such a retrospective analysis would be open to valid criticism. But we agree that the literature supporting the inclusion of anthracyclines in grade 3 follicular lymphoma^{5,6} is

less controversial than the comparable literature in other follicular lymphomas.⁷ We thank him for drawing attention to this important topic. The F2 study (coordinated by M. Federico) is ongoing to determine the influence of grade and the impact of diffuse areas on prognosis.

Peter McLaughlin and Philippe Solal-Céligny,
on behalf of the Scientific Committee of the FLIPI

Correspondence: Philippe Solal-Céligny, Centre J. Bernard, 9 rue Beauverger, 72000 Le Mans, France; e-mail: celigny@roos.fr

References

- Armitage JO, Cavalli F, Zucca E, Longo DL. Follicular lymphoma. In: Text Atlas of Lymphomas. Martin Dunitz Ltd; 1999:13-21.
- Bosga-Bouwer AG, van Imhoff GW, Boonstra R, et al. Follicular lymphoma grade 3b includes 3 cytogenetically defined subgroups with primary t(14;18) 3q27 or other translocations: t(14;18) and 3q27 are mutually exclusive. *Blood*. 2003;101:1145-1154.
- Hornig SJ. Something old, something new, something subjective, something déjà vu. *J Clin Oncol*. 2003;21:1-2.
- Hans CP, Weisenburger DD, Vose JM, et al. A significant diffuse component predicts for inferior survival in grade 3 follicular lymphoma, but cytologic types do not predict survival. *Blood*. 2003;101:2363-2367.
- Wendum D, Sebban C, Gaulard P, et al. Follicular large-cell lymphoma treated with intensive chemotherapy: an analysis of 89 cases included in the LNH87 trial and comparison with the outcome of diffuse large B-cell lymphoma. *J Clin Oncol*. 1997;15:1654-1663.
- Rodriguez J, McLaughlin P, Hagemeister FB, et al. Follicular large cell lymphoma: an aggressive lymphoma that often presents with favorable prognostic features. *Blood*. 1999;93:2002-2007.
- Rigacci L, Federico M, Martinelli M, et al. The role of anthracyclines in combination chemotherapy for the treatment of follicular lymphomas. *Leuk Lymphoma*. 2003;44:1911-1917.

To the editor:

Analysis of total phosphotyrosine levels in CD34⁺ cells from CML patients to predict the response to imatinib mesylate treatment

The achievement of partial or complete cytogenetic remission (CCR) in response to imatinib mesylate in chronic myeloid leukemia (CML) is an indicator of favorable response to the drug.¹ The current assays for prediction of response rely on immunoblot analyses of Bcr-Abl or its substrates Crkl² or Stat5.³ However, these require large numbers of primary cells and are technically demanding. We previously showed that the inhibition of total phosphotyrosine was a reliable indicator of the effect of imatinib mesylate on *BCR-ABL*⁺ cell lines.⁴ We have now further developed this flow cytometric test to assess the response of primary cells to in vitro imatinib treatment.

We analyzed cryopreserved mononuclear cells from 23 patients with CML in chronic phase, 16 of whom had responded to imatinib mesylate, reaching at least a partial cytogenetic response within the first year of therapy. There was no significant difference in the interval between diagnosis and the initiation of imatinib mesylate treatment between the 2 groups. After thawing, CD34⁺ cells were isolated by immunomagnetic separation, kept overnight in culture, and subsequently exposed for 2 hours to doses of imatinib mesylate ranging from 0.1 to 50 μ M, and intracellularly stained as described in Figure 1. The overall tyrosine phosphorylation was determined by flow cytometry and expressed as a percentage of the level of phosphorylation of the nonexposed control cells.

The results showed a significant difference between patients who achieved a complete or partial cytogenetic remission and those who did not (Figure 1). For the most discriminatory concentration of 20 μ M imatinib mesylate, the median level of tyrosine phosphorylation inhibition in cells from the responders was 46%, as opposed to only 32% in the nonresponders ($P = .034$, Mann-Whitney test). None of the nonresponders had more than 45% inhibition of their total cellular phosphotyrosine content, whereas this was achieved in half of the responding patients. The molecular response to imatinib mesylate also correlated with its capacity to inhibit tyrosine phosphorylation in vitro: in none of the patients who failed to exhibit a greater than 2-log reduction in the number of *BCR-ABL* transcripts or a *BCR-ABL/ABL* ratio less than 2% was imatinib able to reduce the cellular phosphotyrosine content below 55% of the pretreatment levels, whereas 75% (6/8) of those with at

Figure 1. Inhibition of tyrosine phosphorylation in primary cells by imatinib mesylate. (A) Median levels of tyrosine phosphorylation in cytogenetic responders (□) and nonresponders (■). 10^4 to 10^5 CD34⁺ cells per data point were treated for 2 hours with a range of imatinib mesylate concentrations, fixed in 1% paraformaldehyde, permeabilized with 0.3% saponin, and incubated with an antiphosphotyrosine (PY99; Santa Cruz, Holly Ditch Farm, United Kingdom) and a secondary fluorescent antibody (goat F(ab')₂ antimouse; Caltag, Silverstone, United Kingdom). The dose of imatinib mesylate which best distinguishes the 2 groups was found to be 20 μ M. (B) Median percentage tyrosine phosphorylation of imatinib mesylate-treated cells from cytogenetic responders (R) and nonresponders (NR) compared with cells not exposed to the drug. The data show a significant difference between the median tyrosine phosphorylation of the 2 groups (Mann-Whitney test, $P = .034$). In about half of the patients responding to imatinib mesylate the tyrosine phosphorylation levels upon in vitro exposure to the inhibitor fell below 55% (dotted line) of that in the nonexposed cells, whereas this level of reduction was not achieved in any of the nonresponders. (C) Illustrative flow cytometric profile of 1 responder (top) and 1 nonresponder (bottom). Cells stained with the isotype control antibody (gray line) are used to adjust the flow cytometer settings, whereas cells not exposed to imatinib mesylate serve as a phosphotyrosine staining baseline for all measurements (black line). Imatinib mesylate-treated cells (shaded curve) from patients who eventually responded to imatinib mesylate show a clearly visible shift to the left, while the shift is significantly smaller for the nonresponders (as shown by the distance between the arrows pointing to the peak fluorescence of the exposed and nonexposed cells). (D) Comparison of CD34 cells used fresh or after 1 round of freezing for the assessment of total phosphotyrosine phosphorylation. From 7 extra patients, CD34⁺ cells from freshly collected blood were divided into 2 aliquots: 1 was placed in culture for the assay on "fresh" cells, and the second was cryopreserved and thawed later on for the same test. Each line depicts 1 paired sample of cells.

least 45% inhibition of phosphotyrosine levels achieved those degrees of molecular remission. No association was found between the Sokal and Hasford scores⁵ and the level of decrease in tyrosine phosphorylation.

The test is able to generate reproducible data from as little as 5×10^4 CD34⁺ cells, can be completed within 1 day from the receipt of the sample in the laboratory, and yields similar results on fresh or cryopreserved cells (Figure 1D). In order to confirm the results of this pilot study, we initiated a prospective analysis on a larger patient group where additional clinical and molecular parameters will be assessed. If validated, this assay may facilitate decisions on whether to opt for imatinib mesylate treatment as opposed to allogeneic stem cell transplantation in the first place.

Beate Schultheis, Richard Szydlo, François X. Mahon, Jane F. Apperley, and Junia V. Melo

Correspondence: Junia V. Melo, Department of Haematology, Imperial College London, Hammersmith Hospital, Ducane Road, London W12 0NN, United Kingdom; e-mail: j.melo@imperial.ac.uk

We thank Efthymios Fidanis for technical assistance, Dr David Marin and Linda Fung for providing clinical information on the patients, Dr Giuseppe Saglio for cells from 1 patient, and Dr Elisabeth Buchdunger for donation of imatinib mesylate.

This work was funded by the Hammersmith Hospital Trust Research Committee and the Leukaemia Research Fund (United Kingdom).

References

1. Kantarjian H, Sawyers C, Hochhaus A, et al. Hematologic and cytogenetic responses to imatinib mesylate in chronic myelogenous leukemia. *N Engl J Med*. 2002;346:645-652.
2. Druker BJ, Talpaz M, Resta DJ, et al. Efficacy and safety of a specific inhibitor of the BCR-ABL tyrosine kinase in chronic myeloid leukemia. *N Engl J Med*. 2001;344:1031-1037.
3. Jacobberger JW, Sramkoski RM, Frisa PS, et al. Immunoreactivity of Stat5 phosphorylated on tyrosine as a cell-based measure of Bcr/Abl kinase activity. *Cytometry A*. 2003;54:75-88.
4. Mahon FX, Belloc F, Cholet C, et al. Detection of resistance to STI571 in patients with BCR-ABL positive acute leukemia and chronic myeloid leukemia (CML) blast crisis [abstract]. *Blood*. 2000;96:471a.
5. Hasford J, Pfirrmann M, Hehlmann R et al. A new prognostic score for survival of patients with chronic myeloid leukemia treated with interferon alfa: Writing Committee for the Collaborative CML Prognostic Factors Project Group. *J Natl Cancer Inst*. 1998;90:850-858.

To the editor:

The interleukin-10 gene promoter polymorphism (–1082) does not correlate with clinical outcome in diffuse large B-cell lymphoma

Interleukin-10 (IL-10) is a cytokine important in suppressing the immune response by inhibiting proinflammatory cells. However, this cytokine is multifunctional and can also stimulate proliferation as has been shown both in normal and tumor B cells.¹ IL-10 has been implicated to play a role in lymphoma development, especially considering the finding of an association between increased IL-10 levels and poor outcome in some lymphoma entities.^{1,2} Furthermore, IL-10 gene promoter polymorphisms have been shown to affect the levels of IL-10 expression, for example, the IL-10_{–1082A} and IL-10_{–1082G} alleles correlate with low and high IL-10 production, respectively.³ Recently, Lech-Maranda et al² reported that the IL-10_{–1082} genotype influenced clinical outcome in patients with diffuse large B-cell lymphoma (DLBCL), where an improved overall survival and a higher rate of complete remission were found for patients with IL-10_{–1082AG/GG} genotypes compared with patients with IL-10_{–1082AA} genotype. They also showed a higher IL-10_{–1082G} allele frequency in DLBCL compared with healthy controls (0.47 vs 0.39). To investigate this further, we screened 244 samples obtained from patients with DLBCL (168 with de novo DLBCL, 67 with DLBCL and previous history of low-grade lymphoma, and 9 with DLBCL and unknown previous

Figure 1. Overall survival for the 244 patients with DLBCL with the IL-10_{–1082AG/GG} (174 cases) or IL-10_{–1082AA} (70 cases) genotype.

history) from Uppsala and Umeå University Hospitals to establish their IL-10_{–1082} genotype and compare it with overall survival. We also assessed the genotype in 195 healthy controls. In the present analysis, the frequency of the IL-10_{–1082G} allele was not significantly different in our patients with DLBCL versus the control group (0.43 vs 0.46; $P = .38$). Thus, the increased IL-10_{–1082G} allele frequency could not be verified in our analysis of DLBCL (Table 1). Furthermore, we did not find any difference in overall survival between the 174 patients with IL-10_{–1082AG/GG} genotype and the 70 patients with IL-10_{–1082AA} (median survival, 39 vs 37 months; log-rank test, $P = .50$; Figure 1). No significant difference in the clinical presentation was indicated between the 2 cohorts considering sex, clinical stage, complete remission rate, and international prognostic index. Although patients in our analysis, in general, were older (χ^2 ; $P = .014$), had B symptoms more often (χ^2 ; $P = .034$), and had elevated serum lactate dehydrogenase (S-LDH; $P = .012$) levels, no difference in survival was shown between patients with IL-10_{–1082AG/GG} and IL-10_{–1082AA} genotype when considering only patients younger than 60 years of age or

Table 1. Allele and genotype frequency of the IL-10_{–1082} polymorphism in 244 DLBCLs and 195 control subjects

	DLBCL cases	Controls	<i>P</i>
Allele frequency			
IL-10 _{–1082G}	0.43	0.46	.38
Genotype frequency (%)			
IL-10 _{–1082GG}	38 (16)	46 (24)	.64
IL-10 _{–1082AG}	136 (56)	89 (46)	
IL-10 _{–1082AA}	70 (29)	60 (31)	

P values were calculated using χ^2 test. The *P* value for the genotype frequency is calculated for IL-10_{–1082AG/GG} versus IL-10_{–1082AA}.

when patients with or without B symptoms or elevated S-LDH levels were analyzed separately. Moreover, no difference in outcome was evident for patients with or without the IL-10_{-1082G} allele when separately analyzing de novo DLBCL and cases with previous history of low-grade lymphoma. In conclusion, we could not confirm the findings in the report by Lech-Maranda et al² of increased frequency of the IL-10_{-1082G} allele and its association with superior outcome in our analysis of patients with DLBCL. We therefore believe that the clinical relevance of the IL-10₋₁₀₈₂ promoter polymorphism may be limited in DLBCL, although larger studies have to be performed to conclude this properly.

Mattias Berglund, Ulf Thunberg, Göran Roos, Richard Rosenquist, and Gunilla Enblad

Correspondence: Mattias Berglund, Departments of Oncology, Radiology, and Clinical Immunology, Rudbeck Laboratory, Uppsala University, SE-751 85 Uppsala, Sweden; e-mail: Mattias.Berglund@genpat.uu.se.

Response:

Different patients, different results: the need for prospective controlled studies

In this issue, Berglund and colleagues report their analysis of the IL-10 gene polymorphism (*IL10*₋₁₀₈₂) in a series of 244 lymphoma patients. This series included 168 de novo diffuse large B-cell lymphomas whereas other cases had a previous history of low-grade lymphoma and should probably be considered as transformed lymphoma. The authors did not find a different distribution in *IL10*₋₁₀₈₂ alleles between the patient and the control populations. Of note, the *IL10*_{-1082G} allele disequilibrium frequency that we previously described only applies to diffuse large B-cell lymphoma and not to other lymphoma subtypes (E.W. and G.S., unpublished results, May 2005). In this context, these authors also failed to observe a prognostic significance for the *IL10*_{-1082 G} allele in diffuse large B-cell lymphoma.

The reasons for this discrepancy are unclear and may include several biases such as patient characteristics and treatment options and results. For instance, the median survival reported by Berglund et al is close to 38 months for the whole population but was

The authors are grateful to Drs Lars Klareskog and Leonid Padyukov and to the Epidemiologic investigation of rheumatoid arthritis (EIRA) group for help with the collection of the healthy control material and to Dr Thomas Axelsson for technical assistance. The single nucleotide polymorphism (SNP) genotyping was performed by the Wallenberg Consortium North (WCN) SNP platform, Department of Medical Sciences, Uppsala University.

References

1. Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A. Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol*. 2001;19:683-765.
2. Lech-Maranda E, Baseggio L, Bienvenu J, et al. Interleukin-10 gene promoter polymorphisms influence the clinical outcome of diffuse large B-cell lymphoma. *Blood*. 2004;103:3529-3534.
3. Turner DM, Williams DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV. An investigation of polymorphism in the interleukin-10 gene promoter. *Eur J Immunogenet*. 1997;24:1-8.

projected to be 70 months in our study. The fact that this study includes older patients is also significant since the prognostic significance of the *IL10*_{-1082 G} allele was more pronounced in patients younger than 60 years than in older patients in our study. Finally, how inherited immune response may actually influence patient outcome may indeed depend of the initial characteristics of the patient, the disease, and the efficiency of treatment. Altogether, these results emphasize the need for prospective and multicentric genomic DNA collection linked to a clinical database, such as the one currently built with Groupe d'Etude des Lymphomes de l'Adulte (GELA) clinical trials, which will be analyzed in the coming years.

Gilles Salles and Ewa Maranda

Correspondence: Centre Hospitalier Leon-SUD, 69495 Pierre-Bénite cedex, France; e-mail: gilles.salles@chu-lyon.fr.

To the editor:

The bone marrow microenvironment influences the differential chemokine receptor expression of normal and neoplastic plasma cells

A number of mechanisms may explain the clinical and biologic differences in behavior between monoclonal gammopathy of undetermined significance (MGUS) and myeloma plasma cells (PCs). These include inherent differences in PC programming, immortalization of PCs at different stages of differentiation, different cells of origin, and potentially the influence of the normal/neoplastic bone marrow (BM) microenvironment. Chemokines and their receptors are known to be paramount to normal cell differentiation and homing. As B cells differentiate into PCs they undergo a coordinated change in chemokine receptor expression and chemokine responsiveness. We therefore hypothesized that the expression of chemokine receptors on myeloma PCs would be altered compared with normal PCs and may explain some of the differences in behavior between normal and malignant populations.

As our previous studies have demonstrated that 2 populations of PCs (normal and neoplastic) can be identified in myeloma and MGUS cases based on expression of CD19 expression¹⁻³ and that MGUS cases with a predominantly neoplastic phenotype have a higher chance of progressing to myeloma than those with a mixed normal and malignant phenotype,³ we also hypothesized that by examining both normal and neoplastic PCs from the same BM in MGUS patients we could determine whether any variation in chemokine receptor profile was due to the marrow microenvironment or to neoplastic transformation.

Normal (CD38⁺CD138⁺CD19⁺CD56⁻) and neoplastic PCs (CD38⁺CD138⁺CD19⁻CD56^{+/+}) were distinguished from other leukocytes using 4-color flow cytometry with CELLQuest version 3.1 on a FACSort (BD Biosciences, Oxford, United Kingdom) as

Figure 1. Flow-cytometric analysis of chemokine receptor expression. (A) Chemokine receptor expression on normal and myeloma PCs. The mean expression level and range is shown. (B) CCR2 expression is significantly lower on CD19⁺ PCs from MGUS patients than that on CD19⁺ PCs from the same bone marrow. The figure shows 5th, 25th, 50th, 75th, and 95th percentiles for expression, measured as mean fluorescence intensity.

previously described.^{1,4} Cells were incubated with CD38 allophycocyanin; CD19 phycoerythrin–cyanine 5.5; CD45 fluorescein isothiocyanate (in-house conjugates) combined with phycoerythrin-conjugated CCR1, CCR2, CCR3, CXCR-5 (R&D, Minneapolis, MN), CCR5, CCR6, CXCR1, CXCR2, CXCR3, or CXCR4 (BD Biosciences, Mountain View, CA); and a minimum of 3000 PCs were studied. Expression levels were recorded as the geometric mean fluorescence intensity and compared using the Student *t* test.

Chemokine receptor expression was assessed on PCs from 20 normal and 19 myeloma BMs. CCR3, CCR5, CXCR2, and CXCR3 were not expressed on PCs (< 3-fold difference in expression with control). In contrast, CCR1, CCR2, CCR6, CXCR1, CXCR4, and CXCR5 were expressed on both types of PCs, in keeping with data from other groups.^{5–9} The differences in chemokine receptor expression between normal and myeloma PCs were small (Figure 1A). CCR1 and CXCR5 showed no difference in expression between the 2 cell types. CCR2, CCR6, and CXCR1 showed decreased expression on myeloma PCs compared with normal PCs (average 3.3, 1.5, and 1.8 fold, respectively). In contrast, CXCR4 was up-regulated on myeloma PCs in comparison to normal PCs (average 2.2 fold). This suggests that PCs have a specific chemokine receptor expression profile; however, differences in the level of expression of some receptors, particularly CCR2 and CXCR1, may explain the abnormal localization of myeloma PCs.

These differences were studied further by assessing normal and neoplastic PCs from the same BM in 20 MGUS patients. CCR6, CXCR1, and CXCR4 showed no significant difference in expression between the 2 cell types, suggesting that feedback loops

between neoplastic PCs and BM stroma also influence normal PC chemokine receptor expression. In contrast, CCR2 expression on CD19⁺ PCs was on average 1.6-fold lower than on their CD19⁺ counterparts (range, 1.1–6.8 fold), suggesting that a key feature of neoplastic PCs is the down-regulation of CCR2 (Figure 1B). This reduced expression is therefore not a consequence of differences in BM microenvironment but rather a function of the neoplastic process or of the stage of differentiation of the originating neoplastic cell. Interfering with chemokines and their receptors that are related to malignant transformation may prove useful as adjunct to chemotherapy approaches.

Mona H. Al Rayes, Andy C. Rawstron, Gareth J. Morgan, and Faith E. Davies

Correspondence: Faith E. Davies, Haemato-oncology Unit, Institute of Cancer Research and Royal Marsden Hospital, Downs Road, Sutton, Surrey, United Kingdom; e-mail: faith.davies@icr.ac.uk.

Supported by the British Society of Haematology (A.C.R., F.E.D.), Leukaemia Research Fund UK (F.E.D., G.J.M.), Department of Health (F.E.D.), International Myeloma Foundation UK (A.C.R.), and The Egyptian Missions Department (M.H.A.R.).

References

- Rawstron AC, Davies FE, DasGupta R, et al. Flow cytometric disease monitoring in multiple myeloma: the relationship between normal and neoplastic PCs predicts outcome after transplantation. *Blood*. 2002;100:3095–3100.
- Ocqueteau M, Orfao A, Almeida J, et al. Immunophenotypic characterization of PCs from monoclonal gammopathy of undetermined significance patients: implications for the differential diagnosis between MGUS and multiple myeloma. *Am J Pathol*. 1998;152:1655–1665.
- Rawstron AC, Fenton JAL, Gonzalez D, et al. High-risk MGUS: identification by immunophenotype, karyotype, and clonal homogeneity [abstract 116]. *Blood*. 2003;102:36.
- Rawstron AC, Fenton JA, Ashcroft J, et al. The interleukin-6 receptor alpha-chain (CD126) is expressed by neoplastic but not normal PCs. *Blood*. 2000;96:3880–3886.
- Hargreaves DC, Hyman PL, Lu T, et al. A coordinated change in chemokine responsiveness guides PC movement. *J Exp Med*. 2001;194:45–56.
- Nakayama T, Hieshima K, Izawa D, Tatsumi Y, Kanamaru A, Yoshie O. Cutting edge: profile of chemokine receptor expression on human PCs accounts for their efficient recruitment to target tissues. *J Immunol*. 2003;170:1136–1140.
- Broek IV, Asosingh K, Vanderkerken K, Straetmans N, Van Camp B, Van Riet I. Chemokine receptor CCR2 is expressed by human multiple myeloma cells and mediates migration to bone marrow stromal cell-produced monocyte chemoattractant proteins MCP-1, -2 and 3. *Br J Cancer*. 2003;88:855–862.
- Durig J, Schmucker U, Duhrsen U. Differential expression of chemokine receptors in B cell malignancies. *Leukemia*. 2001;15:752–756.
- Moller C, Stromberg T, Turemalm M, Nilsson K, Nilsson G. Expression and function of chemokine receptors in human multiple myeloma. *Leukemia*. 2003;17:203–210.

To the editor:

Patients with thalassemia in the United States

Thalassemia is one of the most common monogenic diseases of man, prevalent in tropical and subtropical regions of the world. Population migration during the past decades has led to increasing numbers of these patients being encountered in all parts of the world, including in the United States. A clinical research network, supported by National Institutes of Health (NIH), was established to study β -thalassemia major/intermedia. A report on disease complications was published in July 2004, based on 342 patients (222 in the United States and 120 in Toronto).¹

Our laboratory was established in 2003 to provide molecular diagnosis of hemoglobinopathies and thalassemias. In the past

18 months, we have genotyped 40 β -thalassemia major/intermedia patients of all ethnic backgrounds (Table 1). Only 4 cases are of Mediterranean ancestry, hitherto the source of most β -thalassemia patients in our country. Seventy percent of the cases are 15 years of age or younger. They reside in Massachusetts (18), Rhode Island (3), West Virginia (4), New York (1), Maryland (3), and Georgia (11). Surprisingly, only one patient in this recently diagnosed cohort has the β -thalassemia mutation genotype [IVS1-6 (T>C)/IVS1-110 (G>A)] that corresponds to 1 of the 5 genotypes reported to be most commonly found in the United States.¹

These findings highlight the fact that β -thalassemia is more

Table 1. Forty recently diagnosed β -thalassemia major/intermedia patients and their β -thalassemia mutations

Patient no.	Age, y	Sex	Ethnic background	β -thalassemia mutation on one <i>HBB</i> allele	β -thalassemia mutation on the other <i>HBB</i> allele
1	12	M	African American	Nt-88 (C>T)	Polyadenylation signal (AATAAA>AACAAA)
2*	7	M	Syrian	Nt -87 (C>G)	Nt -87 (C>G)
3*	13	M	Syrian	Nt -87 (C>G)	Nt -87 (C>G)
4*	25	M	Syrian	Nt -87 (C>G)	Nt -87 (C>G)
5	16	F	Greek	Nt -87 (C>G)	Codon 39 (CAG>TAG)
6	14	F	Ghanaian	Nt -29 (A>G)	Nt -29 (A>G)
7	3.5	F	Asian	Nt -28 (A>G)	Codons 41/42 (-CTTT)
8	1.75	F	Chinese	Nt -28 (A>G)	Codons 71/72 (+A)
9	34	F	Pakistani	Cap + 1 (A>C)	IVSI (-1) (AGGT>AAGT)
10	11	M	Afghani	Codon 5 (-CT)	Codon 16 (-C)
11	< 1	F	Pakistani	Codons 8/9 (+G)	Codons 8/9 (+G)
12	16	M	Cambodian	Codon 17 (AAG>TAG)	Codon 17 (AAG>TAG)
13	6	F	Asian	Codon 17 (AAG>TAG)	Codons 41/42 (-CTTT)
14	1	F	Laotian	Hb E; Codon 26 (GAG>AAG)	Codons 41/42 (-CTTT)
15†	1.5	F	Laotian	Hb E; Codon 26 (GAG>AAG)	Codons 41/42 (-CTTT)
16†	7	M	Laotian	Hb E; Codon 26 (GAG>AAG)	Codons 41/42 (-CTTT)
17	4	M	East Indian	IVSI-1 (G>T)	IVSI-1 (G>T)
18	19	F	East Indian	IVSI-1 (G>T)	IVSI-5 (G>C)
19	5	F	Asian	IVSI-1 (G>T)	IVSI-5 (G>C)
20	12	M	Asian	IVSI-1 (G>T)	Codons 41/42 (-CTTT)
21	14	M	East Indian	IVSI-1 (G>T)	IVSII-837 (T>G)
22	< 1	M	Bangladesh	IVSI-5 (G>C)	IVSI-5 (G>C)
23	15	F	Asian	IVSI-5 (G>C)	Codons 41/42 (-CTTT)
24	11	M	Asian	IVSI-5 (G>C)	3' 619-bp deletion
25	26	M	Bangladesh	IVSI-5 (G>C)	HPFH-3, Indian type
26	26	M	Lebanese	IVSI-6 (T>C)	IVSI-6 (T>C)
27	31	F	Greek	IVSI-6 (T>C)	IVSI-6 (T>C)
28	12	M	Lebanese	IVSI-6 (T>C)	IVSI-110 (G>A)
29‡	4	F	Irish/Italian/Portuguese	IVSI-6 (T>C)	Codon 39 (CAG>TAG)
30‡	4	F	Irish/Italian/Portuguese	IVSI-6 (T>C)	Codon 39 (CAG>TAG)
31	7	F	Hispanic	IVSI-6 (T>C)	Codon 39 (CAG>TAG)
32§	8	M	Cuban	IVSI-6 (T>C)	Codon 39 (CAG>TAG)
33§	9	M	Cuban	IVSI-6 (T>C)	Codon 39 (CAG>TAG)
34	29	F	Italian	IVSI-6 (T>C)	IVSII-1 (G>A)
35	3	F	Irish	Codon 91 (-T) ²	No mutation
36	25	F	Irish	Codon 91 (-T) ²	No mutation
37	51	F	Irish	Codon 91 (-T) ²	No mutation
38	53	F	Irish	Codon 91 (-T) ²	No mutation
39¶	< 1	F	Egyptian	IVSII-745 (C>G)	Hb Lepore Boston-Washington
40¶	15	M	Egyptian	IVSII-745 (C>G)	Hb Lepore Boston-Washington

Nt indicates nucleotide; IVS, intron; and HPFH-3, hereditary persistence of fetal hemoglobin 3.

*Patients are the same kindred.

†Patients are siblings.

‡Patients are twins.

§Patients are siblings.

||Dominant β -thalassemia in the same kindred.²

¶Patients are siblings.

common and diverse than generally thought of in the United States; many of the recently diagnosed patients are not of the Mediterranean ancestry any more. They also suggest that carrier screening and genetic counseling may not be used optimally for this disease (12 of our patients are 5 years of age or younger). Additionally, our laboratory has diagnosed 11 cases of clinically significant hemoglobin (Hb) H disease³ from Massachusetts, New York, and Georgia: 1 (-MED/- $\alpha^{3.7}$); 5 (-SEA/- $\alpha^{3.7}$), including 1 Hb E heterozygote; and 5 (-SEA/ $\alpha^{Constant Spring \alpha}$), including 2 Hb E heterozygotes. Efforts devoted to community public education appropriately adapted to different cultures and languages, continuous education for health care providers, and research in disease pathophysiology and treatment for α - and β -thalassemias are needed.

Hong-yuan Luo, Jeanne Boudreaux, Martin H. Steinberg, and David H. K. Chui

Correspondence: David H. K. Chui, Hemoglobin Diagnostic Reference Laboratory, Boston Medical Center, 88 East Newton Street, Boston, MA 02118; e-mail: david.chui@bmc.org.

Supported in part by National Heart, Lung, and Blood Institute (NHLBI) Cooperative Agreement 1U54 HL70819.

References

- Cunningham MJ, Macklin EA, Neufeld EJ, Cohen AR. Complications of β -thalassemia major in North America. *Blood*. 2004;104:34-39.
- Luo H-Y, Tang W, Eung SH, et al. Dominantly inherited β -thalassemia intermedia due to a new single nucleotide deletion in exon 2 of the β -globin gene: Hb Morgantown ($\beta 91$ CTG>CG). *J Clin Pathol*. In press.
- Chui DHK, Fucharoen S, Chan V. Hemoglobin H disease: not necessarily a benign disorder. *Blood*. 2003;101:791-800.

To the editor:

The Marburg I polymorphism of factor VII–activating protease is not associated with venous thrombosis

Table 1. Marburg I polymorphism and the risk of venous thrombosis

FSAP nt 1601	Patients, no. (%)	Controls, no. (%)	OR	95% CI
GG	444 (94.3)	441 (93.6)	1*	NA
GA	26 (5.5)	28 (6.0)	0.9	0.5-1.6
AA	1 (0.2)	2 (0.4)	0.5	0.05-5.5
GA + AA	27 (5.7)	30 (6.4)	0.9	0.5-1.5

There were 471 patients and 471 controls.
nt indicates nucleotide; NA, not applicable.

*Reference category.

In a recently published paper in *Blood*, Hoppe et al¹ showed in 213 patients and 213 controls that carriers of the Marburg I polymorphism of the factor VII–activating protease (FSAP)² had an increased risk of venous thrombosis (odds ratio [OR], 3.5; 95% confidence interval [CI], 1.2-10.0).

FSAP is a serine protease that has 2 functions in hemostasis. It activates factor VII, thereby promoting coagulation,³ but it can also activate single-chain plasminogen activators, thus promoting fibrinolysis.⁴ Recently, a single nucleotide polymorphism (1601G>A) was discovered in the gene coding for FSAP, which results in the substitution of glycine 511 by glutamic acid (FSAP Marburg I) and which is present in 2% to 9% of the white population.^{1,5,6} The Marburg I variant has an impaired prourokinase activating potency, whereas it can still activate factor VII normally.⁷

If true, the finding of Hoppe et al¹ would support the hypothesis that reduced fibrinolysis contributes to the risk of venous thrombosis. However, the authors indicated that their control group might be biased due to the exclusive inclusion of healthy blood donors and that larger studies were needed to validate their results. Therefore, we determined the Marburg I polymorphism in 471 consecutive patients with a first episode of deep venous thrombosis (DVT) and 471 sex- and age-matched healthy controls of the Leiden Thrombophilia Study (LETS), a case-control study on the causes of venous thrombosis.⁸

We determined the Marburg I 1601G>A polymorphism with a 5' nuclease/TaqMan assay (Assay by design; Applied Biosystems, Foster City, CA).⁹ Nucleotide sequences of primers and probes are available on request. An odds ratio with 95% CI was calculated as a measure of the relative risk of thrombosis for carriers of the Marburg I allele (homozygous or heterozygous) compared with homozygous wild-type allele carriers. Factor VII activity was measured previously using Thromborel S reagent (Behringwerke, Warburg, Germany) and factor VII–deficient plasma.¹⁰ In Table 1, the risk of venous thrombosis is shown for the Marburg I polymorphism. Marburg I was found in 30 controls (allele frequency 0.034) and 27 cases (allele frequency 0.030). No association between Marburg I and venous thrombosis was found. Similar results were obtained when the analysis was stratified by sex or age (individuals < 45 years of age versus individuals ≥ 45 years of

age). Factor VII activity was not influenced by the presence of Marburg I.

Our results indicate, in contrast with the finding of Hoppe et al,¹ that the Marburg I allele of FSAP is not a risk factor for venous thrombosis. This difference may be explained by the frequency of the Marburg I allele in the control population studied by Hoppe et al,¹ which was considerably lower (0.012) than that reported in other studies, including ours (0.023-0.043).^{5,6} In thrombosis patients the frequency of the Marburg I allele was similar in the study of Hoppe et al¹ (0.039) and ours (0.030), so the low prevalence in the control group (5/426) seems to be the explanation for the findings by Hoppe et al, either because these were blood donors, in which case the mutant allele would be infrequent in a group selected on health, or because the control group was relatively small. On the other hand we cannot exclude small geographic differences in the Marburg I frequency.

**Rick van Minkelen, Marieke C. H. de Visser, Hans L. Vos,
Rogier M. Bertina, and Frits R. Rosendaal**

Correspondence: Rick van Minkelen, Department of Hematology, Haemostasis and Thrombosis Research Center, C2-R, Leiden University Medical Center, PO box 9600, 2300 RC, Leiden, the Netherlands; e-mail: r.van_minkelen@lumc.nl.

Supported by grant 912-02-036 from the Netherlands Organization for Scientific Research (NWO). The LETS study was supported by grant 89-063 from the Netherlands Heart Foundation.

References

- Hoppe B, Tolou F, Radtke H, et al. Marburg I polymorphism of factor VII–activating protease is associated with idiopathic venous thromboembolism. *Blood*. 2005;105:1549-1551.
- Romisch J. Factor VII activating protease (FSAP): a novel protease in hemostasis. *Biol Chem*. 2002;383:1119-1124.
- Romisch J, Feussner A, Vermohlen S, Stohr HA. A protease isolated from human plasma activating factor VII independent of tissue factor. *Blood Coagul Fibrinolysis*. 1999;10:471-479.
- Romisch J, Vermohlen S, Feussner A, Stohr H. The FVII activating protease cleaves single-chain plasminogen activators. *Haemostasis*. 1999;29:292-299.
- Willeit J, Kiechl S, Weimer T, et al. Marburg I polymorphism of factor VII–activating protease: a prominent risk predictor of carotid stenosis. *Circulation*. 2003;107:667-670.
- Ireland H, Miller GJ, Webb KE, Cooper JA, Humphries SE. The factor VII activating protease G511E (Marburg) variant and cardiovascular risk. *Thromb Haemost*. 2004;92:986-992.
- Roemisch J, Feussner A, Nerlich C, Stoeckl HA, Weimer T. The frequent Marburg I polymorphism impairs the pro-u-rokinase activating potency of the factor VII activating protease (FSAP). *Blood Coagul Fibrinolysis*. 2002;13:433-441.
- Koster T, Rosendaal FR, de Ronde H, et al. Venous thrombosis due to poor anticoagulant response to activated protein C: Leiden Thrombophilia Study. *Lancet*. 1993;342:1503-1506.
- Livak KJ. Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal*. 1999;14:143-149.
- Koster T, Rosendaal FR, Reitsma PH, et al. Factor VII and fibrinogen levels as risk factors for venous thrombosis: a case-control study of plasma levels and DNA polymorphisms—the Leiden Thrombophilia Study (LETS). *Thromb Haemost*. 1994;71:719-722.

Response:

Association of Marburg I polymorphism of factor VII-activating protease with venous thromboembolism is limited to idiopathic events

In this issue, van Minkelen et al report on venous thrombosis and the Marburg I variant of factor VII-activating protease (FSAP), commenting on a study published recently by our group in the journal.¹ They essentially describe 2 findings that appear to be in contrast to our study: (1) The frequency of carriers of *FSAP* Marburg I in the controls of van Minkelen et al was considerably higher (6.4%) than that of our control group (2.3%); and (2) an association of *FSAP* Marburg I with deep vein thrombosis (DVT) could not be confirmed.

In reply to these comments we want to remark the following. (1) Indeed, the frequency of carriers of *FSAP* Marburg I in our controls (5/213; 2.3%)¹ comprising healthy blood donors was lower than that of the Bruneck study cohort (37/810; 4.6%),² the controls of van Minkelen et al (30/471; 6.4%), and the Northwick Park Heart Study cohort (176/2066; 8.5%).³ Different frequencies in the described study populations could be due to low sample size, selection bias, and different geographic distributions. To exclude the possibility that our controls might be biased by negative selection of *FSAP* Marburg I due to its association with atherosclerosis,² we analyzed a larger, independent control group consisting of consecutive patients admitted to our institution for reasons other than venous thromboembolism (non-VTE patients). Main diagnoses of these patients included transient ischemic attack/stroke, family testing for hereditary thrombophilia, pregnancy complications, and myocardial infarction. Considering the positive association of *FSAP* Marburg I with progression of atherosclerosis² and assuming at least no negative association of this variant with VTE (Hoppe et al¹ and van Minkelen et al), the frequency of *FSAP* Marburg I carriers in these non-VTE patients (15/327; 4.6%) will presumably overestimate the normal frequency of this variant in our geographic region.

(2i) The data presented by van Minkelen et al included patients with DVT, making no distinction between idiopathic and secondary events. As we described in the study,¹ the association of *FSAP* Marburg I with VTE (including DVT and/or pulmonary embolism) was nearly exclusively attributable to idiopathic events (Table 1; "Comparison vs controls I"). Secondary VTE due to acquired risk factors (immobilization, surgery, trauma, pregnancy, puerperium, malignancy) was not associated with this variant. (ii) To avoid possible biases by the control group (blood donors), we performed a statistical analysis based on the second, independent control group (non-VTE patients; Table 1; "Comparison vs controls II"). Again, *FSAP* Marburg I was significantly associated with idio-

Table 1. Association of *FSAP* Marburg I with VTE

	No.	Carriers of <i>FSAP</i> Marburg I, no. (%)	Comparison vs controls I, odds ratio (95% CI)	Comparison vs controls II, odds ratio (95% CI)
Idiopathic VTE	103	12 (11.7)	5.5 (1.9,16.0)†*	2.7 (1.2,6.1)‡
Secondary VTE	110	5 (4.5)	2.0 (0.56,7.0)*	0.99 (0.35,2.8)
VTE	213	17 (8.0)	3.6 (1.3,10.0)*‡	1.8 (0.88,3.7)
Controls I	213	5 (2.3)	NA	NA
Controls II	327	15 (4.6)	NA	NA

Controls I indicates healthy blood donors; controls II, non-VTE patients; and NA, not applicable.

*The data presented here (univariate analyses) differ slightly from those of our initial report (logistic regression analyses; independent variables: factor V Leiden, prothrombin 20210G→A, and *FSAP* Marburg I).¹

†*P* = .001 (Fisher exact test).

‡*P* < .05 (Fisher exact test).

pathic but not secondary VTE (odds ratio, 2.7; 95% confidence interval [CI], 1.2-6.1).

The *FSAP* genotypes have no influence on its capacity to activate factor VII in vitro.² Moreover, the physiologic role of *FSAP* and the site where its action takes place remains unclear. Thus, the absence of a detectable difference in circulating factor VII activities with both *FSAP* genotypes as described by van Minkelen et al is not surprising.

Based on the additional data described here and as stated in our initial report,¹ *FSAP* Marburg I is associated with idiopathic VTE.

Berthold Hoppe, Guido A. Heymann, Hartmut Radtke, Holger Kiesewetter, Thomas Dörner, and Abdulgabar Salama

Correspondence: Berthold Hoppe, Institute of Transfusion Medicine, Campus Virchow-Klinikum, Charité—Universitätsmedizin Berlin, Augustenburger Platz 1, 13353 Berlin, Germany; e-mail: berthold.hoppe@charite.de.

References

- Hoppe B, Tolou F, Radtke H, Kiesewetter H, Dörner T, Salama A. Marburg I polymorphism of factor VII-activating protease is associated with idiopathic venous thromboembolism. *Blood*. 2005;105:1549-1551.
- Willeit J, Kiechl S, Weimer T, et al. Marburg I polymorphism of factor VII-activating protease: a prominent risk predictor of carotid stenosis. *Circulation*. 2003;107:667-670.
- Ireland H, Miller GJ, Webb KE, Cooper JA, Humphries SE. The factor VII activating protease G511E (Marburg) variant and cardiovascular risk. *Thromb Haemost*. 2004;92:986-992.