



Universiteit
Leiden
The Netherlands

SWS observations of solid CO₂ in molecular clouds

Graauw, Th. de; Whittet, D.C.B.; Gerakines, P.A.; Bauer, O.H.; Beintema, D.A.; Boogert, A.C.A.; ... ; Salama, A.

Citation

Graauw, T. de, Whittet, D. C. B., Gerakines, P. A., Bauer, O. H., Beintema, D. A., Boogert, A. C. A., ... Salama, A. (1996). SWS observations of solid CO₂ in molecular clouds. *Astron. Ap.*, 315, L345-L348. Retrieved from <https://hdl.handle.net/1887/2253>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/2253>

Note: To cite this publication please use the final published version (if applicable).

SWS observations of solid CO₂ in molecular clouds*

Th. de Graauw^{1,2}, D.C.B. Whittet³, P.A. Gerakines³, O.H. Bauer⁴, D.A. Beintema^{1,5}, A.C.A. Boogert², D.R. Boxhoorn¹, J.E. Chiar³, P. Ehrenfreund⁶, H. Feuchtgruber^{4,5}, F.P. Helmich⁶, A.M. Heras⁵, R. Huygen⁷, D.J.M. Kester¹, D. Kunze⁴, F. Lahuis⁵, K.J. Leech⁵, D. Lutz⁴, P.W. Morris^{5,8}, T. Prusti⁵, P.R. Roelfsema^{1,5}, A. Salama⁵, S.G. Schaeidt^{4,5}, W.A. Schutte⁶, H.W.W. Spoon⁸, A.G.G.M. Tielens⁹, E.A. Valentijn¹, B. Vandenbushe^{5,7}, E.F. van Dishoeck⁶, P.R. Wesselius¹, E. Wieprecht^{4,5}, and C.M. Wright⁴

¹ SRON, P.O. Box 800, 9700 AV Groningen, The Netherlands

² Kapteyn Astronomical Institute, PO Box 800, 9700 AV Groningen, The Netherlands

³ Department of Physics, Applied Physics & Astronomy, Rensselaer Polytechnic Institute, Troy, NY 12180, USA

⁴ Max-Planck-Institut für Extraterrestrische Physik, Postfach 1603, D-85740 Garching, Germany

⁵ ISO Science Operations Centre, Astrophysics Division, ESA, Villafranca del Castillo, P.O. Box 50727, E-28080 Madrid, Spain

⁶ Leiden Observatory, P.O. Box 9513, 2300 RA Leiden, The Netherlands

⁷ Sterrenkundig Instituut, K.U. Leuven, Celestijnenlaan 200B, B-3001 Heverlee, Belgium

⁸ SRON, Sorbonnelaan 2, 3584 CA Utrecht, The Netherlands

⁹ NASA Ames Research Center, Mail Stop 245-6, Moffett Field, CA 94035, USA

Received 1 July 1996 / Accepted 29 August 1996

Abstract. We report absorption features of solid CO₂ stretching and bending modes in several lines of sight, including embedded young stellar objects and the Galactic Center source Sgr A*. The overall CO₂ abundance in ices is $\sim 15\%$ relative to H₂O. Profile shapes are consistent with the presence of grain mantles with distinct polar (H₂O-rich) and nonpolar (CO or CO₂-rich) layers. In addition to the normal isotopic form, we report detection of the stretching mode of ¹³CO₂; the ¹²C/¹³C ratio is consistent with terrestrial and interstellar values.

Key words: ISM: molecules – dust, extinction – infrared: interstellar: lines – stars: pre-main sequence – Galaxy: center

1. Introduction

A key problem in astrophysics is to understand the cosmic evolution of carbon, from its creation by nucleosynthesis in post-main-sequence stars to its inclusion in living systems on the surface of the Earth. Between these extremes, dust grains provide a repository for condensed carbon in interstellar clouds and protoplanetary disks. In molecular clouds, refractory carbon and silicate grains acquire icy mantles, composed primarily

of H₂O, but also containing carbon-bearing species such as CH₄, CH₃OH and CO (Whittet 1993). The relative abundance of CH- and CO-bonded carbon in the ices is an important (but currently poorly constrained) parameter; this will affect chemical evolution when the ices are subject to irradiation and warm-up with the onset of star formation.

Ground-based observations showed that CO is a ubiquitous constituent of grain mantles in cold, dense regions of molecular clouds (Chiar et al. 1995 and references therein). Its abundance may range up to $\sim 50\%$ of the H₂O abundance. The presence of CO on grains naturally led to the prediction that CO₂ might also be present, as CO₂ forms readily in laboratory ice analogues containing CO when subject to energetic processing such as ultraviolet irradiation or particle bombardment (e.g. d’Hendecourt et al. 1986; Sandford et al. 1988). Profiles of the 4.67 μm CO feature in the spectra of several young stellar objects (YSOs) are best fit by laboratory mixtures of CO and CO₂ (Chiar et al. 1995), suggesting that energetic production of CO₂ is, indeed, occurring in the vicinities of these embedded stars.

Direct spectroscopic detection of interstellar CO₂ was previously hindered by the presence of CO₂ absorption in the Earth’s atmosphere. Both the strong stretching-mode resonance at 4.27 μm (2340 cm^{-1}) and the bending-mode resonance at 15.2 μm (660 cm^{-1}) require a satellite platform for detection. Observations of the bending mode with the low resolution spectrometer of the Infrared Astronomical Satellite have been reported (d’Hendecourt & Jourdain de Muizon 1989), but this instrument had insufficient spectral resolution to provide a very sensitive diagnostic. Following the launch of the Infrared Space Observatory (ISO; Kessler et al. 1996) it is possible to observe

Send offprint requests to: D.C.B. Whittet

* Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries France, Germany, the Netherlands and the United Kingdom) and with the participation of ISAS and NASA.




Fig. 1. Optical depth spectra of the ¹²CO₂ stretching mode feature in (a) GL 2136, (b) GL 2591, (c) GL 4176 and (d) Sgr A*. Observational data are represented by ‘x’ symbols. Best-fitting models based on laboratory data (thick lines) are also shown; polar and nonpolar components of the models are plotted separately as continuous and dashed lines, respectively.

both vibrational modes at appropriate resolving power. This Letter reports first detection of CO₂ in molecular clouds observed by ISO; an accompanying paper (van Dishoeck et al. 1996) reports a search for gas phase CO₂ in four of the five sources included here.

2. Observations

The ISO short wavelength spectrometer (SWS; de Graauw et al. 1996) was used to obtain spectra in the regions of the CO₂ stretching and bending modes (4.1–4.4 μm and 14–16 μm, respectively). Four of the infrared sources included in this study, GL 2136, GL 2591, GL 4176 and NGC 7538 IRS9, are thought to be YSOs embedded in molecular clouds. For comparison, we also discuss SWS data for the Galactic Center source Sgr A*, reported by Lutz et al. (1996): although often used as a probe of dust in the diffuse ISM, the presence of absorption by CO₂ and H₂O ices implies the presence of molecular material in this line of sight. The four YSOs were observed in SWS grating mode AOT06, giving a mean resolving power $\lambda/\Delta\lambda \sim 2000$. Sgr A* was observed in mode AOT01 (full grating scan, speed 4), yielding resolution a factor ~ 2 lower. The instrument, its calibration, and data reduction techniques are described in detail elsewhere in this volume.

3. Results and discussion

The stretching mode is detected in all five sources. The bending mode is seen in GL 2136, NGC 7538 IRS9 and Sgr A* (but is too weak in Sgr A* for detailed profile analysis). Only upper limits could be set on bending modes in GL 2591 and GL 4176. Optical depth spectra are shown in Fig. 1 (stretching mode) and Fig. 2 (bending mode), together with fits described below. Note that the stretching mode in NGC 7538 IRS9 (not shown) is saturated. Optical depths were calculated by fitting local continua through adjacent regions of the flux spectra (4.10–4.18 and 4.32–4.37 μm for the stretch mode; 14.6–14.8 and 15.8–16.0 μm for the bend mode). Positions, widths (FWHM) and peak optical depths are listed in Table 1. Weak absorptions at 4.39 μm due to the ¹³CO₂ stretching mode were also detected in GL 2136 ($\tau_{\max} \approx 0.04$) and NGC 7538 IRS9 ($\tau_{\max} \approx 0.1$). The ¹²CO₂/¹³CO₂ ratio is ~ 40 – 70 , comparable with terrestrial and interstellar ¹²C/¹³C ratios.

Gas phase absorption lines may cause fine structure in the solid state profiles. In the case of the stretch mode, the *P* and *R* branches of the ro-vibrational spectrum of gaseous CO₂ cannot be easily distinguished from the solid state feature (van Dishoeck et al. 1996). In the case of the bend mode, the unresolved *Q* branch causes excess absorption near 14.98 μm in the




Fig. 2. Optical depth spectra of the ¹²CO₂ bending modes in (a) GL 2136 and (b) NGC 7538 IRS9. Lines and symbols have the same meaning as in Fig. 1. Note that the model spectra do not attempt to fit narrow substructure in the profiles.

short-wavelength wing of the solid state feature: this structure is clearly seen in GL 2136 and NGC 7538 IRS9 (Fig. 2). Estimated gas phase CO₂ column densities are only a few percent of the solid state CO₂ column densities (van Dishoeck et al. 1996).

Solid state CO₂ column densities (Table 2) were calculated from the formula $N = \int \tau(\nu) d\nu / A$, where A is the band strength. The integration is carried out over the entire profile of the observed feature. A is only weakly dependent on the composition of the matrix containing the CO₂ (Gerakines et al. 1995), thus we effectively measure $N(\text{CO}_2)$ summed over all possible matrices. Also listed in Table 2 are results for H₂O and CO in ices, estimated from data in the literature. Note that $N(\text{CO})$ towards Sgr A* is based on a tentative detection in IRS12 contrasted with non-detections in IRS3 and IRS7 (McFadzean et al. 1989); all three sources lie in the SWS beam.

The five objects in our sample have $N(\text{CO}_2)/N(\text{H}_2\text{O})$ ratios in the range 0.12–0.16. It is interesting that the Galactic Center source, presumably obscured by foreground clouds, yields essentially the same result as the embedded YSOs. The relative CO abundance is lower, varying between 0.13 (NGC 7538 IRS9) and < 0.03 (GL 2591). No objects in the present study have solid CO abundances approaching the highest values seen in some dark clouds (Chiar et al. 1995). Together, CO and CO₂ account

Table 1. Positions, widths and optical depths of the observed ¹²CO₂ features. For the bending mode, the positions of the subpeaks are given in GL 2136 and NGC 7538 IRS9. Units for ν_0 and $\Delta\nu$ are cm⁻¹. A colon indicates an uncertain value.

Source	Stretching mode			Bending mode		
	ν_0	$\Delta\nu$	τ_{max}	ν_0	$\Delta\nu$	τ_{max}
GL 2136	2342.2	19.3	2.14	655.9 662.4	24.5	0.28
GL 2591	2344.8	25.8	0.71	—	—	<0.1
GL 4176	2345.3	19.7	0.37	—	—	<0.1
N 7538 IRS9	2345:	>20	>3	655.8 662.0	19.7	0.78
Sgr A*	2342.5	17.2	0.72	656.2	19:	0.07

Table 2. Solid state column densities of ¹²CO₂ from our results, together with values for H₂O and CO from the literature. All values are in units of 10¹⁷ cm⁻².

Source	$N(\text{CO}_2)$	$N(\text{H}_2\text{O})$	$N(\text{CO})$	Reference
GL 2136	6.1	50	1.8	1, 2
GL 2591	2.7	17	<0.5	3, 4
GL 4176	1.2	9	<0.5	5
N 7538 IRS9	12	80	10	2, 6
Sgr A*	1.5	12	≲1.5	7

References: 1. Schutte et al. (1996a); 2. Tielens et al. (1991); 3. Smith et al. (1989); 4. Lacy et al. (1984); 5. Ehrenfreund et al. (in preparation); 6. Schutte et al. (1996b); 7. McFadzean et al. (1989).

for a substantial fraction of the inventory of detected molecules in interstellar ices. This may be contrasted with the rather low abundance of CH bonded carbon in known species such as CH₃OH (5–10%) and CH₄ (~1%) (Whittet 1993; Boogert et al. 1996). Our current inventory is no doubt far from complete.

Laboratory spectra for various ice mixtures containing CO and CO₂ at 10–80 K (Ehrenfreund et al. 1996) were used to fit the observed CO₂ profiles. Model spectra were calculated using Mie theory for power-law size distributions of spherical particles in the small particle limit. The fitting routine selects the best fitting mixture or combination of mixtures by χ^2 minimization from a suite of 47 calculated spectra. The method is the same as previously used to fit ground-based solid CO spectra (Chiar et al. 1995). Results are summarized in Table 3 and plotted with the observed spectra in Figs. 1 and 2. Best fits were obtained in every case by combining a polar mixture (dominated by H₂O) with a nonpolar mixture (dominated by CO₂) at temperatures 10–30 K, consistent with studies of solid CO. However, it is generally the nonpolar ices that contribute greatest optical depth in the CO feature, whereas polar ices dominate the CO₂ features. Hence, the distribution of CO and CO₂ between polar and nonpolar ices must be different. This is expected if CO in the more volatile nonpolar ices is partially desorbed in conditions that promote

Table 3. Summary of best-fitting laboratory mixtures.

Source	Mixtures	T (K)	τ_{\max}
	Stretching mode:		
GL 2136	H ₂ O:CO ₂ :CO = 100:8:8	30	2.02
	H ₂ O:CO ₂ = 1:10	10	0.89
GL 2591	H ₂ O:CO ₂ :CO = 100:8:8	30	0.63
	H ₂ O:CO ₂ = 1:10	10	0.42
GL 4176	H ₂ O:CO ₂ :CO = 100:5:10	10	0.32
	H ₂ O:CO ₂ = 1:10	10	0.18
Sgr A*	H ₂ O:CO ₂ :CO = 100:8:8	30	0.60
	CO:CO ₂ = 100:50	10	0.30
	Bending mode:		
GL 2136	H ₂ O:CO ₂ :CO = 100:5:10	10	0.20
	H ₂ O:CO ₂ = 1:10	10	0.19
N 7538 IRS9	H ₂ O:CO ₂ = 100:20	10	0.53
	H ₂ O:CO ₂ = 1:10	10	0.46

CO₂ formation, whereas CO in the polar ices is converted more efficiently into CO₂.

The bending modes in GL 2136 and NGC 7538 IRS9 show considerable structure, with subpeaks at 15.10 and 15.25 μm and a shoulder at 15.4–15.6 μm (Fig. 2). We interpret this structure in terms of a broad ($\sim 0.5 \mu\text{m}$) underlying component, peaking around $\sim 15.2 \mu\text{m}$, with separate narrow component(s) responsible for the subpeaks. The broad component is well fit by essentially the same combination of polar and nonpolar mixtures that fit the stretching mode (Table 3). The subpeaks might arise in a minor phase (containing $< 15\%$ of the total CO₂) which is indistinguishable in the stretching region. Two classes of model might explain them. First, bulk spectra of *pure* CO₂ show splitting of the bending mode (this vibration is doubly degenerate and splits when the axial symmetry is broken). However, particle shape influences the positions of the subpeaks (Ehrenfreund et al. 1996), and good fits to the observed positions are only obtained for a very restrictive set of parameters (oblate or prolate spheroids with axial ratios 0.1 ± 0.05). Alternatively, the narrow subpeaks might reflect the presence of traces of CO₂ in an environment which allows multiple trapping sites: e.g., a CO:O₂:CO₂ = 100:50:8 mixture shows subpeaks at 15.12 and 15.19 μm , near those in the interstellar spectra. However, the subpeaks are very sensitive to matrix conditions and concentrations (see Fig. 3 of Ehrenfreund et al. 1996). Thus, both explanations for the substructure require very special conditions, either in grain population or in mantle composition. It is clear that, in principle, much information on interstellar ices is contained within these features.

Acknowledgements. We are indebted to the SRON–MPE SWS teams and the SIDT. This work is supported by NASA (grants NAGW–3144 and NAGW–4039) and by the Netherlands Organization for Scientific Research (NWO). MPE is supported by DARA grants 50QI8610 8 and 50QI9402 3.

References

- Boogert A.C.A., Schutte W.A., Tielens A.G.G.M., et al., 1996, A&A this volume
- Chiar J.E., Adamson A.J., Kerr T.H., Whittet D.C.B., 1995, ApJ 455, 234
- de Graauw Th., Haser L.N., Beintema D.A., et al., 1996a, A&A this volume
- d’Hendecourt L.B., Allamandola L.J., Grim R.J.A., Greenberg J.M., 1986, A&A 158, 119
- d’Hendecourt L.B., de Muizon M.J., 1989, A&A 223, L5
- Ehrenfreund P., Boogert A., Gerakines P.A., Schutte W.A., Jansen D., van Dishoeck E.F., 1996, A&A this volume
- Gerakines P.A., Schutte W.A., Greenberg J.M., van Dishoeck E.F., 1995, A&A 296, 810
- Kessler M.F., et al., 1996, A&A this volume
- Lacy J.H., Baas F., Allamandola L.J., et al., 1984, ApJ 276, 533
- Lutz D., Feuchtgruber H., Genzel R., et al., 1996, A&A this volume
- McFadzean A.D., Whittet D.C.B., Longmore A.J., Bode M.F., Adamson A.J., 1989, MNRAS 241, 873
- Sandford S.A., Allamandola L.J., Tielens A.G.G.M., Valero, G.J., 1988, ApJ 329, 498
- Schutte W.A., Gerakines P.A., Geballe T.R., van Dishoeck E.F., Greenberg J.M., 1996a, A&A 309, 633
- Schutte W.A., Tielens A.G.G.M., Whittet D.C.B., et al., 1996b, A&A this volume
- Smith R.G., Sellgren K., Tokunaga A.T., 1989, ApJ 344, 413
- Tielens A.G.G.M., Tokunaga A.T., Geballe T.R., Baas, F., 1991, ApJ 381, 181
- van Dishoeck E.F., Helmich F.P., de Graauw Th., et al., 1996, A&A this volume
- Whittet D.C.B., 1993, in *Dust and Chemistry in Astronomy*, eds. T.J. Millar & D.A. Williams, (Cambridge University Press), p. 9