



Universiteit  
Leiden  
The Netherlands

## **Circulating gut-associated antigens of *Schistosoma mansoni* : biological, immunological, and molecular aspects**

Dam, G.J. van

### **Citation**

Dam, G. J. van. (1995, February 9). *Circulating gut-associated antigens of Schistosoma mansoni : biological, immunological, and molecular aspects*. Retrieved from <https://hdl.handle.net/1887/41317>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/41317>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/41317> holds various files of this Leiden University dissertation.

**Author:** Dam, G.J. van

**Title:** Circulating gut-associated antigens of *Schistosoma mansoni* : biological, immunological, and molecular aspects

**Issue Date:** 1995-02-09

**Circulating gut-associated antigens  
of *Schistosoma mansoni*:  
biological, immunological, and  
molecular aspects**



**Circulating gut-associated antigens  
of *Schistosoma mansoni*:  
biological, immunological,  
and molecular aspects**

PROEFSCHRIFT

ter verkrijging van de graad van Doctor  
aan de Rijksuniversiteit te Leiden,  
op gezag van de Rector Magnificus Dr. L. Leertouwer,  
hoogleraar in de faculteit der Godgeleerdheid,  
volgens besluit van het College van Dekanen  
te verdedigen op donderdag 9 februari 1995  
te klokke 16.15 uur

door

Govert Jan van Dam

geboren te Gouda in 1960

**Promotiecommissie**

Promotor	Prof. Dr. A.M. Deelder
Co-promotor	Dr. J.P. Rotmans
Referent	Prof. Dr. A.E. Butterworth (Cambridge University, Cambridge, UK)
Overige leden	Prof. Dr. B.J. Bogitsh (Vanderbilt University, Nashville, USA) Prof. Dr. M.R. Daha Prof. Dr. J.P. Kamerling Prof. Dr. W.J.M. Spaan

The work described in this thesis was performed at the Department of Parasitology, University of Leiden (Head: Prof. Dr. A.M. Deelder). The financial support for the cost of printing given by the Hubrecht Janssen Fonds, and the Leids Universiteits Fonds is gratefully acknowledged.

*our little systems have their day  
they have their day and cease to be  
they are but broken lights of Thee  
but Thou O Lord art more than they*

*Alfred Tennyson*

*aan mijn vader en moeder  
aan Betty  
aan Rieke, Paul en Jaap*

Cover illustration: gut-associated antigens of *Schistosoma mansoni*, visualized by immunofluorescence microscopy (McAb 99-1G3-A) on a section of adult male worms fixed in Rossman's fixative

CIP-GEGEVENS KONINKLIJKE BIBLIOTHEEK, DEN HAAG

Dam, Govert Jan van

Circulating gut-associated antigens of *Schistosoma mansoni*:

biological, immunological, and molecular aspects /

Govert Jan van Dam. - [S.l. : s.n.] - III.

Proefschrift Leiden. - Met lit. opg. - Met samenvatting in het Nederlands.

ISBN 90-9007796-0

Trefw.: schistosomiasis / biochemie / immunologie.



---

## Contents

Preface	ix
1. <i>Schistosoma</i> gut-associated antigens	1
2. <i>Schistosoma</i> carbohydrates in host-parasite (immunological) interactions	19
2.1. Glycoconjugates in general - short review	19
2.2. Glycoconjugates of <i>Schistosoma</i>	26
3. <i>Schistosoma</i> : Analysis of monoclonal antibodies reactive with the circulating antigens CAA and CCA	47
4. <i>Schistosoma mansoni</i> : analysis of monoclonal antibodies reactive with gut-associated antigens	73
5. Application of the FITC-anti-FITC-gold system to ultrastructural localization of antigens	91
6. Detection of IgM antibodies directed against the gut-associated circulating cathodic antigen in sera from <i>Schistosoma mansoni</i> infected patients	101
7. The immunologically reactive O-linked polysaccharide chains derived from circulating cathodic antigen isolated from the human blood fluke <i>Schistosoma mansoni</i> have Lewis x as repeating unit	117
8. The immunologically reactive part of the immunopurified circulating anodic antigen from <i>Schistosoma mansoni</i> is a threonine-linked polysaccharide consisting of $\rightarrow 6$ )-[ $\beta$ -D-GlcpA-(1 $\rightarrow$ 3)]- $\beta$ -D-GalpNAc-(1 $\rightarrow$ repeating units	151
9. <i>Schistosoma mansoni</i> excretory antigen CCA shares carbohydrate epitopes with human granulocytes and evokes host antibodies mediating complement-dependent lysis of granulocytes	173
10. <i>Schistosoma mansoni</i> circulating anodic antigen but not circulating cathodic antigen interacts with complement component C1q	189

---

11. <i>Schistosoma mansoni</i> ; <i>in vitro</i> and <i>in vivo</i> excretion of CAA and CCA by developing schistosomula and adult worms	205
12. General discussion	225
Summary	237
Samenvatting	241
Appendices	245
I. List of abbreviations	246
II. List of figures	247
III. List of tables	250
IV. Structure of polysaccharides from CAA	252
V. Structure of polysaccharides from CCA	253
Acknowledgements	255
Curriculum vitae	257

---

## **Preface**

---



## Preface

Human schistosomiasis (bilharzia) is one of the major parasitic diseases in the world, affecting 200 million people predominantly in third world countries. In areas where the disease is highly prevalent it causes important health problems, and it also has socially-economic effects on the population. As the parasite is transmitted through surface water, increasing irrigation measures may also contribute to expansion of the disease. Laboratory research on schistosomiasis is mainly directed at the (immunological) interaction of the parasite and the host and the improvement of diagnostic methods. Besides having fundamental scientific interest and/or using *Schistosoma* infection as an immunological model, research efforts may also lead to improved therapeutic strategies or, finally, to the development of an effective vaccine.

Schistosomiasis is caused by the presence of the blood-fluke *Schistosoma* (Trematoda) in the blood-vessels of mammalian hosts. The current method for diagnosis of schistosomiasis in developing countries is the parasitological examination of urine and faeces for the presence of *Schistosoma* eggs. An alternative method which is now increasingly used is based on the detection of *Schistosoma* antigens in the circulatory system or the urine of the host. The gut of the parasite is an important source of these antigens since many gut-associated antigens are excreted into the circulation of the host following digestion of food (e.g. blood cells, proteins) by the parasite. Two major gut-associated antigens which have been thoroughly studied with regard to diagnostic detectability, are the circulating anodic antigen (CAA) and the circulating cathodic antigen (CCA).

**Chapter 1** of this thesis includes a general introduction to *Schistosoma mansoni* gut-associated antigens with particular emphasis on CAA and CCA, and deals also shortly with host-parasite interactions. In addition, a literature overview is presented dealing with localization of the antigens in the parasite, detection in the host circulation, as well as development of the host immune response. As important components of these antigens are carbohydrate structures, in **Chapter 2** a short general overview is given on glycoconjugates with respect to their structure and techniques used for structural analysis. Examples given are mostly from the field of parasitology and schistosomiasis research.

To improve the detection of CAA and CCA a large panel of monoclonal antibodies (McAbs) not only directed against CAA and CCA, but also against various other *Schistosoma* antigens was developed. An analysis of anti-CAA and anti-CCA McAbs generated during the ongoing research of many years is

presented in **Chapter 3**. To detect other gut-associated antigens as potential candidates for immunodiagnosis, a study was undertaken to analyze McAbs reactive with *Schistosoma mansoni* gut-associated antigens other than CAA, CCA, or well-studied gut proteases. The outcome of this study, described in **Chapter 4**, corroborated the important role of CCA. **Chapter 5** describes a technical improvement in the detection or ultrastructural localization of antigens using an anti-FITC McAb. As an example the detection of CCA in *S. mansoni* adult worms is used. In a routinely used serodiagnostic assay (immunofluorescence on sections of adult worms) human IgM antibodies are measured which are directed against gut-associated antigens (predominantly against CCA). To more specifically study the IgM response against CCA and to simplify this detection an ELISA system was developed with the use of immunopurified CCA, as described in **Chapter 6**.

The immunopurification of sufficiently large amounts of CAA and CCA allowed for their structural analysis. The elucidation of the primary structures of the carbohydrate parts, most relevant immunodiagnostically and immunologically, is described in **Chapters 7** and **8**. Because the carbohydrate structure of CCA was found to resemble a major granulocyte surface antigen, the possible role of anti-CCA antibodies in granulocytotoxicity was investigated in **Chapter 9**. In **Chapter 10** a possible function for CAA is described, as it was shown to interact with the first complement component C1q, which may result in an interference with cellular cytotoxicity mechanisms against the parasite. Finally, to contribute to the study of the possible physiological role of CAA and CCA, the *in vivo* and *in vitro* excretion patterns of CAA and CCA by newly transformed and developing schistosomula as well as by 7-week old adult worms were investigated and discussed with respect to clearance mechanisms in **Chapter 11**. This thesis concludes with a short general discussion in **Chapter 12**.