



Universiteit
Leiden
The Netherlands

Circulating gut-associated antigens of *Schistosoma mansoni* : biological, immunological, and molecular aspects

Dam, G.J. van

Citation

Dam, G. J. van. (1995, February 9). *Circulating gut-associated antigens of Schistosoma mansoni : biological, immunological, and molecular aspects*. Retrieved from <https://hdl.handle.net/1887/41317>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/41317>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/41317> holds various files of this Leiden University dissertation.

Author: Dam, G.J. van

Title: Circulating gut-associated antigens of *Schistosoma mansoni* : biological, immunological, and molecular aspects

Issue Date: 1995-02-09

Chapter 2

***Schistosoma* carbohydrates in host–parasite (immunological) interactions**

Chapter 2

***Schistosoma* carbohydrates in host–parasite (immunological) interactions**

2.1. Glycoconjugates in general – short review.


Structural characteristics of glycoconjugates

Glycoconjugates are molecules which contain one or more carbohydrate structures, like glycoproteins, glycolipids, proteoglycans or lipopolysaccharides. Glycosylation of proteins, the attachment of one or more carbohydrate chains (glycans) to a protein backbone, is a very common phenomenon shared by almost all organisms. The biological roles of these carbohydrate units include protection against proteolytic attack, induction and maintenance of the 3-dimensional conformation in a biologically active form, facilitation of the extracellular secretion, direction and modulation of the immune response, provision of ligand structures for cell–adhesive molecules (see recent reviews [46,98,181] and papers cited therein).

The major linkages described between the protein and the oligosaccharide or polysaccharide side-chains are N-glycosidic (to asparagine) or O-glycosidic (to serine or threonine). The major types of N-linked carbohydrate chains are summarized in Table 1, and seven types of core structures found in O-linked glycans of the mucin-type in Table 2. These core structures may be extended with poly-*N*-acetylglucosamine chains, containing repeating units consisting of Gal and GlcNAc [111]. These polysaccharide side-chains may be substituted with *e.g.* Fuc [165] and/or Neu5Ac (in terminal position) [109], and have been shown to carry various antigenic determinants such as I/i, ABO blood group and Lewis antigens (reviewed in *e.g.* [43,45]). Lewis x (Le^x, also called SSEA-1 or X-hapten [82], being the dominant epitope in lacto-*N*-fucopentaose-III, LNFP-III [75,93,174,182]) and sialyl-Le^x, present on granulocytes and monocytes, may serve as ligands for adhesion molecules (L-selectin and P-selectin) on endothelial cells and platelets [22,77,166,170], thus playing a critical role in the recruitment of leukocytes to inflammatory sites [84]. Besides

these roles in normal physiological processes, Le^X-containing glycoproteins and glycolipids are also considered to be onco-developmental antigens [43,59] and in some human neoplasms, the expression of Le^X or sialyl-Le^X appears to be associated with increased metastasis or tumor relapse [121,152,155].

Table 1. Examples of oligomannose, *N*-acetylactosamine, hybrid and xylose-containing types of asparagine-linked carbohydrate chains.^a










^a structures described *e.g.* in [4,81,95,117,180]

Proteoglycans are a class of glycoconjugates found in *e.g.* animal connective tissues and have as a common structural characteristic that they contain several linear polysaccharide chains (named glycosaminoglycans) generally composed of disaccharide repeating units of a uronic acid (GlcA or IdoA) and GlcNAc or GalNAc. Glycosaminoglycans are highly negatively charged at neutral pH and

most of them are sulphated, *e.g.* chondroitin sulphate and dermatan sulphate. The glycosaminoglycans are usually covalently bound to the protein backbone by Xyl–Ser linkages, but also GalNAc–Ser/Thr or GlcNAc–Asn linkages are found, dependent on the type of proteoglycan [169]. Functions of proteoglycans in which the carbohydrate chains are essential are *e.g.* interaction with complement components and inhibition of complement activation [145], regulatory functions (anticoagulant) in hemostasis [8], and stability of extracellular matrix [111].

Table 2. Core structures found in mucin–type serine– or threonine–linked oligosaccharides and polysaccharides.^a

	core 1		core 5
	core 2		core 6
	core 3		core 7
	core 4		

^a structures described *e.g.* in [64,172]

Glycoconjugates of parasites

The function and role of glycosylated structures in the interaction of pathogenic organisms like unicellular and multicellular parasites is becoming increasingly apparent. Glycoconjugates are involved in parasite membrane protection [44,150,191], specific interaction with some host cell–types [18,63,118,136,151,191], physical stability and protection of cercarial tegument [17,38,123,153], or masking of antigens the recognition whereof by the host immune system may result in damage to the parasite [38,160]. Since

the carbohydrate chains of the glycoconjugates appear to be directly involved in these interactions, the elucidation of the glycan structure may yield new insights into parasite glycoconjugate structure/effect relationships and into the biosynthetic pathways responsible for their formation. Several studies have indicated that parasite glycans may be highly immunogenic and thus can be exploited in vaccine development, as it has been shown that antibodies against these structures can mediate protective immunity [56,78,90,91,134,173].

The structural analysis of parasite glycoconjugates has often been hampered by the small amounts of purified material available. For characterization, peptides and proteins which are scarce can be obtained in theoretically unlimited amounts by recombinant DNA-technology, if the amino acid sequence or DNA code is known. However, only careful and laborious purification can yield sufficient material needed for carbohydrate structural analysis. For this reason many research groups have used (partially) purified antigen preparations and/or immunochemical or histochemical methods for studying these structures. These methods comprise specific antibody and lectin binding, to the structures themselves and/or in combination with various chemical and enzymatic carbohydrate degradation methods. A short description of the various techniques used follows below.

Structural analysis of glycoconjugate glycans

Biochemical techniques. To the biochemist, a variety of analytical techniques is available for the identification of the primary structure of carbohydrates isolated from glycoconjugates (reviewed in *e.g.* [74,98,99]). Among the techniques often used are monosaccharide analysis, enzymatic degradation methods, mass spectrometry (MS) and nuclear magnetic resonance (NMR) spectroscopy. A major advantage of MS is the very small amount of sample needed (as low as femtomoles [76]) while NMR spectroscopy needs more material (usually in the nanomole range), but is in contrast to MS non-destructive, allowing multiple analyses to be performed. Each technique yields specific information, and combination of the methods is mostly essential for a complete elucidation of a molecular structure. As an example, NMR spectroscopy can identify the type and sequence of sugar residues (Gal, Man, etc.) as well as the type of linkages, while MS can give size, composition (hexose, pentose, etc.) and linkage information. Enzymatic degradation methods, *e.g.* sequential exoglycosidase digestion [92,188], make use of the specificity of naturally occurring enzymes (purified or recombinant), sometimes in combination with sequential lectin-affinity chromatography [21,116]. These methods may also yield information on the identity of the monosaccharide residue as well as on the sequence and type of linkages. The introduction of radioactive (^3H , ^{14}C) or fluorescent labels makes detection highly sensitive, and are therefore widely applied in the analysis of minute amounts of parasite carbohydrates.

Lectin-based techniques. Lectins are sugar-binding proteins or glycoproteins of non-immune origin without enzymatic activity to the carbohydrate bound [94]. Various lectins show agglutination properties towards cells *in vitro* or precipitate glycoconjugates [36,50]. Many highly purified lectins of well-defined carbohydrate specificity (those applied in the analysis of schistosomal antigens are summarized in Table 3) are easily available, which has led to the widespread adoption of lectin techniques [4]. These techniques employ conjugates of lectins with a variety of fluorescent labels or enzymes, biotin-conjugates as well as immobilized lectins for affinity chromatography [21,48,51,104,116]. Because they do not usually enter cells, lectins can be used as probes to provide information about the location, abundance and function of glycoconjugates at cell surfaces [37,85,102,108,123,159,161,176,177]. The specific and reversible interactions between lectins and mono- or oligosaccharides form the basis for important affinity chromatographical methods used in isolation and fractionation of glycoconjugates and glycopeptides [21,105,116,131,167].

Table 3. Various lectins used for the analysis of carbohydrates in schistosomal antigens.^a

Lectin	Carbohydrate specificity
Con A <i>Canavalia ensiformis</i> (jack bean) (Concanavalin A)	α -D-Man > α -D-Glc > α -D-GlcNAc
DBA <i>Dolichos biflorus</i> (horse gram)	α -D-GalNAc
GS-I <i>Griffonia simplicifolia</i>	α -D-Gal, α -D-GalNAc
GS-II <i>Griffonia simplicifolia</i>	α/β -D-GlcNAc
HPA <i>Helix pomatia</i> (edible snail)	α -D-GalNAc
LBA <i>Phaseolus lunatus</i> (lima bean)	α -D-GalNAc > Gal
LcH <i>Lens culinaris</i> (lentil)	α -D-Man, α -D-Glc, α -D-GlcNAc
Lotus <i>Lotus tetragonolobus purpureas</i>	α -L-Fuc
MAA <i>Maackia amurensis</i>	Neu5Ac- α (2-3)-Gal
PNA <i>Arachis hypogaea</i> (peanut)	D-Gal- β (1-3)-D-GalNAc > β -D-Gal
PWA <i>Phytolacca americana</i> (pokeweed)	D-GlcNAc- β (1-4)-D-GlcNAc- β - (1-4)-D-GlcNAc
RCA-I <i>Ricinus communis</i> I – 120 kDa (castor bean)	β -D-Gal
RCA-II <i>Ricinus communis</i> II – 60 kDa (castor bean)	β -D-Gal, α/β -D-GalNAc
SBA <i>Glycine max</i> (soybean)	α/β -D-GalNAc
SJA <i>Sophora japonica</i>	α -D-GalNAc
UEA-I <i>Ulex europaeus</i>	α -L-Fuc(1→x) ^b
WFA <i>Wisteria floribunda</i> (Japanese wisteria)	α/β -D-GalNAc
WGA <i>Triticum vulgaris</i> (wheat germ)	β -D-GlcNAc- β (1→4)-D-GlcNAc

^a data based on e.g. [5,51,104,177].

^b x may be any aglycon

Antibody-based techniques. There are close parallels between lectins and carbohydrate-specific monoclonal antibodies (McAbs), resulting in a similar spectrum of applications and methods. A major difference is that, in general, antibody binding to carbohydrate epitopes involves more than one sugar residue, and is not easily inhibited by a single monosaccharide or even by a disaccharide [13,179]. The maximum size of an antigen-binding site of an antibody is about 6 sugar or amino acid residues [19,24,127,179,186]. For the determination of overall carbohydrate recognition patterns in humoral immune responses polyclonal antibodies present in immune sera are often used. Another frequent application of antibodies is the immunohistochemical and immunocytochemical localization of antigens [25,26,71,72,96,135,163].

Partial destruction and inhibition. The above described techniques may also be applied to glycoprotein preparations which are partially degraded with respect to their carbohydrate or peptide portion. Methods generally used for degradation include chemical methods like sodiumperiodate treatment [5,91,126,187], or mild acid hydrolysis (for desialylation [137,175] and defucosylation [91,137,143,175,183]), as well as enzymatic methods [52,91,98]. The inhibitory effect of these treatments on the binding of antibodies or lectins yields information about the type of mono- or oligosaccharide involved. Inhibition can also be achieved by mono- or oligosaccharides themselves [19,90,100,105,157], by other antigenic preparations [57,124,144,179], or by other lectins, McAbs or polyclonal antibodies [58,100,144].

2.2. Glycoconjugates of *Schistosoma*

Parasites of the genus *Schistosoma* are complex organisms shown to contain and synthesize numerous glycoproteins and glycolipids, which include some completely unique carbohydrate structures. This chapter will give an overview, dealing in particular with glycoprotein carbohydrates which might interact with the immune system of the definitive host. As the methods employed in glycan analysis are summarized above, these will only be shortly referred to in the paragraphs below.

Schistosomula and adult worms

Purified lectins with defined specificity have been employed in a number of studies to determine which glycans in worm tissues and secretions were accessible for the host immune system. A summary of the lectins recognizing

carbohydrate structures on the tegument or in the gut of the schistosome is given in Table 4. The host is exposed to parasite gut-associated antigens when these enter the circulation after regular regurgitation of the undigested contents of the gut.

Table 4. Lectin binding to *Schistosoma* adult worms and schistosomula.^{a,b,c}

Tegument	Gut	Reference
Con A, SBA, DBA, PNA, RCA-I, UEA-I	Con A, RCA-I, SBA, DBA, UEA-I, PNA	Beisler <i>et al.</i> (1984) [5] ^d
Con A, RCA, SBA ±, PNA, UEA-I, Lotus	Con A, RCA, SBA, PNA, UEA-I, Lotus	Linder and Huldts (1982) [103]
Con A, LcH, SBA, RCA-I, RCA-II	n.d. ^e	Hayunga and Sumner (1986) [69]
PNA, DBA, Con A, SBA	n.d.	MacGregor <i>et al.</i> (1985) [108]
Con A, RCA-II, PNA, SBA, Lotus	n.d.	Simpson and Smithers (1980) [161]
Con A, UEA-I, SBA	n.d.	Murrell <i>et al.</i> (1978) [122]

^a as various reports indicated that the binding of WGA is largely aspecific [5,105,122,159,161], this lectin has been omitted in this Table

^b unless otherwise indicated, studies were performed in *Schistosoma mansoni*

^c shaded lectins were also investigated but showed no binding

^d studied in *Schistosoma japonicum*

^e n.d. = not done

In some studies, the reduction in binding of several lectins to the schistosomulum surface in the course of development of the parasite indicated a decline in the number of exposed carbohydrate epitopes [138,139,159]. On the other hand, Con A-agarose chromatography showed the presence of 19 and 45 kDa surface glycoproteins in two-day old but not in younger schistosomula [138], while PNA-binding demonstrated a 170 kDa glycoprotein abundantly present on 4-week-old worms, but absent in pre-liver stages of the parasite [107]. The loss of lectin binding sites is increased after a short stay in the circulation of the host and appears to be due to antigen shedding rather than masking of carbohydrate epitopes [139].

As a more general phenomenon, antigen shedding has also been suggested by a decrease in surface-bound anti-carbohydrate antibodies with the development

of the parasite, resulting in an absence of antibodies recognizing lung schistosomula [97]. Shedding of surface antigens is one of the evasion mechanisms employed by the parasite to escape the host immune attack occurring through antibodies [10,97], or via the alternative complement activation route [112,113,154]. On the other hand, the parasite may evade the host immune response by masking of surface antigens through acquisition of host molecules on the tegument, as described by Clegg, Smithers and Terry in 1971 [20], as well as by others [49,156]. This phenomenon has been clearly linked with a decrease in binding to the schistosomula of host antibodies or complement components [73,146], while other studies also indicate some intrinsic changes in the tegument which render the parasite insusceptible to immune attack [14,31,120].

The presence of Con A-binding epitopes on tegument antigens as well as on gut antigens has been consistently shown (Table 4). The highest binding affinity of Con A is to the Man residue (Table 3), for which reason it is generally used to indicate the presence of N-linked carbohydrate structures (see Table 1) [21,116]. Binding of SBA and PNA to the parasite surface indicates the presence of GalNAc- and Gal-GalNAc-containing oligosaccharides, respectively (Table 3). These findings have been confirmed more recently by studies of Cummings and co-workers [128-131,167,168], using glycoconjugates synthesized by the worms after incubation *in vitro* with radiolabeled monosaccharides and combinations of the above described direct and indirect analytical methods (*e.g.* lectin-affinity chromatography, exo- and endo-glycosidase treatment). They were able to characterize a number of unique carbohydrate structures which are summarized in Table 5.

The presence of Le^x on the surface of schistosomula as detected by McAbs prepared from mice immunized with *S. mansoni* eggs [66,91], was described by Ko *et al.* (1990) [90]. Using sera of *S. mansoni* infected patients these investigators also found a positive reaction on sections of embryonic mouse head with a distribution similar to that of the SSEA-1 epitope (is identical to Le^x), from which they concluded that anti-Le^x antibodies are generated during human schistosomiasis infection. Recently, Köster and Strand (1994) described another McAb recognizing the Le^x epitope and showed that the epitope was expressed at the surface of schistosomula (immediately after transformation) and adult worms, as well as at the gut epithelium of adult worms [96].

In contrast with these reports on the presence of the Le^x trisaccharide in *Schistosoma* [90,128], several studies failed to show binding of the Fuc-specific lectins UEA-I and/or Lotus to schistosomula and adult worms (Table 4) [103,138,139,159,161], although both lectins were shown to recognize Le^x [163,167]. A possible explanation is that, due to conformational changes, steric hindrance, or epitope masking by other (glyco)proteins, the

Table 5. Schistosome carbohydrate structures described by Cummings and co-workers.

Carbohydrate structure	Reference
<i>one-day cultured male worms</i>	
O-linked terminal GalNAc	[128]
O-linked terminal GlcNAc	
O-linked Gal-GalNAc	
N-linked oligomannose-type oligosaccharides, ranging in size from Man ₇ GlcNAc ₂ to Man ₉ GlcNAc ₂	[130]
N-linked tri- and diantennary N-acetyllactosamine-type oligosaccharides, containing Man, Fuc, GlcNAc, and GalNAc ^a	[131]
N-linked poly-Le ^x , (→3)-β-Gal-(1→4)-[α-Fuc-(1→3)]-β-GlcNAc-(1→) _n	[167]
N-linked biantennary, complex-type oligosaccharides, containing on both antennae the terminal sequence β-GalNAc-(1→4)-[±α-Fuc-(1→3)]-β-GlcNAc-(1→2)-α-Man-(1→	[168]
glycosphingolipid with novel core β-GalNAc-(1→4)-Glc-ceramide	[110]
<i>two-day cultured mechanically transformed schistosomula</i>	
O-linked GalNAc	[129]
O-linked GlcNAc	
N-linked oligomannose-type oligosaccharides, ranging in size from Man ₆ GlcNAc ₂ to Man ₉ GlcNAc ₂	
N-linked complex-type oligosaccharides, containing Man, Fuc, GlcNAc, and GalNAc ^b	

^a Neu5Ac could not be detected; GalNAc was β-linked in a terminal position

^b Neu5Ac could not be detected

Fuc-containing epitopes are accessible to the lectins only in (partly) purified antigens but not if the epitopes are still present in worm tissues¹.

An interesting excursion on the interaction of fucosylated (Le^x and Le^y) oligosaccharides has recently been described by Harn and co-workers [182]. These authors show that these carbohydrates induce proliferation of human peripheral blood mononuclear cells (which include B cells) [65], as well as interleukin 10 production by isolated B cells (B220⁺) of *S. mansoni*-infected

¹ Recently, in our laboratory using an IFA on sections of adult male *Schistosoma* worms fixed in Rossman's fixative, we found a strong reactivity of Lotus and a weak reactivity of UEA-I in the gut of the parasites (unpublished results). In Chapter 7 of this thesis it is demonstrated that circulating antigen CCA contains Le^x repeating units and that Lotus but not UEA-I binds to immunopurified CCA in ELISA [178]. In contrast, Carlier *et al.* (1980) have shown only a very weak binding of Lotus to partly purified M antigen (CCA), which might be due to a TCA-precipitation step in the preparation procedure causing a partial degradation of the Fuc-containing epitope [15].

and not of non-infected mice [182]. Moreover, they were able to show the presence of antibodies against these fucosylated oligosaccharides in the cerebrospinal fluid of schistosomiasis patients with cerebral disorders [47]. Based on these very new phenomena, the authors suggest a role of Le^X- and/or Le^Y-containing oligosaccharides in the immunoregulation of the helper T cell response in schistosomiasis and maybe other chronic infectious diseases [182].

The presence of sialic acid (Neu5Ac) in schistosome glycoconjugates is a controversial issue. A number of researchers could not detect its presence in *in vitro* synthesized glycoproteins [128-131,168], while others found a minor amount in adult schistosome carcasses which they attributed to the presence of ingested erythrocytes [149]. In contrast, using immunochemical methods of lectin (PNA) or dye binding after neuraminidase treatment of the parasites, it was shown that some glycoconjugates in lung-stage schistosomula and adult worms contain sialic acid, as opposed to newly transformed schistosomula [115,159,161]. In the cercarial glycocalyx (surface coat), although negatively charged, Neu5Ac could not be detected [123].

The above described lectin-based studies demonstrate a large variety of carbohydrate structures on the surface of adult worms and schistosomula. Indeed, antigenic carbohydrate structures dominate at the surface of the schistosomulum indicated by absorption of more than 90 % of the schistosomulum-specific antibodies in infected mouse serum by an egg antigen preparation from which the protein antigens had been removed [134]. The cross-reaction between the carbohydrate structures of schistosome egg antigens and schistosomula surface antigens has formed the basis of a theory on the slow development of immunity observed in heavily-exposed children [9,10,12]. This experimentally well-supported hypothesis (recently reviewed in *e.g.* [11]) involves the development in young children of IgM and IgG2 antibodies which block the protective immune mechanisms of antibody-dependent cellular cytotoxicity (ADCC) by eosinophils and other cells. Polysaccharide antigens released by the large quantity of eggs deposited in the tissues elicit a predominantly thymus-independent antibody response resulting in especially IgM and IgG2 antibodies [3]. These so-called "blocking" antibodies may cross-react with carbohydrate epitopes on schistosomula surface glycoproteins, and not only fail to mediate ADCC reactions but also block "effector" antibodies like IgG1 and IgE. As the infection progresses and the child reaches adulthood the balance in the antibody response may switch to a predominantly protective type of response which results in the development of (partial) immunity.

Experimental and epidemiological evidence for this theory came from the groups of Capron, Butterworth, and recently of the group of Dessein. Capron and co-workers described a rat IgG2c McAb (isotype comparable with IgG3 in mice

[30,119,142] or IgG2 in man [3,79,89,158]) which inhibited a rat IgG2a McAb-mediated, eosinophil-dependent killing of schistosomula [54,55]. Mouse IgM McAbs recognizing cross-reactive egg polysaccharides and schistosomulum surface antigens as well as human IgM against the schistosomulum surface could effectively block the eosinophil-dependent killing of schistosomula by sera from infected humans [40,86]. Also in murine schistosomiasis blocking antibodies could be characterized [134,189,190] and it was shown that egg antigens elicit thymus-independent responses of the IgM and IgG3 isotypes [114]. The inhibitory effect on protection against schistosome infections of mice IgM antibodies which bind to egg and schistosomulum surface antigens might be limited to anti-carbohydrate antibodies as a number of studies indicate the presence of protective IgM responses or McAbs (in passive transfer experiments) which recognize egg and/or schistosomulum antigens [53,80,88,147,164].

After purification of individual IgG subclasses from schistosomiasis mansoni patient sera it was shown that IgG1 and IgG3 antibodies mediated eosinophil killing of schistosomula, that IgG4 consistently blocked killing, and that IgG2 would either mediate or block killing, depending on the state of activation of the eosinophils [87]. Epidemiological support for the blocking antibodies theory came mainly from the Kenyan reinfection studies of Butterworth *et al.* (1987,1988), who found that blocking antibodies of the IgG2 and IgM isotypes in the sera of young children prevented the expression of immunity [9,10]. Resistance to reinfection after treatment is associated with the presence of IgG isotypes other than IgG2 which recognize similar schistosomulum antigens [10], although Omer-Ali *et al.* (1989) obtained indications that, in general, antibodies against carbohydrate epitopes on the surface of schistosomula of *S. haematobium* do not have a major protective role in man [132]. Finally, Demeure *et al.* (1993) recently described that, in a population cross-section comprising all age-groups, anti-schistosomal IgG2 levels were predictive for susceptibility to reinfection [29].

The carbohydrate epitope originally defined by the protective rat IgG2a McAb produced by Capron and co-workers [33,54] (already mentioned above), is present on a 38 kDa schistosomula surface molecule and expressed in various molecular mass components in the different parasite stages of *S. mansoni* [32,33,35]. Interestingly, this epitope has been shown to be additionally synthesized by the freshwater snail *Biomphalaria glabrata* and may be implied in molecular mimicry between schistosomes and their invertebrate intermediate hosts [35]. Moreover, the epitope is present in products of other snails, including *Bulinus truncatus*, intermediate host for *S. haematobium* [35], as well as in keyhole limpet hemocyanin (KLH), synthesized by a marine mollusc *Megathura crenulata* [57]. The major immunogenicity of the 38 kDa carbohydrate moieties in mammalian hosts was confirmed in mouse and human

schistosomiasis [86,133]. The molecular structure of this carbohydrate epitope has not been elucidated (Dr. J.P. Kamerling, personal communication).

The above described role of schistosomulum surface carbohydrates in the development of a humoral immune response should be extended to the role of the excretory and secretory schistosome antigens. Indeed, in schistosome infections, high antibody titers are observed against parasite gut-associated antigens, which are regularly released into the circulation of the host [27,28,42,125]. Radiolabeling experiments showed that essentially all components released by the worm were recognized by antibodies in infected human sera, in contrast with most of the major (non-secreted) membrane and tegumental proteins [101]. It was therefore suggested that the adult worm is protected against immune attack by direction of the host's antibody response against released (glyco)proteins rather than against parasite surface antigens [101]. In passive transfer experiments, it was demonstrated that after administration of a mouse IgM McAb directed against the parasite gut epithelium, larger numbers of lung-stage schistosomula could be isolated than when an irrelevant IgM McAb or physiological salt was given [1]. The antigen recognized by this McAb was also found in high concentrations in *S. mansoni* eggs, was excreted in the urine of schistosomiasis patients and experimental animals [2,148] and appeared to have a polysaccharide nature, of which the antigenicity was readily destroyed by periodate, but not by protein-denaturing treatments [1]. In addition, a number of highly glycosylated gut-associated glycoproteins was described (among these CAA and CCA), which are summarized in Chapter 1 of this thesis. Although the immune response against those antigens is high, there appears to be so far no specific interaction with immune mechanisms, besides from those described in this thesis (Chapter 9,10).

Egg

The localization of glycans in schistosome eggs has been studied using lectins in similar procedures as for the carbohydrate structures found in schistosomula and adult worms (see above). A summary of the results of two studies is given in Table 6.

The cross-reactivity of carbohydrate epitopes on egg antigens with those on schistosomula surface antigens in relation to the occurrence of blocking antibodies has been discussed above. These epitopes were present on a 38 kDa schistosomulum surface molecule and also found on molecules of different M_r present in other schistosome life stages (miracidia, adult worm) or even in the intermediate snail host. A number of mouse IgM and IgG3 McAbs, which also reacted with antigens of about 38 kDa at the surface of *S. mansoni*

Table 6. Lectin binding to *Schistosoma* eggs.^{a,b}

Miracidium	Vitelline membrane ^c	Vitelline space ^d	Reference
Con A, SBA ±, PNA, UEA-I ±	Con A, RCA-I	Con A, SBA, DBA, PNA, UEA-I	Beisler <i>et al.</i> (1984) [5] ^e
RCA, SBA, PNA, UEA-I, Lotus ^f	^f	^f	Linder and Hyltd, (1982) [103] ^{f,g}

^a as various reports indicated that the binding of WGA is largely aspecific [5,105,122,159,161], this lectin has been omitted in this Table

^b shaded lectins were also investigated but showed no binding

^c inside of egg shell

^d containing hatching fluid (with miracidia excreta) which leaks through pores in the eggshell

^e studied in *Schistosoma japonicum*

^f no differentiation within egg was made

^g studied in *Schistosoma mansoni*

schistosomula recognized a well-characterized polysaccharide egg antigen designated κ_3 [6,39,40]. Cross-reacting antigens were additionally found in a crude cercarial antigen preparation but not in adult worm homogenate. The IgM but not the IgG3 McAbs inhibited the killing of schistosomula by eosinophils in the presence of sera from infected humans [7,40]. However, passive transfer of the IgM McAbs to vaccinated mice at the time of challenge failed to produce *in vivo* blocking [6]. The polysaccharide κ_3 showed a molecular mass in the range of >750 – 70 kDa and was resistant to boiling and protease degradation, but sensitive to 50 mM periodate. It partially bound to Con A and remained unaffected by 0.1 M NaOH or 0.1 M HCl treatment. Cross-reactivity to other schistosome species was shown by rabbit anti- κ_3 antibodies which bound to the surface of *S. japonicum*, *S. haematobium*, and *S. bovis* schistosomula while also an egg antigen of *S. japonicum* was recognized [39].

Besides the blocking of cellular-dependent immune mechanisms, another effect of antibodies directed against egg carbohydrates was described by Simpson *et al.* (1990) who showed that the administration of a mouse anti-egg carbohydrate IgM McAb to mice intravenously injected with 2000 *S. mansoni* eggs resulted in larger lung granulomas than in mice which received no or an irrelevant McAb [162].

S. mansoni-infected T cell-deprived mice suffer from severe hepatocyte damage most likely caused by an egg-derived cationic glycoprotein antigen named ω_1 . Together with another cationic glycoprotein, α_1 , these antigens were recently purified from *S. mansoni* eggs and characterized with respect to their biochemical, immunological as well as hepatotoxic properties [41]. ω_1 is a monomeric glycoprotein with a pI > 9.0 and a molecular mass of 31 kDa. The antigen α_1 consists of two immunologically cross-reactive dimers with a pI

ranging from 7.5 to 8.5 and molecular masses of 41 kDa and 36 kDa, each of which is composed of one unique and one common subcomponent. Since both antigens bound to Con A they might contain N-linked carbohydrate chains (Tables 3 and 1). The immunodominance of the carbohydrate epitopes was shown by the observation that antigenic activity against monospecific antisera was readily lost after oxidation with 50 mM periodate, but not affected by protease treatment. Mild acid hydrolysis destroyed the carbohydrate epitopes of w_1 but not of α_1 , while α_1 was more sensitive to NaOH treatment, suggesting the presence of O-linked glycans. Antibody-binding was abrogated after the antigens were heated at 100°C for 5 min.

Using an ELISA with sera from infected humans and mice, it has been shown that purified w_1 is *S. mansoni*-specific and is a better marker than α_1 or unfractionated egg antigens to distinguish *S. mansoni* infection from other schistosome infections [41]. Passive transfer of monospecific anti- w_1 sera into *S. mansoni*-infected, T cell-deprived mice completely prevented the occurrence of microvesicular hepatocyte damage in these animals, while monospecific anti- α_1 serum had no hepatoprotective activity [41].

Another common carbohydrate epitope identified by a mouse McAb (named 128C3/3, produced after immunization with cercarial antigens [171]) was found to be present on *S. mansoni*, *S. haematobium*, *S. japonicum* egg glycoproteins and glycolipids as well as on *S. mansoni* cercarial and adult worm antigens [96,184,185]. In addition, sera of both infected humans and mice contain large amounts of antibodies with a similar epitope specificity [184]. In an immunopurified glycolipid fraction (using the same McAb) the major immunogens were shown to be unique glycosphingolipids with repeating trisaccharide units containing internal Fuc residues substituted onto a novel glucosylceramide core structure [100]. The core structure was independently described by Makaanu *et al.* (1992) [110]. The largest antigen identified in the glycolipid preparation was interpreted to have the following structure¹:




Figure 1. Chemical structure of a major carbohydrate antigen recognized by McAb 128C3/3.

¹ The anomeric configuration of the constituent monosaccharides is undetermined due to the analytical technique used.

Other smaller components have also been recognized by McAb 128C3/3 in which one or more Fuc(1→3) side chains were absent. The immunodominance of L-Fuc in the binding of the McAb to egg glycoproteins was demonstrated by inhibition with monomeric L-Fuc, while the McAb also bound to immobilized L-Fuc [100].

The same McAb 128C3/3 bound to a common epitope present on various polypeptides of the cercarial glycocalyx as well as in the tegument of cercariae and schistosomula [23]. The glycocalyx components were shed after transformation of cercariae to schistosomula, but two components of 170 and 180 kDa continued to be present in cercariae, in newly transformed schistosomula and in 36 h *in vitro* cultured schistosomula. The lectins Con A (to Man, Glc) and Lotus (to Fuc) also bound to these antigens as well as to a number of other high and low molecular mass components [23]. Inhibition studies later showed that in the binding of McAb 128C3/3 Fuc was dominant [100]. Remarkably, 128C3/3 and Con A but not Lotus were reactive with a 38 kDa glycoprotein from schistosomula which is possibly the same as the antigen recognized by the partially protective McAbs described by different groups, *e.g.* Dissous *et al.* (1982,1985) [33,34], Phillips *et al.* (1982,1986) [83,192], Bickle *et al.* (1987,1988) [6,40], or Harn *et al.* (1984,1987) [66,67]. Nevertheless, the epitopes recognized by these McAbs need not be the same, even though each of the McAbs bound to a carbohydrate moiety. Finally, the antigen recognized by McAb 128C3/3 appears to be present in the serum of *Schistosoma*-infected patients and an immunodiagnostic antigen-capture ELISA using this McAb has been described [68].

A number of egg glycoproteins (major serological antigens, MSA_{1,2,3}) showing immunodiagnostic potential were isolated from *S. mansoni* eggs by successive steps of Con A-affinity and ion-exchange chromatography [62,140,141]. MSA₃ bound weakly to Con A and appeared to be poorly glycosylated. MSA₁ and MSA₂ bound to Con A with higher affinity and PAS-staining, specific for carbohydrates, was also more intense. MSA₁ may be a dimer composed of two glycoproteins of 50 kDa each, while MSA₂ was found to be a heterogeneous lipoglycoprotein with a molecular mass of M_r 450 kDa. The M_r of MSA₃ was 80 kDa [140]. The major source of the antigens appeared to be the hatching fluid, containing miracidial excretory or surface coat products [62,140]. MSA₁ appeared to be both stage- and species-specific, while MSA₂ and MSA₃ showed partial cross-reactivity with cercarial but not with adult worm antigen, as well as with egg antigens from *S. haematobium* and *S. japonicum* [140]. Some evidence for strain variation of the antigen was provided by Hamburger *et al.* (1982) [60], who purified a glycoprotein from an Egyptian strain of *S. mansoni* (named major egg glycoprotein, MEG) using the same procedure as for MSA₁ from a Puerto Rican strain [141]. The glycoprotein from the Egyptian strain differed in molecular mass (70 kDa) and its reactivity with antibodies in

antigen-competitive radioimmunoassay was similar but not identical to that of MSA₁. In particular, MEG showed a high degree of cross-reaction with egg antigens from *S. haematobium* [60]. The major immunogenic epitopes of MEG appeared to be carbohydrates because various biochemical procedures as heating to 100°C for 1 h, treatment with 10 % TCA, 0.1 M NaOH, or 0.1 M HCl had no effect on the serologic reactivity of MEG, but treatment with 65 mM periodate resulted in a drastic loss of molecular mass and of serologic reactivity. Pronase treatment of MEG caused a limited fragmentation of the molecule and only a minor decrease in serum antibody recognition [61].

Minor carbohydrate-rich fragments of soluble egg antigens (<13 kDa and >10 kDa, isolated by ultrafiltration followed by dialysis) contained the same carbohydrate epitopes as MEG, but showed no immunopathological activity [106]. Having shown that a number of lectins (Con A, PNA, RCA II, Fuc) bound to MEG as well as to the smaller fragments, it was suggested that MEG consists of a polypeptide backbone to which several polysaccharide chains are bound [105,106].

Cercaria

A number of cercarial antigens showing cross-reacting carbohydrate epitopes with antigens on other life-stages have already been mentioned and described above. The results of localization studies employing lectins are summarized in Table 7, similar to the egg and worm carbohydrates.

As shown in Table 7, there appeared to be a considerable heterogeneity between the various studies in the types of lectins bound to cercarial structures. A number of these differences might be explained by species-specific expression of carbohydrate structures, *e.g.* between Beisler *et al.* (1984) [5], and Linder (1985) [102], but the discordant binding of Lotus to the cercarial heads is difficult to interpret. The components present in the excretion products of cercariae (as described by [102]) might partly be produced by the penetration glands causing overlap of the lectin-binding patterns, suggesting that enzymes used for skin penetration are glycosylated.

Using Bolton-Hunter radiolabeling of cercarial tegument structures, Hayunga *et al.* (1986) [70] identified a number of Con A-binding high and low molecular mass glycoproteins (180, 150, 43 and 30 kDa). These components were retained by 24 h cultured schistosomula, and supplemented by two other glycoproteins (66 and 57 kDa) [70].

Table 7. Lectin binding to *Schistosoma* cercariae and newly transformed schistosomula (≤ 3 h.).^{a,b,c}

Head	Tail	Penetration glands	Excreta	Reference
Con A, SBA, DBA, UEA-I \pm , PNA, RCA-I	Con A, SBA \pm , DBA, RCA-I	SBA, UEA-I	n.d. ^d	Beisler <i>et al.</i> (1984) [5] ^e
Con A, GS-I, RCA-I, SBA, HPA, WFA, UEA-I, Lotus, PNA, DBA	Con A, GS-I, RCA-I, SBA, HPA, WFA, UEA-I, Lotus, PNA, DBA	n.d.	Con A, GS-I, RCA-I, SBA, HPA, WFA, UEA-I \pm , Lotus, PNA, DBA	Linder (1985) [102]
Con A, SBA, UEA-I ^{f,g}	n.d.	n.d.	n.d.	Murrell <i>et al.</i> (1978) [122] ^g
Lotus	Lotus \pm	n.d.	n.d.	Nanduri <i>et al.</i> (1991) [123]
Con A, PNA, Lotus	n.d.	n.d.	n.d.	Payares and Simpson (1985) [138] ^h

^a as various reports indicated that the binding of WGA is largely aspecific [5,105,122,159,161], this lectin has been omitted in this Table

^b unless otherwise indicated, studies were performed in *Schistosoma mansoni*

^c shaded lectins were also investigated but showed no binding

^d n.d. = not done

^e studied in *Schistosoma japonicum*

^f 2 h schistosomula were weakly positive for SBA and UEA-I

^g no differentiation within cercariae was made

^h surface proteins of newly transformed schistosomula

Nanduri *et al.* (1991) [123] used lectin-affinity chromatography on immobilized Lotus lectin to isolate Fuc-containing carbohydrate structures from cercarial body and tail glycocalyx. Monosaccharide composition analysis of body glycocalyx showed that Fuc is the major sugar residue present, while in the tail Glc is predominant. Alkaline treatment indicated that the glycans are attached to the protein backbone via an O-glycosidic bond, both for bodies and tails. After reductive alkaline treatment, two glycan chains of 10.5 kDa and 5.6 kDa appeared to account for most of the glycocalyx components of both bodies and tails. The larger glycan was retarded on an anion-exchange column, thus being negatively charged, but Neu5Ac could not be demonstrated. Both body and tail glycocalyx were highly sensitive to mild acid hydrolysis, which for the body supports the presence of high amounts of Fuc [123]. These results confirm a

previous study by Caulfield *et al.* (1988) [16], who found a predominance of Fuc residues in the carbohydrate fraction and a relatively high amount of Thr and Ser in the amino acid fraction of the cercarial glycocalyx. Additionally, using a McAb which can be inhibited by monomeric L-Fuc (128C3/3, already been discussed in detail above) several smaller surface glycoproteins of cercariae have been identified [23]. The same McAb recognizes carbohydrate epitopes present on a variety of *S. mansoni*, *S. haematobium*, *S. japonicum* egg glycoproteins and glycolipids as well as *S. mansoni* cercarial and adult worm antigens [184,185]. These observations can be used as a clear illustration of a common feature in carbohydrate analysis, namely that the same glycan structure may be expressed throughout the schistosome life-cycle, and exhibit different biological roles, but that yet the overall glycolipid and glycoprotein structures are not conserved.

References

- Appriou M, Ben Younes R, Tribouley-Duret J, Tribouley J. Étude par la réaction d'immunofluorescence des anticorps dirigés contre les antigènes de l'épithélium intestinal de *Schistosoma mansoni*. IV. Étude du pouvoir bloquant sur l'immunité d'un anticorps monoclonal de classe IgM. *Annales de Parasitologie Humaine et Comparée* 1989; 64:456-468.
- Appriou M, Tribouley-Duret J, Tribouley J. Étude par la réaction d'immunofluorescence des anticorps dirigés contre les antigènes de l'épithélium intestinal de *Schistosoma mansoni*. III. Étude de la réactivité d'un anticorps monoclonal. *Annales de Parasitologie Humaine et Comparée* 1986; 61:435-446.
- Barrett DJ, Ayoub EM. IgG2 subclass restriction of antibody to pneumococcal polysaccharides. *Clinical and Experimental Immunology* 1986; 63:127-134.
- Beeley JG. *Laboratory techniques in biochemistry and molecular biology*, Vol. 16. Elsevier, Amsterdam, 1985:
- Beisler GK, Nakao M, Matsuda H, Tanaka H. Characterization of the carbohydrates of *Schistosoma japonicum* adult worm, egg and cercaria by analysis of lectin binding and antibody reaction. *Japanese Journal of Experimental Medicine* 1984; 54:263-273.
- Bickle QD, Andrews BJ. Characterization of *Schistosoma mansoni* monoclonal antibodies which block *in vitro* killing: failure to demonstrate blockage of immunity *in vivo*. *Parasite Immunology* 1988; 10:151-168.
- Bickle QD, Andrews BJ, Taylor MG. *Schistosoma mansoni*: characterization of two protective monoclonal antibodies. *Parasite Immunology* 1986; 8:95-107.
- Bourin MC. La Thrombomoduline: un nouveau protéoglycane. Relation structure-fonction. *Annales de Biologie Clinique* 1991; 49:199-207.
- Butterworth AE, Bensted-Smith R, Capron A, Capron M, Dalton PR, Dunne DW, Grzych J-M, Kariuki HC, Khalife J, Koech DK, Mugambi M, Ouma JH, Arap Siongok TK, Sturrock RF. Immunity in human schistosomiasis mansoni: prevention by blocking antibodies of the expression of immunity in young children. *Parasitology* 1987; 94:281-300.
- Butterworth AE, Dunne DW, Fulford AJC, Capron M, Khalife J, Capron A, Koech DK, Ouma JH, Sturrock RF. Immunity in human schistosomiasis mansoni: cross-reactive IgM and IgG2 anti-carbohydrate antibodies block the expression of immunity. *Biochimie* 1988; 70:1053-1063.
- Butterworth AE, Dunne DW, Fulford AJC, Thorne KJI, Gachuhi K, Ouma JH, Sturrock RF. Human immunity to *Schistosoma mansoni* - observations on mechanisms, and implications for control. *Immunological Investigations* 1992; 21:391-407.
- Butterworth AE, Hagan P. Immunity in human schistosomiasis. *Parasitology Today* 1987; 3:11-16.
- Campbell JH, Pappenheimer AM. Quantitative studies of the specificity of anti-pneumococcal polysaccharide antibodies,

- types III and VII. 2. Inhibition of precipitin reactions with oligosaccharides isolated from hydrolysates of S3 and S8. *Immunochemistry* **1966**; **3**:213–222.
14. Capron A, Dessaint JP, Capron M, Ouma JH, Butterworth AE. Immunity to schistosomes: progress toward vaccine. *Science* **1987**; **238**:1065–1072.
 15. Carlier Y, Bout D, Strecker G, Debray H, Capron A. Purification, immunochemical, and biological characterization of the *Schistosoma* circulating M antigen. *Journal of Immunology* **1980**; **124**:2442–2450.
 16. Caulfield JP, Cianci CM, McDiarmid SS, Suyemitsu T, Schmid K. Ultrastructure, carbohydrate, and amino acid analysis of two preparations of the cercarial glycocalyx of *Schistosoma mansoni*. *Journal of Parasitology* **1987**; **73**:514–522.
 17. Caulfield JP, Yuan HC, Cianci CM, Hein A. *Schistosoma mansoni*: development of the cercarial glycocalyx. *Experimental Parasitology* **1988**; **65**:10–19.
 18. Chang CS, Chang K-P. Monoclonal antibody affinity purification of a *Leishmania* membrane glycoprotein and its inhibition of *Leishmania*-macrophage binding. *Proceedings of the National Academy of Sciences of the United States of America* **1986**; **83**:100.
 19. Chen HT, Kabat EA. Immunochemical studies on blood groups. The combining site specificities of mouse monoclonal hybridoma anti-A and anti-B. *Journal of Biological Chemistry* **1985**; **260**:13208–13217.
 20. Clegg JA, Smithers SR, Terry RJ. Acquisition of human antigens by *Schistosoma mansoni* during cultivation *in vitro*. *Nature (London)* **1971**; **232**:653–654.
 21. Cummings RD, Kornfeld S. Fractionation of asparagine-linked oligosaccharides by serial lectin-agarose affinity chromatography: a rapid sensitive and specific technique. *Journal of Biological Chemistry* **1982**; **257**:11235–11240.
 22. Cummings RD, Smith DF. The selectin family of carbohydrate-binding proteins – structure and importance of carbohydrate ligands for cell adhesion. *BioEssays* **1992**; **14**:849–856.
 23. Dalton JP, Lewis SA, Aronstein WS, Strand M. *Schistosoma mansoni*: immunogenic glycoproteins of the cercarial glycocalyx. *Experimental Parasitology* **1987**; **63**: 215–226.
 24. Danielson SJ, Gray GR. Immunochemical characterization of polylysine conjugates containing reductively aminated cellulose oligosaccharides. *Glycoconjugate Journal* **1986**; **3**:363–377.
 25. De Water R, Franssen JAM, Deelder AM. Ultrastructural localization of the circulating anodic antigen in the digestive tract of *Schistosoma mansoni* using monoclonal antibodies in an immunogold labeling procedure. *American Journal of Tropical Medicine and Hygiene* **1986**; **35**:549–558.
 26. De Water R, Franssen JAM, Deelder AM. Ultrastructural localization of the circulating cathodic antigen in the digestive tract of various life-cycle stages of *Schistosoma mansoni*. *Zeitschrift für Parasitenkunde* **1986**; **72**:635–646.
 27. Deelder AM, Kornelis D. Immunodiagnosis of recently acquired *Schistosoma mansoni* infection. A comparison of various immunological techniques. *Tropical and Geographical Medicine* **1981**; **33**:36–41.
 28. Deelder AM, Van Zeyl RJM, Fillié YE, Rotmans JP, Duchenne W. Recognition of gut-associated antigens by immunoglobulin M in the indirect fluorescent antibody test for schistosomiasis mansoni. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **1989**; **83**:364–367.
 29. Demeure CE, Rihet P, Abel L, Ouattara M, Bourgeois A, Dessein AJ. Resistance to *Schistosoma mansoni* in humans – influence of the IgE/IgG4 balance and IgG2 in immunity to reinfection after chemotherapy. *Journal of Infectious Diseases* **1993**; **168**:1000–1008.
 30. Der Balian GP, Slack J, Clevinger BL, Bazin H, Davie JM. Subclass restriction of murine antibodies. III. Antigens that stimulate IgG3 in mice stimulate IgG2c in rats. *Journal of Experimental Medicine* **1980**; **152**:209–218.
 31. Dessein AJ, Samuelson JC, Butterworth AE, Hogan M, Sherry BA, Vadas MA, David JR. Immune evasion by *Schistosoma mansoni*: loss of susceptibility to antibody or complement-dependent eosinophil attack by schistosomula cultured in medium free of macromolecules. *Parasitology* **1981**; **82**: 357–374.
 32. Dissous C, Capron A. *Schistosoma mansoni*: antigenic community between schistosomula surface and adult worm incubation products as a support for concomitant immunity. *FEBS Letters* **1983**; **162**:355–359.
 33. Dissous C, Grzych J-M, Capron A. *Schistosoma mansoni* surface antigen defined by a rat monoclonal IgG2a. *Journal of Immunology* **1982**; **129**:2232–2234.
 34. Dissous C, Grzych J-M, Capron A. Biochemical studies on the 30–40 kDa *Schistosoma mansoni* surface antigens. *Molecular and Biochemical Parasitology* **1985**; **16**:277–288.
 35. Dissous C, Grzych J-M, Capron A. *Schistosoma mansoni* shares a protective

- epitope with freshwater and marine snails. *Nature (London)* **1986**; **323**:443–445.
36. Dixon HFB. *Nature (London)* **1981**; **292**:192–192.
 37. Dunn TS, Yoshino TP. Tegumental surface modulation in *Schistosoma mansoni* primary sporocysts in response to ligand binding. *Parasite Immunology* **1991**; **13**:121–135.
 38. Dunne DW. Schistosome carbohydrates. *Parasitology Today* **1990**; **6**:45–48.
 39. Dunne DW, Bickle QD. Identification and characterisation of a polysaccharide-containing antigen from *Schistosoma mansoni* eggs which cross-reacts with the surface of schistosomula. *Parasitology* **1987**; **94**:255–268.
 40. Dunne DW, Bickle QD, Butterworth AE, Richardson BA. The blocking of human antibody-dependent, eosinophil-mediated killing of *Schistosoma mansoni* schistosomula by monoclonal antibodies which cross-react with a polysaccharide-containing egg antigen. *Parasitology* **1987**; **94**:269–280.
 41. Dunne DW, Jones FM, Doenhoff MJ. The purification, characterization, serological activity and hepatotoxic properties of two cationic glycoproteins ($\alpha 1$ and $\Omega 1$) from *Schistosoma mansoni* eggs. *Parasitology* **1991**; **103**:225–236.
 42. Evengård B, Hammarström L, Smith CIE, Linder E. Early antibody responses in human schistosomiasis. *Clinical and Experimental Immunology* **1990**; **80**:69–76.
 43. Feizi T. Demonstration by monoclonal antibodies that carbohydrate structures of glycoproteins and glycolipids are onco-developmental antigens. *Nature (London)* **1985**; **314**:53–57.
 44. Ferrante A, Allison AC. Alternative pathway activation of complement by African trypanosomes lacking a glycoprotein coat. *Parasite Immunology* **1983**; **5**:491.
 45. Fukuda M. *Biochemical and Biophysical Acta* **1985**; **780**:119–150.
 46. Fukuda M. Lysosomal membrane glycoproteins: structure, biosynthesis, and intracellular trafficking. *Journal of Biological Chemistry* **1991**; **266**:21327–21330.
 47. Galvao dos Reis M, Secor E, De Souza Andrade Filho A, Cardoso Silva M, Silveira Santos SR, Vasilaski G, Almeida dos Reis E, Velupillai P, Harn DA. Antibodies against defined oligosaccharide in spinal cord fluid: a study among patients with cerebral disorder. In: *Abstract book of the International Symposium on Schistosomiasis*. Instituto Oswaldo Cruz, Rio de Janeiro, **1993**:148.
 48. Gerard C. Purification of glycoproteins. *Methods in Enzymology* **1990**; **182**:529–539.
 49. Goldring OL, Clegg JA, Smithers SR, Terry RJ. Acquisition of human blood group antigens by *Schistosoma mansoni*. *Clinical and Experimental Immunology* **1976**; **26**:181–187.
 50. Goldstein IJ, Hughes RC, Monsigny M, Osawa T, Sharon N. What should be called a lectin? *Nature (London)* **1980**; **285**:66–66.
 51. Goldstein IJ, Poretz RD. In: Liener IE, Sharon N, Goldstein IJ, eds. *The lectins: properties, functions, and applications in biology and medicine*. Academic Press Inc., London, **1986**:33–245.
 52. Gooi HC, Thorpe SJ, Hounsell EF, Rumpold H, Kraft D, Förster O, Feizi T. Marker of peripheral blood granulocytes and monocytes of man recognized by two monoclonal antibodies VEP8 and VEP9 involves the trisaccharide 3-fucosyl-N-acetyl-lactosamine. *European Journal of Immunology* **1983**; **13**:306–312.
 53. Gregoire RJ, Shi MH, Rekosch DM, Loverde PT. Protective monoclonal antibodies from mice vaccinated or chronically infected with *Schistosoma mansoni* that recognize the same antigens. *Journal of Immunology* **1987**; **139**:3792–3801.
 54. Grzych J-M, Capron M, Bazin H, Capron A. *In vitro* and *in vivo* effector function of rat IgG2a monoclonal anti-*Schistosoma mansoni* antibodies. *Journal of Immunology* **1982**; **129**:2739–2743.
 55. Grzych J-M, Capron M, Dissous C, Capron A. Blocking activity of rat monoclonal antibodies in experimental schistosomiasis. *Journal of Immunology* **1984**; **133**:998–1004.
 56. Grzych J-M, Capron M, Lambert P-H, Dissous C, Torres S, Capron A. An anti-idiotypic vaccine against experimental schistosomiasis. *Nature (London)* **1985**; **316**:74–76.
 57. Grzych J-M, Dissous C, Capron M, Torres S, Lambert P-H, Capron A. *Schistosoma mansoni* shares a protective carbohydrate epitope with keyhole limpet haemocyanin. *Journal of Experimental Medicine* **1987**; **165**:865–878.
 58. Haavik S, Codrington JF, Davison PF. Development and characterization of monoclonal antibodies against a mucin-type glycoprotein. *Glycobiology* **1992**; **2**:217–224.
 59. Hakomori S-I, Nudelman E, Lavery SB, Kannagi R. Novel fucolipids accumulating in human adenocarcinoma. I. Glycolipids with

- di- or trifucosylated type 2 chains. *Journal of Biological Chemistry* 1984; 259:4672-4680.
60. Hamburger J, Lustigman S, Arap Siongok TK, Ouma JH, Mahmoud AA. Characterization of a purified glycoprotein from *Schistosoma mansoni* eggs: specificity, stability, and the involvement of carbohydrate and peptide moieties in its serologic activity. *Journal of Immunology* 1982; 128:1864-1869.
61. Hamburger J, Lustigman S, Arap Siongok TK, Ouma JH, Mahmoud AA. Analysis and preliminary purification of glycoproteins isolated from eggs in the urine of patients with *Schistosoma haematobium* infection. *Journal of Immunology* 1982; 129:1711-1714.
62. Hamburger J, Pelley RP, Warren KS. *Schistosoma mansoni* soluble egg antigens: determination of the stage and species specificity of their serologic reactivity by radioimmunoassay. *Journal of Immunology* 1976; 117:1561-1566.
63. Handman E, Goding JW. The *Leishmania* receptor for macrophages is a lipid-containing glycoconjugate. *EMBO Journal* 1985; 4:329.
64. Hanisch F-G, Chai W, Rosankiewicz JR, Lawson AM, Stoll MS, Feizi T. *European Journal of Biochemistry* 1993; 217:645-655.
65. Harn DA, David JR, Galvao dos Reis M, Almeida dos Reis E, Velupillai P. Fucosylated oligosaccharides induce proliferation of PBMC from *Schistosoma mansoni* infected patients. In: *Abstract book of the International Symposium on Schistosomiasis*. Instituto Oswaldo Cruz, Rio de Janeiro, 1993:147.
66. Harn DA, Mitsuyama M, David JR. *Schistosoma mansoni*. Anti-egg monoclonal antibodies protect against cercarial challenge *in vivo*. *Journal of Experimental Medicine* 1984; 159:1371-1387.
67. Harn DA, Quinn JJ, Cianci CM, Ko AI. Evidence that a protective membrane epitope is involved in early but not late phase immunity in *Schistosoma mansoni*. *Journal of Immunology* 1987; 138:1571-1580.
68. Hassan MM, Badawi MA, Strand M. Circulating schistosomal antigen in diagnosis and assessment of cure in individuals infected with *Schistosoma mansoni*. *American Journal of Tropical Medicine and Hygiene* 1992; 46:737-744.
69. Hayunga EG, Sumner MP. Characterization of surface glycoproteins on *Schistosoma mansoni* adult worms by lectin affinity chromatography. *Journal of Parasitology* 1986; 72:283-291.
70. Hayunga EG, Sumner MP. Expression of lectin-binding surface glycoproteins during the development of *Schistosoma mansoni* schistosomula. *Journal of Parasitology* 1986; 72:913-920.
71. Holy JM, O'Leary K, Oaks JA, Tracy JW. Immunocytochemical localization of the major Glutathione S-transferases in adult *Schistosoma mansoni*. *Journal of Parasitology* 1989; 75:181-190.
72. Hong Z, Loverde PT, Thakur A, Hammarskjold ML, Rekosh D. *Schistosoma mansoni*: a Cu/Zn superoxide dismutase is glycosylated when expressed in mammalian cells and localizes to a subtegumental region in adult schistosomes. *Experimental Parasitology* 1993; 76:101-114.
73. Horta MFM, Ramalho-Pinto FJ. Role of human Decay-accelerating factor in the evasion of *Schistosoma mansoni* from the complement-killing *in vitro*. *Journal of Experimental Medicine* 1991; 174:1399-1406.
74. Hounsell EF. A general strategy for glycoprotein oligosaccharide analysis. In: Hounsell EF, ed. *Methods in Molecular Biology, Vol. 14: Glycoprotein analysis in biomedicine*. Humana Press Inc., Totowa, 1993:1-15.
75. Hounsell EF, Jones NJ, Gooi HC, Feizi T, Donald ASR, Feeney J. 500 MHz ¹H-N.M.R. and conformational studies of fucosyl-oligosaccharides recognised by monoclonal antibodies with specificities related to Le-a, Le-b and SSEA-1. *Carbohydrate Research* 1988; 178:67-78.
76. Huberty MC, Vath JE, Yu W, Martin SA. Site-specific carbohydrate identification in recombinant proteins using MALD-TOF MS. *Analytical Chemistry* 1993; 65:2791-2800.
77. Imai Y, Lasky LA, Rosen SD. Further characterization of the interaction between L-selectin and its endothelial ligands. *Glycobiology* 1992; 2:373-381.
78. Jasmer DP, Perryman LE, Conder GA, Crow S, Mcguire T. Protective immunity to *Haemonchus contortus* induced by immunoaffinity isolated antigens that share a phylogenetically conserved carbohydrate gut surface epitope. *Journal of Immunology* 1993; 151:5450-5460.
79. Jefferis R, Kumararatne DS. Selective IgG subclass deficiency: quantification and clinical relevance. *Clinical and Experimental Immunology* 1990; 81:357-367.
80. Jwo J, Loverde PT. The ability of fractionated sera from animals vaccinated with irradiated cercariae of *Schistosoma mansoni* to transfer immunity to mice. *Journal of Parasitology* 1989; 75:252-260.
81. Kamerling JP. Xylose-containing carbohydrate chains derived from N-glycoproteins.

- Pure & Applied Chemistry* 1991; 63: 465-472.
82. Kannagi R, Nudelman E, Lavery SB, Hakomori S-I. A series of human erythrocyte glycosphingolipids reacting to the monoclonal antibody directed to a developmentally regulated antigen, SSEA-1. *Journal of Biological Chemistry* 1982; 257:14865-14874.
 83. Kelly C, Simpson AJG, Fox E, Phillips SM, Smithers SR. The identification of *Schistosoma mansoni* surface antigens recognized by protective monoclonal antibodies. *Parasite Immunology* 1986; 8:193-198.
 84. Kerr MA, Stocks SC. The role of CD15-(Lex)-related carbohydrates in neutrophil adhesion. *Histochemical Journal* 1992; 24:811-826.
 85. Kéry V. Lectin-carbohydrate interactions in immunoregulation. *International Journal of Biochemistry* 1991; 23:631-640.
 86. Khalife J, Capron M, Capron A, Grzych J-M, Butterworth AE, Dunne DW, Ouma JH. Immunity in human schistosomiasis mansoni. Regulation of protective immune mechanisms by IgM blocking antibodies. *Journal of Experimental Medicine* 1986; 164: 1626-1640.
 87. Khalife J, Dunne DW, Richardson BA, Mazza G, Thorne KJI, Capron A, Butterworth AE. Functional role of human IgG subclasses in eosinophil-mediated killing of schistosomula of *Schistosoma mansoni*. *Journal of Immunology* 1989; 142:4422-4427.
 88. King CH, El Ibiary S, El Nawawi M, Sawyer J, Griffin A, El Hawey A, Mahmoud AAF. Intensity of *Schistosoma mansoni* infection in a human population is inversely correlated with antibody response to Smw68, a protective parasite antigen. *Journal of Infectious Diseases* 1989; 160:686-691.
 89. Knutsen AP. Patients with IgG subclass and or selective antibody deficiency to polysaccharide antigens. Initiation of a controlled clinical trial of intravenous immune globulin. *Journal of Allergy and Clinical Immunology* 1989; 84:640-647.
 90. Ko AI, Dräger UC, Harn DA. A *Schistosoma mansoni* epitope recognized by a protective monoclonal antibody is identical to the stage-specific embryonic antigen 1. *Proceedings of the National Academy of Sciences of the United States of America* 1990; 87:4159-4163.
 91. Ko AI, Harn DA. Characterization of protective and non-protective surface membrane carbohydrate epitopes of *Schistosoma mansoni*. *Memorias Do Instituto Oswaldo Cruz* 1987; 82 Suppl 4:115-119.
 92. Kobata A. The carbohydrates of glycoproteins. In: Ginsberg V, Robbins PW, eds. *Biology of Carbohydrates, Vol. 2*. Wiley, Chichester, 1984:87.
 93. Kobata A, Ginsburg V. Oligosaccharides from human milk. II. Isolation and characterization of a new pentasaccharide, lacto-N-fucopentaose III. *Journal of Biological Chemistry* 1969; 244:5496-5502.
 94. Kocourek J, Horejsi V. Defining a lectin. *Nature (London)* 1981; 290:188-188.
 95. Kornfeld R, Kornfeld S. In: Lennarz WJ, ed. *The biochemistry of glycoproteins and proteoglycans*. Plenum, New York, 1980:1-34.
 96. Köster B, Strand M. *Schistosoma mansoni*: immunolocalization of two different fucose-containing carbohydrate epitopes. *Parasitology* 1994; 108:433-446.
 97. Langley JG, Dunne DW. Temporal variation in the carbohydrate and peptide surface epitopes in antibody-dependent, eosinophil-mediated killing of *Schistosoma mansoni* schistosomula. *Parasite Immunology* 1992; 14:185-200.
 98. Lee KB, Loganathan D, Merchant ZM, Linhardt RJ. Carbohydrate Analysis of Glycoproteins. A Review. *Applied Biochemistry and Biotechnology* 1990; 23:53-80.
 99. Lennarz W, Hart GW. *Guide to techniques in glycobiology. Series: Methods in Enzymology, vol. 230*. Academic Press, Inc., San Diego, 1994.
 100. Lavery SB, Weiss JB, Salyan MEK, Roberts CE, Hakomori S-I, Magnani JL, Strand M. Characterization of a series of novel fucose-containing glycosphingolipid immunogens from eggs of *Schistosoma mansoni*. *Journal of Biological Chemistry* 1992; 267:5542-5551.
 101. Lewis SA, Strand M. Characterization of proteins and immunogens released by adult *Schistosoma mansoni*. *Journal of Parasitology* 1991; 77:263-271.
 102. Linder E. *Schistosoma mansoni*: visualization with fluorescent lectins of secretions and surface carbohydrates of living cercariae. *Experimental Parasitology* 1985; 59: 307-312.
 103. Linder E, Hult G. Distribution of exposed and hidden carbohydrates of *Schistosoma mansoni* adult worms demonstrated by selective binding of fluorochrome-conjugated lectins. *Parasitology* 1982; 85:503-509.
 104. Lis H, Sharon N. *Annual Review of Biochemistry* 1986; 55:35.

105. Lustigman S, Hamburger J. *Schistosoma mansoni*: radiometric assay of lectin binding specificities of the major egg glycoprotein and its carbohydrate-rich fragment. *Experimental Parasitology* 1985; 59:59-67.
106. Lustigman S, Mahmoud AA, Hamburger J. Glycopeptides in soluble egg antigen of *Schistosoma mansoni*: isolation, characterization, and elucidation of their immunochemical and immunopathological relation to the major egg glycoprotein (MEG). *Journal of Immunology* 1985; 134:1961-1967.
107. MacGregor AN, Kusel JR. Isolation and characterization of a surface membrane glycoprotein from adult *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* 1989; 34:237-244.
108. MacGregor AN, Stott DI, Kusel JR. Lectin binding to glycoproteins in the surface membrane of *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* 1985; 16:163-172.
109. Maemura K, Fukuda M. Poly-*N*-acetyl-lactosaminyl O-glycans attached to leukosialin - the presence of sialyl Le-x structures in O-glycans. *Journal of Biological Chemistry* 1992; 267:24379-24386.
110. Makaanu CK, Damian RT, Smith DF, Cummings RD. The human blood fluke *Schistosoma mansoni* synthesizes a novel type of glycosphingolipid. *Journal of Biological Chemistry* 1992; 267:2251-2257.
111. Margolis RK, Margolis RU. Structure and Localization of Glycoproteins and Proteoglycans. In: Margolis RU, Margolis RK, eds. *Neurobiology of Glycoconjugates*. Plenum Publishing Corporation, New York, 1989:85-126.
112. Marikovsky M, Arnon R, Fishelson Z. Proteases secreted by transforming schistosomula of *Schistosoma mansoni* promote resistance to killing by complement. *Journal of Immunology* 1988; 141:273-278.
113. Marikovsky M, Parizade M, Arnon R, Fishelson Z. Complement regulation on the surface of cultured schistosomula and adult worms of *Schistosoma mansoni*. *European Journal of Immunology* 1990; 20:221-227.
114. Mazza G, Dunne DW, Butterworth AE. Antibody isotype responses to the *Schistosoma mansoni* schistosomulum in the CBA/N mouse induced by different stages of the parasite life cycle. *Parasite Immunology* 1990; 12:529-543.
115. McDiarmid SS, Podesta RB. Identification of a sialic acid containing glycocalyx on the surface of *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* 1984; 10:33-43.
116. Merkle RK, Cummings RD. Lectin affinity chromatography of glycopeptides. *Methods in Enzymology* 1987; 138:232-259.
117. Montreuil J. Primary structure of glycoprotein glycans (basis for the molecular biology of glycoproteins). *Advances in Carbohydrate Chemistry and Biochemistry* 1980; 37:157-223.
118. Moody SF. Molecular variation in *Leishmania*. *Acta Tropica (Basel)* 1993; 53:185-204.
119. Moreno C, Esdaine J. Immunoglobulin isotype in the murine response to polysaccharide antigens. *European Journal of Immunology* 1983; 13:262-264.
120. Moser G, Wassom DL, Sher A. Studies of the antibody-dependent killing of schistosomula of *Schistosoma mansoni* employing haptenic target antigens. I. Evidence that the loss in susceptibility to immune damage undergone by developing schistosomula involves a change unrelated to the masking of parasite antigens by host molecules. *Journal of Experimental Medicine* 1980; 152:41-53.
121. Muramatsu T. Carbohydrate signals in metastasis and prognosis of human carcinomas. *Glycobiology* 1993; 3:294-296.
122. Murrell KD, Taylor DW, Vannier WE, Dean DA. *Schistosoma mansoni*: analysis of surface membrane carbohydrates using lectins. *Experimental Parasitology* 1978; 46:247-255.
123. Nanduri J, Dennis JE, Rosenberry TL, Mahmoud AAF, Tartakoff AM. Glycocalyx of bodies versus tails of *Schistosoma mansoni* cercariae. Lectin-binding, size, charge, and electron microscopic characterization. *Journal of Biological Chemistry* 1991; 266:1341-1347.
124. Nash TE, Deelder AM. Comparison of four schistosome excretory-secretory antigens: phenol-sulfuric test active peak, cathodic circulating antigen, gut-associated proteoglycan, and circulating anodic antigen. *American Journal of Tropical Medicine and Hygiene* 1985; 34:236-241.
125. Nash TE, Ottesen EA, Cheever AW. Antibody response to a polysaccharide antigen present in the schistosome gut. II. Modulation of antibody response. *American Journal of Tropical Medicine and Hygiene* 1978; 27:944-950.
126. Nash TE, Prescott B, Neva FA. The characteristics of a circulating antigen in schistosomiasis. *Journal of Immunology* 1974; 112:1500-1507.
127. Newman BA, Liao J, Gruezo F, Sugii S, Kabat EA, Torii M, Clevinger BL, Davie JM, Schilling J, Bond M, Hood L. Immunochemical studies of mouse monoclonal antibodies to dextran B1355S-II. Combining site specificity,

- sequence, idotype and affinity. *Molecular Immunology* **1986**; **23**:413-424.
128. Nyame K, Cummings RD, Damian RT. *Schistosoma mansoni* synthesizes glycoproteins containing terminal O-linked N-acetylglucosamine residues. *Journal of Biological Chemistry* **1987**; **262**:7990-7995.
129. Nyame K, Cummings RD, Damian RT. Characterization of the N- and O-linked oligosaccharides in glycoproteins synthesized by *Schistosoma mansoni* schistosomula. *Journal of Parasitology* **1988**; **74**:562-572.
130. Nyame K, Cummings RD, Damian RT. Characterization of the high mannose asparagine-linked oligosaccharides synthesized by *Schistosoma mansoni* adult male worms. *Molecular and Biochemical Parasitology* **1988**; **28**:265-274.
131. Nyame K, Smith DF, Damian RT, Cummings RD. Complex-type asparagine-linked oligosaccharides in glycoproteins synthesized by *Schistosoma mansoni* adult males contain terminal β -linked N-acetylgalactosamine. *Journal of Biological Chemistry* **1989**; **264**:3235-3243.
132. Omer-Ali P, Hagan P, Wilkins HA, Simpson AJG. Antibody to schistosomulum surface carbohydrate epitopes in subjects infected with *Schistosoma haematobium*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **1989**; **83**:358-361.
133. Omer-Ali P, Magee AI, Kelly C, Simpson AJG. A major role for carbohydrate epitopes preferentially recognized by chronically infected mice in the determination of *Schistosoma mansoni* schistosomulum surface antigenicity. *Journal of Immunology* **1986**; **137**:3601-3607.
134. Omer-Ali P, Smithers SR, Bickle QD, Phillips SM, Harn DA, Simpson AJG. Analysis of the anti-*Schistosoma mansoni* surface antibody response during murine infection and its potential contribution to protective immunity. *Journal of Immunology* **1988**; **140**:3273-3279.
135. Page AP, Hamilton AJ, Maizels RM. *Toxocara canis*: monoclonal antibodies to carbohydrate epitopes of secreted (TES) antigens localize to different secretion-related structures in infective larvae. *Experimental Parasitology* **1992**; **75**:56-71.
136. Palatnikdesousa CB, Dutra HS, Borojevic R. *Leishmania donovani* surface glycoconjugate Gp36 is the major immunogen component of the fucose-mannose ligand (FML). *Acta Tropica (Basel)* **1993**; **53**:59-72.
137. Patterson SD, Bell K. The carbohydrate side chains of the major plasma serpins of horse and wallaby - analyses of enzymatic and chemically treated (including "Smith degradation") protein blots by lectin binding. *Biochemistry International* **1990**; **20**:429-436.
138. Payares G, Simpson AJG. *Schistosoma mansoni* surface glycoproteins. Analysis of their expression and antigenicity. *European Journal of Biochemistry* **1985**; **153**:195-201.
139. Pearce EJ, Basch PF, Sher A. Evidence that the reduced surface antigenicity of developing *Schistosoma mansoni* schistosomula is due to antigen shedding rather than host molecule acquisition. *Parasite Immunology* **1986**; **8**:79-94.
140. Pelley RP, Pelley RJ. *S. mansoni* soluble egg antigens - IV. Biochemistry and immunology of major serological antigens with particular emphasis on MSA₁. In: Van den Bossche H, ed. *Biochemistry of parasites and host-parasite relationships*. Elsevier/North-Holland Biomedical Press, Amsterdam, **1976**:283-290.
141. Pelley RP, Pelley RJ, Hamburger J, Peters PA, Warren KS. *Schistosoma mansoni* soluble egg antigens I. Identification and purification of three major antigens, and the employment of radioimmunoassay for further characterization. *Journal of Immunology* **1976**; **117**:1553-1560.
142. Perlmutter RM, Hansburg D, Briles DE, Nicolotti RA, Davie JM. Subclass restriction of murine anti-carbohydrate antibodies. *Journal of Immunology* **1978**; **121**:566-572.
143. Prenner C, Mach L, Glössl J, März L. The antigenicity of the carbohydrate moiety of an insect glycoprotein, honeybee (*Apis mellifera*) venom phospholipase A2. The role of α 1,3-fucosylation of the asparagine-bound N-acetylglucosamine. *Biochemical Journal* **1992**; **284**:377-380.
144. Pujol FH, Liprandi F, Rodriguez M, Cesari IM. Production of a mouse monoclonal antibody against the alkaline phosphatase of adult *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* **1990**; **40**:43-52.
145. Quigg RJ. Inhibition of the alternative pathway of complement by glomerular chondroitin sulphate proteoglycan. *Immunology* **1992**; **76**:373-377.
146. Rasmussen KR, Kemp WM. *Schistosoma mansoni*: interactions of adult parasites with the complement system. *Parasite Immunology* **1987**; **9**:235-248.
147. Richter D, Incani RN, Harn DA. Isotype responses to candidate vaccine antigens in protective sera obtained from mice vaccinated with irradiated cercariae of *Schistosoma mansoni*. *Infection and Immunity* **1993**; **61**:3003-3011.
148. Ripert C, Combe A, Daulouede S, Appriou M, Tribouley-Duret J, Tribouley J, Moyou-Somo

- R, Same-Ekobo A, Ambassa P. Detection with a monoclonal antibody of an antigen characteristic of the genus *Schistosoma* excreted in the urine. *Tropical Medicine and Parasitology* 1988; 39:131-135.
149. Robertson NP, Cain GD. Isolation and characterization of glycosaminoglycans from *Schistosoma mansoni*. *Comparative Biochemistry and Physiology* 1985; 82B: 299-306.
150. Rovis L, Musoke AJ, Moloo SK. Failure of trypanosomal membrane antigens to induce protection against tsetse-transmitted *Trypanosoma vivax* or *T. brucei* in goats and rabbits. *Acta Tropica (Basel)* 1984; 41:227.
151. Russell DG. The macrophage-attachment glycoprotein gp63 is the predominant C3-acceptor site on *Leishmania mexicana* promastigotes. *European Journal of Biochemistry* 1987; 164:213.
152. Saitoh O, Wang W-C, Lotan R, Fukuda M. Differential glycosylation and cell surface expression of lysosomal membrane glycoproteins in sublines of a human colon cancer exhibiting distinct metastatic potentials. *Journal of Biological Chemistry* 1992; 267:5700-5711.
153. Samuelson JC, Caulfield JP. The cercarial glycocalyx of *Schistosoma mansoni*. *Journal of Cell Biology* 1985; 100:1423-1434.
154. Samuelson JC, Caulfield JP. Cercarial glycocalyx of *Schistosoma mansoni* activates human complement. *Infection and Immunity* 1986; 51:181-186.
155. Sheinfeld J, Reuter VE, Sarkis AS, Cordon-Cardo C. Blood group antigens in normal and neoplastic urothelium. *Journal of Cellular Biochemistry* 1992; Suppl. 16: 50-55.
156. Sher A. Acquisition of murine major histocompatibility complex gene products by schistosomula of *Schistosoma mansoni*. *Journal of Experimental Medicine* 1978; 148:46-57.
157. Sherblom AP, Sathyamoorthy N, Decker JM, Muchmore AV. IL-2, a lectin with specificity for high mannose glycopeptides. *Journal of Immunology* 1989; 143:939-944.
158. Siber GR, Schur PH, Aisenberg AC, Weitzman SA, Schiffman G. Correlation between serum IgG2 concentrations and the antibody response to bacterial polysaccharide antigens. *The New England Journal of Medicine* 1980; 303:178-182.
159. Simpson AJG, Correa-Oliveira R, Smithers SR, Sher A. The exposed carbohydrates of schistosomula of *Schistosoma mansoni* and their modification during maturation *in vivo*. *Molecular and Biochemical Parasitology* 1983; 8:191-205.
160. Simpson AJG, Omer-Ali P, Yi X, Smithers SR. Schistosome glycoconjugates as antigens and their relevance in experimental and human schistosomiasis. *Biochemical Society Transactions* 1988; 16:264-265.
161. Simpson AJG, Smithers SR. Characterization of the exposed carbohydrates on the surface membrane of adult *Schistosoma mansoni* by analysis of lectin binding. *Parasitology* 1980; 81:1-15.
162. Simpson AJG, Yi X, Lillywhite JE, Omer-Ali P, Kelly CG, Mott R, Smithers SR, Correa-Oliveira R, Katz N, Rumjanek F, Hagan P. Dissociation of antibody responses during human schistosomiasis and evidence for enhancement of granuloma size by anti-carbohydrate IgM. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1990; 84:808-814.
163. Sinn HP, Brown SA, Oberle E, Thompson JS. Analysis of the lewis-x epitope in human pancreas and pancreatic adenocarcinomas. *International Journal of Pancreatology* 1992; 11:125-135.
164. Smith MA, Clegg JA, Snary D, Trejdosiewicz AJ. Passive immunization of mice against *Schistosoma mansoni* with an IgM monoclonal antibody. *Parasitology* 1982; 84:83-91.
165. Spooncer E, Fukuda M, Klock JC, Oates JE, Dell A. Isolation and characterization of polyfucosylated lactosaminoglycan from human granulocytes. *Journal of Biological Chemistry* 1984; 259:4792-4801.
166. Springer TA, Lasky LA. Sticky sugars for selectins. *Nature (London)* 1991; 349: 196-197.
167. Srivatsan J, Smith DF, Cummings RD. The human blood fluke *Schistosoma mansoni* synthesizes glycoproteins containing the Lewis X antigen. *Journal of Biological Chemistry* 1992; 267:20196-20203.
168. Srivatsan J, Smith DF, Cummings RD. *Schistosoma mansoni* synthesizes novel biantennary Asn-linked oligosaccharides containing terminal β -linked *N*-acetyl-galactosamine. *Glycobiology* 1992; 2: 445-452.
169. Stelma FF, Talla I, Verlé P, Niang M, Gryseels B. Morbidity due to heavy *Schistosoma mansoni* infections in a recently established focus in northern Senegal. *American Journal of Tropical Medicine and Hygiene* 1994; 50:575-579.
170. Stöcks SC, Kerr MA. Stimulation of neutrophil adhesion by antibodies recognizing CD15 (Lex) and CD15-expressing carcino-embryonic antigen-related glycoprotein NCA-160. *Biochemical Journal* 1992; 288: 23-27.

171. Strand M, McMillan A, Pan X. *Schistosoma mansoni*: reactivity with infected human sera and monoclonal antibody characterization of a glycoprotein in different developmental stages. *Experimental Parasitology* 1982; 54:145–156.
172. Strous GJ, Dekker J. Mucin-type glycoproteins. *Critical Reviews in Biochemistry and Molecular Biology* 1992; 27:57–92.
173. Taylor DW. Schistosome vaccines. *Experientia* 1991; 47:152–157.
174. Tetteroo P, Geurts van Kessel A. Expression of CD15 (FAL) on myeloid cells and chromosomal localization of the gene. *Histochemical Journal* 1992; 24:777–782.
175. Tretter V, Altmann F, Kubelka V, März L, Becker WM. Fucose α 1,3-linked to the core region of glycoprotein N-glycans creates an important epitope for IgE from honeybee venom allergic individuals. *International Archives of Allergy and Immunology* 1993; 102:259–266.
176. Tsuda M, Kurokawa T, Takeuchi M, Sugino Y. Changes in cell surface structure by viral transformation studied by binding of lectins differing in sugar specificity. *Japanese Journal of Cancer Research* 1975; 66: 513–521.
177. Uchikawa R, Loker ES. Lectin-binding properties of the surfaces of *in vitro*-transformed *Schistosoma mansoni* and *Echinostoma paraensei* sporocysts. *Journal of Parasitology* 1991; 77:742–748.
178. Van Dam GJ, Bergwerff AA, Thomas-Oates JE, Rotmans JP, Kamerling JP, Vliegenthart JFG, Deelder AM. The immunologically reactive O-linked polysaccharide chains derived from Circulating Cathodic Antigen isolated from the human blood fluke *Schistosoma mansoni* have Lewis x as repeating unit. *European Journal of Biochemistry* 1994; 225:467–482.
179. Van Dam GJ, Verheul AFM, Zigterman GJWJ, De Reuver MJ, Snippe H. Estimation of the avidity of antibodies in polyclonal antisera against *Streptococcus pneumoniae* type 3 by inhibition ELISA. *Molecular Immunology* 1989; 26:269–274.
180. Van Kuik JA, Van Halbeek H, Kamerling JP, Vliegenthart JFG. Primary structure of the low-molecular weight carbohydrate chains of *Helix pomatia* α -hemocyanin. *Journal of Biological Chemistry* 1985; 260: 13984–13988.
181. Varki A. Biological roles of oligosaccharides: all of the theories are correct. *Glycobiology* 1993; 3:97–130.
182. Velupillai P, Harn DA. Oligosaccharide-specific induction of interleukin 10 production by B220+ cells from schistosome-infected mice: a mechanism for regulation of CD4+ T-cell subsets. *Proceedings of the National Academy of Sciences of the United States of America* 1994; 91:18–22.
183. Watanabe K, Lain RA, Hakomori S-I. On neutral fucoglycolipids having long, branched carbohydrate chains: H-active and I-active glycosphingolipids of human erythrocytes. *Biochemistry* 1975; 14:2725–2733.
184. Weiss JB, Magnani JL, Strand M. Identification of *Schistosoma mansoni* glycolipids that share immunogenic carbohydrate epitopes with glycoproteins. *Journal of Immunology* 1986; 136: 4275–4282.
185. Weiss JB, Strand M. Characterization of developmentally regulated epitopes of *Schistosoma mansoni* egg glycoprotein antigens. *Journal of Immunology* 1985; 135: 1421–1429.
186. Wessels MR, Pozsgay V, Kasper DL, Jennings HJ. Structure and immunochemistry of an oligosaccharide repeating unit of the capsular polysaccharide of type III group B *Streptococcus*. A revised structure for the type III group B streptococcal polysaccharide. *Journal of Biological Chemistry* 1987; 262:8262–8267.
187. Woodward MP, Young Jr WW, Bloodgood RA. Detection of monoclonal antibodies specific for carbohydrate epitopes using periodate oxidation. *Journal of Immunological Methods* 1985; 78:143–153.
188. Yamashita K, Mizuochi T, Kobata A. Analysis of oligosaccharides by gel filtration. *Methods in Enzymology* 1982; 83:105.
189. Yi X, Omer-Ali P, Kelly C, Simpson AJG, Smithers SR. IgM antibodies recognizing carbohydrate epitopes shared between schistosomula and miracidia of *Schistosoma mansoni* which block *in vitro* killing. *Journal of Immunology* 1986; 137:3946–3954.
190. Yi X, Simpson AJG, De Rossi R, Smithers SR. The presence of antibody in mice chronically infected with *Schistosoma mansoni* which blocks *in vitro* killing of schistosomula. *Journal of Immunology* 1986; 137: 3955–3958.
191. Zamze S. Glycosylation in parasitic protozoa of the Trypanosomatidae family. *Glyco-conjugate Journal* 1991; 8:443–447.
192. Zodda DM, Phillips SM. Monoclonal antibody-mediated protection against *Schistosoma mansoni* infection in mice. *Journal of Immunology* 1982; 129: 2326–2328.