



Universiteit
Leiden
The Netherlands

Circulating gut-associated antigens of *Schistosoma mansoni* : biological, immunological, and molecular aspects

Dam, G.J. van

Citation

Dam, G. J. van. (1995, February 9). *Circulating gut-associated antigens of Schistosoma mansoni : biological, immunological, and molecular aspects*. Retrieved from <https://hdl.handle.net/1887/41317>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/41317>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/41317> holds various files of this Leiden University dissertation.

Author: Dam, G.J. van

Title: Circulating gut-associated antigens of *Schistosoma mansoni* : biological, immunological, and molecular aspects

Issue Date: 1995-02-09

Chapter 12

General discussion



Chapter 12

General discussion

Schistosoma antigens have long been a topic of intensive research. An important reason for this undoubtedly is the interesting life-cycle involving many host-parasite interactions. In particular in the final hosts, *e.g.* humans and cattle, these interactions often may be dangerous for the parasite, as they take place in the blood, the schistosome habitat. The blood is equipped with a number of physiological defense mechanisms (especially immunological and hemostatic ones) against injury and infection, which have forced the parasite to develop its own efficient defense system. It is in the interest of man, as a final host, to overcome these parasite evasion mechanisms by manipulation and education of his own defense systems to recognize and destroy the parasite. For this reason most of the research has been focused on tegumental surface antigens since they are the major targets for the host immune response. However, another source of large amounts of parasite antigens presented to the host is the schistosome gut, a very active organ for the digestion of host blood cells and nutrients, as well as for excretion and regurgitation of undigested compounds.

In the early report of Berggren and Weller (1967) a negatively charged antigen was described which generally is assumed to be identical with CAA, although localization studies had not yet been performed [6]. CCA was first described independently by Carlier *et al.* (1975) [16] and Deelder *et al.* (1976) [26], but not until 20 years after the first description of the antigens, reliable, accurate and reproducible assay-systems for the detection of CAA and CCA were developed and widely applied [23–25,71]. Essential to the success of these diagnostic techniques was the development and application of monoclonal antibodies. Moreover, the use of these CAA- and CCA-specific McAbs has facilitated structural analysis of the antigens, as well as studies on the clearance in experimental animal infection and on the immunopathological involvement in schistosome infections in experimental animals and humans.

Considering circulating antigens as diagnostic markers, a number of other antigens have also been described as potentially useful targets for immunodetection and immunodiagnosis of schistosomiasis. Hayunga *et al.* (1986) developed an assay for detection of cercarial antigen in serum, which could



determine a 100-worm infection in mice, as early as 1 week after infection [39]. Application of this assay would facilitate an early and accurate diagnosis, *e.g.* of infected travellers. However, only a very limited field study using this assay has been performed, which showed that chronic schistosomiasis infections also could be detected [40].

Strand and co-workers described that a schistosome egg antigen containing carbohydrate epitopes, which were cross-reactive with antigens of various life-cycle stages, could be detected in serum of patients using a McAb [37,74]. After praziquantel treatment of the patients antigen levels were shown to decrease [37]. A McAb-based inhibition passive hemagglutination assay for detection of a polysaccharide egg antigen in urine has been used by Ripert *et al.* in several epidemiological studies [47,55,56].

Recently, a highly sensitive and specific ELISA for the detection of circulating soluble egg antigen was developed within our laboratory [53]. In this assay two different McAbs were utilized in a mixture both for coating and detection. The correlation with egg output and serum CAA was very high. The McAbs were shown to be highly reactive with two different repetitive carbohydrate epitopes of soluble egg antigen and also showed a strong reactivity with antigens present on the cercarial and schistosomular surface, as well as recognizing antigens in the parenchyma (flame cells, excretory system) of adult worms [8,52].

An attempt to define and apply circulating antigens other than those described above but still potentially relevant for immunodiagnosis using McAbs present in our laboratory was unsuccessful. Remarkably, it was found that a major portion of the selected McAbs, showing diverse reaction patterns in fluorescence assays and in other tests, was reactive with a non-repetitive epitope on CCA [68], which suggests that the antigen contains a large number of different epitopes.

Over many years, a large number of anti-CAA and anti-CCA McAbs ($n=25$ and $n=55$, respectively) has been generated, the various reaction patterns of which are described in Chapter 3. These McAbs displayed a strong isotype restriction with only IgM, IgG3, and IgG1 McAbs. In all, 60% of the McAbs against CAA were of the IgG1, and 80% of the anti-CCA McAbs were of the IgM isotype. In the mouse, a carbohydrate antigen-driven isotype restriction in the form of IgM and IgG3 is well-known [14,49,54], but anti-protein antibodies (needing T-cell help) are generally IgG1 [2,14,50], which isotype corresponds to the human IgG4 [1,30,44]. In this light, the IgG1 preference for McAbs against the carbohydrate antigen CAA is remarkable.

The major theme of this thesis are the results on the use of the CAA- and CCA-specific McAbs for the purification and subsequent molecular analysis of the antigens themselves. A purification procedure was optimized and immunopurified preparations of CAA and CCA were obtained in sufficient



amounts for molecular characterization. In a collaborative study, it was found that CAA contains a novel and unique polysaccharide structure consisting of multiple repeats of a $\rightarrow 6$ [GlcA- β (1 \rightarrow 3)]GalNAc- β (1 \rightarrow) disaccharide unit [7]. This structure may explain the nearly absolute specificity of the CAA-based assays for the diagnosis of schistosomiasis. CCA-detection, especially in urine, was described to be less specific, necessitating a higher cut-off value in the assays [43,71]. The elucidation of the main carbohydrate structures of this antigen as an *O*-linked polysaccharide having the Lewis x trisaccharide as a repeating unit provided a (partial) explanation for these observations. Glycoproteins or glycolipids carrying a single or multiple Lewis x determinant ($\rightarrow 3$)Gal β (1 \rightarrow 4)[Fuc α (1 \rightarrow 3)]GlcNAc β (1 \rightarrow , Le^x) are frequently found on a variety of human cell-types, including granulocytes [32,60] and carcinoma cells [22,35,59]. Current and preliminary investigations indeed indicate an association between a positive reaction in the urine CCA-ELISA and the presence of leukocytes in the urine of *Schistosoma*-negative individuals (Deelder *et al.*, unpublished results).

Another interesting area of research is the role of these schistosome gut-associated antigens in the host-parasite interaction. It is inconceivable that the parasite would certainly not excrete the antigens in such relatively large amounts in order to facilitate immuno-detection. Since the digestive tract of the parasite is the source of the antigens, it is easy to imagine that they play a role in enzymatic digestive processes [51], but the highly glycosylated nature of the antigens suggests a protective function for the gastrodermis [75]. Indeed, CCA shares some structural characteristics with mucins [17,67], compounds known to provide lubrication and physical protection for epithelial cell surfaces [38,64]. Besides this possible "innate" physical protection, a few specific interactions of gut antigens with the host immune system had been described previously [3,29] and/or were further investigated during the course of the present research.

It has been shown that antigens originating from the schistosome gut caused complement consumption in the absence of specific antibodies, although the activation pathway (classical or alternative) could not be identified [3,70]. Therefore, it could be envisaged, that the antigens play a role in avoiding complement-mediated damage by activation of the complement-system in the lumen rather than directly at the vulnerable surface of the gut. Using immunopurified CAA, and CAA-specific McAbs it could be shown that CAA binds to C1q with characteristics similar to the C1q-C1q-receptor interaction (Chapter 10 and [69]). By acting as a C1q-receptor and binding the C1q present locally, CAA may interfere with the binding of C1q-immune complexes with C1q-receptor-containing cells such as monocytes, neutrophils and platelets [21], thereby preventing activation of these cells. In the same study it was shown that immunopurified CCA binds to isolated C1q-heads, while CAA binds to C1q-stalks (similar to the C1q-receptor). In ELISA's, measuring C3 and/or C4 deposition as a measure for complement activation, the purified antigens did not



activate the complement system, neither by the classical nor by the alternative pathway (unpublished results).

Following characterization of the carbohydrate structure of CAA as a glycosaminoglycan-like polyanion containing GlcA-GalNAc repeating units, it is tempting to speculate on the biological effects (apart from the C1q-interaction described above). A number of polyanions (*e.g.* heparin, chondroitin-sulfate) are known for their anticoagulant activities which probably reside in the glycosaminoglycan-region of the molecule [10-12]. Indeed, in a number of studies anticoagulant activities were found in schistosome antigen preparations and preliminary characterizations of the active component do not exclude CAA as a possible candidate [31,41,65,66]. The mechanism of action, however, remains controversial. One study described the inhibition of the activation of Hageman factor (XII) by schistosome adult worm antigens, without inhibiting the activated factor XII itself [31], while another study showed that the activity of factor XII after activation was specifically inhibited by a similar antigen preparation [65,66]. A third study reported that anticoagulation probably occurred only in the factor XII-independent system, the extrinsic pathway [41]. In addition, Robertson and Cain (1985) isolated a number of glycosaminoglycans from *Schistosoma*, predominantly from the tegumental fraction, suggesting that these components may prevent entrapment of the schistosome by the host's blood-clotting process [57]. However, the observation that galactosaminoglucuronoglycan structures did not influence the major blood clotting parameters may argue against the possible anticoagulation effect of CAA [48].

Finally, Chiang and Caulfield (1989) described that polyanions like suramin, heparin, and dextran sulfate caused displacement of human lipoproteins from the schistosomulum tegument [18,19]. Opposed to the other above described phenomena, this possible effect of the polyanion CAA might prove to be a disadvantage for the parasite, because the surface-bound lipoproteins are thought to mask parasite tegument antigens, which otherwise are recognized by host antibodies [18,19].

The major O-linked polysaccharide chains on CCA are composed of multiple repeating Le^x units (Chapter 7 and [67]). In a pool of glycoproteins isolated from adult schistosome worm-pairs, Srivatsan *et al.* (1992) have recently demonstrated a similar structure with at least four repeating units of the Le^x determinant attached to N-linked carbohydrate chains [62]. Due to the antigen preparation procedure the source of the antigen could not be located, while CCA clearly originates from the gut of the parasite [27]. Besides this, we found that CCA consists of about 80% carbohydrates, and electrophoresis resulted in an antigen smear of 70 - 200 kDa [28,68]. The anti-Le^x McAb used by Srivatsan *et al.* reacted clearly with two bands (one band at 200 kDa and another (broad) band at about 90 kDa) on a Western blot of isolated glycoproteins [62]. In our hands, treatment of CCA with α -fucosidase did not result in a reduction of

McAb-binding (unpublished results), while Srivatsan *et al.*, on the other hand, found a 60% reduction in binding of the antigens to an affinity-column prepared with antibodies from an infected hamster, indicating that only part of their antigens was not susceptible to α -fucosidase. Besides these differences, there are also structural similarities since anti-Le^x McAbs bound to both antigen preparations. It is likely from these results that the schistosome worms are able to synthesize a series of different antigens containing the Le^x or poly-Le^x epitopes, both on *N*-linked and *O*-linked carbohydrate chains.

The finding that the major polysaccharide structures on CCA contained Le^x as a repeating unit produces some interesting hypotheses. Le^x- and, to a much larger extent, sialylated Le^x-containing glycoproteins, play an important role in granulocyte and monocyte adhesion to endothelial cells and platelets [34,45,46,61,63,73,76], thereby recruiting granulocytes to sites of inflammation [13,61]. In this context, it could be hypothesized that the excretion of relatively large amounts of a poly-Le^x containing component by the parasite induces local anti-inflammatory and anti-thrombogenic effects [61,62]. Inflammatory reactions as well as blood coagulation are host protection mechanisms, which potentially are very harmful to the schistosome, and the interference of CCA with these mechanisms may thus be one of the important parasite's survival strategies.

Circulating human granulocytes are rich in Le^x and carry in relatively high abundance branched *N*-linked polysaccharides having Le^x repeating units [60]. Deelder *et al.* (1989) [29] demonstrated that the predominant IgM response in humans, as measured by immunofluorescence assay, against *Schistosoma mansoni* gut-associated antigens was actually directed against CCA. In addition, a mild to moderate neutropenia in chronic schistosomiasis patients has been observed [9], which might be caused by an inhibitory factor in the sera of these patients delaying the maturation of neutrophils in the bone marrow and spleen [9,58]. From these three observations, we proposed the hypothesis that excretion of CCA evokes high titres of anti-poly-Le^x antibodies, which are also directed against identical host carbohydrate structures on *e.g.* neutrophils, thereby causing complement mediated antibody-dependent lysis of these cells. In this context, Ko *et al.* (1990) [42] found that a murine protective IgM McAb, raised against *S. mansoni* eggs, recognized the Le^x determinant (also called SSEA-1, stage-specific embryonic antigen-1), and showed binding to the surface of live schistosomula. In the present study (Chapter 9) it is shown that McAbs, as well as affinity-purified human IgM antibodies directed against CCA, recognized human granulocytes, and, in the presence of complement, caused lysis of the cells.

Recently, another interesting effect of carbohydrate structures similar or identical to CCA (*i.e.* fucosylated (Le^x and Le^y) oligosaccharides) on the host immune system was described by Harn and co-workers [36,72]. These authors showed that Le^x and Le^y carbohydrates (conjugated to human serum albumin) induced



proliferation of human peripheral blood mononuclear cells (including B cells) [36], as well as interleukin 10 production by isolated B cells (B220⁺) of infected but not of non-infected mice [72]. Moreover, they were able to show the presence of antibodies in the cerebrospinal fluid of schistosomiasis patients with cerebral disorders against these fucosylated oligosaccharides [33]. Based on these very new phenomena, the authors suggested that Le^x- and/or Le^y-containing oligosaccharides played a role in the immunoregulation of the helper T cell response in schistosomiasis and maybe other chronic infectious diseases [72]. In addition to the above mentioned highly local (anti-inflammatory and anti-thrombogenic) or indirect (induction of anti-Le^x antibodies) effects, this hypothesis describes a more general effect of excreted CCA.

Appriou *et al.* (1989) described an IgM McAb recognizing a carbohydrate epitope of a schistosome gut-associated antigen [3]. The structural characteristics of this antigen, its localization, and presence in different developmental stages, including eggs, indicated a similarity with CCA [4]. Indeed, comparisons between the 'Appriou-McAb' and the anti-CAA and anti-CCA McAbs, which were performed in our laboratory, confirmed the presence of common epitopes with CCA and not with CAA. Some anti-CCA McAbs, predominantly those which also recognized an egg antigen, showed inhibition of the 'Appriou-McAb' in an immunofluorescence assay (Deelder *et al.*, unpublished results). The 'Appriou-McAb' displayed an inhibitory effect on immunity by passive transfer experiments in mice, both in a secondary and in a primary infection indicating interference with innate immune systems [3]. These investigators hypothesized that inhibition was due to a masking by the antibodies of the complement-activating surface structures on schistosomula. However, IgM deposition on the schistosomulum-surface would also activate the complement system (via the classical route). If the McAb described by Appriou *et al.* would also recognize the main carbohydrate chains of CCA which have a poly-Le^x structure, a weakening of the immune system by complement-dependent lysis of granulocytes, which carry multiple Le^x epitopes on their surface, might also be hypothesized as a possible inhibition mechanism (see also Chs. 7, 9, and above). In contrast to the results of the passive immunization described above, we could not demonstrate an inhibitory effect on immunity, using several different anti-CCA McAbs in passive transfer experiments (unpublished data). Moreover, Ko *et al.* (1990) described a McAb which recognized the Le^x (SSEA-1) epitope but which showed a host-protective effect after passive immunization of mice [42].

The kinetics of CAA and CCA relative to the physiology of the schistosome was investigated by determining the excretion patterns of *in vitro* and *in vivo* developing schistosomula and adult worms (Chapter 11). The observation that the feeding of red blood cells had no effect on the *in vitro* production of the antigens revealed no direct enzymatic role in digestion, and agreed with nature of the antigens themselves (*i.e.* mucin-like molecules). In the growing parasite, CAA

and CCA appeared to be continuously excreted at a low level but showed a steep increase after about 10–15 days which coincides with the rapid development of the gut at that stage [5,15,20]. This supports the potential gut-protective role already indicated above.

In summary, the research described in this thesis focused mainly on two dominant *Schistosoma* gut-associated antigens, CAA and CCA, which are major targets in highly specific and sensitive McAb-based immunodiagnostic assays for schistosomiasis. After purification of the antigens by specific McAbs, the molecular structures of the immunologically dominant carbohydrate parts were elucidated, pointing to distinctive biological functions. Finally, it could thus be shown that CAA and CCA are indeed involved in specifically manipulating parts of the host defense system.

References

1. Aalberse RC, Van der Gaag R, Van Leeuwen J. Serological aspects of IgG4 antibodies. I. Prolonged immunization results in an IgG4-restricted response. *Journal of Immunology* 1983; 32:722–726.
2. Abraham KM, Teale JM. Isotype restriction during infection of mice with the cestode *Mesocestoides corti*: role of immune suppression. *Journal of Immunology* 1987; 138:1699–1704.
3. Appriou M, Ben Younes R, Tribouley-Duret J, Tribouley J. Étude par la réaction d'immunofluorescence des anticorps dirigés contre les antigènes de l'épithélium intestinal de *Schistosoma mansoni*. IV. Étude du pouvoir bloquant sur l'immunité d'un anticorps monoclonal de classe IgM. *Annales de Parasitologie Humaine et Comparée* 1989; 64:456–468.
4. Appriou M, Tribouley-Duret J, Tribouley J. Étude par la réaction d'immunofluorescence des anticorps dirigés contre les antigènes de l'épithélium intestinal de *Schistosoma mansoni*. III. Étude de la réactivité d'un anticorps monoclonal. *Annales de Parasitologie Humaine et Comparée* 1986; 61:435–446.
5. Basch PF. Cultivation of *Schistosoma mansoni* in vitro. I. Establishment of cultures from cercariae and development until pairing. *Journal of Parasitology* 1981; 67:179–185.
6. Berggren WL, Weller TH. Immunoelectrophoretic demonstration of specific circulating antigen in animals infected with *Schistosoma mansoni*. *American Journal of Tropical Medicine and Hygiene* 1967; 16:606–612.
7. Bergwerff AA, Van Dam GJ, Rotmans JP, Deelder AM, Kamerling JP, Vliegthart JFG. The immunologically reactive part of immunopurified circulating anodic antigen from *Schistosoma mansoni* is a threonine-linked polysaccharide consisting of $\rightarrow 6$ -[β -D-Glc pA-(1 \rightarrow 3)]- β -D-GalpNAc-(1 \rightarrow repeating units. *Journal of Biological Chemistry* 1994; in press
8. Bogers JJPM, Nibbeling HAM, Van Marck EAE, Deelder AM. Immunofluorescent visualization of the excretory and gut system of *Schistosoma mansoni* by confocal laser scanning microscopy. *American Journal of Tropical Medicine and Hygiene* 1994; 50:612–619.
9. Borojevic R, Santos da Silva C, Carvalho EA. Chronic schistosomiasis mansoni: splenic myelopoiesis and inhibition of neutrophil granulocytopenesis mediated by the sera of infected patients. *Journal of Infectious Diseases* 1983; 148:422–426.
10. Bourin MC. La Thrombomoduline: un nouveau protéoglycane. Relation structure-fonction. *Annales de Biologie Clinique* 1991; 49: 199–207.
11. Bourin MC, Lindahl U. Glycosaminoglycans and the regulation of blood coagulation. *Biochemical Journal* 1993; 289:313–330.
12. Bourin MC, Öhlin AK, Lane DA, Stenflos J, Lindahl U. Relationship between anticoagulant activities and polyanionic properties of rabbit thrombomodulin. *Journal of Biological Chemistry* 1988; 263:8044–8052.
13. Brandley BK, Swiedler SJ, Robbins PW. Carbohydrate ligands of the LEC cell adhesion molecules. *Cell* 1990; 63:861–863.



14. Briles DE, Claflin JL, Schroer K, Forman C. Mouse IgG3 antibodies are highly protective against infection with *Streptococcus pneumoniae*. *Nature (London)* 1981; 294: 88–89.
15. Brink LH, McLaren DJ, Smithers SR. *Schistosoma mansoni*: a comparative study of artificially transformed schistosomula and schistosomula recovered after cercarial penetration of isolated skin. *Parasitology* 1977; 74:73–86.
16. Carlier Y, Bout D, Bina JC, Camus D, Figueiredo JFM, Capron A. Immunological studies in human schistosomiasis. I. Parasitic antigen in urine. *American Journal of Tropical Medicine and Hygiene* 1975; 24:949–954.
17. Carlier Y, Bout D, Strecker G, Debray H, Capron A. Purification, immunochemical, and biological characterization of the *Schistosoma* circulating M antigen. *Journal of Immunology* 1980; 124:2442–2450.
18. Chiang Ch–P, Caulfield JP. The binding of human low–density lipoproteins to the surface of schistosomula of *Schistosoma mansoni* is inhibited by polyanions and reduces the binding of anti–schistosomal antibodies. *American Journal of Pathology* 1989; 134:1007–1018.
19. Chiang Ch–P, Caulfield JP. Human lipoprotein binding to schistosomula of *Schistosoma mansoni*. Displacement by polyanions, parasite antigen masking, and persistence in young larvae. *American Journal of Pathology* 1989; 135:1015–1024.
20. Clegg JA. In vitro cultivation of *Schistosoma mansoni*. *Experimental Parasitology* 1965; 16:133–147.
21. Cooper NR. The classical complement pathway: activation and regulation of the first complement component. *Advances in Immunology* 1985; 37:151–216.
22. Cordon–Cardo C, Reuter VE, Lloyd KO, Sheinfeld J, Fair WR, Old LJ, Melamed MR. Blood group–related antigens in human urothelium: enhanced expression of precursor, Le–x, and Le–y determinants in urothelial carcinoma. *Cancer Research* 1988; 48:4113–4120.
23. De Jonge N, Kreamsner PG, Krijger FW, Schommer G, Fillié YE, Kornelis D, Van Zeyl RJM, Van Dam GJ, Feldmeier H, Deelder AM. Detection of the schistosome circulating cathodic antigen by enzyme immunoassay using biotinylated monoclonal antibodies. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1990; 84:815–818.
24. De Jonge N, Schommer G, Krijger FW, Feldmeier H, Zwingenberger K, Steiner A, Bienze U, Deelder AM. Presence of circulating anodic antigen in serum of *Schistosoma intercalatum*–infected patients from Gabon. *Acta Tropica (Basel)* 1989; 46:115–120.
25. Deelder AM, De Jonge N, Boerman OC, Fillié YE, Hilberath GW, Rotmans JP, Gerritse MJ, Schut DWOA. Sensitive determination of circulating anodic antigen in *Schistosoma mansoni* infected individuals by an enzyme–linked immunosorbent assay using monoclonal antibodies. *American Journal of Tropical Medicine and Hygiene* 1989; 40:268–272.
26. Deelder AM, Klappe HTM, Van den Aardweg GJM, Van Meerbeke EHEM. *Schistosoma mansoni*: demonstration of two circulating antigens in infected hamsters. *Experimental Parasitology* 1976; 40:189–197.
27. Deelder AM, Kornelis D, Van Marck EAE, Eveleigh PC, Van Egmond JG. *Schistosoma mansoni*: characterization of two circulating polysaccharide antigens and the immunological response to these antigens in mouse, hamster, and human infections. *Experimental Parasitology* 1980; 50:16–32.
28. Deelder AM, Van Dam GJ, Kornelis D, Fillié YE, Van Zeyl RJM. *Schistosoma*: analysis of monoclonal antibodies reactive with the circulating antigens CAA and CCA. *submitted 1994*.
29. Deelder AM, Van Zeyl RJM, Fillié YE, Rotmans JP, Duchenne W. Recognition of gut–associated antigens by immunoglobulin M in the indirect fluorescent antibody test for schistosomiasis mansoni. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1989; 83:364–367.
30. Falconer AE, Friedmann PS, Bird P, Calvert JE. Abnormal immunoglobulin G subclass production in response to keyhole limpet haemocyanin in atopic patients. *Clinical and Experimental Immunology* 1992; 89:495–499.
31. Foster CB, Flanigan TP, Destigter KK, Blanton R, Dumenco LL, Gallagher C, Ratnoff OD. Inhibition of the activation of Hageman factor (factor–XII) by extracts of *Schistosoma mansoni*. *Journal of Laboratory and Clinical Medicine* 1992; 120:735–739.
32. Fox N, Damjanov I, Knowles BB, Solder D. Immunohistochemical localization of the mouse stage–specific embryonic antigen 1 in human tissues and tumors. *Cancer Research* 1983; 43:669–678.
33. Galvao dos Reis M, Secor E, De Souza Andrade Filho A, Cardoso Silva M, Silveira Santos SR, Vasilaski G, Almeida dos Reis E, Velupillai P, Harn DA. Antibodies against defined oligosaccharide in spinal cord fluid: a study among patients with cerebral disorder. In: *Abstract book of the International Symposium on Schistosomiasis*. Instituto Oswaldo Cruz, Rio de Janeiro, 1993:148.
34. Hakomori S–I. LeX and related structures as adhesion molecules. *Histochemical Journal* 1992; 24:771–776.



35. Hakomori S-I, Nudelman E, Lavery SB, Kannagi R. Novel fucolipids accumulating in human adenocarcinoma. I. Glycolipids with di- or trifucosylated type 2 chains. *Journal of Biological Chemistry* 1984; 259:4672-4680.
36. Harn DA, David JR, Galvao dos Reis M, Almeida dos Reis E, Velupillai P. Fucosylated oligosaccharides induce proliferation of PBMC from *Schistosoma mansoni* infected patients. In: *Abstract book of the International Symposium on Schistosomiasis*. Instituto Oswaldo Cruz, Rio de Janeiro, 1993: 147.
37. Hassan MM, Badawi MA, Strand M. Circulating schistosomal antigen in diagnosis and assessment of cure in individuals infected with *Schistosoma mansoni*. *American Journal of Tropical Medicine and Hygiene* 1992; 46:737-744.
38. Hayes DF, Silberstein DS, Rodrigue SW, Kufe DW. DF3 antigen, a human epithelial cell mucin, inhibits adhesion of eosinophils to antibody-coated targets. *Journal of Immunology* 1990; 145:962-970.
39. Hayunga EG, Møllegård I, Duncan Jr JF, Sumner MP, Stek Jr M, Hunter Jr KW. Development of circulating antigen assay for rapid detection of acute schistosomiasis. *Lancet* 1986; 2:716-718.
40. Hayunga EG, Møllegård I, Duncan Jr JF, Sumner MP, Stek Jr M, Hunter Jr KW. Early diagnosis of *Schistosoma mansoni* in mice using assays directed against cercarial antigens isolated by hydrophobic chromatography. *Journal of Parasitology* 1987; 73:351-362.
41. Klufft C, Trumpi-Kalshoven MM, Deelder AM. Factor XII-independent activator generation in human plasma. In: Davidson JF, Cēpelák V, Samama M, Desnoyers PC, eds. *Progress in Chemical Fibrinolysis and Thrombolysis, Vol IV*. Churchill Livingstone, Edinburgh, 1979: 362-367.
42. Ko AI, Dräger UC, Harn DA. A *Schistosoma mansoni* epitope recognized by a protective monoclonal antibody is identical to the stage-specific embryonic antigen 1. *Proceedings of the National Academy of Sciences of the United States of America* 1990; 87:4159-4163.
43. Krijger FW, Van Lieshout L, Deelder AM. A simple technique to pretreat urine and serum samples for quantitation of schistosome circulating anodic and cathodic antigen. *Acta Tropica (Basel)* 1994; 56:55-63.
44. Lal RB, Dhawan RR, Tarrand JJ, Ayoub EM, Ottesen EA. Lack of IgG4 antibody response to carbohydrate antigens in patients with lymphatic filariasis. *Immunology* 1991; 74:333-337.
45. Larsen E, Patabrica T, Sajer S, Gilbert GE, Wagner DD, Furie BC, Furie B. PADGEM-dependent adhesion of platelets to monocytes and neutrophils is mediated by a lineage-specific carbohydrate, LNF III (CD15). *Cell* 1990; 63:467-474.
46. Lowe JB, Stoolman LM, Nair RP, Larsen RD, Berhend TL, Marks RM. ELAM-1-dependent cell adhesion to vascular endothelium determined by a transfected human fucosyltransferase cDNA. *Cell* 1990; 63:475-484.
47. Martin-Prevel Y, Bertheau F, Bouyssou M, Ripert C, Pinder M. An epidemiological study of a *Schistosoma intercalatum* focus in south-east Gabon. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1992; 86:401-405.
48. Matera M, Scalia S. Anticoagulant activity of galactosaminoglucuronoglycan: a pharmacological study. *Drugs Under Experimental and Clinical Research* 1991; 17:65-73.
49. Mazza G, Dunne DW, Butterworth AE. Antibody isotype responses to the *Schistosoma mansoni* schistosomulum in the CBA/N mouse induced by different stages of the parasite life cycle. *Parasite Immunology* 1990; 12:529-543.
50. Moreno C, Esdaine J. Immunoglobulin isotype in the murine response to polysaccharide antigens. *European Journal of Immunology* 1983; 13:262-264.
51. Nash TE. Localization of the circulating antigen within the gut of *Schistosoma mansoni*. *American Journal of Tropical Medicine and Hygiene* 1974; 23:1085-1087.
52. Nour el Din MSA, Kornelis D, Van Zeyl RJM, Deelder AM. Immunologic characterization of two monoclonal antibodies reactive with repetitive carbohydrate epitopes of circulating *Schistosoma mansoni* egg antigen. *American Journal of Tropical Medicine and Hygiene* 1994; 50:487-498.
53. Nour el Din MSA, Nibbeling R, Rotmans JP, Polderman AM, Krijger FW, Deelder AM. Quantitative determination of circulating soluble egg antigen in urine and serum of *Schistosoma mansoni*-infected individuals using a combined two-site enzyme-linked immunosorbent assay. *American Journal of Tropical Medicine and Hygiene* 1994; 50:585-594.
54. Perlmutter RM, Hansburg D, Briles DE, Nicolotti RA, Davie JM. Subclass restriction of murine anti-carbohydrate antibodies. *Journal of Immunology* 1978; 121:566-572.
55. Raccurt CP, Lardillier-Rey D, Appriou M, Tribouley J, Ripert C. Dépistage de la bilharziose intestinale selon la cinétique d'élimination urinaire d'un antigène polysaccharidique. *Cahiers Santé* 1992; 2:91-97.



56. Ripert C, Combe A, Daulouede S, Appriou M, Tribouley-Duret J, Tribouley J, Moyou-Somo R, Same-Ekobo A, Ambassa P. Detection with a monoclonal antibody of an antigen characteristic of the genus *Schistosoma* excreted in the urine. *Tropical Medicine and Parasitology* 1988; 39:131-135.
57. Robertson NP, Cain GD. Isolation and characterization of glycosaminoglycans from *Schistosoma mansoni*. *Comparative Biochemistry and Physiology* 1985; 82B: 299-306.
58. Santos da Silva C, Carvalho EA, Goncalvez MS, Borojevic R. Experimental murine schistosomiasis mansoni: inhibition of neutrophil granulocyte inflammatory reaction. *Brazilian Journal of Medical and Biological Research* 1988; 21:273-279.
59. Sheinfeld J, Reuter VE, Melamed MR, Fair WR, Morse M, Sogani PC, Herr HW, Whitmore WF, Cordon-Cardo C. Enhanced bladder cancer detection with the Lewis-x antigen as a marker of neoplastic transformation. *Journal of Urology* 1990; 143:285-288.
60. Spooncer E, Fukuda M, Klock JC, Oates JE, Dell A. Isolation and characterization of polyfucosylated lactosaminoglycan from human granulocytes. *Journal of Biological Chemistry* 1984; 259:4792-4801.
61. Springer TA, Lasky LA. Sticky sugars for selectins. *Nature (London)* 1991; 349: 196-197.
62. Srivatsan J, Smith DF, Cummings RD. The human blood fluke *Schistosoma mansoni* synthesizes glycoproteins containing the Lewis X antigen. *Journal of Biological Chemistry* 1992; 267:20196-20203.
63. Stocks SC, Kerr MA. Stimulation of neutrophil adhesion by antibodies recognizing CD15 (Lex) and CD15-expressing carcinoembryonic antigen-related glycoprotein NCA-160. *Biochemical Journal* 1992; 288:23-27.
64. Strous GJ, Dekker J. Mucin-type glycoproteins. *Critical Reviews in Biochemistry and Molecular Biology* 1992; 27:57-92.
65. Tsang VCW, Damian RT. Demonstration and mode of action of an inhibitor for activated Hageman factor (factor XIIa) of the intrinsic blood coagulation pathway from *Schistosoma mansoni*. *Blood* 1977; 49:619-633.
66. Tsang VCW, Hubbard WJ, Damian RT. Coagulation factor XIIa (activated Hageman factor) inhibitor from adult *Schistosoma mansoni*. *American Journal of Tropical Medicine and Hygiene* 1977; 26:243-247.
67. Van Dam GJ, Bergwerff AA, Thomas-Oates JE, Rotmans JP, Kamerling JP, Vliegthart JFG, Deelder AM. The immunologically reactive O-linked polysaccharide chains derived from Circulating Cathodic Antigen isolated from the human blood fluke *Schistosoma mansoni* have Lewis x as repeating unit. *European Journal of Biochemistry* 1994; 225:467-482.
68. Van Dam GJ, Cornelis D, Van Zeyl RJM, Rotmans JP, Deelder AM. *Schistosoma mansoni*: analysis of monoclonal antibodies reactive with gut-associated antigens. *Parasitology Research* 1993; 79:55-62.
69. Van Dam GJ, Seino J, Rotmans JP, Daha MR, Deelder AM. *Schistosoma mansoni* circulating anodic antigen but not circulating cathodic antigen interacts with complement component C1q. *European Journal of Immunology* 1993; 23:2807-2812.
70. Van Egmond JG, Deelder AM, Daha MR. *Schistosoma mansoni*: complement activation by antigenic preparations. *Experimental Parasitology* 1981; 51:188-194.
71. Van Lieshout L, De Jonge N, El Masry NA, Mansour MM, Krijger FW, Deelder AM. Improved diagnostic performance of the circulating antigen assay in human schistosomiasis by parallel testing for circulating anodic and cathodic antigens in serum and urine. *American Journal of Tropical Medicine and Hygiene* 1992; 47:463-469.
72. Velupillai P, Harn DA. Oligosaccharide-specific induction of interleukin 10 production by B220+ cells from schistosome-infected mice: a mechanism for regulation of CD4+ T-cell subsets. *Proceedings of the National Academy of Sciences of the United States of America* 1994; 91:18-22.
73. Walz G, Aruffo A, Kolanus W, Bevilacqua M, Seed B. Recognition by ELAM-1 of the sialyl-Le-x determinant on myeloid and tumor cells. *Science* 1990; 250:1132-1135.
74. Weiss JB, Strand M. Characterization of developmentally regulated epitopes of *Schistosoma mansoni* egg glycoprotein antigens. *Journal of Immunology* 1985; 135:1421-1429.
75. Wilson RA, Barnes PE. Synthesis of macromolecules by the epithelial surfaces of *Schistosoma mansoni*: an autoradiographic study. *Parasitology* 1979; 78:295-310.
76. Zhou Q, Moore KL, Smith DF, Varki A, McEver RP, Cummings RD. The selectin GMP-140 binds to sialylated, fucosylated lactosaminoglycans on both myeloid and nonmyeloid cells. *Journal of Cell Biology* 1991; 115:557-564.