

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/43419> holds various files of this Leiden University dissertation.

**Author:** Schaap, M.M.

**Title:** The use of transcriptomics data in detecting non-genotoxic carcinogens

**Issue Date:** 2016-10-04

## Curriculum Vitae

Mirjam Martha Schaap was born on the 7th of Juli 1978 in Zwolle. After graduating from secondary education at the Carolus Clusius College in Zwolle in 1994, she started her post-secondary education at MBO 't Knooppunt in Zwolle to become pharmacy technician. After completion, she studied Biological and Medical Research at the University of Applied Sciences Hogeschool IJsselland in Deventer. During this study, a minor internship project was completed at the Nottingham Trent University, United Kingdom under supervision of P.L.R Bonner. Her major internship project was carried out at the Laboratory of Health Protection Research at the National Institute for Public Health and the Environment (RIVM) under the supervision of Dr. E van Schothorst and Dr. B. Hoebee. In 2001, she graduated and started as research technician at the Center of Health Protection (RIVM). In 2010, she started her PhD study at the Leiden University Medical Center while stationed at the Center of Health Protection at the RIVM under the direct supervision of Prof. Dr. H. van Steeg and Dr. M. Luijten. Since February 2014, Mirjam has been working as scientific advisor at the Biosecurity Office and the Center for Environmental Health (cGM) at the RIVM.



## List of publications

**Schaap MM**, Wackers PF, Zwart EP, van Benthem J, Jacobs MN, Colacci A, Kienhuis AS, van Steeg H, Luijten M. Improving the comparison approach to categorize (non-)genotoxic carcinogens: a case study. *Issues in Toxicology* No. 28, *Toxicogenomics in Predictive Carcinogenicity*, Edited by Michael D. Waters and Russell S. Thomas, The Royal Society of Chemistry 2016

Grinberg M, Stöber RM, Edlund K, Rempel E, Godoy P, Reif R, Widera A, Madjar K, Schmidt-Heck W, Marchan R, Sachinidis A, Spitkovsky D, Hescheler J, Carmo H, Arbo MD, van de Water B, Wink S, Vinken M, Rogiers V, Escher S, Hardy B, Mitic D, Myatt G, Waldmann T, Mardinoglu A, Damm G, Seehofer D, Nüssler A, Weiss TS, Oberemm A, Lampen A, **Schaap MM**, Luijten M, van Steeg H, Thasler WE, Kleinjans JC, Stierum RH, Leist M, Rahnenführer J, Hengstler JG. Toxicogenomics directory of chemically exposed human hepatocytes. *Arch Toxicol.* 2014 Dec;88(12):2261-87.

**Schaap MM**, Wackers PF, Zwart EP, Huijskens I, Jonker MJ, Hendriks G, Breit TM, van Steeg H, van de Water B, Luijten M. A novel toxicogenomics-based approach to categorize (non-)genotoxic carcinogens. *Arch Toxicol.* 2014 89(12):2413-27

Schmeits PC, **Schaap MM**, Luijten M, van Someren E, Boorsma A, van Loveren H, Peijnenburg AA, Hendriksen PJ. Detection of the mechanism of immunotoxicity of cyclosporine A in murine in vitro and in vivo models. *Arch Toxicol.* 2014 89(12):2325-37

Melis JP, Derks KW, Pronk TE, Wackers P, **Schaap MM**, Zwart E, van Ijcken WF, Jonker MJ, Breit TM, Pothof J, van Steeg H, Luijten M. In vivo murine hepatic microRNA and mRNA expression signatures predicting the (non-)genotoxic carcinogenic potential of chemicals. *Arch Toxicol.* 2014 Apr;88(4):1023-34.

van Kesteren PCE, Zwart PE, **Schaap MM**, Pronk TE, van Herwijnen MHM, Kleinjans JCS, Bokkers BGH, Godschalk RWL, Zeilmaker MJ, van Steeg H, Luijten M. Benzo[a]pyrene-induced transcriptomic responses in primary hepatocytes and in vivo liver: Toxicokinetics is essential for in vivo–in vitro comparisons. *Arch Toxicol.* 2013 Mar;87(3):505-15.

Melis JP, Speksnijder EN, Kuiper RV, Salvatori DC, **Schaap MM**, Maas S, Robinson J, Verhoef A, van Benthem J, Luijten M, van Steeg H. Detection of genotoxic and non-genotoxic carcinogens in Xpc(-/-)p53(+/-) mice. *Toxicol Appl Pharmacol*. 2013 Jan 15; 266(2): 289-97.

**Schaap MM**, Zwart EP, Wackers PF, Huijskens I, van de Water B, Breit TM, van Steeg H, Jonker MJ, Luijten M. Dissecting modes of action of non-genotoxic carcinogens in primary mouse hepatocytes. *Arch Toxicol*. 2012 Nov;86(11):1717-27.

**Schaap MM**, Zwart EP, van den Dungen MW, Braakhuis HM, White PA, Steeg Hv, Benthem Jv, Luijten M. Proliferating primary hepatocytes from the pUR288 lacZ plasmid mouse are valuable tools for genotoxicity assessment in vitro. *Environ Mol Mutagen*. 2012 Jun;53(5):376-83.

Mahabir AG, **Schaap MM**, Pennings JL, van Benthem J, Hendriksen CF, van Steeg H. Comparison of clastogen-induced gene expression profiles in wild-type and DNA repair-deficient Rad54/Rad54B cells. *BMC Genomics*. 2010 Jan 12;11:24.

Jonker MJ, Bruning O, van Iterson M, **Schaap MM**, van der Hoeven TV, Vrieling H, Beems RB, de Vries A, van Steeg H, Breit TM, Luijten M. Finding transcriptomics biomarkers for in vivo identification of (non-)genotoxic carcinogens using wild-type and Xpa/p53 mutant mouse models. *Carcinogenesis*. 2009 Oct; 30(10):1805-12.

Mahabir AG, Zwart E, **Schaap M**, van Benthem J, de Vries A, Hernandez LG, Hendriksen CF, van Steeg H. lacZ mouse embryonic fibroblasts detect both clastogens and mutagens. *Mutat Res*. 2009 Jun 18;666(1-2):50-6.

Mahabir AG, **Schaap M**, Theunissen P, van Benthem J, Essers J, de Vries A, Hendriksen CF, van Steeg H. DNA-repair-deficient Rad54/Rad54B mice are more sensitive to clastogens than wild-type mice. *Toxicol Lett*. 2008 Dec 15;183(1-3):112-7

Baken KA, Pennings JL, Jonker MJ, **Schaap MM**, de Vries A, van Steeg H, Breit TM, van Loveren H. Overlapping gene expression profiles of model compounds provide opportunities for immunotoxicity screening. *Toxicol Appl Pharmacol*. 2008 Jan 1;226(1):46-59.

Bruins W, Jonker MJ, Bruning O, Pennings JL, **Schaap MM**, Hoogervorst EM, van Steeg H, Breit TM, de Vries A. Delayed Expression of Apoptotic and Cell Cycle Control Genes in Carcinogen-Exposed Bladders of Mice Lacking p53.S389 Phosphorylation. *Carcinogenesis*. 2007 Aug;28(8):1814-23

Wijnhoven SW, Speksnijder EN, Liu X, Zwart E, vanOostrom CT, Beems RB, Hoogervorst EM, **Schaap MM**, Attardi LD, Jacks T, van Steeg H, Jonkers J, de Vries A. Dominant-negative but not gain-of-function effects of a p53.R270H mutation in mouse epithelium tissue after DNA damage. *Cancer Res*. 2007 May 15;67(10):4648-56.

Kooter IM, Pennings J, Fokkens P, Leseman D, Boere J, Gerlofs M, Cassee F, Schalk JA, Orzechowski T, **Schaap M**, Breit T, Dormans J, van Oostrom C, de Vries A, van Steeg H. Ozone induces clear cellular and molecular responses in the mouse lung independently of the transcription-coupled repair status. *J Appl Physiol*. 2007 Mar;102(3):1185-92

Wijnhoven SW, Zwart E, Speksnijder EN, Beems RB, Olive KP, Tuveson DA, Jonkers J, **Schaap MM**, van den Berg J, Jacks T, van Steeg H, de Vries A. Mice expressing a mammary gland-specific R270H mutation in the p53 tumor suppressor gene mimic human breast cancer development. *Cancer Res*. 2005 Sep 15; 65(18):8166-73.

Van Schothorst EM, Franssen-van Hal N, **Schaap MM**, Pennings J, Hoebee B, Keijer J. Adipose gene expression patterns of weight gain suggest counteracting steroid hormone synthesis. *Obes Res*. 2005 Jun;13(6):1031-41.



