



Universiteit
Leiden
The Netherlands

Celiac disease : from basic insight to therapy development

Stepniak, D.T.

Citation

Stepniak, D. T. (2006, December 14). *Celiac disease : from basic insight to therapy development*. Retrieved from <https://hdl.handle.net/1887/5435>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/5435>

Note: To cite this publication please use the final published version (if applicable).

List of publications

- **Stepniak D**, Wiesner M., de Ru A, Moustakas AK, Papadopoulos GK, van Veelen PA, Koning F. Large-scale characterization of natural ligands provides insight into the unique binding-properties of HLA-DQ2. (submitted).
- **Stepniak D**, Koning F. Enzymatic gluten detoxification: the proof of pudding is in the eating! *Trends in Biotechnol.* 2006 Oct;24(10):433-4.
- **Stepniak D**, Spaenij-Dekking L, Mitea C, Moester M, de Ru A, Baak-Pablo R, van Veelen P, Edens L, Koning F. Highly efficient gluten degradation with a newly identified prolyl endoprotease: implications for celiac disease. *Am J Physiol Gastrointest Liver Physiol.* 2006 Jun;67(6):460-8.
- **Stepniak D**, Koning F. Celiac disease – sandwiched between innate and adaptive immunity. Review. *Hum Immunol* 2006 Jun;67(6):460-8.
- Monsuur A, **Stepniak D**, Diosdado B, Wapenaar MC, Mearin ML, Koning F, Wijmenga C. Genetic and functional analysis of pyroglutamyl-peptidase I in celiac disease. *Eur J Gastroenterol Hepatol.* 2006 Jun;18(6):637-644.
- Diosdado B*, **Stepniak DT***, Monsuur AJ, Franke L, Wapenaar MC, Mearin ML, Koning F, Wijmenga C. No genetic association of the human prolyl endopeptidase gene in the Dutch celiac disease population. *Am J Physiol Gastrointest Liver Physiol.* 2005 Sep;289(3):G495-500.
- **Stepniak D**, Vader LW, Kooy Y, van Veelen PA, Moustakas A, Papandreu NA, Eliopoulos E, Drijfhout JW, Papadopoulos GK, Koning F. T-cell recognition of HLA-DQ2-bound gluten peptides can be influenced by an N-terminal proline at p-1. *Immunogenetics.* 2005 Apr;57(1-2):8-15.
- Vader W, **Stepniak D**, Kooy Y, Mearin L, Thompson A, van Rood JJ, Spaenij L, Koning F. The HLA-DQ2 gene dose effect in celiac disease is directly related to the magnitude and breadth of gluten-specific T cell responses. *Proc Natl Acad Sci U S A.* 2003 Oct 14;100(21):12390-5.
- Vader LW, **Stepniak DT**, Bunnik EM, Kooy YM, de Haan W, Drijfhout JW, Van Veelen PA, Koning F. Characterization of cereal toxicity for celiac disease patients based on protein homology in grains. *Gastroenterology.* 2003 Oct;125(4):1105-13.
- Campo N, Tjalsma H, Buist G, **Stepniak D**, Meijer M, Veenhuis M, Westermann M, Muller JP, Bron S, Kok J, Kuipers OP, Jongbloed JD. Subcellular sites for bacterial protein export. *Mol Microbiol.* 2004 Sep;53(6):1583-99.
- Zimecki M, **Stepniak D**, Szynol A, Kruzel ML. Lactoferrin regulates proliferative response of human peripheral blood mononuclear cells to phytohemagglutinin and mixed lymphocyte reaction. *Arch Immunol Ther Exp.* 2001;49(2):147-54.

* Equal contribution