



Universiteit
Leiden
The Netherlands

Optimization of care in orthopaedics and neurosurgery

Hofstede, S.N.

Citation

Hofstede, S. N. (2016, September 14). *Optimization of care in orthopaedics and neurosurgery*. Retrieved from <https://hdl.handle.net/1887/43013>

Version: Not Applicable (or Unknown)

License:

Downloaded from: <https://hdl.handle.net/1887/43013>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/43013> holds various files of this Leiden University dissertation.

Author: Hofstede, S.N.


Title: Optimization of care in orthopaedics and neurosurgery

Issue Date: 2016-09-14



Part 1

**Implementation of evidence based guidelines of non-surgical
care in osteoarthritis and sciatica care**



Chapter 2

Designing a strategy to implement optimal conservative treatments in patients with knee or hip osteoarthritis in orthopedic practice: a study protocol of the BART-OP study

Stefanie N. Hofstede, Thea P.M. Vliet Vlieland, Cornelia H.M van den Ende, Perla J. Marang-van de Mheen, Rob G.H.H. Nelissen, Leti van Bodegom-Vos

ABSTRACT

Background

National and international evidence-based guidelines for hip and knee osteoarthritis recommend to start with (a combination of) conservative treatments, followed by surgical intervention if a patient does not respond sufficiently to conservative treatment options. Despite these recommendations, there are strong indications that conservative treatments are not optimally used in orthopedic practice. Our study aims to quantify the use of conservative treatments in Dutch orthopedic practice and to explore the barriers and facilitators for the use of conservative treatments that should be taken into account in a strategy to improve the embedding of conservative treatments in hip and knee osteoarthritis in orthopedic practice.

Methods

This study consists of three phases. First, current use of conservative treatments in patients with hip and knee osteoarthritis will be explored using an internet-based survey among at least 100 patients to identify the underused conservative treatments. Second, barriers and facilitators for the use of conservative treatments in orthopedic practice will be identified using semi-structured interviews among 10 orthopedic surgeons and 5 patients. The interviews will be followed by an internet based survey among approximately 450 orthopedic surgeons and at least 100 patients in which the identified barriers and facilitators will be ranked by importance. Finally, an implementation strategy will be developed based on the results of the previous phases using intervention mapping.

Discussion

The developed strategy is likely to result in an optimal and standardized use of conservative treatment options in hip and knee osteoarthritis in orthopedic practice, because it is focused on identified barriers and facilitators. In addition, the results of this study can be used as an example for optimizing the use of conservative care in other patient groups. In a subsequent study, the developed implementation strategy will be assessed on its effectiveness, feasibility and costs.

BACKGROUND

Osteoarthritis (OA) is a degenerative joint disease primarily characterized by progressive loss of articular cartilage. It leads to pain and loss of function [1]. Approximately 10% of men and 18% of women older than 60 years have OA [2]. Symptomatic OA of the knee and the hip have the highest prevalence within the group of arthritis. Due to the ageing society and obesity, the prevalence of hip and knee OA is still increasing [3].

In 2009, 154 patients per 100,000 persons received a Total Hip Arthroplasty (THA) or Total Knee Arthroplasty (TKA), and 118 patients per 100,000 persons received a TKA in Western countries [4]. However, the lifespan of a prosthesis is limited. The revision rate after a TKA or THA is 12.9% after ten years [5], and revision arthroplasty is less successful than primary TKA or THA [6]. Therefore, it is important to delay the primary TKA or THA, by optimizing the use of conservative treatment options, especially in young people.

National and international evidence-based guidelines for hip and knee OA recommend to start with (a combination of) conservative treatments [7-11]. Conservative treatments include pharmacological options, (e.g., the use of analgesics, non-steroidal anti-inflammatory drugs and steroid injection therapy) and non-pharmacological options (e.g., physical therapy, patient education and weight loss interventions). Conservative treatments aim to prevent progression and reduce symptoms such as joint pain and impairment of functions [11]. If persons do not respond (sufficiently) to conservative treatment options, joint replacement (i.e. THA or TKA) can be considered. Despite the recommendation in guidelines to start with conservative treatments and only use surgical intervention if a patient does not respond sufficiently to conservative treatment options, the use of conservative treatments in daily practice is suboptimal [12-15]. For example a study showed that conservative treatments were not fully exploited in 81% of the patients who were referred to specialized knee/hip OA outpatient clinics [12]. Information about conservative treatments patients receive in orthopedic practice is lacking. Furthermore, surgery rates are rising [16]. TKA and THA in patients with OA increased with 196% and 50% respectively between 1995 and 2005 in the Netherlands [16]. In addition, large variation exists in preoperative status (e.g., disease severity) across different centers in Europe and Australia, which suggests differences in the timing of surgery [17,18]. Optimal use of conservative treatments could reduce these differences.

A few models of care were developed to optimize the use of conservative treatments. In Australia, a clinical pathway model and clinician and patient toolkits were developed to support implementation of nonsurgical management of hip and knee OA [19]. However, in Australia, rheumatologists play a leading role, while in the Netherlands the orthopedic surgeon is responsible for OA treatment in hospital care. In the Netherlands,

a stepped-care strategy (SCS), based on (inter)national guidelines [20,21] is developed to facilitate the use of conservative treatments in three steps in primary care [22,23]. The first step consists of education, life style advice, and acetaminophen. If the treatment options in the first step are not sufficient, treatment options in the second step can be considered (exercise therapy, dietary therapy, and non-steroidal anti-inflammatory drugs). Multidisciplinary care, intra-articular injections, and transcutaneous electrical nerve stimulation are treatment options in the third step and could be considered if treatment options in step one or two are ineffective. After implementation of the SCS, most recommended conservative treatments seem to be well used, except dietary therapy [23]. Both studies provide evidence to promote the use of conservative treatments in primary care or in a setting where the rheumatologists play a leading role, but strategies for the optimization of conservative treatments in orthopedic care are still lacking. Information about the current use of conservative treatments, and barriers and facilitators influencing the adoption of conservative treatments in orthopedic practice is needed to develop a tailored implementation strategy focused on orthopedic care.

In the Netherlands, patients with OA are usually treated by the general practitioner. According to guidelines patients should be referred to the orthopedic surgeon if they do not respond sufficiently to conservative treatment options. In orthopedic practice, the decision will be made to start/ continue conservative treatments or to perform a surgery depending on previous received treatments and disease severity. The leading role of an orthopedic surgeon could result in other barriers and facilitators compared to a setting where the rheumatologists play a leading role. This subsequently results in another strategy to improve the embedding of conservative therapies in hip and knee OA in orthopedic practice. While rheumatologists and general practitioners only provide conservative treatments in OA, orthopedic surgeons can provide both conservative treatments and surgical interventions. It is unclear to what extent factors such as lack of information about conservative treatment options, increasing number of orthopedic surgeons [16], or patient preferences play a role. It is important to explore these factors for the development of a tailored implementation strategy, so that orthopedic surgeons will provide underused treatment options in primary care, such as dietary therapy. Part of this implementation strategy could be the SCS or a clinical pathway model as used in previous implementations.

Objective

The BART-OP study (Beating osteoARthritis in the Orthopedic Practice) aims to quantify the use of conservative treatments in Dutch orthopedic practice before THA or TKA and to explore the barriers and facilitators for the use of conservative treatments that should be taken into account in a strategy to improve the embedding of conservative treatments in hip and knee OA in orthopedic practice.

To reach the aim of this study, we formulated the following research questions:

1. What is the current use of conservative treatments, before patients receive a surgery, in orthopedic practice?
2. Which barriers and facilitators influence the use of conservative treatments in orthopedic practice?
3. What is an appropriate tailored implementation strategy for the embedding of conservative treatments in orthopedic practice?

In a subsequent study, the developed implementation strategy will be assessed on its effectiveness, feasibility and costs.

METHODS

This study consists of three phases to be executed in one year:

- A. The analysis of current use of conservative treatments, before patients receive a surgery in orthopedic practice (months 1 to 9).
- B. Identification of barriers and facilitators for non-optimal conservative treatments, using two steps (months 1 to 9).
 - i. Barriers and facilitators for non-optimal conservative treatments are explored with interviews among orthopedic surgeons and patients.
 - ii. Identified barriers and facilitators are ranked by importance in a survey among a representative sample of orthopedic surgeons and patients.
- C. The development of the implementation strategy based on the results of phases A and B (months 9 to 12).

The study design, study population, analysis and outcome measures are described per study phase.

Phase A. The analysis of current use of conservative treatments before patients receive a surgery in orthopedic practice

Study design

To analyze the current use of conservative treatments, before patients undergo THA or TKA in orthopedic practice, an internet-based survey among patients will be performed. The survey will include questions about which conservative treatment options are used before surgery. This information is needed to be able to focus the implementation strategy on the right conservative treatments. The content of the survey will be developed based on the Dutch guideline of OA of the hip and knee [11]. Reminders to non-responders will be sent after two weeks and again after four weeks.

Study population

The survey will be sent to a sample of at least 100 patients living in different regions of the Netherlands. Inclusion criteria for patients are: age ≥ 18 years, a doctor's diagnosis of hip or knee OA, and who have had a TKA or THA no longer than 12 months ago or are on the waiting list for surgery within the next 3 months. Patients with an inability to understand written Dutch will be excluded from the study. We will sample these patients using advertisements in local newspapers, and at websites or newsletters of patient associations.

Analysis

Descriptive statistics will be used to describe the current use of conservative treatment options in orthopedic practice. Independent t-tests or Mann Whitney U tests for continuous variables and Chi square tests or Fisher's exact tests for proportions are used to analyze differences in the frequency of use between different regions or other conditions.

Outcome measures

The main outcome measure is the percentage of patients in whom the conservative treatment options are applied optimally before they undergo surgery, as described in the guideline. These results will help us to focus the implementation strategy, developed in phase C, on the right conservative treatments.

Phase B. Identification of barriers and facilitators for non-optimal treatment

Study design

Two steps will be taken to identify barriers and facilitators associated with the non-optimal use of conservative treatments. First, semi-structured interviews among orthopedic surgeons and patients will be performed to explore all relevant barriers and facilitators for non-optimal conservative therapy. The interview questions will be based on the Theoretical Domains Interview framework (TDI) [24]. The TDI framework includes 12 theoretical construct domains derived from 33 psychological theories, and covering 128 explanatory constructs that enhance implementation of evidence-based practice [24]. In addition, barriers and facilitators reported in a previous study about the use of the SCS to optimize hip and knee OA in primary care [25] are included in the interview questions. Second, an internet-based survey will be held among a selection of orthopedic surgeons ($n \approx 400$) and sample of patients ($n \geq 100$) to rank barriers and facilitators identified in the interviews on importance. The survey will include questions to determine which of these barriers and facilitators are associated with the use of conservative treatments.

Study population

For the semi-structured interviews, we anticipate interviewing 10 orthopedic surgeons involved in hip and knee surgery and 5 patients who have had a THA or TKA no longer than 12 months ago (≥ 18 years, and able to understand oral Dutch). If we do not reach data saturation after these interviews (three consecutive interviews without new barriers or facilitators [26]), we will continue interviewing until data saturation is reached. To obtain contrasting views on barriers and facilitators, we will apply purposive sampling. First, we will purposively select orthopedic surgeons and patients from Dutch regions with high surgery rates and from Dutch regions with relatively low surgery rates based on the report of Van Beek et al. (2010) about variation in clinical practice [27]. In addition, we will select orthopedic surgeons in such a way as to ensure diversity of hospital type (public hospitals and academic hospitals). It is important to include orthopedic surgeons of public and academic hospitals, because this may reveal other facilitators and barriers. For the internet based survey, Dutch orthopedic surgeons listed in the registry of the Dutch Orthopedic Association (NOV) or the Dutch medical address book will be approached for participation. Inclusion criteria are: involved in hip or knee OA, and access to email address. Patients ($n \geq 100$) are recruited using advertisements in local newspapers. Included are patients: ≥ 18 years who have had total hip or knee surgery no longer than 12 months ago, or are on the waiting list for receiving a THA or TKA. Patients with an inability to understand oral Dutch will be excluded from the study.

Analysis

The semi-structured interviews will be audio-taped and transcribed in full for analysis. The interviews will be analyzed by two researchers using open coding to ensure that we find all barriers and facilitators for the non-optimal use of conservative therapy. This qualitative analysis will be executed using the software package ATLAS.ti (ATLAS.ti Scientific Software Development GmbH, Berlin, Germany) for this qualitative analysis.

The subsequent survey data will allow us to rank the importance of barriers and facilitators and their relationship with the use of conservative treatments. These relationships will be assessed using multiple regression analysis. We will use SPSS 20.0 for analysis.

Outcome measures

A list of the most relevant barriers and facilitators for the optimal use of conservative treatments in orthopedic practice before patients with hip or knee OA receive THA or TKA.

Phase C. The development of the implementation strategy

Study design

The results of the previous phases will be used to develop a tailored implementation strategy for the optimal use of conservative treatments in orthopedic practice in patients with hip or knee OA. The results of phase A will show at which type(s) of conservative treatment the strategy should be aimed. Phase B results will show the most relevant barriers and facilitators that should be taken into account in the development of the strategy. From literature, it is known that, in general, multifaceted strategies are more effective than single strategies [28,29]. Assuming this, and our expectation that several barriers on different theoretical domains will be found, it is very likely that the developed implementation strategy includes several components directed at different levels (i.e. knowledge or social influences). Furthermore, it is expected that the strategy components will include educational outreach, an interactive educational strategy, and/or patient-specific strategies, because these facets seem to be promising for implementation [28].

In the development process, the project team will use the intervention mapping approach of Bartholomew et al. [30]. This method begins with the creation of matrices, in which the performance objectives are set against the most important factors that hinder or facilitate the adoption of conservative treatments. Subsequently, the project team will brainstorm about the strategy components needed to achieve the performance objective in the presence of the barrier or facilitator mentioned in the matrix. The cells of the matrices are then gradually filled with implementation strategy components [30]. Next, the project team will translate the formulated strategy components into practical strategies.

Analysis

The study group meeting will be summarized. The project members will receive a summary of the meeting and the formulated implementation strategy and will be asked whether the summary and implementation strategy is consistent with the conclusions reached in the meeting.

Outcome measures

A tailored implementation strategy for the embedding of conservative treatments in orthopedic practice in patients with hip or knee OA.

Ethical approval

This study protocol was presented to the Medical Ethical Committee of the Leiden University Medical Center (CME P13.087/NV/nv). An exemption was obtained, as ethical approval for this type of study is not required under Dutch law.

DISCUSSION

The goal of this study is to develop a tailored implementation strategy to optimize the use of conservative treatments in hip and knee OA in patients referred to the orthopedic surgeon.

Several studies have been performed to develop and test implementation strategies, including identification of barriers that prevent implementation [31-33]. They all conclude that a prior inventory of barriers to develop a tailored implementation strategy is useful and can confirm whether barriers differ in different settings. Prior inventory thereby reduces the number of costly trials evaluating different implementation strategies [28,32,34]. Although previous studies already explored barriers for the use of conservative treatments, these studies were performed in other settings, and not focused on orthopedic care. Furthermore, the uptake of several implementation activities was poor, since only 9% of the participating GPs were present at the seminar [23]. It was very difficult to reach all GPs in seminars. This could be easier in orthopedic practice. Orthopedic surgeons may have more interest in OA, because it is part of their specialization whereas for GPs it is one of the many health problems in their daily practice. This highlights the importance of optimizing the use of conservative treatments in orthopedic practice as well, so that patients will receive optimal treatment options in orthopedic practice if conservative care was suboptimal in their primary care trajectory. Our study and the study performed in primary care together will provide useful information for the development of interventions based on the full spectrum of barriers and facilitators in primary care and orthopedic practice. This is important because a multidisciplinary approach is likely to be more effective to obtain optimal conservative therapy [35].

A strength of this study is the purposive sampling of orthopedic surgeons of regions with low and high surgery rates, because they could have contrasting views on barriers and facilitators. We think that this will reveal most barriers and facilitators for the implementation of the optimal use of conservative treatments in hip and knee OA in orthopedic practice. A limitation may be the selection of patients. Patients will be recruited via advertisement, which can lead to selection bias, because patients who respond to the advertisements may perceive other barriers and facilitators as most important compared to non-responders. Furthermore, the use of an internet-based survey could also induce selection bias. Knee and hip OA increases with age [3], but not all elderly persons do have internet or an email address. This can lead to the selection of younger persons compared to the average age of OA patients, while elderly persons may perceive other barriers and facilitators as most important. We will assess the impact of selection bias by comparing elderly respondents with younger ones. If they perceive the same barriers and facilitators we can conclude that the impact of this type of selection bias does not influence our results.

The developed strategy is likely to result in an optimal and standardized use of conservative treatment options in hip and knee OA in orthopedic practice. In addition, the results of this study can be used as an example for optimizing the use of conservative care in other patient groups. In a subsequent study, the developed implementation strategy will be assessed on its effectiveness, feasibility and costs.

REFERENCES

1. Woolf AD, Pfleger B: Burden of major musculoskeletal conditions. *Bull World Health Organ* 2003, 81: 646-656.
2. World Health Organization. Chronic diseases and health promotion: Chronic rheumatic conditions. Available at: <http://www.who.int/chp/topics/rheumatic/en/>. Accessed 5-11-2013.
3. Pereira D, Peleteiro B, Araujo J, Branco J, Santos RA, Ramos E: The effect of osteoarthritis definition on prevalence and incidence estimates: a systematic review. *Osteoarthritis Cartilage* 2011, 19: 1270-1285.
4. OECD: Hip and knee replacement. In *Health at a Glance 2011: OECD Indicators*. OECD Publishing; 2011.
5. Labeck G, Thaler M, Janda W, Agreiter M, Stockl B: Revision rates after total joint replacement: cumulative results from worldwide joint register datasets. *J Bone Joint Surg Br* 2011, 93: 293-297.
6. Bennell KL, Hunter DJ, Hinman RS: Management of osteoarthritis of the knee. *BMJ* 2012, 345: e4934.
7. Zhang W, Nuki G, Moskowitz RW, Abramson S, Altman RD, Arden NK et al.: OARSI recommendations for the management of hip and knee osteoarthritis: part III: Changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage* 2010, 18: 476-499.
8. Hochberg MC, Altman RD, April KT, Benkhalti M, Guyatt G, McGowan J et al.: American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)* 2012, 64: 465-474.
9. Conaghan PG, Dickson J, Grant RL: Care and management of osteoarthritis in adults: summary of NICE guidance. *BMJ* 2008, 336: 502-503.
10. Fernandes L, Hagen KB, Bijlsma JW, Andreassen O, Christensen P, Conaghan PG et al.: EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis. *Ann Rheum Dis* 2013, 72: 1125-1135.
11. Netherlands Orthopaedic Association (NOV). Guideline diagnostics and treatment of hip- and knee osteoarthritis. Utrecht: Quality Institute for Health Care (CBO); 2007.
12. Snijders GF, den Broeder AA, van Riel PL, Straten VH, de Man FH, van den Hoogen FH et al.: Evidence-based tailored conservative treatment of knee and hip osteoarthritis: between knowing and doing. *Scand J Rheumatol* 2011, 40: 225-231.
13. Shrier I, Feldman DE, Gaudet MC, Rossignol M, Zukor D, Tanzer M et al.: Conservative non-pharmacological treatment options are not frequently used in the management of hip osteoarthritis. *J Sci Med Sport* 2006, 9: 81-86.
14. De Haan MN, Guzman J, Bayley MT, Bell MJ: Knee osteoarthritis clinical practice guidelines -- how are we doing? *J Rheumatol* 2007, 34: 2099-2105.
15. Hunter DJ: Quality of osteoarthritis care for community-dwelling older adults. *Clin Geriatr Med* 2010, 26: 401-417.
16. Otten R, van Roermund PM, Picavet HS: Trends in the number of knee and hip arthroplasties: considerably more knee and hip prostheses due to osteoarthritis in 2030 (in Dutch). *Ned Tijdschr Geneesk* 2010, 154: A1534.
17. Dieppe P, Judge A, Williams S, Ikwueke I, Guenther KP, Floeren M et al.: Variations in the pre-operative status of patients coming to primary hip replacement for osteoarthritis in European orthopaedic centres. *BMC Musculoskelet Disord* 2009, 10: 19.
18. Ackerman IN, Dieppe PA, March LM, Roos EM, Nilsson AK, Brown GC et al.: Variation in age and physical status prior to total knee and hip replacement surgery: a comparison of centers in Australia and Europe. *Arthritis Rheum* 2009, 61: 166-173.
19. Brand C, Cox S: Systems for implementing best practice for a chronic disease: management of osteoarthritis of the hip and knee. *Intern Med J* 2006, 36: 170-179.
20. Smink AJ, van den Ende CH, Vliet Vlieland TP, Swierstra BA, Kortland JH, Bijlsma JW et al.: "Beating osteoARThritis": development of a stepped care strategy to optimize utilization and timing of non-surgical treatment modalities for patients with hip or knee osteoarthritis. *Clin Rheumatol* 2011, 30: 1623-1629.
21. van den Ende CM, Bierma-Zeinstra SM, Vliet Vlieland TP, Swierstra BA, Voorn TB, Dekker J: Conservative treatment of hip and knee osteoarthritis: a step-by-step treatment strategy (in Dutch). *Ned Tijdschr Geneesk* 2010, 154:A1574.
22. Smink AJ, Bierma-Zeinstra SM, Dekker J, Vliet Vlieland TP, Bijlsma JW, Swierstra BA et al.: Agreement of general practitioners with the guideline-based stepped-care strategy for patients with osteoarthritis of the hip or knee: a cross-sectional study. *BMC Fam Pract* 2013, 14: 33.
23. Smink AJ, Dekker J, Vliet Vlieland TP, Swierstra BA, Kortland JH, Bijlsma JW et al.: Health care use of patients with osteoarthritis of the hip or knee after implementation of a stepped care strategy: an observational study. *Arthritis Care Res (Hoboken)* 2013, 65.
24. Michie S, Johnston M, Abraham C, Lawton R,

- Parker D, Walker A: Making psychological theory useful for implementing evidence based practice: a consensus approach. *Qual Saf Health Care* 2005, 14: 26-33.
25. Cuperus N, Smink AJ, Bierma-Zeinstra SM, Dekker J, Schers HJ, de Boer F et al.. Patient reported barriers and facilitators to use a self-management booklet for hip and knee osteoarthritis in primary care: Results of a qualitative interview study. *BMC Family Practice* 2013, 14:181.
 26. Francis JJ, Johnston M, Robertson C, Glidewell L, Entwistle V, Eccles MP et al.: What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychol Health* 2010, 25: 1229-1245.
 27. Van Beek E, Lemmers K, Van Schooten G, Vlieger E. Reduction of variation in clinical practice: budgetary effects of more precise indications (in Dutch). *PLEXUS* 2010.
 28. Prior M, Guerin M, Grimmer-Somers K: The effectiveness of clinical guideline implementation strategies-a synthesis of systematic review findings. *J Eval Clin Pract* 2008, 14: 888-897.
 29. Legare F, Turcotte S, Stacey D, Ratto S, Kryworuchko J, Graham ID: Patients' perceptions of sharing in decisions: a systematic review of interventions to enhance shared decision making in routine clinical practice. *Patient* 2012, 5: 1-19.
 30. Bartholomew L, Parcel G, Kok G, Gottlieb N: *Planning health promotion programs: an intervention mapping approach*. New York: John Wiley & Sons Inc; 2006.
 31. Grol R: Personal paper. Beliefs and evidence in changing clinical practice. *BMJ* 1997, 315: 418-421.
 32. Grol R, Grimshaw J: From best evidence to best practice: effective implementation of change in patients' care. *Lancet* 2003, 362: 1225-1230.
 33. Baker R, Camosso-Stefinovic J, Gillies C, Shaw EJ, Cheater F, Flottorp S et al.: Tailored interventions to overcome identified barriers to change: effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* 2010, 3: CD005470.
 34. Grimshaw JM, Thomas RE, MacLennan G, Fraser C, Ramsay CR, Vale L et al.: Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health Technol Assess* 2004, 8: iii-72.
 35. Mazieres B, Scmidely N, Hauselmann HJ, Martin-Mola E, Serni U, Verbruggen AA et al.: Level of acceptability of EULAR recommendations for the management of knee osteoarthritis by practitioners in different European countries. *Ann Rheum Dis* 2005, 64: 1158-1164.

