



Universiteit
Leiden
The Netherlands

Topologies and convergence structures on vector lattices of operators

Deng, Y.

Citation

Deng, Y. (2021, February 2). *Topologies and convergence structures on vector lattices of operators*. Retrieved from <https://hdl.handle.net/1887/3135024>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3135024>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3135024> holds various files of this Leiden University dissertation.

Author: Deng, Y.

Title: Topologies and convergence structures on vector lattices of operators

Issue Date: 2021-02-02

Chapter 3

Convergence structures and locally solid topologies on vector lattices of operators

Abstract

For vector lattices E and F , where F is Dedekind complete and supplied with a locally solid topology, we introduce the corresponding locally solid absolute strong operator topology on the order bounded operators $\mathcal{L}_{\text{ob}}(E, F)$ from E into F . Using this, it follows that $\mathcal{L}_{\text{ob}}(E, F)$ admits a Hausdorff uo-Lebesgue topology whenever F does.

For each of order convergence, unbounded order convergence, and—when applicable—convergence in the Hausdorff uo-Lebesgue topology, there are both a uniform and a strong convergence structure on $\mathcal{L}_{\text{ob}}(E, F)$. Of the six conceivable inclusions within these three pairs, only one is generally valid. On the orthomorphisms of a Dedekind complete vector lattice, however, five are generally valid, and the sixth is valid for order bounded nets. The latter condition is redundant in the case of sequences of orthomorphisms on a Banach lattice, as a consequence of a uniform order boundedness principle for orthomorphisms that we establish.

We also show that, in contrast to general order bounded operators, the orthomorphisms preserve not only order convergence of nets, but unbounded order convergence and—when applicable—convergence in the Hausdorff uo-Lebesgue topology as well.

3.1 Introduction and overview

Let X be a non-empty set. A *convergence structure* on X is a non-empty collection \mathcal{C} of pairs $((x_\alpha)_{\alpha \in A}, x)$, where $(x_\alpha)_{\alpha \in A}$ is a net in X and $x \in X$, such that:

- (1) when $((x_\alpha)_{\alpha \in A}, x) \in \mathcal{C}$, then also $((y_\beta)_{\beta \in B}, x) \in \mathcal{C}$ for every subnet $(y_\beta)_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$;
- (2) when a net $(x_\alpha)_{\alpha \in A}$ in X is constant with value x , then $((x_\alpha)_{\alpha \in A}, x) \in \mathcal{C}$.

One can easily vary on this definition. For example, one can allow only sequences. There does not appear to be a consensus in the literature about the notion of a convergence structure; [8] uses filters, for example. Ours is sufficient for our merely descriptive purposes, and close in spirit to what may be the first occurrence of such a definition in [22] for sequences. Although we shall not pursue this in the present paper, let us still mention that the inclusion of the subnet criterion in the definition makes it possible to introduce an associated topology on X in a natural way. Indeed, define a subset S of X to be \mathcal{C} -closed when $x \in S$ for all pairs $((x_\alpha)_{\alpha \in A}, x) \in \mathcal{C}$ such that $(x_\alpha)_{\alpha \in A} \subseteq S$. Then the collection of the complements of the \mathcal{C} -closed subsets of X is a topology on X .

The convergent nets in a topological space, together with their limits, are the archetypical example of a convergence structure. In the context of vector lattices, there are other ones that are rarely of a topological nature. For example, the order convergence nets with their order limits form a convergence structure, and likewise there is a convergence structure for unbounded order convergence. Taken together with the (topological) structure for convergence in a Hausdorff uo-Lebesgue topology, when this exists, there are three natural and related convergence structures on a vector lattice to consider.

Suppose that E and F are vector lattices, where F is Dedekind complete. The above then yields three convergence structures on the vector lattice $\mathcal{L}_{\text{ob}}(E, F)$ of order bounded operators from E into F , but there are also three others that are derived from those in F . For example, one can consider all pairs $((T_\alpha)_{\alpha \in A}, T)$, where $(T_\alpha)_{\alpha \in A}$ is a net in $\mathcal{L}_{\text{ob}}(E, F)$ and $T \in \mathcal{L}_{\text{ob}}(E)$, such that $(T_\alpha x)_{\alpha \in A}$ is order convergent to Tx in F for all $x \in E$. These pairs also form a convergence structure on $\mathcal{L}_{\text{ob}}(E, F)$. Likewise, the pointwise unbounded order convergence in F and—when applicable—the pointwise convergence in a Hausdorff uo-Lebesgue topology on F yield convergence structures on $\mathcal{L}_{\text{ob}}(E, F)$. Motivated by the terminology for operators between Banach spaces, we shall speak of *uniform* and *strong* convergence structures on $\mathcal{L}_{\text{ob}}(E)$ —with the obvious meanings.

The present paper is primarily concerned with the possible inclusions between the uniform and strong convergence structure for each of order convergence, unbounded order convergence, and—when applicable—convergence in a Hausdorff uo-Lebesgue topology. We consider these inclusions for $\mathcal{L}_{\text{ob}}(E, F)$, but also for the orthomorphisms $\text{Orth}(E)$ on a Dedekind complete vector lattice. This special interest in $\text{Orth}(E)$ stems from representation theory. When a group acts as order automorphisms on a Dedekind complete vector lattice E , then the Boolean lattice of all invariant bands in E can be retrieved from the commutant of the group action in $\text{Orth}(E)$. This commutant, therefore, plays the role of the von Neumann algebra which is the commutant of a unitary action of a group on a Hilbert space. It has been known long since that more than one topology on a von Neumann algebras is needed to understand it and its role in representation theory on Hilbert spaces, and the same holds

true for the convergence structures as related to these commutants in an ordered context. Using these convergence structures, it is, for example, possible to obtain ordered versions of von Neumann's bicommutant theorem. We shall report separately on this. Apart from its intrinsic interest, the material on $\text{Orth}(E)$ in the present paper is an ingredient for these next steps.

This paper is organised as follows.

Section 3.2 contains the basic notations, definitions, conventions, and references to earlier results.

In Section 3.3, we show how, given a vector lattice E , a Dedekind complete vector lattice F , and a (not necessarily Hausdorff) locally solid linear topology τ_F on F , a locally solid linear topology can be introduced on $\mathcal{L}_{\text{ob}}(E, F)$ that deserves to be called the absolute strong operator topology that is generated by τ_F . This is a preparation for Section 3.4, where we show that regular vector sublattices of $\mathcal{L}_{\text{ob}}(E, F)$ admit a Hausdorff uo-Lebesgue topology when F admits one.

For each of order convergence, unbounded order convergence, and—when applicable—convergence in a Hausdorff uo-Lebesgue topology, there are two conceivable implications between uniform and strong convergence of a net of order bounded operators. In Section 3.5, we show that only one of these six is generally valid. Section 3.9 will make it clear that the five failures are, perhaps, not as ‘only to be expected’ as one might think at first sight.

In Section 3.6, we review some material concerning orthomorphism and establish a few auxiliary result for use in the present paper and in future ones. It is shown here that a Dedekind complete vector lattice and its orthomorphisms have the same universal completion.

Section 3.7 briefly digresses from the main line of the paper. It is shown that orthomorphisms preserve not only the order convergence of nets, but also the unbounded order convergence and—when applicable—the convergence in a Hausdorff uo-Lebesgue topology. None of this is true for arbitrary order bounded operators.

In Section 3.8, we return to the main line, and we specialise the results in Sections 3.3 and 3.4 to the orthomorphisms. When restricted to $\text{Orth}(E)$, the absolute strong operator topologies from Section 3.3 are simply strong operator topologies.

Section 3.9 on orthomorphisms is the companion of Section 3.5, but the results are quite in contrast. For each of order convergence, unbounded order convergence, and—when applicable—convergence in a Hausdorff uo-Lebesgue topology, both implications between uniform and strong convergence of a net of orthomorphisms *are* valid, with an order boundedness condition on the net being necessary only for order convergence. For sequences of orthomorphisms on Banach lattices, this order boundedness condition is redundant as a consequence of a uniform order boundedness principle for orthomorphisms that is also established in this section.

3.2 Preliminaries

In this section, we collect a number of definitions, notations, conventions and earlier results.

All vector spaces are over the real numbers; all vector lattices are supposed to be Archimedean. We write E^+ for the positive cone of a vector lattice E . For a non-empty subset S of E , we let I_S and B_S denote the ideal of E and the band in E , respectively, that are generated by S ; we write S^\vee for $\{s_1 \vee \cdots \vee s_n : s_1, \dots, s_n \in S\}$.

Let E be a vector lattice, and let $x \in E$. We say that a net $(x_\alpha)_{\alpha \in \mathcal{A}}$ in E is *order convergent* to $x \in E$ (denoted by $x_\alpha \xrightarrow{o} x$) when there exists a net $(y_\beta)_{\beta \in \mathcal{B}}$ in E such that $y_\beta \downarrow 0$ and with the property that, for every $\beta_0 \in \mathcal{B}$, there exists an $\alpha_0 \in \mathcal{A}$ such that $|x - x_\alpha| \leq y_{\beta_0}$ whenever α in \mathcal{A} is such that $\alpha \geq \alpha_0$. We explicitly include this definition to make clear that the index sets \mathcal{A} and \mathcal{B} need not be equal.

Let $(x_\alpha)_{\alpha \in \mathcal{A}}$ be a net in a vector lattice E , and let $x \in E$. We say that (x_α) is *unbounded order convergent* to x in E (denoted by $x_\alpha \xrightarrow{uo} x$) when $|x_\alpha - x| \wedge y \xrightarrow{o} 0$ in E for all $y \in E^+$. Order convergence implies unbounded order convergence to the same limit. For order bounded nets, the two notions coincide.

Let E and F be vector lattices. The order bounded operators from E into F will be denoted by $\mathcal{L}_{\text{ob}}(E, F)$. We write E^\sim for $\mathcal{L}_{\text{ob}}(E, \mathbb{R})$. A linear operator $T : E \rightarrow F$ between two vector lattices E and F is *order continuous* when, for every net $(x_\alpha)_{\alpha \in \mathcal{A}}$ in E , the fact that $x_\alpha \xrightarrow{o} 0$ in E implies that $Tx_\alpha \xrightarrow{o} 0$ in F . An order continuous linear operator between two vector lattices is automatically order bounded; see [7, Lemma 1.54], for example. The order continuous linear operators from E into F will be denoted by $\mathcal{L}_{\text{oc}}(E, F)$. We write E_{oc}^\sim for $\mathcal{L}_{\text{oc}}(E, \mathbb{R})$.

Let F be a vector sublattice of a vector lattice E . Then F is a *regular vector sublattice* of E when the inclusion map from F into E is order continuous. Ideals are regular vector sublattices. For a net in a regular vector sublattice F of E , its uo -convergence in F and in E are equivalent; see [28, Theorem 3.2].

When E is a vector space, a *linear topology* on E is a (not necessarily Hausdorff) topology that provides E with the structure of a topological vector space. When E is a vector lattice, a *locally solid linear topology* on E is a linear topology on E such that there exists a base of (not necessarily open) neighbourhoods of 0 that are solid subsets of E . For the general theory of locally solid linear topologies on vector lattices we refer to [6]. When E is a vector lattice, a *locally solid additive topology* on E is a topology that provides the additive group E with the structure of a (not necessarily Hausdorff) topological group, such that there exists a base of (not necessarily open) neighbourhoods of 0 that are solid subsets of E .

A topology τ on a vector lattice E is an *o-Lebesgue topology* when it is a (not necessarily Hausdorff) locally solid linear topology on E such that, for a net $(x_\alpha)_{\alpha \in \mathcal{A}}$ in E , the fact that $x_\alpha \xrightarrow{o} 0$ in E implies that $x_\alpha \xrightarrow{\tau} 0$. A vector lattice need not admit a Hausdorff o-Lebesgue topology. A topology τ on a vector lattice E is a *uo-Lebesgue topology* when it is a (not necessarily Hausdorff) locally solid linear topology on E such that, for a net $(x_\alpha)_{\alpha \in \mathcal{A}}$ in E , the fact that $x_\alpha \xrightarrow{uo} 0$ in E implies that $x_\alpha \xrightarrow{\tau} 0$. Since order convergence implies unbounded order convergence, a uo -Lebesgue topology is an o -Lebesgue topology. A vector lattice E need not admit a Hausdorff uo -Lebesgue topology, but when it does, then this topology is unique (see [11, Propositions 3.2, 3.4, and 6.2] or [44, Theorems 5.5 and 5.9]) and we denote it by $\widehat{\tau}_E$.

Let E be a vector lattice, let F be an ideal of E , and suppose that τ_F is a (not necessarily Hausdorff) locally solid linear topology on F . Take a non-empty subset S of F . Then there exists a unique (possibly non-Hausdorff) locally solid linear topology $u_S\tau_F$ on E such that, for a net $(x_\alpha)_{\alpha \in A}$ in E , $x_\alpha \xrightarrow{u_S\tau_F} 0$ if and only if $|x_\alpha| \wedge |s| \xrightarrow{\tau_F} 0$ for all $s \in S$; see [20, Theorem 3.1] for this, which extends earlier results in this vein in, e.g., [11] and [44]. This topology $u_S\tau_F$ is called the unbounded topology on E that is generated by τ_F via S . Suppose that E admits a Hausdorff uo-Lebesgue topology $\widehat{\tau}_E$. The uniqueness of such a topology then implies that $u_E\widehat{\tau}_E = \widehat{\tau}_E$. In the sequel we shall use this result from [11] and [44] a few times.

Finally, the characteristic function of a set S will be denoted by χ_S , and the identity operator on a vector space will be denoted by I .

3.3 Absolute strong operator topologies on $\mathcal{L}_{\text{ob}}(E, F)$

Let E and F be vector lattices, where F is Dedekind complete. In this section, we start by showing how topologies can be introduced on vector sublattices of $\mathcal{L}_{\text{ob}}(E, F)$ that can be regarded as absolute strong operator topologies; see Corollary 3.3.5 and Remark 3.3.7, below. Once this is known to be possible, it is easy to relate this to o-Lebesgue topologies and uo-Lebesgue topologies on regular vector sublattices of $\mathcal{L}_{\text{ob}}(E, F)$. In particular, we shall see that every regular vector sublattice of $\mathcal{L}_{\text{ob}}(E, F)$ admits a (necessarily unique) Hausdorff uo-Lebesgue topology when F admits a Hausdorff o-Lebesgue topology; see Corollary 3.4.5, below.

When restricted to the orthomorphisms on a Dedekind complete vector lattice, the picture simplifies; see Section 3.8. In particular, the restrictions of absolute strong operator topologies are then simply strong operator topologies.

The construction in the proof of the following result is an adaptation of that in the proof of [20, Theorem 3.1]. The latter construction is carried out under minimal hypotheses and uses neighbourhood bases at zero as in [44, proof of Theorem 2.3] rather than Riesz pseudo-norms. Such an approach enables one to also understand various ‘pathologies’ in the literature from one central result; see [20, Example 3.10]. It is for this reason of maximum flexibility that we also choose such a neighbourhood approach here.

Theorem 3.3.1. *Let E and F be vector lattices, where F is Dedekind complete, and let τ_F be a (not necessarily Hausdorff) locally solid additive topology on F . Take a non-empty subset S of E . There exists a unique (possibly non-Hausdorff) additive topology $\text{ASOT}_S\tau_F$ on $\mathcal{L}_{\text{ob}}(E, F)$ such that, for a net $(T_\alpha)_{\alpha \in A}$ in $\mathcal{L}_{\text{ob}}(E, F)$, $T_\alpha \xrightarrow{\text{ASOT}_S\tau_F} 0$ if and only if $|T_\alpha||s| \xrightarrow{\tau_F} 0$ for all $s \in S$.*

Let I_S be the ideal of E that is generated by S . For a net $(T_\alpha)_{\alpha \in A}$ in $\mathcal{L}_{\text{ob}}(E, F)$, $T_\alpha \xrightarrow{\text{ASOT}_S\tau_F} 0$ if and only if $|T_\alpha||x| \xrightarrow{\tau_F} 0$ for all $x \in I_S$; and also if and only if $|T_\alpha|x \xrightarrow{\tau_F} 0$ for all $x \in I_S$.

Furthermore:

- (1) *for every $x \in I_S$, the map $T \mapsto Tx$ is an $\text{ASOT}_S\tau_F$ -continuous map from $\mathcal{L}_{\text{ob}}(E, F)$ into F ;*
- (2) *the topology $\text{ASOT}_S\tau_F$ on $\mathcal{L}_{\text{ob}}(E, F)$ is a locally solid additive topology;*

- (3) when τ_F is a Hausdorff topology on F , the following are equivalent for an additive subgroup \mathcal{G} of $\mathcal{L}_{\text{ob}}(E, F)$:
- (a) the restriction $\text{ASOT}_S \tau_F|_{\mathcal{G}}$ of $\text{ASOT}_S \tau_F$ to \mathcal{G} is a Hausdorff topology on \mathcal{G} ;
 - (b) I_S separates the points of \mathcal{G} .
- (4) the following are equivalent for a linear subspace \mathcal{V} of $\mathcal{L}_{\text{ob}}(E, F)$:
- (a) for all $T \in \mathcal{V}$ and $s \in S$, $|\varepsilon T||s| \xrightarrow{\tau_F} 0$ as $\varepsilon \rightarrow 0$ in \mathbb{R} ;
 - (b) the restriction $\text{ASOT}_S \tau_F|_{\mathcal{V}}$ of $\text{ASOT}_S \tau_F$ to \mathcal{V} is a (possibly non-Hausdorff) linear topology on \mathcal{V} .

Proof. Suppose that τ_F is a (not necessarily Hausdorff) locally solid additive topology on F .

It is clear from the required translation invariance of $\text{ASOT}_S \tau_F$ that it is unique, since the nets that are $\text{ASOT}_S \tau_F$ -convergent to zero are prescribed.

For its existence, we take a τ_F -neighbourhood base $\{U_\lambda\}_{\lambda \in \Lambda}$ of zero in F that consists of solid subsets of F . For $x \in I_S$ and $\lambda \in \Lambda$, we set

$$V_{\lambda, x} := \{T \in \mathcal{L}_{\text{ob}}(E, F) : |T||x| \in U_\lambda\}.$$

The $V_{\lambda, x}$ are solid subsets of $\mathcal{L}_{\text{ob}}(E, F)$ since the U_λ are solid subsets of F .

Set

$$\mathcal{N}_0 := \{V_{\lambda, x} : \lambda \in \Lambda, x \in I_S\}.$$

We shall now verify that \mathcal{N}_0 satisfies the necessary and sufficient conditions in [31, Theorem 3 on p. 46] to be a base of neighbourhoods of zero for an additive topology on $\mathcal{L}_{\text{ob}}(E, F)$.

Take $V_{\lambda_1, x_1}, V_{\lambda_2, x_2} \in \mathcal{N}_0$. There exists a $\lambda_3 \in \Lambda$ such that $U_{\lambda_3} \subseteq U_{\lambda_1} \cap U_{\lambda_2}$, and it is easy to verify that then $V_{\lambda_3, |x_1| \vee |x_2|} \subseteq V_{\lambda_1, x_1} \cap V_{\lambda_2, x_2}$. Hence \mathcal{N}_0 is a filter base.

It is clear that $V_{\lambda, x} = -V_{\lambda, x}$.

Take $V_{\lambda, x} \in \mathcal{N}_0$. There exists a $\mu \in \Lambda$ such that $U_\mu + U_\mu \subseteq U_\lambda$, and it is easy to see that then $V_{\mu, x} + V_{\mu, x} \subseteq V_{\lambda, x}$.

An appeal to [31, Theorem 3 on p. 46] now yields that \mathcal{N}_0 is a base of neighbourhoods of zero for an additive topology on $\mathcal{L}_{\text{ob}}(E, F)$ that we shall denote by $\text{ASOT}_S \tau_F$. It is a direct consequence of its definition that, for a net $(T_\alpha)_{\alpha \in A}$ in $\mathcal{L}_{\text{ob}}(E, F)$, $T_\alpha \xrightarrow{\text{ASOT}_S \tau_F} 0$ if and only if $|T_\alpha||x| \xrightarrow{\tau_F} 0$ for all $x \in I_S$. Using the fact that τ_F is a locally solid additive topology on F , it is routine to verify that the latter condition is equivalent to the condition that $|T|x \xrightarrow{\tau_F} 0$ for all $x \in I_S$, as well as to the condition that $|T_\alpha||s| \xrightarrow{\tau_F} 0$ for all $s \in S$.

We turn to the statements in the parts (1)–(4).

For part (1), suppose that $(T_\alpha)_{\alpha \in A}$ is a net in $\mathcal{L}_{\text{ob}}(E, F)$ such that $T_\alpha \xrightarrow{\text{ASOT}_S \tau_F} 0$. Then $|T_\alpha||x| \xrightarrow{\tau_F} 0$ for all $x \in I_S$. Since $|T_\alpha x| \leq |T_\alpha||x|$, the fact that τ_F is locally solid implies that then also $T_\alpha x \xrightarrow{\tau_F} 0$ for all $x \in I_S$.

Since the topology $\text{ASOT}_S \tau_F$ is a locally solid additive topology on $\mathcal{L}_{\text{ob}}(E, F)$ by construction, part (2) is clear.

For part (3), we recall from [31, p. 48, Theorem 4] that an additive topology on a group is Hausdorff if and only if the intersection of the elements of a neighbourhood base of zero is trivial. Using this for F in the second step, and invoking [20, Proposition 2.1] in the third,

we see that

$$\begin{aligned}
 \bigcap_{\lambda \in \Lambda, x \in I_S} (V_{\lambda, x} \cap \mathcal{G}) &= \{ T \in \mathcal{L}_{\text{ob}}(E, F) : |T||x| \in \bigcap_{\lambda \in \Lambda} U_\lambda \text{ for all } x \in I_S \} \cap \mathcal{G} \\
 &= \{ T \in \mathcal{L}_{\text{ob}}(E, F) : |T||x| = 0 \text{ for all } x \in I_S \} \cap \mathcal{G}. \\
 &= \{ T \in \mathcal{L}_{\text{ob}}(E, F) : Tx = 0 \text{ for all } x \in I_S \} \cap \mathcal{G} \\
 &= \{ T \in \mathcal{G} : Tx = 0 \text{ for all } x \in I_S \}.
 \end{aligned}$$

Another appeal to [31, p. 48, Theorem 4] then completes the proof of part (3).

We prove that part (4a) implies part (4b). It is clear that $\text{ASOT}_S \tau_F|_{\mathcal{V}}$ is an additive topology on \mathcal{V} . From what we have already established, we know that the assumption implies that also $|\varepsilon T||x| \xrightarrow{\tau_F} 0$ as $\varepsilon \rightarrow 0$ in \mathbb{R} for all $T \in \mathcal{V}$ and $x \in I_S$. Fix $\lambda \in \Lambda$ and $x \in I_S$, and take $T \in \mathcal{V}$. Since $|\varepsilon T||x| \xrightarrow{\tau_F} 0$ as $\varepsilon \rightarrow 0$ in \mathbb{R} , there exists a $\delta > 0$ such that $|\varepsilon T||x| \in U_\lambda$ whenever $|\varepsilon| < \delta$. That is, $\varepsilon T \in V_{\lambda, x} \cap \mathcal{V}$ whenever $|\varepsilon| < \delta$. Hence $V_{\lambda, x} \cap \mathcal{V}$ is an absorbing subset of \mathcal{V} . Furthermore, since $V_{\lambda, x}$ is a solid subset of $\mathcal{L}_{\text{ob}}(E, F)$, it is clear that $\varepsilon T \in V_{\lambda, x} \cap \mathcal{V}$ whenever $T \in V_{\lambda, x} \cap \mathcal{V}$ and $\varepsilon \in [-1, 1]$. We conclude from [5, Theorem 5.6] that $\text{ASOT}_S \tau_F|_{\mathcal{V}}$ is a linear topology on \mathcal{V} .

We prove that part (4b) implies part (4a). Take $T \in \mathcal{V}$. Then $\varepsilon T \xrightarrow{\text{ASOT}_S \tau_F|_{\mathcal{V}}} 0$ as $\varepsilon \rightarrow 0$ in \mathbb{R} . By construction, this implies that (and is, in fact, equivalent to) the fact that $|\varepsilon T||s| \xrightarrow{\tau_F} 0$ for all $s \in S$. \square

Remark 3.3.2. It is clear from the convergence criteria for nets that the topologies $\text{ASOT}_{S_1} \tau_F$ and $\text{ASOT}_{S_2} \tau_F$ are equal when $I_{S_1} = I_{S_2}$. One could, therefore, work with ideals from the very start, but it seems worthwhile to keep track of a smaller set of presumably more manageable ‘test vectors’. See also the comments preceding Theorem 3.4.3, below.

Remark 3.3.3. Suppose that $(T_\alpha)_{\alpha \in A}$ is a net in $\mathcal{L}_{\text{ob}}(E, F)$ such that $T_\alpha \xrightarrow{\text{ASOT}_S \tau_F} 0$. It is easy to see that then $|T_\alpha|x \xrightarrow{\tau_F} 0$ uniformly on every order bounded subset of I_S , so that then also $T_\alpha x \xrightarrow{\tau_F} 0$ uniformly on every order bounded subset of I_S . When τ_F is a Fatou topology on F (in particular: when τ_F is an o-Lebesgue topology on F ; see [6, Lemma 4.2]), then, conversely, the fact that $T_\alpha x \xrightarrow{\tau_F} 0$ uniformly on every order bounded subset of I_S implies that $T_\alpha \xrightarrow{\text{ASOT}_S \tau_F} 0$. This follows readily from the Riesz-Kantorovich formula for the modulus of an operator.

Definition 3.3.4. The topology $\text{ASOT}_S \tau_F$ in Theorem 3.3.1 is called the *absolute strong operator topology that is generated by τ_F via S* . We shall comment on this nomenclature in Remark 3.3.7, below.

The following result, which can also be obtained using Riesz pseudo-norms, is clear from Theorem 3.3.1.

Corollary 3.3.5. *Let E and F be vector lattices, where F is Dedekind complete, and let τ_F be a (not necessarily Hausdorff) locally solid linear topology on F . Take a vector sublattice \mathcal{E} of $\mathcal{L}_{\text{ob}}(E, F)$ and a non-empty subset S of E .*

There exists a unique additive topology $\text{ASOT}_S\tau_F$ on \mathcal{E} such that, for a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , $T_\alpha \xrightarrow{\text{ASOT}_S\tau_F} 0$ if and only if $|T_\alpha||s| \xrightarrow{\tau_F} 0$ for all $s \in S$.

Let I_S be the ideal of E that is generated by S . For a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , $T_\alpha \xrightarrow{\text{ASOT}_S\tau_F} 0$ if and only if $|T_\alpha||x| \xrightarrow{\tau_F} 0$ for all $x \in I_S$; and also if and only if $|T_\alpha|x \xrightarrow{\tau_F} 0$ for all $x \in I_S$.

Furthermore:

- (1) for every $x \in I_S$, the map $T \mapsto Tx$ is an $\text{ASOT}_S\tau_F\text{-}\tau_F$ continuous map from \mathcal{E} into F ;
- (2) the additive topology $\text{ASOT}_S\tau_F$ on the group \mathcal{E} is, in fact, a locally solid linear topology on the vector lattice \mathcal{E} . When τ_F is a Hausdorff topology on F , then $\text{ASOT}_S\tau_F$ is a Hausdorff topology on \mathcal{E} if and only if I_S separates the points of \mathcal{E} .

Remark 3.3.6. Although in the sequel of this paper we shall mainly be interested in the nets that are convergent in a given topology, let us still remark that is possible to describe an explicit $\text{ASOT}_S\tau_F$ -neighbourhood base of zero in \mathcal{E} . Take a τ_F -neighbourhood base $\{U_\lambda\}_{\lambda \in \Lambda}$ of zero in F that consists of solid subsets of F . For $\lambda \in \Lambda$ and $x \in I_S$, set

$$V_{\lambda,x} := \{ T \in \mathcal{E} : |T||x| \in U_\lambda \}.$$

Then $\{ V_{\lambda,x} : \lambda \in \Lambda, x \in I_S \}$ is an $\text{ASOT}_S\tau_F$ -neighbourhood base of zero in \mathcal{E} .

Remark 3.3.7. It is not difficult to see that $\text{ASOT}_S\tau_F$ is the weakest locally solid linear topology $\tau_{\mathcal{E}}$ on \mathcal{E} such that, for every $x \in I_S$, the map $T \mapsto Tx$ is a $\tau_{\mathcal{E}}\text{-}\tau_F$ continuous map from \mathcal{E} into F . It is also the weakest linear topology $\tau'_{\mathcal{E}}$ on \mathcal{E} such that, for every $x \in I_S$, the map $T \mapsto |T|x$ is a $\tau'_{\mathcal{E}}\text{-}\tau_F$ continuous map from \mathcal{E} into F . The latter characterisation is our motivation for the name ‘absolute strong operator topology’.

Take $F = \mathbb{R}$ and $S = E$. Then $\text{ASOT}_E\tau_{\mathbb{R}}$ is what is commonly known as the absolute weak*-topology on E^* . There is an unfortunate class of ‘weak’ and ‘strong’ here that appears to be unavoidable.

Remark 3.3.8. For comparison with Remark 3.3.7, and in order to make clear the role of the local solidness of the topologies in the present section, we mention the following, which is an easy consequence of [5, Theorem 5.6], for example. Let E and F be vector spaces, where F is supplied with a (not necessarily) Hausdorff linear topology τ_F . Take a linear subspace \mathcal{E} of the vector space of all linear maps from E into F , and take a non-empty subset S of E . Then there exists a unique (not necessarily Hausdorff) linear topology $\text{SOT}_S\tau_F$ on \mathcal{E} such that, for a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , $T_\alpha \xrightarrow{\text{SOT}_S\tau_F} 0$ if and only if $T_\alpha s \xrightarrow{\tau_F} 0$ for all $s \in S$. The subsets of \mathcal{E} of the form $\bigcap_{i=1}^n \{ T \in \mathcal{E} : Ts_i \in V_{\lambda_i} \}$, where the s_i run over S and the V_{λ_i} run over a balanced τ_F -neighbourhood base $\{V_\lambda : \lambda \in \Lambda\}$ of zero in F , are an $\text{SOT}_S\tau_F$ -neighbourhood base of zero in \mathcal{E} . When τ_F is Hausdorff, then $\text{SOT}_S\tau_F$ is Hausdorff if and only if S separates the points of \mathcal{E} . This strong operator topology $\text{SOT}_S\tau_F$ on \mathcal{E} that is generated by τ_F via S , is the weakest linear topology $\tau_{\mathcal{E}}$ on \mathcal{E} such that, for every $s \in S$, the map $T \mapsto Ts$ is $\tau_{\mathcal{E}}\text{-}\tau_F$ -continuous.

3.4 o-Lebesgue topologies and uo-Lebesgue topologies on vector lattices of operators

In order to arrive at results concerning o-Lebesgue topologies and uo-Lebesgue topologies on regular vector sublattices of operators, we need a preparatory result for which we are not aware of a reference. Given its elementary nature, we refrain from any claim to originality. It will re-appear at several places in the sequel.

Lemma 3.4.1. *Let E and F be vector lattices, where F is Dedekind complete, and let \mathcal{E} be a regular vector sublattice of $\mathcal{L}_{\text{ob}}(E, F)$. Suppose that $(T_\alpha)_{\alpha \in \mathcal{A}}$ is net in \mathcal{E} such that $T_\alpha \xrightarrow{0} 0$ in \mathcal{E} . Then $T_\alpha x \xrightarrow{0} 0$ for all $x \in E$.*

Proof. By the regularity of \mathcal{E} , we also have that $T_\alpha \xrightarrow{0} 0$ in $\mathcal{L}_{\text{ob}}(E, F)$. Hence there exists a net $(S_\beta)_{\beta \in \mathcal{B}}$ in $\mathcal{L}_{\text{ob}}(E, F)$ such that $S_\beta \downarrow 0$ in $\mathcal{L}_{\text{ob}}(E, F)$ and with the property that, for every $\beta_0 \in \mathcal{B}$, there exists an $\alpha_0 \in \mathcal{A}$ such that $|T_\alpha| \leq S_{\beta_0}$ for all $\alpha \in \mathcal{A}$ such that $\alpha \geq \alpha_0$. We know from [7, Theorem 1.18], for example, that $S_\beta x \downarrow 0$ for all $x \in E^+$. Since $|T_\alpha x| \leq |T_\alpha| x$ for $x \in E^+$, it then follows easily that $T_\alpha x \xrightarrow{0} 0$ for all $x \in E^+$. Hence $T_\alpha x \xrightarrow{0} 0$ for all $x \in E$. \square

We can now show that the o-Lebesgue property of a locally solid linear topology on the Dedekind complete codomain is inherited by the associated absolute strong operator topology on a regular vector sublattice of operators.

Proposition 3.4.2. *Let E and F be vector lattices, where F is Dedekind complete. Suppose that F admits an o-Lebesgue topology τ_F . Take a regular vector sublattice \mathcal{E} of $\mathcal{L}_{\text{ob}}(E, F)$ and a non-empty subset S of E . Then $\text{ASOT}_S \tau_F$ is an o-Lebesgue topology on \mathcal{E} . When τ_F is a Hausdorff topology on F , then $\text{ASOT}_S \tau_F$ is a Hausdorff topology on \mathcal{E} if and only if I_S separates the points of \mathcal{E} .*

Proof. In view of Corollary 3.3.5, we merely need to show that, for a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , the fact that $T_\alpha \xrightarrow{0} 0$ in \mathcal{E} implies that $T_\alpha \xrightarrow{\text{ASOT}_S \tau_F} 0$. Take $s \in S$. Since also $|T_\alpha| \xrightarrow{0} 0$ in \mathcal{E} , Lemma 3.4.1 implies that $|T_\alpha||s| \xrightarrow{0} 0$ in F . Using that τ_F is an o-Lebesgue topology on F , we find that $|T_\alpha||s| \xrightarrow{\tau_F} 0$. Since this holds for all $s \in S$, Corollary 3.3.5 shows that $T_\alpha \xrightarrow{\text{ASOT}_S \tau_F} 0$ in \mathcal{E} . \square

We conclude by showing that every regular vector sublattice of $\mathcal{L}_{\text{ob}}(E, F)$ admits a (necessarily unique) Hausdorff uo-Lebesgue topology when the Dedekind complete codomain F admits a Hausdorff o-Lebesgue topology. It is the unbounded topology that is associated to (in general multiple) absolute strong operator topologies on the vector sublattice. Our most precise result in this direction is the following. The convergence criterion in part (2) is a ‘minimal one’ that is convenient when one wants to show that a net is convergent, whereas the criteria in part (3) exploits the known convergence of a net to its maximum.

Theorem 3.4.3. *Let E and F be vector lattices, where F is Dedekind complete. Suppose that F admits an o-Lebesgue topology τ_F . Take a regular vector sublattice \mathcal{E} of $\mathcal{L}_{\text{ob}}(E, F)$, a non-empty subset \mathcal{S} of \mathcal{E} , and a non-empty subset S of E .*

Then $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ is a *uo-Lebesgue topology* on \mathcal{E} .

We let I_S denote the ideal of E that is generated by S , and $I_{\mathcal{S}}$ the ideal of \mathcal{E} that is generated by \mathcal{S} . For a net $(T_\alpha)_{\alpha \in A}$ in \mathcal{E} , the following are equivalent:

- (1) $T_\alpha \xrightarrow{u_{\mathcal{S}}\text{ASOT}_S\tau_F} 0$;
- (2) $(|T_\alpha| \wedge |T|)|s| \xrightarrow{\tau_F} 0$ for all $T \in \mathcal{S}$ and $s \in S$;
- (3) $(|T_\alpha| \wedge |T|)x \xrightarrow{\tau_F} 0$ for all $T \in I_{\mathcal{S}}$ and $x \in I_S$.

Suppose that τ_F is actually a Hausdorff *o-Lebesgue topology* on F . Then the following are equivalent:

- (1) $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ is a (necessarily unique) Hausdorff *uo-Lebesgue topology* on \mathcal{E} ;
- (2) I_S separates the points of \mathcal{E} and $I_{\mathcal{S}}$ is order dense in \mathcal{E} .

In that case, the Hausdorff *uo-Lebesgue topology* $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ on \mathcal{E} is the restriction of the (necessarily unique) Hausdorff *uo-Lebesgue topology* on $\mathcal{L}_{\text{ob}}(E, F)$, i.e., of $u_{\mathcal{L}_{\text{ob}}(E, F)}\text{ASOT}_E\tau_F$, and the criteria in (1), (2), and (3) are also equivalent to:

- (4) $(|T_\alpha| \wedge |T|)x \xrightarrow{\tau_F} 0$ for all $T \in \mathcal{L}_{\text{ob}}(E, F)$ and $x \in E$.

Proof. It is clear from Proposition 3.4.2 and [20, Proposition 4.1] that $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ is a *uo-Lebesgue topology* on \mathcal{E} . The two convergence criteria for nets follow from the combination of those in [20, Theorem 3.1] and in Corollary 3.3.5.

According to [20, Proposition 4.1], $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ is a Hausdorff topology on \mathcal{E} if and only if $\text{ASOT}_S\tau_F$ is a Hausdorff topology on \mathcal{E} and $I_{\mathcal{S}}$ is order dense in \mathcal{E} . An appeal to Proposition 3.4.2 then completes the proof of the necessary and sufficient conditions for $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ to be Hausdorff.

Suppose that τ_F is actually also Hausdorff, that I_S separates the points of \mathcal{E} , and that $I_{\mathcal{S}}$ is order dense in \mathcal{E} . From what we have already established, it is clear that $u_{\mathcal{L}_{\text{ob}}(E, F)}\text{ASOT}_E\tau_F$ is a (necessarily unique) Hausdorff *uo-Lebesgue topology* on $\mathcal{L}_{\text{ob}}(E, F)$. Since the restriction of a Hausdorff *uo-Lebesgue topology* on a vector lattice to a regular vector sublattice is a (necessarily unique) Hausdorff *uo-Lebesgue topology* on the vector sublattice (see [44, Proposition 5.12]), the criterion in part (4) follows from that in part (3) applied to $u_{\mathcal{L}_{\text{ob}}(E, F)}\text{ASOT}_E\tau_F$. \square

Remark 3.4.4. Take a τ_F -neighbourhood base $\{U_\lambda\}_{\lambda \in \Lambda}$ of zero in F that consists of solid subsets of F . For $\lambda \in \Lambda$, $\tilde{T} \in I_{\mathcal{S}}$, and $x \in I_S$, set

$$V_{\lambda, \tilde{T}, x} := \{T \in \mathcal{E} : (|T| \wedge |\tilde{T}|)|x| \in U_\lambda\}.$$

As a consequence of the constructions of unbounded and absolute strong operator topologies, $\{V_{\lambda, \tilde{T}, x} : \lambda \in \Lambda, \tilde{T} \in I_{\mathcal{S}}, x \in I_S\}$ is then a $u_{\mathcal{S}}\text{ASOT}_S\tau_F$ -neighbourhood base of zero in \mathcal{E} .

The following is a less precise consequence of Theorem 3.4.3 that will be sufficient in many situations.

Corollary 3.4.5. Let E and F be vector lattices, where F is Dedekind complete. Suppose that F admits a Hausdorff *o-Lebesgue topology* τ_F .

Take a regular vector sublattice \mathcal{E} of $\mathcal{L}_{\text{ob}}(E, F)$. Then \mathcal{E} admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_{\mathcal{E}}$. This topology equals $\mathfrak{u}_{\mathcal{E}}\text{ASOT}_E\tau_F$, and is also equal to the restriction to \mathcal{E} of the Hausdorff uo-Lebesgue topology $\mathfrak{u}_{\mathcal{L}_{\text{ob}}(E, F)}\text{ASOT}_E\tau_F$ on $\mathcal{L}_{\text{ob}}(E, F)$.

For a net $(T_{\alpha})_{\alpha \in A}$ in \mathcal{E} , the following are equivalent:

- (1) $T_{\alpha} \xrightarrow{\widehat{\tau}_{\mathcal{E}}} 0$;
- (2) $(|T_{\alpha}| \wedge |T|)x \xrightarrow{\tau_F} 0$ for all $T \in \mathcal{E}$ and $x \in E$;
- (3) $(|T_{\alpha}| \wedge |T|)x \xrightarrow{\tau_F} 0$ for all $T \in \mathcal{L}_{\text{ob}}(E, F)$ and $x \in E$.

Remark 3.4.6. There can, sometimes, be other ways to see that a given regular vector sublattice of $\mathcal{L}_{\text{ob}}(E, F)$ admits a Hausdorff uo-Lebesgue topology. For example, suppose that F_{oc}^{\sim} separates the points of F . For $x \in E$ and $\varphi \in F_{\text{oc}}^{\sim}$, the map $T \mapsto \varphi(Tx)$ defines an order continuous linear functional on $\mathcal{L}_{\text{oc}}(E, F)$, and it is then clear that the order continuous dual of $\mathcal{L}_{\text{oc}}(E, F)$ separates the points of $\mathcal{L}_{\text{oc}}(E, F)$. Hence $\mathcal{L}_{\text{oc}}(E, F)$ can also be supplied with a Hausdorff uo-Lebesgue topology as in [20, Theorem 5.2] which, in view of its uniqueness, coincides with the one as supplied by Corollary 3.4.5.

3.5 Comparing uniform and strong convergence structures on $\mathcal{L}_{\text{ob}}(E, F)$

Suppose that E and F are vector lattices, where F is Dedekind complete. As explained in Section 3.1, there exist a uniform and a strong convergence structure on $\mathcal{L}_{\text{ob}}(E, F)$ for each of order convergence, unbounded order convergence, and—when applicable—convergence in the Hausdorff uo-Lebesgue topology. In this section, we investigate what the relation is between the members of each of these three pairs. We shall show that only one of the six conceivable implications is valid in general, and that the others are not even generally valid for uniformly bounded sequences of order continuous operators on Banach lattices. Whilst the failures of such general implications may, perhaps, not come as too big a surprise, the positive results for orthomorphisms (see Theorems 3.9.4, 3.9.7, 3.9.9, and 3.9.12, below) may serve to indicate that they are less evident than one would think at first sight.

For monotone nets in $\mathcal{L}_{\text{ob}}(E, F)$, however, the following result shows that then even all four (or six) convergence structures on $\mathcal{L}_{\text{ob}}(E, F)$ are equal.

Proposition 3.5.1. *Let E and F be vector lattices, where F is Dedekind complete, and let $(T_{\alpha})_{\alpha \in A}$ be a monotone net in $\mathcal{L}_{\text{ob}}(E, F)$. The following are equivalent:*

- (1) $T_{\alpha} \xrightarrow{0} 0$ in $\mathcal{L}_{\text{ob}}(E, F)$;
- (2) $T_{\alpha} \xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E, F)$;
- (3) $T_{\alpha}x \xrightarrow{0} 0$ in F for all $x \in E$;
- (4) $T_{\alpha}x \xrightarrow{\text{uo}} 0$ in F for all $x \in E$.

Suppose that, in addition, F admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_F$, so that $\mathcal{L}_{\text{ob}}(E, F)$ also admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E, F)}$ by Corollary 3.4.5. Then (1)–(4) are also equivalent to:

- (5) $T_{\alpha} \xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E, F)}} 0$;

(6) $T_\alpha x \xrightarrow{\widehat{\tau}_F} 0$ for all $x \in E$.

Proof. We may suppose that $T_\alpha \downarrow 0$ and that $x \in E^+$. For order bounded nets in a vector lattice, order convergence and unbounded order convergence are equivalent. Passing to an order bounded tail of $(T_\alpha)_{\alpha \in A}$, we thus see that the parts (1) and (2) are equivalent. Similarly, the parts (3) and (4) are equivalent. The equivalence of the parts (1) and (3) is well known; see [6, Theorem 1.67], for example.

Suppose that F admits a Hausdorff uo-Lebesgue topology $\widehat{\tau}_F$. In that case, it follows from [20, Lemma 7.2] that the parts (2) and (5) are equivalent, as are the parts (4) and (6). \square

When $(T_\alpha)_{\alpha \in A}$ is a not necessarily monotone net in $\mathcal{L}_{\text{ob}}(E, F)$ such that $T_\alpha \xrightarrow{o} 0$, then Lemma 3.4.1 shows that $T_\alpha x \xrightarrow{o} 0$ in F for all $x \in E$. We shall now give five examples to show that each of the remaining five conceivable implications between a corresponding uniform and strong convergence structures on $\mathcal{L}_{\text{ob}}(E, F)$ is not generally valid. In each of these examples, we can even take $E = F$ to be a Banach lattice, and for the net $(T_\alpha)_{\alpha \in A}$ we can even take a uniformly bounded sequence $(T_n)_{n=1}^\infty$ of order continuous operators on E .

Example 3.5.2. We give an example of a uniformly bounded sequence $(T_n)_{n=1}^\infty$ of positive order continuous operators on a Dedekind complete Banach lattice E with a strong order unit, such that $T_n x \xrightarrow{o} 0$ in E for all $x \in E$ but $T_n \not\xrightarrow{o} 0$ in $\mathcal{L}_{\text{ob}}(E)$ because the sequence is not even order bounded in $\mathcal{L}_{\text{ob}}(E)$.

We choose $\ell_\infty(\mathbb{N})$ for $E = F$. For $n \geq 1$, we set $T_n := S^n$, where S is the right shift operator on E . The T_n are evidently positive and of norm one. A moment's thought shows that they are order continuous. Furthermore, it is easy to see that $T_n x \xrightarrow{o} 0$ in E for all $x \in E$. We shall now show that $\{T_n : n \geq 1\}$ is not order bounded in $\mathcal{L}_{\text{ob}}(E)$. For this, we start by establishing that the T_n are mutually disjoint. Let $(e_i)_{i=1}^\infty$ be the standard sequence of unit vectors in E . Take $m \neq n$ and $i \geq 1$. Since e_i is an atom, the Riesz-Kantorovich formula for the infimum of two operators shows that

$$0 \leq (T_m \wedge T_n)e_i = \inf\{te_{m+i} + (1-t)e_{n+i} : 0 \leq t \leq 1\} \leq \inf\{e_{m+i}, e_{n+i}\} = 0.$$

Hence $(T_m \wedge T_n)$ vanishes on the span of the e_i . Since this span is order dense in E , and since $T_n \wedge T_m \in \mathcal{L}_{\text{oc}}(E)$, it follows that $T_n \wedge T_m = 0$.

We can now show that $(T_n)_{n=1}^\infty$ is not order bounded in $\mathcal{L}_{\text{ob}}(E)$. Indeed, suppose that $T \in \mathcal{L}_{\text{ob}}(E)$ is an upper bound for all T_n . Set $e := \bigvee_{i=1}^\infty e_i$. Then, for all $N \geq 1$,

$$Te \geq \left(\bigvee_{n=1}^N T_n \right) e = \left(\sum_{n=1}^N T_n \right) e \geq Ne_{N+1}.$$

This shows that Te cannot be an element of ℓ_∞ . We conclude from this contradiction that $(T_n)_{n=1}^\infty$ is not order bounded in $\mathcal{L}_{\text{ob}}(E)$.

Example 3.5.3. We give an example of a uniformly bounded sequence $(T_n)_{n=1}^\infty$ of positive order continuous operators on a Dedekind complete Banach lattice E with a strong order unit, such that $T_n \xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E)$ but $T_n x \not\xrightarrow{\text{uo}} 0$ for some $x \in E$.

We choose $\ell_\infty(\mathbb{Z})$ for $E = F$. For $n \geq 1$, we set $T_n := S^n$, where S is the right shift operator on E . Just as in Example 3.5.2, the T_n are positive order continuous operators on E of norm one that are mutually disjoint. Since disjoint sequences in vector lattices are unbounded order convergent to zero (see [28, Corollary 3.6]), we have $T_n \xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E)$. On the other hand, if we let e be the two-sided sequence that is constant 1, then $T_n e = e$ for all $n \geq 1$. Hence $(T_n e)_{n=1}^\infty$ is not unbounded order convergent to zero in E .

For our next example, we require a preparatory lemma.

Lemma 3.5.4. *Let μ be the Lebesgue measure on the Borel σ -algebra \mathcal{B} of $[0, 1]$, and let $1 \leq p \leq \infty$. Take a Borel subset S of $[0, 1]$, and define the positive operator $T_S : L_p([0, 1], \mathcal{B}, \mu) \rightarrow L_p([0, 1], \mathcal{B}, \mu)$ by setting*

$$T_S(f) := \int_S f \, d\mu \cdot \chi_S$$

for $f \in L_p([0, 1], \mathcal{B}, \mu)$. Then $T_S \wedge I = 0$.

Proof. Take an $n \geq 1$, and choose disjoint a partition $[0, 1] = \bigcup_{i=1}^n A_i$ of $[0, 1]$ into Borel sets A_i of measure $1/n$. Let e denote the constant function 1. Then

$$\begin{aligned} (T_S \wedge I)e &= \sum_{i=1}^n (T_S \wedge I)\chi_{A_i} \\ &\leq \sum_{i=1}^n (T_S \chi_{A_i}) \wedge \chi_{A_i} \\ &\leq \sum_{i=1}^n (\mu(A_i) \chi_S) \wedge \chi_{A_i} \\ &\leq \sum_{i=1}^n \mu(A_i) \chi_{A_i} \\ &= \frac{1}{n} e. \end{aligned}$$

Since n is arbitrary, we see that $(T_S \wedge I)e = 0$. Because $0 \leq T_S \wedge I \leq I$, $T_S \wedge I$ is order continuous. From the fact that the positive order continuous operator $T_S \wedge I$ vanishes on the weak order unit e of $L_p([0, 1], \mathcal{B}, \mu)$, we conclude that $T_S \wedge I = 0$. \square

Example 3.5.5. We give an example of a uniformly bounded sequence $(T_n)_{n=1}^\infty$ of order continuous operators on a separable reflexive Banach lattice E with a weak order unit, such that $T_n x \xrightarrow{\text{uo}} 0$ in E for all $x \in E$ but $T_n \not\xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E)$ because even $T_n \xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}} 0$ in $\mathcal{L}_{\text{ob}}(E)$.

Let μ be the Lebesgue measure on the Borel σ -algebra \mathcal{B} of $[0, 1]$, and let $1 \leq p \leq \infty$. For E we choose $L_p([0, 1], \mathcal{B}, \mu)$, so that E is reflexive for $1 < p < \infty$. For $n \geq 1$, we let \mathcal{B}_n be the sub- σ -algebra of \mathcal{B} that is generated by the intervals $S_{n,i} := [(i-1)/2^n, i/2^n]$ for $i = 1, \dots, 2^n$, and we let $\mathbb{E}_n : E \rightarrow E$ be the corresponding conditional expectation. By [9, Theorem 10.1.5], \mathbb{E}_n is a positive norm one projection. A moment's thought shows that every open subset of $[0, 1]$ is the union of the countably infinitely many $S_{n,i}$ that are

contained in it, so that it follows from [9, Theorem 10.2.3] that $\mathbb{E}_n f \rightarrow f$ almost everywhere as $n \rightarrow \infty$. By [28, Proposition 3.1], we can now conclude that $\mathbb{E}_n f \xrightarrow{\text{uo}} f$ for all $f \in E$.

On the other hand, it is not true that $\mathbb{E}_n \xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}} I$. To see this, we note that, by [9, Example 10.1.2], every \mathbb{E}_n is a linear combination of operators as in Lemma 3.5.4. Hence $\mathbb{E}_n \perp I$ for all n . Since $\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}$ is a locally solid linear topology, a possible $\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}$ -limit of the \mathbb{E}_n is also disjoint from I , hence cannot be I itself.

On setting $T_n := \mathbb{E}_n - I$ for $n \geq 1$, we have obtained a sequence of operators as desired.

Example 3.5.6. We give an example of a *uniformly bounded sequence* $(T_n)_{n=1}^\infty$ of positive order continuous operators on a Dedekind complete Banach lattice E with a strong order unit that admits a Hausdorff uo-Lebesgue topology, such that $T_n \xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}} 0$ in $\mathcal{L}_{\text{ob}}(E)$ but $T_n x \not\xrightarrow{\widehat{\tau}_E} 0$ in E for some $x \in E$.

We choose E , the $T_n \in \mathcal{L}_{\text{ob}}(E)$, and $e \in E$ as in Example 3.5.3. There are several ways to see that E admits a Hausdorff uo-Lebesgue topology. This follows most easily from the fact that E is atomic (see [44, Lemma 7.4]) and also from [20, Theorem 6.3] in the context of measure spaces. By Corollary 3.4.5, $\mathcal{L}_{\text{ob}}(E)$ then also admits such a topology. Since we already know from Example 3.5.3 that $T_n \xrightarrow{\text{uo}} 0$, we also have that $T_n \xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}} 0$. On the other hand, the fact that $T_n e = e$ for $n \geq 1$ evidently shows that $(T_n e)_{n=1}^\infty$ is not $\widehat{\tau}_E$ -convergent to zero in E .

Example 3.5.7. We note that Example 3.5.5 also gives an example of a *uniformly bounded sequence* $(T_n)_{n=1}^\infty$ of order continuous operators on a separable reflexive Banach lattice E with a weak order unit that admits a Hausdorff uo-Lebesgue topology, such that $T_n x \xrightarrow{\widehat{\tau}_E} 0$ in E for all $x \in E$ but $T_n \not\xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}} 0$ in $\mathcal{L}_{\text{ob}}(E)$.

3.6 Orthomorphisms

In this section, we review some material concerning orthomorphism and establish a few auxiliary result for use in the present paper and in future ones.

Let E be a vector lattice. We recall from [7, Definition 2.41] that an operator on E is called an *orthomorphism* when it is a band preserving order bounded operator. An orthomorphism is evidently disjointness preserving, it is order continuous (see [7, Theorem 2.44]), and its kernel is a band (see [7, Theorem 2.48]). We denote by $\text{Orth}(E)$ the collection of all orthomorphism on E . Even when E is not Dedekind complete, the supremum and infimum of two orthomorphisms S and T in E always exists in $\mathcal{L}_{\text{ob}}(E)$. In fact, we have

$$\begin{aligned} [S \vee T](x) &= S(x) \vee T(x) \\ [S \wedge T](x) &= S(x) \wedge T(x) \end{aligned} \tag{3.1}$$

for $x \in E^+$ and

$$|Tx| = |T||x| = |T(|x|)| \tag{3.2}$$

for $x \in E$; see [7, Theorems 2.43 and 2.40]. Consequently, $\text{Orth}(E)$ is a unital vector lattice algebra for every vector lattice E . Even more is true: according to [7, Theorem 2.59], $\text{Orth}(E)$ is an (obviously Archimedean) f -algebra for every vector lattice E , so it is commutative by [7, Theorem 2.56]. Furthermore, for every vector lattice E , when $T \in \text{Orth}(E)$ and $T : E \rightarrow E$ is injective and surjective, then the linear map $T^{-1} : E \rightarrow E$ is again an orthomorphism. We refer to [37, Theorem 3.1.10] for a proof of this result of Huijsmans' and de Pagter's.

It follows easily from equation (3.1) that, for every vector lattice E , the identity operator is a weak order unit of $\text{Orth}(E)$. When E is Dedekind complete, $\text{Orth}(E)$ is the band in $\mathcal{L}_{\text{ob}}(E)$ that is generated by the identity operator on E ; see [7, Theorem 2.45].

Let E be a vector lattice, let $T \in \mathcal{L}_{\text{ob}}(E)$, and let $\lambda \geq 0$. Using [7, Theorem 2.40], it is not difficult to see that the following are equivalent:

- (1) $-\lambda I \leq T \leq \lambda I$;
- (2) $|T|$ exists in $\mathcal{L}_{\text{ob}}(E)$, and $|T| \leq \lambda I$;
- (3) $|Tx| \leq \lambda|x|$ for all $x \in E$.

The set of all such T is a unital subalgebra $\mathcal{Z}(E)$ of $\text{Orth}(E)$ consisting of ideal preserving order bounded operators on E . It is called the *ideal centre* of E .

Let E be a vector lattice, and define the *stabiliser* of E , denoted by $\mathcal{S}(E)$, as the set of linear operators on E that are ideal preserving. It is not required that these operators be order bounded, but this is nevertheless always the case. In fact, $\mathcal{S}(E)$ is a unital subalgebra of $\text{Orth}(E)$ for every vector lattice E (see [47, Proposition 2.6]), so that we have the chain

$$\mathcal{Z}(E) \subseteq \mathcal{S}(E) \subseteq \text{Orth}(E)$$

of unital algebras for every vector lattice E . For every Banach lattice E , we have

$$\mathcal{Z}(E) = \mathcal{S}(E) = \text{Orth}(E);$$

see [47, Corollary 4.2], so that the identity operator on E is then even an order unit of $\text{Orth}(E)$.

For every Banach lattice E , $\text{Orth}(E)$ is a unital Banach subalgebra of the bounded linear operators on E in the operator norm. This follows easily from the facts that bands are closed and that a band preserving operator on a Banach lattice is automatically order bounded; see [7, Theorem 4.76].

Let E be a Banach lattice. Since the identity operator is an order unit of $\text{Orth}(E)$, we can introduce the order unit norm $\|\cdot\|_I$ with respect to I on $\text{Orth}(E)$ by setting

$$\|T\|_I := \inf\{\lambda \geq 0 : |T| \leq \lambda I\}$$

for $T \in \text{Orth}(E)$. Then $\|T\| = \|T\|_I$ for all $T \in \text{Orth}(E)$; see [47, Proposition 4.1]. Since we already know that $\text{Orth}(E)$ is complete in the operator norm, it follows that $\text{Orth}(E)$, when supplied with $\|\cdot\| = \|\cdot\|_I$, is a unital Banach lattice algebra that is also an AM-space. When E is a Dedekind complete Banach lattice, then evidently $\|T\| = \|T\|_I = \| |T| \|_I = \| |T| \| = \|T\|_r$ for $T \in \text{Orth}(E)$. Hence $\text{Orth}(E)$ is then also a unital Banach lattice subalgebra of the Banach lattice algebra of all order bounded operators on E in the regular norm.

Let E be Banach lattice. It is clear from the above that $(\text{Orth}(E), \|\cdot\|) = (\text{Orth}(E), \|\cdot\|_I)$ is a unital Banach f -algebra in which its identity element is also a (positive) order unit. The following result is, therefore, applicable with $\mathcal{A} = \text{Orth}(E)$ and $e = I$. It shows, in particular, that $\text{Orth}(E)$ is isometrically Banach lattice algebra isomorphic to a $C(K)$ -space. Both its statement and its proof improve on the ones in [15, Proposition 2.6], [41, Proposition 1.4], and [30].

Theorem 3.6.1. *Let \mathcal{A} be a unital f -algebra such that its identity element e is also a (positive) order unit, and such that it is complete in the submultiplicative order unit norm $\|\cdot\|_e$ on \mathcal{A} . Let \mathcal{B} be a (not necessarily unital) associative subalgebra of \mathcal{A} . Then $\overline{\mathcal{B}}^{\|\cdot\|_e}$ is a Banach f -subalgebra of \mathcal{A} . When $e \in \overline{\mathcal{B}}$, then there exist a compact Hausdorff space K , uniquely determined up to homeomorphism, and an isometric surjective Banach lattice algebra isomorphism $\psi : \overline{\mathcal{B}}^{\|\cdot\|_e} \rightarrow C(K)$.*

Proof. Since $(\mathcal{A}, \|\cdot\|_I)$ is an AM-space with order unit e , there exist a compact Hausdorff space K' and an isometric surjective lattice homomorphism $\psi' : \mathcal{A} \rightarrow C(K')$ such that $\psi'(e) = 1$; see [37, Theorem 2.1.3] for this result of Kakutani's, for example. Via this isomorphism, the f -algebra multiplication on $C(K')$ provides the vector lattice \mathcal{A} with a multiplication that makes \mathcal{A} into an f -algebra with e as its positive multiplicative identity element. Such a multiplication is, however, unique; see [7, Theorem 2.58]. Hence ψ' also preserves multiplication, and we conclude that $\psi' : \mathcal{A} \rightarrow C(K')$ is an isometric surjective Banach lattice algebra isomorphism.

We now turn to \mathcal{B} . It is clear that $\overline{\mathcal{B}}^{\|\cdot\|_e}$ is Banach subalgebra of \mathcal{A} . After moving to the $C(K')$ -model for \mathcal{A} that we have obtained, [23, Lemma 4.48] shows that $\overline{\mathcal{B}}^{\|\cdot\|_e}$ is also a vector sublattice of \mathcal{A} . Hence $\overline{\mathcal{B}}^{\|\cdot\|_e}$ is a Banach f -subalgebra of \mathcal{A} . When $e \in \overline{\mathcal{B}}^{\|\cdot\|_e}$, we can then apply the first part of the proof to \mathcal{B} , and obtain a compact Hausdorff space K and an isometric surjective Banach lattice algebra isomorphism $\psi : \overline{\mathcal{B}}^{\|\cdot\|_e} \rightarrow C(K)$. The Banach-Stone theorem (see [12, Theorem VI.2.1], for example) implies that K is uniquely determined up to homeomorphism. \square

We now proceed to show that E and $\text{Orth}(E)$ have isomorphic universal completions. We start with a preparatory lemma.

Proposition 3.6.2. *Let E be a Dedekind complete vector lattice, and let $x \in E$. Let I_x be the principal ideal of E that is generated by x , let B_x be the principal band in E that is generated by x , let $P_x : E \rightarrow B_x$ be the corresponding order projection, and let \mathcal{I}_{P_x} be the principal ideal of $\mathcal{L}_{\text{ob}}(E)$ that is generated by P_x . For $T \in \mathcal{I}_{P_x}$, set $\psi_x(T) := T|x|$. Then $\psi_x(T) \in I_x$, and:*

- (1) *the map $\psi_x : \mathcal{I}_{P_x} \rightarrow I_x$ is a surjective vector lattice isomorphism such that $\psi_x(P_x) = |x|$;*
- (2) *$\mathcal{I}_{P_x} = P_x \mathcal{L}(E)$.*

Proof. Take $T \in \mathcal{I}_{P_x}$. There exists a $\lambda \geq 0$ such that $|T| \leq \lambda P_x$, and this implies that $|Ty| \leq \lambda P_x|y|$ for all $y \in E$. This shows that $T|x| \in I_x$, so that ψ_x maps \mathcal{I}_{P_x} into I_x ; it also shows that $T(B_x^d) = \{0\}$. Suppose that $T|x| = 0$. Since the kernel of T is a band in E , this implies that T vanishes on B_x . We already know that it vanishes on B_x^d . Hence $T = 0$, and we conclude that ψ_x is injective. We show that ψ_x is surjective. Let $y \in I_x$. Take a

$\lambda > 0$ such that $0 \leq |y/\lambda| \leq |x|$. An inspection of the proof of [7, Theorem 2.49] shows that there exists a $T \in \mathcal{Z}(E)$ with $T|x| = y/\lambda$. Since $\lambda TP_x \in \mathcal{J}_{P_x}$ and $(\lambda TP_x)|x| = y$, we see that ψ_x is surjective. Finally, it is clear from equation (3.1) that ψ_x is a vector lattice homomorphism. This completes the proof of part (1).

We turn to part (2). It is clear that $\mathcal{J}_{P_x} \supseteq P_x \mathcal{Z}(E)$. Take $T \in \mathcal{J}_{P_x} \subseteq \mathcal{Z}(E)$. Then also $P_x T \in \mathcal{J}_{P_x}$. Since $\psi_x(T) = \psi_x(P_x T)$, the injectivity of ψ_x on \mathcal{J}_{P_x} implies that $T = P_x T \in P_x \mathcal{Z}(E)$. \square

The first part of Proposition 3.6.2 is used in the proof of our next result.

Proposition 3.6.3. *Let E be a Dedekind complete vector lattice. Then there exist an order dense ideal I of E and an order dense ideal \mathcal{J} of $\text{Orth}(E)$ such that I and \mathcal{J} are isomorphic vector lattices.*

Proof. Choose a maximal disjoint system $\{x_\alpha : \alpha \in \mathcal{A}\}$ in E . For each $\alpha \in \mathcal{A}$, let I_{x_α} , B_{x_α} , $P_{x_\alpha} : E \rightarrow B_{x_\alpha}$, $\mathcal{J}_{P_{x_\alpha}}$, and the vector lattice isomorphism $\psi_{x_\alpha} : \mathcal{J}_{P_{x_\alpha}} \rightarrow I_{x_\alpha}$ be as in Proposition 3.6.2.

Since the x_α are mutually disjoint, it is clear that the ideal $\sum_{\alpha \in \mathcal{A}} I_{x_\alpha}$ of E is, in fact, an internal direct sum $\bigoplus_{\alpha \in \mathcal{A}} I_{x_\alpha}$. Since the disjoint system is maximal, $\bigoplus_{\alpha \in \mathcal{A}} I_{x_\alpha}$ is an order dense ideal of E .

It follows easily from equation (3.1) that the P_{x_α} are also mutually disjoint. They even form a maximal disjoint system in $\text{Orth}(E)$. To see this, suppose that $T \in \text{Orth}(E)$ is such that $|T| \wedge P_{x_\alpha} = 0$ for all $\alpha \in \mathcal{A}$. Then $(|T|x_\alpha) \wedge x_\alpha = (|T| \wedge P_{x_\alpha})x_\alpha = 0$ for all $\alpha \in \mathcal{A}$. Since $|T|$ is band preserving, this implies that $|T|x_\alpha = 0$ for all $\alpha \in \mathcal{A}$. The fact that the kernel of $|T|$ is a band in E then yields that $|T| = 0$. Just as for E , we now conclude that the ideal $\sum_{\alpha \in \mathcal{A}} \mathcal{J}_{P_{x_\alpha}}$ of $\text{Orth}(E)$ is an internal direct sum $\bigoplus_{\alpha \in \mathcal{A}} \mathcal{J}_{P_{x_\alpha}}$ that is order dense in $\text{Orth}(E)$.

Since $\bigoplus_{\alpha \in \mathcal{A}} \psi_{x_\alpha} : \bigoplus_{\alpha \in \mathcal{A}} \mathcal{J}_{P_{x_\alpha}} \rightarrow \bigoplus_{\alpha \in \mathcal{A}} I_{x_\alpha}$ is a vector lattice isomorphism by Proposition 3.6.2, the proof is complete. \square

It is generally true that a vector lattice and an order dense vector sublattice of it have isomorphic universal completions; see [6, Theorems 7.21 and 7.23]. Proposition 3.6.3 therefore implies the following.

Corollary 3.6.4. *Let E be a Dedekind complete vector lattice. Then the universal completions of E and of $\text{Orth}(E)$ are isomorphic vector lattices.*

The previous result enables us to relate the countable sup property of E to that of $\text{Orth}(E)$. We recall that vector lattice E has the countable sup property when, for every non-empty subset S of E that has a supremum in E , there exists an at most countable subset of S that has the same supremum in E as S . In parts of the literature, such as in [36] and [51], E is then said to be order separable. We also recall that a subset of a vector lattice is said to be an *order basis* when the band that it generates is the whole vector lattice.

Proposition 3.6.5. *Let E be a Dedekind complete vector lattice. The following are equivalent:*

- (1) $\text{Orth}(E)$ has the countable sup property;
- (2) E has the countable sup property and an at most countably infinite order basis.

Proof. It is proved in [33, Theorem 6.2] that, for an arbitrary vector lattice F , F^u has the countable sup property if and only if F has the countable sup property as well as an at most countably infinite order basis. Since $\text{Orth}(E)$ has a weak order unit I , we see that $\text{Orth}(E)^u$ has the countable sup property if and only if $\text{Orth}(E)$ has the countable sup property. On the other hand, since $\text{Orth}(E)^u$ and E^u are isomorphic by Corollary 3.6.4, an application of this same result to E shows that $\text{Orth}(E)^u$ has the countable sup property if and only if E has the countable sup property and an at most countably infinite order basis. \square

We conclude by giving some estimates for orthomorphisms in Proposition 3.6.7 that will be used in the sequel. As a preparation, we need the following extension of [7, Exercise 1.3.7].

Lemma 3.6.6. *Let E be a vector lattice with the principal projection property. Take $x, y \in E$. For $\lambda \in \mathbb{R}$, let P_λ denote the order projection in E onto the band generated by $(x - \lambda y)^+$. Then $\lambda P_\lambda y \leq P_\lambda x$. When $x, y \in E^+$ and $\lambda \geq 0$, then $x \leq \lambda y + P_\lambda x$.*

Proof. The first inequality follows from the fact that

$$0 \leq P_\lambda(x - \lambda y)^+ = P_\lambda(x - \lambda y) = P_\lambda x - \lambda P_\lambda y.$$

For the second inequality, we note that $x - \lambda y \leq (x - \lambda y)^+ = P_\lambda(x - \lambda y)^+$ for all x, y , and λ . When $x, y \in E^+$ and $\lambda \geq 0$, then $(x - \lambda y)^+ \leq x^+ = x$, so that

$$x \leq \lambda y + P_\lambda(x - \lambda y)^+ \leq \lambda y + P_\lambda x.$$

\square

Proposition 3.6.7. *Let E be a Dedekind complete vector lattice, and let $T \in \text{Orth}(E)^+$. For $\lambda > 0$, let \mathcal{P}_λ be the order projection in $\text{Orth}(E)$ onto the band generated by $(T - \lambda I)^+$ in $\text{Orth}(E)$. There exists a unique order projection P_λ in E such that $\mathcal{P}_\lambda(S) = P_\lambda S$ for all $S \in \text{Orth}(E)$. Furthermore:*

- (1) $\lambda P_\lambda \leq P_\lambda T \leq T$;
- (2) $T \leq \lambda I + P_\lambda T$;
- (3) $(P_\lambda T x) \wedge y \leq \frac{1}{\lambda} T y$ for all $x, y \in E^+$.

Proof. Since $0 \leq \mathcal{P}_\lambda \leq I_{\text{Orth}(E)}$, it follows from [6, Theorem 2.62] that there exists a unique $P_\lambda \in \text{Orth}(E)$ with $0 \leq P_\lambda \leq I$ such that $\mathcal{P}_\lambda(S) = P_\lambda S$ for all $S \in \text{Orth}(E)$. The fact that \mathcal{P}_λ is idempotent implies that P_λ is also idempotent. Hence P_λ is an order projection.

The inequalities in the parts (1) and (2) are then a consequence of those in Lemma 3.6.6. For part (3), we note that $(P_\lambda T x) \wedge y$ is in the image of the projection P_λ . Since order projections are vector lattice homomorphisms, we have, using part (1) in the final step, that

$$(P_\lambda T x) \wedge y = P_\lambda((P_\lambda T x) \wedge y) = (P_\lambda^2 T x) \wedge P_\lambda y \leq P_\lambda y \leq \frac{1}{\lambda} T y.$$

\square

3.7 Continuity properties of orthomorphisms

Orthomorphisms preserve order convergence of nets. In this short section, we show that they also preserve unbounded order convergence and, when applicable, convergence in the (necessarily unique) Hausdorff uo-Lebesgue topology.

Before doing so, let us note that this is in contrast to the case of general order bounded operators. Surely, there exist order bounded operators that are not order continuous. For the remaining two convergence structures, we consider ℓ_1 with its standard basis $(e_n)_{n=1}^\infty$. It follows from [28, Corollary 3.6] that $e_n \xrightarrow{\text{uo}} 0$. There are several ways to see that ℓ_1 admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_{\ell_1}$. This follows from the fact that its norm is order continuous (see [44, p. 993]), from the fact that it is atomic (see [44, Lemma 7.4]), and from a result in the context of measure spaces (see [20, Theorem 6.2]).

The latter two results also show that $e_n \xrightarrow{\widehat{\tau}_{\ell_1}} 0$ which is, of course, also a consequence of the fact that $e_n \xrightarrow{\text{uo}} 0$. Define $T : \ell_1 \rightarrow \ell_1$ by setting $Tx := (\sum_{n=1}^\infty x_n) e_1$ for $x = \sum_{n=1}^\infty x_n e_n \in \ell_1$. Since $Te_n = e_1$ for all $n \geq 1$, the order continuous positive operator T on ℓ_1 preserves neither uo-convergence nor $\widehat{\tau}_{\ell_1}$ -convergence of sequences in ℓ_1 .

Proposition 3.7.1. *Let E be a Dedekind complete vector lattice, and let $T \in \text{Orth}(E)$. Suppose that $(x_\alpha)_{\alpha \in \mathcal{A}}$ is a net in E such that $x_\alpha \xrightarrow{\text{uo}} 0$ in E . Then $Tx_\alpha \xrightarrow{\text{uo}} 0$ in E .*

Proof. Using equation (3.2), one easily sees that we may suppose that T and the x_α are positive. For $n \geq 1$, we let \mathcal{P}_n be the order projection in $\text{Orth}(E)$ onto the band generated by $(T - nI)^+$ in $\text{Orth}(E)$. According to Proposition 3.6.7, there exists a unique order projection P_n in E such that $\mathcal{P}_n(S) = P_n S$ for all $S \in \text{Orth}(E)$. Take $e \in E^+$. By applying part (2) of Proposition 3.6.7 in the first step and its part (3) in the third, we see that, for $\alpha \in \mathcal{A}$ and $n \geq 1$,

$$\begin{aligned} (Tx_\alpha) \wedge e &\leq (nx_\alpha + P_n Tx_\alpha) \wedge e \\ &\leq n(x_\alpha \wedge e) + P_n Tx_\alpha \wedge e \\ &\leq n(x_\alpha \wedge e) + \frac{1}{n} Te. \end{aligned} \tag{3.3}$$

This implies that, for $n \geq 1$,

$$0 \leq \inf_\alpha \sup_{\beta \geq \alpha} [(Tx_\beta) \wedge e] \leq n \inf_\alpha \sup_{\beta \geq \alpha} [x_\beta \wedge e] + \frac{1}{n} Te.$$

Since $x_\alpha \wedge e \xrightarrow{o} 0$ in E , it now follows from [28, Remark 2.2] that

$$0 \leq \inf_\alpha \sup_{\beta \geq \alpha} [(Tx_\beta) \wedge e] \leq \frac{1}{n} Te$$

for all $n \geq 1$. Hence $\inf_\alpha \sup_{\beta \geq \alpha} [(Tx_\beta) \wedge e] = 0$, and we conclude that $(Tx_\alpha) \wedge e \xrightarrow{o} 0$ in E . Since $e \in E^+$ was arbitrary, the proof is complete. \square

For the case of a Hausdorff uo-Lebesgue topology, we need the following preparatory result that has some independent interest. Lemma 3.9.11 is of the same flavour.

Proposition 3.7.2. *Let E be a Dedekind complete vector lattice that admits a (not necessarily Hausdorff) locally solid linear topology τ_E , and let $T \in \text{Orth}(E)$. Suppose that $(x_\alpha)_{\alpha \in \mathcal{A}}$ is a net in E such that $x_\alpha \xrightarrow{\tau_E} 0$ in E . Then $Tx_\alpha \xrightarrow{u_E \tau_E} 0$ in E .*

Proof. As in the proof of Proposition 3.7.1, we may suppose that T and the x_α are positive. For $n \geq 1$, we let \mathcal{P}_n be the order projection in $\text{Orth}(E)$ onto the band generated by $(T - nI)^+$ in $\text{Orth}(E)$ again, so that again there exists a unique order projection P_n in E such that $\mathcal{P}_n(S) = P_n S$ for all $S \in \text{Orth}(E)$. Fix $e \in E^+$. Take a solid τ_E -neighbourhood U of 0 in E , and choose a τ_E -neighbourhood V of 0 such that $V + V \subseteq U$. Take an $n_0 \geq 1$ such that $Te/n_0 \in V$. As $x_\alpha \xrightarrow{\tau_E} 0$, there exists an $\alpha_0 \in \mathcal{A}$ such that $n_0 x_\alpha \in V$ for all $\alpha \geq \alpha_0$. Continuing the chain of inequalities in equation (3.3) for n_0 for one more step, we see that, for all $\alpha \geq \alpha_0$,

$$\begin{aligned} (Tx_\alpha) \wedge e &\leq n_0(x_\alpha \wedge e) + \frac{1}{n_0}Te \\ &\leq n_0 x_\alpha + \frac{1}{n_0}Te \\ &\in V + V \subseteq U \end{aligned} \tag{3.4}$$

The solidness of V then implies that $(Tx_\alpha) \wedge e \in U$ for all $\alpha \geq \alpha_0$. Since U and e were arbitrary, we conclude that $Tx_\alpha \xrightarrow{u_E \tau_E} 0$. \square

Since the unbounded topology $u_E \widehat{\tau}_E$ that is generated by a Hausdorff uo-Lebesgue topology $\widehat{\tau}_E$ equals $\widehat{\tau}_E$ again, the following is now clear.

Corollary 3.7.3. *Let E be a Dedekind complete vector lattice that admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_E$, and let $T \in \text{Orth}(E)$. Suppose that $(x_\alpha)_{\alpha \in \mathcal{A}}$ is a net in E such that $x_\alpha \xrightarrow{\widehat{\tau}_E} 0$ in E . Then $Tx_\alpha \xrightarrow{\widehat{\tau}_E} 0$ in E .*

3.8 Topologies on $\text{Orth}(E)$

Let E be a Dedekind complete vector lattice, and suppose that τ_E is a (not necessarily Hausdorff) locally solid additive topology on E . Take a non-empty subset S of E . According to Theorem 3.3.1, there exists a unique additive topology $\text{ASOT}_S \tau_E$ on $\mathcal{L}_{\text{ob}}(E)$ such that, for a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in $\mathcal{L}_{\text{ob}}(E)$, $T_\alpha \xrightarrow{\text{ASOT}_S \tau_E} 0$ if and only if $|T_\alpha||s| \xrightarrow{\tau_E} 0$ for all $s \in S$. When $(T_\alpha)_{\alpha \in \mathcal{A}} \subseteq \text{Orth}(E)$, equation (3.2) and the local solidness of τ_E imply that this convergence criterion is also equivalent to the one that $T_\alpha s \xrightarrow{\tau_E} 0$ for all $s \in S$. Hence on subsets of $\text{Orth}(E)$, an absolute strong operator topology that is generated by a locally solid additive topology on E coincides with the corresponding strong operator topology. In order to remind ourselves of the connection with the topology on the enveloping vector lattice $\mathcal{L}_{\text{ob}}(E)$ of $\text{Orth}(E)$, we

shall keep writing $\text{ASOT}_S \tau_F$ when considering the restriction of this topology to subsets of $\text{Orth}(E)$, rather than switch to, e.g., $\text{SOT}_S \tau_F$.

The above observation can be used in several results in Section 3.3. For the ease of reference, we include the following consequence of Corollary 3.3.5.

Corollary 3.8.1. *Let E be a Dedekind complete vector lattice, and let τ_E be a (not necessarily Hausdorff) locally solid linear topology on E . Take a vector sublattice \mathcal{E} of $\text{Orth}(E)$ and a non-empty subset S of E .*

There exists a unique locally solid linear topology $\text{ASOT}_S \tau_E$ on \mathcal{E} such that, for a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , $T_\alpha \xrightarrow{\text{ASOT}_S \tau_E} 0$ if and only if $T_\alpha s \xrightarrow{\tau_E} 0$ for all $s \in S$.

Let I_S be the ideal of E that is generated by S . For a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , $T_\alpha \xrightarrow{\text{ASOT}_S \tau_E} 0$ if and only if $T_\alpha x \xrightarrow{\tau_E} 0$ for all $x \in I_S$.

When τ_E is a Hausdorff topology on F , then $\text{ASOT}_S \tau_E$ is a Hausdorff topology on \mathcal{E} if and only if I_S separates the points of \mathcal{E} .

According to the next result, there is an intimate relation between the existence of Hausdorff o-Lebesgue topologies and uo-Lebesgue topologies on E and on $\text{Orth}(E)$.

Proposition 3.8.2. *Let E be a Dedekind complete vector lattice. The following are equivalent:*

- (1) *E admits a Hausdorff o-Lebesgue topology;*
- (2) *$\text{Orth}(E)$ admits a Hausdorff o-Lebesgue topology;*
- (3) *E admits a (necessarily unique) Hausdorff uo-Lebesgue topology;*
- (4) *$\text{Orth}(E)$ admits a (necessarily unique) Hausdorff uo-Lebesgue topology.*

Proof. As E and $\text{Orth}(E)$ are Dedekind complete, they are not just order dense vector sublattices of their universal completions but even order dense ideals; see [7, p.126–127]. Since these universal completions are isomorphic vector lattices by Corollary 3.6.4, the proposition follows from a double application of [20, Theorem 4.9.(3)]. \square

For a Dedekind complete vector lattice E , $\text{Orth}(E)$, being a band in $\mathcal{L}_{\text{ob}}(E)$, is a regular vector sublattice of $\mathcal{L}_{\text{ob}}(E)$. A regular vector sublattice \mathcal{E} of $\text{Orth}(E)$ is, therefore, also a regular vector sublattice of $\mathcal{L}_{\text{ob}}(E)$, and Proposition 3.4.2 then shows how o-Lebesgue topologies on \mathcal{E} can be obtained from an o-Lebesgue topology on E as (absolute) strong operator topologies. In particular, this makes the fact that part (1) of Proposition 3.8.2 implies its part (2) more concrete. The fact that part (1) implies part (2) is made more concrete as a special case of the following consequence of Theorem 3.4.3.

Theorem 3.8.3. *Let E be a Dedekind complete vector lattice. Suppose that E admits an o-Lebesgue topology τ_E . Take a regular vector sublattice \mathcal{E} of $\text{Orth}(E)$, a non-empty subset \mathcal{S} of \mathcal{E} , and a non-empty subset S of E .*

Then $\text{u}_{\mathcal{S}} \text{ASOT}_S \tau_E$ is a uo-Lebesgue topology on \mathcal{E} .

We let I_S denote the ideal of E that is generated by S , and $I_{\mathcal{S}}$ the ideal of \mathcal{E} that is generated by \mathcal{S} . For a net $(T_\alpha)_{\alpha \in \mathcal{A}}$ in \mathcal{E} , the following are equivalent:

- (1) *$T_\alpha \xrightarrow{\text{u}_{\mathcal{S}} \text{ASOT}_S \tau_E} 0$;*
- (2) *$|T_\alpha s| \wedge |Ts| \xrightarrow{\tau_E} 0$ for all $T \in \mathcal{S}$ and $s \in S$;*

(3) $|T_\alpha x| \wedge |Tx| \xrightarrow{\tau_E} 0$ for all $T \in I_{\mathcal{S}}$ and $x \in I_{\mathcal{S}}$.

Suppose that τ_E is actually a Hausdorff o -Lebesgue topology $\widehat{\tau}_E$ on E . Then the following are equivalent:

(1) $\mathfrak{u}_{\mathcal{S}}\text{ASOT}_{\mathcal{S}}\widehat{\tau}_E$ is a (necessarily unique) Hausdorff uo -Lebesgue topology on \mathcal{E} ;

(2) $I_{\mathcal{S}}$ separates the points of \mathcal{E} and $I_{\mathcal{S}}$ is order dense in \mathcal{E} .

In that case, the Hausdorff uo -Lebesgue topology $\mathfrak{u}_{\mathcal{S}}\text{ASOT}_{\mathcal{S}}\tau_E$ on \mathcal{E} is the restriction of the (necessarily unique) Hausdorff uo -Lebesgue topology on $\mathcal{L}_{\text{ob}}(E, F)$, i.e., of $\mathfrak{u}_{\mathcal{L}_{\text{ob}}(E, F)}\text{ASOT}_E\tau_E$, and the criteria in (1), (2), and (3) are also equivalent to:

(4) $(|T_\alpha| \wedge |T|)x \xrightarrow{\widehat{\tau}_E} 0$ for all $T \in \mathcal{L}_{\text{ob}}(E)$ and $x \in E$.

3.9 Comparing uniform and strong convergence structures on $\text{Orth}(E)$

Let E and F be vector lattices, where F is Dedekind complete, and let $(T_\alpha)_{\alpha \in A}$ be a net in $\mathcal{L}_{\text{ob}}(E, F)$. In Section 3.5, we studied the relation between uniform and strong convergence of $(T_\alpha)_{\alpha \in A}$ for order convergence, unbounded order convergence, and—when applicable—convergence in a Hausdorff uo -Lebesgue topology. In the present section, we consider the case where $(T_\alpha)_{\alpha \in A}$ is actually contained in $\text{Orth}(E)$. As we shall see, the relation between uniform and strong convergence is now much more symmetrical than in the general case of Section 3.5; see Theorem 3.9.4 (and Theorem 3.9.7), Theorem 3.9.9, and Theorem 3.9.12, below.

These positive results might, perhaps, lead one to wonder whether some of the three uniform convergence structures under consideration might actually even be identical for the orthomorphisms. This, however, is not the case. There even exist sequences of positive orthomorphisms on separable reflexive Banach lattices with weak order units showing that the two ‘reverse’ implications in question are not generally valid. For this, we consider $E := L_p([0, 1])$ for $1 < p < \infty$. In that case, $\text{Orth}(E)$ can canonically be identified with $L_\infty([0, 1])$ as an f -algebra; see [7, Example 2.67], for example. The uo -convergence of a net in the regular vector sublattice $L_\infty([0, 1])$ of $L_0([0, 1])$ coincides with that in $L_0([0, 1])$ which, according to [28, Proposition 3.1], is simply convergence almost everywhere in the case of sequences. According to [20, Theorem 6.3], the convergence of a net in the Hausdorff uo -Lebesgue topology of $L_\infty([0, 1])$ is equal to the convergence in measure. For $n \geq 1$, set $f_n := n\chi_{[0, 1/n]}$. Then $f_n \xrightarrow{uo} 0$ in $L_\infty([0, 1])$, but it is not true that $f_n \xrightarrow{o} 0$ in $L_\infty([0, 1])$ since the f_n are not even order bounded in $L_\infty([0, 1])$. Using $\chi_{[(k-1)2^{-n}, k2^{-n}]}$ for $n \geq 1$ and $k = 1, \dots, 2^n$, one easily finds a sequence that is convergent to zero in measure, but that is not convergent in any point of $[0, 1]$.

We now start with uniform and strong order convergence for nets of orthomorphisms. For this, we need a few preparatory results. The first one is on general order continuous operators.

Lemma 3.9.1. *Let E be a Dedekind complete vector lattice, let $(T_\alpha)_{\alpha \in A}$ be a decreasing net in $\mathcal{L}_{\text{oc}}(E)^+$, and let F be an order dense vector sublattice of E . The following are equivalent:*

(1) $T_\alpha x \xrightarrow{o} 0$ in E for all $x \in F$;

(2) $T_\alpha x \xrightarrow{0} 0$ in E for all $x \in E$.

Proof. We need to show only that part (1) implies part (2). Suppose that $T_\alpha x \xrightarrow{0} 0$ in E for all $x \in F$. By passing to a tail, we may suppose that there exists a $T \in \mathcal{L}_{\text{oc}}(E)^+$ such that $T_\alpha \leq T$ for $\alpha \in \mathcal{A}$. Take $x \in E^+$. Since $(T_\alpha x)_{\alpha \in \mathcal{A}}$ is directed downwards and E is Dedekind complete, there exists a $y \in E^+$ such that $T_\alpha x \downarrow y$ in E . The order denseness of F in E implies that there exists a net $(x_\beta)_{\beta \in \mathcal{B}} \subseteq F^+$ with $x_\beta \uparrow x$ in E . For each $\alpha \in \mathcal{A}$ and $\beta \in \mathcal{B}$, we have

$$\begin{aligned} y &\leq T_\alpha x = T_\alpha(x - x_\beta) + T_\alpha x_\beta \\ &\leq T(x - x_\beta) + T_\alpha x_\beta. \end{aligned}$$

For each fixed $\beta \in \mathcal{B}$, the assumption then implies that

$$y \leq T(x - x_\beta) + \inf_{\alpha} T_\alpha x_\beta = T(x - x_\beta).$$

The order continuity of T then shows that

$$0 \leq y \leq \inf_{\beta} T(x - x_\beta) = 0,$$

and so $y = 0$. We conclude that $T_\alpha x \downarrow 0$ in E for every $x \in E^+$, and the statement in part (2) follows. \square

Proposition 3.9.2. *Let E be a Dedekind complete vector lattice, let $(T_\alpha)_{\alpha \in \mathcal{A}}$ be a decreasing net in $\text{Orth}(E)^+$, and let S be a non-empty subset of E . The following are equivalent:*

- (1) $T_\alpha s \xrightarrow{0} 0$ in E for all $s \in S$;
- (2) $T_\alpha x \xrightarrow{0} 0$ in E for all $x \in B_S$.

In particular, if E has a positive weak order unit e , then $T_\alpha x \xrightarrow{0} 0$ in E for all $x \in E$ if and only if $T_\alpha e \downarrow 0$ in E .

Proof. We need to show only that part (1) implies part (2). Take $y \in I_S^+$. There exist $s_1, \dots, s_n \in S$ and $\lambda_1, \dots, \lambda_n \geq 0$ such that $0 \leq y \leq \sum_{i=1}^n \lambda_i |s_i|$. Hence $0 \leq T_\alpha y \leq \sum_{i=1}^n \lambda_i T_\alpha |s_i| = \sum_{i=1}^n \lambda_i |T_\alpha s_i|$ for $\alpha \in \mathcal{A}$, and the assumption then implies that $T_\alpha y \downarrow 0$ in E . Since orthomorphisms preserve bands, we have $T_\alpha y \in B_S$ for all α , and the fact that B_S is an ideal of E now shows that $T_\alpha y \downarrow 0$ in B_S . It follows that $T_\alpha y \xrightarrow{0} 0$ in B_S for all $y \in I_S$. Since the restriction of each T_α to the regular vector sublattice B_S of E is again order continuous, and since I_S is an order dense vector sublattice of the vector lattice B_S , Lemma 3.9.1 implies that $T_\alpha y \xrightarrow{0} 0$ in B_S for all $y \in B_S$. The fact that B_S is a regular vector sublattice of E then yields that $T_\alpha y \xrightarrow{0} 0$ in E for all $y \in B_S$. \square

Lemma 3.9.3. *Let E be a Dedekind complete vector lattice, and let \mathcal{S} be a subset of $\text{Orth}(E)$ that is bounded above in $\mathcal{L}_{\text{ob}}(E)$. Then, for $x \in E^+$,*

$$\left(\bigvee_{T \in \mathcal{S}} T \right) x = \bigvee_{T \in \mathcal{S}} Tx.$$

Proof. Using [6, Theorem 1.67.(b)] in the second step, we see that, for $x \in E^+$,

$$\left(\bigvee_{T \in \mathcal{S}} T \right) x = \left(\bigvee_{T^\vee \in \mathcal{S}^\vee} T^\vee \right) x = \bigvee_{T^\vee \in \mathcal{S}^\vee} T^\vee x.$$

By equation (3.1), this equals

$$\bigvee_{y^\vee \in (\mathcal{S}x)^\vee} y^\vee = \bigvee_{y \in \mathcal{S}x} y = \bigvee_{T \in \mathcal{S}} Tx.$$

□

We can now establish our main result regarding uniform and strong order convergence for nets of orthomorphisms.

Theorem 3.9.4. *Let E be a Dedekind complete vector lattice, and let $(T_\alpha)_{\alpha \in \mathcal{A}}$ be a net in $\text{Orth}(E)$ that is order bounded in $\mathcal{L}_{\text{ob}}(E)$. Let S be a non-empty subset of E with $B_S = E$. The following are equivalent:*

- (1) $T_\alpha \xrightarrow{o} 0$ in $\text{Orth}(E)$;
- (2) $T_\alpha \xrightarrow{o} 0$ in $\mathcal{L}_{\text{ob}}(E)$;
- (3) $T_\alpha s \xrightarrow{o} 0$ in E for all $s \in S$;
- (4) $T_\alpha x \xrightarrow{o} 0$ in E for all $x \in E$.

In particular, when E has a weak order unit e , then $T_\alpha \xrightarrow{o} 0$ in $\mathcal{L}_{\text{ob}}(E)$ if and only if $T_\alpha e \xrightarrow{o} 0$ in E .

Before proceeding with the proof, we remark that, since $\text{Orth}(E)$ is a projection band in $\mathcal{L}_{\text{ob}}(E)$, the order boundedness of the net could equivalently have been required in $\text{Orth}(E)$.

Proof. Since the net $(T_\alpha)_{\alpha \in \mathcal{A}}$ is supposed to be order bounded in the regular vector sublattice $\text{Orth}(E)$, the equivalence of the parts (1) and (2) follows from [28, Corollary 2.12]. Lemma 3.4.1 shows that part (2) implies part (4), and evidently part (4) implies part (3). The proof will be completed by showing that part (3) implies part (1). Suppose that $T_\alpha s \xrightarrow{o} 0$ in E for all $s \in S$ or, equivalently, that $|T_\alpha||s| = |T_\alpha s| \xrightarrow{o} 0$ in E for all $s \in S$. For $\alpha \in \mathcal{A}$, set $\tilde{T}_\alpha := \bigvee_{\beta \geq \alpha} |T_\beta|$ in $\mathcal{L}_{\text{ob}}(E)$. Since Lemma 3.9.3 shows that $\tilde{T}_\alpha|s| = \bigvee_{\beta \geq \alpha} |T_\beta||s|$ for $\alpha \in \mathcal{A}$ and $s \in S$, we see that $\tilde{T}_\alpha|s| \downarrow 0$ in E for all $s \in S$. Proposition 3.9.2 then yields that $\tilde{T}_\alpha x \xrightarrow{o} 0$ for all $x \in B_{|s|} = E$. Using that $\tilde{T}_\alpha \downarrow$, it follows that $\tilde{T}_\alpha \downarrow 0$ in $\mathcal{L}_{\text{ob}}(E)$. Since $|T_\alpha| \leq \tilde{T}_\alpha$ for $\alpha \in \mathcal{A}$, we see that $|T_\alpha| \xrightarrow{o} 0$ in $\mathcal{L}_{\text{ob}}(E)$, as required. □

In view of Lemma 3.4.1, the most substantial part of Theorem 3.9.4 is the fact that the parts (3) and (4) imply the parts (1) and (2). For this to hold in general, the assumption that $(T_\alpha)_{\alpha \in \mathcal{A}}$ be order bounded is actually necessary. To see this, let Γ be an uncountable set that is supplied with the counting measure, and consider $E := \ell_p(\Gamma)$ for $1 \leq p \leq \infty$. Set

$$\mathcal{A} := \{(n, S) : n \geq 1, S \subset \Gamma \text{ is at most countably infinite}\}$$

and, for $(n_1, S_1), (n_2, S_2) \in \mathcal{A}$, say that $(n_1, S_1) \leq (n_2, S_2)$ when $n_1 \leq n_2$ and $S_1 \subseteq S_2$. For $(n, S) \in \mathcal{A}$, define $T_{(n,S)} \in \mathcal{Z}(E) = \text{Orth}(E)$ by setting

$$T_{(n,S)}x := n\chi_{\Gamma \setminus S}x$$

for all $x : \Gamma \rightarrow \mathbb{R}$ in E . Take an $x \in E$. Then the net $(T_{(n,S)}x)_{(n,S) \in \mathcal{A}}$ has a tail that is identically zero, namely $(T_{(n,S)}x)_{(n,S) \geq (1, \text{supp } x)}$. Hence $T_{(n,S)}x \xrightarrow{0} 0$ in E for all $x \in E$. We claim that $(T_{(n,S)})_{(n,S) \in \mathcal{A}}$ is not order convergent in $\text{Orth}(E)$, and not even in $\mathcal{L}_{\text{ob}}(E)$. For this, it is sufficient to show that it does not have any tail that is order bounded in $\mathcal{L}_{\text{ob}}(E)$. Suppose, to the contrary, that there exist an $n_0 \geq 1$, an at most countably infinite subset S_0 of Γ , and a $T \in \mathcal{L}_{\text{ob}}(E)$ such that $T_{(n,A)} \leq T$ for all $(n,A) \in \mathcal{A}$ with $n \geq n_0$ and $A \supseteq A_0$. As Γ is uncountable, we can choose an $x_0 \in \Gamma \setminus A_0$; we let e_{x_0} denote the corresponding atom in E . Then, in particular, $T_{(n,A_0)}e_{x_0} \leq Te_{x_0}$ for all $n \geq n_0$. Hence $Te_{x_0} \geq ne_{x_0}$ for all $n \geq n_0$, which is impossible.

We now consider uniform and strong order convergence in the case where E is a Dedekind complete Banach lattice. In that case, a version of Theorem 3.9.4 can be obtained for sequences where the order boundedness of the sequence need to be a part of the hypotheses because it is automatic; see Theorem 3.9.7, below. Our results are based on the following ordered version of the uniform boundedness principle for orthomorphisms.

Proposition 3.9.5. *Let E be a Dedekind complete Banach lattice, and let $\{T_\alpha : \alpha \in \mathcal{A}\}$ be a non-empty subset of $\text{Orth}(E)$. The following are equivalent:*

- (1) $\{T_\alpha : \alpha \in \mathcal{A}\}$ is an order bounded subset of $\mathcal{L}_{\text{ob}}(E)$;
- (2) for each $x \in E$, $\{T_\alpha x : \alpha \in \mathcal{A}\}$ is an order bounded subset of E .

As in Theorem 3.9.4, the order boundedness of the net could equivalently have been stated in $\text{Orth}(E)$.

Proof. It is trivial that part (1) implies part (2). We give two proofs for the fact that part (2) implies part (1).

The first proof is as follows. The fact that $|T_\alpha||x| = |T_\alpha x|$ implies that we may suppose that the T_α are positive. Suppose, to the contrary, that $\{T_\alpha : \alpha \in \mathcal{A}\}$ is not an order bounded subset of $\mathcal{L}_{\text{ob}}(E)$. Using that $\text{Orth}(E) = \mathcal{Z}(E)$, it is easy to see that, for every $n \geq 1$, there exists an $\alpha_n \in \mathcal{A}$ such that $(T_{\alpha_n} - 2^n I)^+ > 0$. For $n \geq 1$, we let B_n be the band generated by $(T_{\alpha_n} - 2^n I)^+$ in $\text{Orth}(E)$, and we let P_n be the corresponding non-zero order projection onto B_n . According to Proposition 3.6.7, there exists a unique order projection P_n in E such that $P_n S = P_n S$ for all $S \in \text{Orth}(E)$. Furthermore, $T_{\alpha_n} \geq 2^n P_n$. As $P_n \neq 0$, we can choose an $x_n \in E^+$ such that $\|P_n x_n\| = 1/2^n$. Since $\bigvee_{n=1}^m P_n x_n \leq \sum_{n=1}^\infty P_n x_n$ for all $m \geq 1$, we can set $e := \bigvee_{n=1}^\infty P_n x_n \in E^+$. By assumption, we can choose an upper bound x of $\{T_{\alpha_n} e : n \geq 1\}$ in E^+ . Then

$$x \geq T_{\alpha_n} e \geq 2^n P_n e \geq 2^n P_n (P_n x_n) = 2^n P_n x_n$$

for $n \geq 1$. Again by assumption, we can choose an upper bound y of $\{T_{\alpha_n} x : n \geq 1\}$ in E^+ , and then

$$y \geq T_{\alpha_n} x \geq 2^n P_n x \geq 2^n P_n (2^n P_n x_n) = 4^n P_n x_n$$

for $n \geq 1$. This implies that $\|y\| \geq 4^n \|P_n x_n\| = 2^n$ for all n . This contradiction completes the first proof.

The second proof, which uses somewhat ‘heavier’ auxiliary results, is as follows. The fact that the T_α are pointwise order bounded implies that they are pointwise norm bounded. Hence, by the uniform boundedness principle, the T_α are bounded in the uniform norm on the bounded operators on E . Since they are in $\text{Orth}(E) = \mathcal{Z}(E)$, where (see Section 3.6) the operator norm agrees with the order unit norm with respect to the strong order unit I of $\mathcal{Z}(E)$, the T_α are also order bounded in $\mathcal{Z}(E)$. \square

As a side result, we note the following consequence of Proposition 3.9.5. It is an ordered analogue of the familiar result for a sequence of bounded operators on a Banach space.

Corollary 3.9.6. *Let E be a Dedekind complete Banach lattice, and let $(T_n)_{n=1}^\infty$ be a sequence in $\text{Orth}(E)$. Suppose that the sequence $(T_n x)_{n=1}^\infty$ is order convergent in E for all $x \in E$. Then $\{T_n : n \geq 1\}$ is an order bounded subset of $\mathcal{L}_{\text{ob}}(E)$. For $x \in E$, define $T : E \rightarrow E$ by setting*

$$Tx := o - \lim_{n \rightarrow \infty} T_n x.$$

Then $T \in \text{Orth}(E)$.

Proof. It is clear that T is linear. Since order convergent sequences are order bounded, Proposition 3.9.5 shows that there exist an $S \in \text{Orth}(E)$ such that $|T_n| \leq |S|$ for $n \geq 1$. As $\text{Orth}(E) = \mathcal{Z}(E)$, there exists a $\lambda \geq 0$ such that $|T_n| \leq \lambda I$ for $n \geq 1$. Using equation (3.2), one then easily sees that $|Tx| \leq \lambda|x|$ for $x \in E$. Hence $T \in \mathcal{Z}(E) = \text{Orth}(E)$. \square

Using Theorem 3.9.4 and the order boundedness statement in Corollary 3.9.6, the following is easily established. As announced above, there is no order boundedness in the hypotheses.

Theorem 3.9.7. *Let E be a Dedekind complete Banach lattice, and let $(T_n)_{n=1}^\infty$ be a sequence in $\text{Orth}(E)$. Let S be a non-empty subset of E such that $I_S = E$. The following are equivalent:*

- (1) $T_n \xrightarrow{o} 0$ in $\text{Orth}(E)$;
- (2) $T_n \xrightarrow{o} 0$ in $\mathcal{L}_{\text{ob}}(E)$;
- (3) $T_n s \xrightarrow{o} 0$ in E for all $s \in S$;
- (4) $T_n x \xrightarrow{o} 0$ in E for all $x \in E$.

In particular, when E has a strong order unit e , then $T_n \xrightarrow{o} 0$ in $\text{Orth}(E)$ if and only if $T_n e \xrightarrow{o} 0$ in E .

Remark 3.9.8. In Theorem 3.9.7, the condition that $I_S = E$ cannot be relaxed to $B_S = E$. To see this, we choose $E := c_0$ and set $e := \bigvee_{n \geq 1} e_i / i^2$, where $(e_i)_{i=1}^\infty$ is the standard unit basis of E . It is clear that $B_e = E$. For $n \geq 1$, there exists a unique $T_n \in \text{Orth}(E)$ such that, for $i \geq 1$, $T_n e_i = n e_i$ when $i = n$, and $T_n e_i = 0$ when $i \neq n$. It is clear that $T_n e \xrightarrow{o} 0$ in E . However, a consideration of $T_n(\bigvee_{i \geq 1} e_i / i)$ for $n \geq 1$ shows that $(T_n)_{n=1}^\infty$ fails to be order bounded in $\text{Orth}(E)$, hence cannot be order convergent in $\text{Orth}(E)$.

We continue our comparison of uniform and strong convergence structures on the orthomorphisms by considering unbounded order convergence. In that case, the result is as follows.

Theorem 3.9.9. *Let E be a Dedekind complete vector lattice, and let $(T_\alpha)_{\alpha \in A}$ be a net in $\text{Orth}(E)$. Let S be a non-empty subset of E such that $B_S = E$. The following are equivalent:*

- (1) $T_\alpha \xrightarrow{\text{uo}} 0$ in $\text{Orth} E$;
- (2) $T_\alpha \xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E)$;
- (3) $T_\alpha s \xrightarrow{\text{uo}} 0$ in E for all $s \in S$;
- (4) $T_\alpha x \xrightarrow{\text{uo}} 0$ in E for all $x \in E$.

In particular, when E has a weak order unit e , then $T_\alpha \xrightarrow{\text{uo}} 0$ in $\text{Orth}(E)$ if and only if $T_\alpha e \xrightarrow{\text{uo}} 0$ in E .

Proof. Since $\text{Orth}(E)$ is a regular vector sublattice of $\mathcal{L}_{\text{ob}}(E)$, the equivalence of the parts (1) and (2) is clear from [28, Theorem 3.2]

We prove that part (2) implies part (4). Suppose that $T_\alpha \xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E)$, so that, in particular, $|T_\alpha| \wedge I \xrightarrow{0} 0$ in $\mathcal{L}_{\text{ob}}(E)$. Take $x \in E$. Using equation (3.1) in the second step, and Lemma 3.4.1 in the third, we have

$$(|T_\alpha||x|) \wedge |x| = (|T_\alpha||x|) \wedge (I|x|) = (|T_\alpha| \wedge I)|x| \xrightarrow{0} 0.$$

Since the net $(|T_\alpha||x|)_{\alpha \in A}$ is contained in the band $B_{|x|}$, it now follows from [20, Proposition 7.4] that $|T_\alpha||x| \xrightarrow{\text{uo}} 0$ in E . As $|T_\alpha||x| = |T_\alpha x|$, we conclude that $T_\alpha x \xrightarrow{\text{uo}} 0$ in E .

It is clear that part (4) implies part (3).

We prove that part (3) implies part (2). Suppose that $T_\alpha s \xrightarrow{\text{uo}} 0$ in E for all $s \in S$, so that also $|T_\alpha||s| = |T_\alpha s| \xrightarrow{\text{uo}} 0$ in E for $s \in S$. Using equation (3.1) again, we have

$$(|T_\alpha| \wedge I)|s| = (|T_\alpha||s|) \wedge |s| \xrightarrow{0} 0$$

in E for all $s \in S$. In view of the order boundedness of $(|T_\alpha| \wedge I)_{\alpha \in A}$, Theorem 3.9.4 then yields that $|T_\alpha| \wedge I \xrightarrow{0} 0$ in $\mathcal{L}_{\text{ob}}(E)$. As I is a weak order unit of $\text{Orth}(E)$, [29, Lemma 3.2] (or the more general [20, Proposition 7.4]) shows that $T_\alpha \xrightarrow{\text{uo}} 0$ in $\mathcal{L}_{\text{ob}}(E)$. \square

We now consider uniform and strong convergence of nets of orthomorphisms for the Hausdorff uo-Lebesgue topology. Let E be a Dedekind complete vector lattice. Suppose that E admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_E$. We recall from Theorem 3.8.3 that $\text{Orth}(E)$ then also admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_{\text{Orth}(E)}$, and that this topology equals $u_{\text{Orth}(E)}\text{ASOT}_E \widehat{\tau}_E$. Furthermore, for a net $(T_\alpha)_{\alpha \in A}$ in $\text{Orth}(E)$, we have that $T_\alpha \xrightarrow{\widehat{\tau}_{\text{Orth}(E)}} 0$ if and only if $|T_\alpha x| \wedge |Tx| \xrightarrow{\widehat{\tau}_E} 0$ for all $T \in \text{Orth}(E)$ and $x \in E$.

We need two preparatory results.

Lemma 3.9.10. *Let E be a vector lattice that admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_E$. Suppose that E has a positive weak order unit e . Let $(x_\alpha)_{\alpha \in \mathcal{A}}$ be a net in E . Then $x_\alpha \xrightarrow{\widehat{\tau}_E} 0$ in E if and only if $|x_\alpha| \wedge e \xrightarrow{\widehat{\tau}_E} 0$ in E .*

Proof. We need to show only the “if”-part. Suppose that $|x_\alpha| \wedge e \xrightarrow{\widehat{\tau}_E} 0$ in E . For each $x \in E$, there exists a net $(y_\beta)_{\beta \in \mathcal{B}}$ in I_e such that $y_\beta \xrightarrow{0} x$, and then certainly $y_\beta \xrightarrow{\widehat{\tau}_E} x$. Hence $\overline{I_e}^{\widehat{\tau}_E} = E$. An appeal to [44, Proposition 9.8] then shows that $x_\alpha \xrightarrow{u_E \widehat{\tau}_E} 0$. Since $u_E \widehat{\tau}_E = \widehat{\tau}_E$, we are done. \square

Our second preparatory result is in the same vein as Proposition 3.7.2.

Lemma 3.9.11. *Let E be a vector lattice with the principal projection property that admits a (not necessarily Hausdorff) o-Lebesgue topology τ_E , and let $(T_\alpha)_{\alpha \in \mathcal{A}}$ be a net in $\text{Orth}(E)$. Let S be a non-empty subset of E such that $B_S = E$. Suppose that $T_\alpha s \xrightarrow{\tau_E} 0$ for all $s \in S$. Then $T_\alpha x \xrightarrow{u_E \tau_E} 0$ for all $x \in E$.*

Proof. Using equation (3.2), it follows easily that $T_\alpha x \xrightarrow{\tau_E} 0$ for all $x \in I_S$. Take an $x \in E$, and let U be a solid τ_E -neighbourhood U of 0. Choose a τ_E -neighbourhood V of 0 such that $V + V \subseteq U$. There exists a net $(x_\beta)_{\beta \in \mathcal{B}}$ in I_S such that $x_\beta \xrightarrow{0} x$ in E , and then we can choose a $\beta_0 \in \mathcal{B}$ such that $|x - x_{\beta_0}| \in V$. As $|T_\alpha| |x_{\beta_0}| = |T_\alpha x_{\beta_0}| \xrightarrow{\tau_E} 0$, there exists an $\alpha_0 \in \mathcal{A}$ such that $|T_\alpha| |x_{\beta_0}| \in V$ for all $\alpha \geq \alpha_0$. For all $\alpha \geq \alpha_0$, we then have

$$\begin{aligned} 0 &\leq (|T_\alpha x|) \wedge |x| = (|T_\alpha| \wedge I)|x| \\ &\leq (|T_\alpha| \wedge I)|x_{\beta_0}| + (|T_\alpha| \wedge I)|x - x_{\beta_0}| \\ &\leq |T_\alpha| |x_{\beta_0}| + |x - x_{\beta_0}| \\ &\in V + V \subseteq U. \end{aligned}$$

As U is solid, we see that $(|T_\alpha x|) \wedge |x| \in U$ for $\alpha \geq \alpha_0$, and we conclude that $(|T_\alpha x|) \wedge |x| \xrightarrow{\tau_E} 0$. Since $|T_\alpha x| \in B_{|x|}$ for $\alpha \in \mathcal{A}$, it then follows from [44, Proposition 9.8] that $|T_\alpha x| \wedge |y| \xrightarrow{\tau_E} 0$ in E for all $y \in B_{|x|}$. As $B_{|x|}$ is a projection band in E , this holds, in fact, for all $y \in E$. \square

Theorem 3.9.12. *Let E be a Dedekind complete vector lattice. Suppose that E admits a (necessarily unique) Hausdorff uo-Lebesgue topology $\widehat{\tau}_E$, so that $\text{Orth}(E)$ and $\mathcal{L}_{\text{ob}}(E)$ also admit (necessarily unique) Hausdorff uo-Lebesgue topologies $\widehat{\tau}_{\text{Orth}(E)}$ and $\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}$, respectively. Let $(T_\alpha)_{\alpha \in \mathcal{A}}$ be a net in $\text{Orth}(E)$. Let S be a non-empty subset S of E such that $B_S = E$. The following are equivalent:*

- (1) $T_\alpha \xrightarrow{\widehat{\tau}_{\text{Orth}(E)}} 0$ in $\text{Orth}(E)$;
- (2) $T_\alpha \xrightarrow{\widehat{\tau}_{\mathcal{L}_{\text{ob}}(E)}} 0$ in $\mathcal{L}_{\text{ob}}(E)$;
- (3) $T_\alpha s \xrightarrow{\widehat{\tau}_E} 0$ in E for all $s \in S$;
- (4) $T_\alpha x \xrightarrow{\widehat{\tau}_E} 0$ in E for all $x \in E$.

In particular, when E has a weak order unit e , then $T_\alpha \xrightarrow{\widehat{\tau}_{\text{Orth}(E)}} 0$ in $\text{Orth}(E)$ if and only if $T_\alpha e \xrightarrow{\widehat{\tau}_E} 0$ in E .

Proof. The equivalence of the parts (1) and (2) follows from the final part of Theorem 3.4.3.

We prove that part (1) implies part (4). Suppose that $T_\alpha \xrightarrow{\widehat{\tau}_{\text{Orth}(E)}} 0$ in $\text{Orth}(E)$. Take an $x \in E$. Then certainly $|T_\alpha x| \wedge |x| = |T_\alpha x| \wedge |Ix| \xrightarrow{\widehat{\tau}_E} 0$. The net $(T_\alpha x)_{\alpha \in \mathcal{A}}$ is contained in the band $B_{|x|}$. Since, by [44, Proposition 5.12], the regular vector sublattice $B_{|x|}$ of E also admits a (necessarily unique) Hausdorff uo-Lebesgue topology (namely, the restriction of $\widehat{\tau}_E$ to $B_{|x|}$), it then follows from Lemma 3.9.10 that $T_\alpha x \xrightarrow{\widehat{\tau}_E} 0$ in E .

We prove that part (4) implies part (1). Suppose that $T_\alpha x \xrightarrow{\widehat{\tau}_E} 0$ for all $x \in E$. Since $\widehat{\tau}_E$ is locally solid, we then also have $|T_\alpha x| \wedge |Tx| \xrightarrow{\widehat{\tau}_E} 0$ for all $T \in \text{Orth}(E)$ and $x \in E$. Hence $T_\alpha \xrightarrow{\widehat{\tau}_{\text{Orth}(E)}} 0$ in $\text{Orth}(E)$.

It is clear that part (4) implies part (3).

Since $u_E \widehat{\tau}_E = \widehat{\tau}_E$, Lemma 3.9.11 shows that part (3) implies part (4). \square

