



Universiteit
Leiden
The Netherlands

Cold gas in distant galaxies

Boogaard, L.A.

Citation

Boogaard, L. A. (2021, February 25). *Cold gas in distant galaxies*. Retrieved from <https://hdl.handle.net/1887/3147175>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3147175>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/3147175> holds various files of this Leiden University dissertation.

Author: Boogaard, L.A.

Title: Cold gas in distant galaxies

Issue date: 2021-02-25

Cold gas in distant galaxies

Cold gas in distant galaxies

Koud gas in verre sterrenstelsels

Proefschrift

ter verkrijging van
de graad van Doctor aan de Universiteit Leiden,
op gezag van Rector Magnificus prof. dr. ir. H. Bijl,
volgens besluit van het College voor Promoties
te verdedigen op donderdag 25 februari 2021
klokke 16.15 uur

door

Lein Adriaan Boogaard

geboren te Oegstgeest
in 1992

Promotor: Prof. dr. P. P. van der Werf
Co-promotor: Dr. R.J. Bouwens

Promotiecommissie: Prof. dr. H. J. A. Röttgering
Prof. dr. J. Schaye
Prof. dr. S. Viti
Dr. J. A. Hodge
Dr. J. Brinchmann
Dr. F. Walter
Prof. dr. I. R. Smail

Universidade do Porto (Portugal)
MPIA, Heidelberg (Germany)
Durham University (UK)

Voor mijn ouders

&

Voor Elisabeth

Copyright © 2021 L. A. Boogaard

Printed by: Gildeprint

Cover design: Arjen Wiersma

Cover images: The Hubble Ultra Deep Field, *credit: NASA, ESA, and S. Beckwith (STScI) and the HUDF Team (front)*, and a 3D rendering of the ALMA Spectroscopic Survey of the HUDF 3 mm datacube (*back*).

An electronic copy of this thesis can be found at <https://openaccess.leidenuniv.nl>

ISBN 978 94 6419 120 2

Contents

1	Introduction	1
1.1	From effect to cause	2
1.2	The theory	4
1.2.1	Cosmology, galaxy formation and the baryon cycle	4
1.2.2	Star formation and the cold interstellar medium	6
1.2.3	The light from galaxies across the electromagnetic spectrum	11
1.3	The instruments	12
1.3.1	Multi Unit Spectroscopic Explorer (MUSE)	13
1.3.2	Atacama Large Millimeter Array (ALMA)	13
1.3.3	Other facilities	13
1.4	The state of the art	14
1.4.1	Star formation in galaxies across cosmic time	14
1.4.2	Molecular gas in distant galaxies	15
1.4.3	The need for molecular deep fields	17
1.5	The ALMA Spectroscopic Survey of the HUDF	18
1.5.1	Motivation	18
1.5.2	Observing strategy	19
1.5.3	Data products: two cubes and two images	19
1.6	The thesis	23
1.6.1	This thesis	23
1.6.2	Related science with ASPECS	25
1.7	The future	26
1.7.1	The cosmic baryon cycle	26
1.7.2	Science and facilities	27
2	Constraining the low-mass end of the M_*-SFR relation at $z < 1$	29
2.1	Introduction	30
2.2	Observations and methods	32
2.2.1	Observations, data reduction, and spectral line fitting	33
2.2.2	Sample selection	34
2.2.3	Stellar masses	37
2.2.4	Star formation rates	38

2.3	Consistency of SFR indicators	39
2.4	Bayesian model	42
2.4.1	Definition	42
2.4.2	Execution	44
2.4.3	Model and data limitations	45
2.5	Star formation sequence	45
2.5.1	Global sample	45
2.5.2	Low-mass sample ($\log M_*[M_\odot] < 9.5$)	49
2.5.3	The effect of redshift bins (2D)	49
2.6	Discussion	51
2.6.1	Comparison with the literature	51
2.6.2	The MS slope — a quantitative comparison to models	55
2.6.3	Implications of a shallow slope	57
2.7	Summary and conclusions	58
Appendix 2.A	Simulations	60
2.A.1	Selection function and completeness	60
2.A.2	Transformation	61
3	The nature and physical properties of gas-mass selected galaxies	65
3.1	Introduction	66
3.2	Observations	67
3.2.1	ALMA Spectroscopic Survey	67
3.2.2	MUSE HUDF Survey	68
3.2.3	Multi-wavelength data (UV–radio) and MAGPHYS	70
3.2.4	X-ray photometry	71
3.3	The ASPECS-LP sample	72
3.3.1	Identification of the line search sample	72
3.3.2	Additional sources with MUSE redshift priors at $z < 2.9$	75
3.3.3	Full sample redshift distribution	78
3.4	Physical properties	79
3.4.1	Star formation rates from MAGPHYS and [O II]	79
3.4.2	Metallicities	81
3.4.3	Molecular gas properties	81
3.5	Results: Global sample properties	83
3.5.1	Stellar mass and SFR distributions	83
3.5.2	AGN fraction	85
3.5.3	Obscured and unobscured star formation rates	85
3.5.4	Metallicities at $1.0 < z < 1.42$	86
3.6	Discussion	87
3.6.1	Sensitivity limit to molecular gas reservoirs	87
3.6.2	Molecular gas across the galaxy main sequence	89
3.6.3	Evolution of molecular gas content in galaxies	92
3.7	Summary	93
Appendix 3.A	Source description and redshift identifications	95

Appendix 3.B	MAGPHYS fits for all CO-detected galaxies	104
4	CO excitation, [C I] and ISM conditions in galaxies at $z = 1 - 3$	105
4.1	Introduction	106
4.2	Observations and ancillary data	108
4.2.1	ALMA Spectroscopic Survey Data Reduction	108
4.2.2	ASPECS Sample	110
4.2.3	Very Large Array Observations (VLASPECS)	111
4.2.4	Multi-wavelength data and SED fitting	112
4.3	Methods	112
4.3.1	Spectral line analysis	112
4.3.2	Deriving line luminosities and molecular gas masses	115
4.4	Results	115
4.4.1	Observed emission lines from CO and [C I]	115
4.5	CO excitation	117
4.5.1	Individual sources	117
4.5.2	Stacked line fluxes	120
4.5.3	LVG modeling	122
4.5.4	Dust-continuum versus low- J CO	125
4.6	Atomic carbon	129
4.6.1	Atomic carbon abundances	129
4.6.2	PDR modeling	132
4.7	Discussion	134
4.7.1	Modest excitation in mid- J lines at $z = 1.0 - 1.6$	134
4.7.2	Increasing excitation with redshift	136
4.7.3	The low- J excitation	138
4.7.4	Broader implications of the flux-limited survey	140
4.7.5	Implications for the cosmic molecular gas density	140
4.8	Summary and Conclusions	141
Appendix 4.A	Similar widths for the low- J and high- J CO lines	143
Appendix 4.B	Spectral line fits	144
5	Line-luminosity functions and the cosmic density of molecular gas	153
5.1	Introduction	154
5.2	Observations	156
5.2.1	ALMA data	156
5.2.2	Ancillary data	158
5.3	Analysis and Results	158
5.3.1	Line search at 1.2 mm	158
5.3.2	Line fluxes	160
5.3.3	Line identification and redshifts	160
5.3.4	Line luminosities and molecular gas masses	163
5.3.5	Luminosity functions and ρ_{H_2}	166
5.4	Discussion	166

5.4.1	CO luminosity functions	166
5.4.2	[C I] and [C II] luminosity functions	169
5.4.3	ρ_{H_2} vs redshift	170
5.5	Conclusions	172
Appendix 5.A	Tabulated luminosity functions	173
Appendix 5.B	Cosmic variance	175
Appendix 5.C	Identification of line candidates without near-infrared counterparts	178
6	The average molecular gas content of star-forming galaxies at $z = 3 - 4$	181
6.1	Introduction	182
6.2	Observations and sample selection	183
6.2.1	Parent sample selection and physical properties	183
6.2.2	Measurement of systemic redshifts	185
6.2.3	Final systemic redshift sample	188
6.3	Results	190
6.3.1	Velocity offsets	190
6.3.2	ALMA Stacking	191
6.4	Discussion	194
6.4.1	Molecular gas masses	194
6.4.2	Low metallicity driving a high molecular gas mass-to-light ratio . .	196
6.4.3	Contribution to the cosmic molecular gas density	201
6.4.4	Implications for observing cold gas in low metallicity galaxies at high redshift	201
6.5	Summary and conclusions	203
Appendix 6.A	Table	205
Appendix 6.B	Spectra	208
	Bibliography	211
	Publication list	221
	Nederlandse samenvatting	227
	Curriculum Vitae	233
	Acknowledgements	235