

Measures and matching for number systems Maggioni, M.

Citation

Maggioni, M. (2021, March 4). *Measures and matching for number systems*. Retrieved from https://hdl.handle.net/1887/3147169

Version:	Publisher's Version
License:	<u>Licence agreement concerning inclusion of doctoral thesis in the</u> <u>Institutional Repository of the University of Leiden</u>
Downloaded from:	https://hdl.handle.net/1887/3147169

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <u>https://hdl.handle.net/1887/3147169</u> holds various files of this Leiden University dissertation.

Author: Maggioni, M. Title: Measures and matching for number systems Issue Date: 2021-03-04

Chapter 5

Matching and Measure for Random Systems

This chapter is based on: [DKM20].

Abstract

We extend the notion of matching for interval maps to random matching for pseudoskew products on the interval. For a certain family of piecewise affine interval random systems the property of random matching implies that any invariant density is piecewise constant. Furthermore, we introduce a one-parameter family of random dynamical systems that produce signed binary expansions of numbers in the interval [-1, 1]. We provide results on minimal weight expansions by proving that the frequency of the digit 0 in the associated signed binary expansions never exceeds 1/2. We do this by showing that the family has random matching for Lebesgue almost every parameter.

§5.1 Motivation and context

The research on optimal algorithms that raise elements of a group into some power has a long and rich history. It is a matter of fact that the computation of powers of elements in a group is the basis of many public key cryptosystems, where the group chosen is either the multiplicative group of a finite field \mathbb{F}_q or the group of points on an elliptic curve. The optimality of the algorithms refers to the ability of computing high powers in a short amount of time. One way to reduce the time complexity is given by the so-called binary method, introduced in [K69], and based on the binary expansion of the power. More precisely, if x is an element of a given group, and $a = \sum_{i=0}^{n} d_i 2^i \in \mathbb{N}$ for some digits $d_i \in \{0, 1\}$, then

$$x^a = \prod_{i=0}^n x^{d_i 2^i},$$

and the power x^a is computed by taking the product of the repeated squarings. While the number of squarings is given by the length n of the binary expansion of a, the number of multiplications equals the number of non-zero bits d_i in the expansion or its Hamming weight. Clearly, the lower it is, the less multiplications are required and the faster the algorithm is. To increase the number of zero bits, [B51] introduced a signed binary representation, i.e., a binary representation with digits in the set $\{-1, 0, 1\}$. This signed binary method was later adopted in several methods in elliptic cryptosystems, see e.g. [CMO98, HP06] and the references therein.

For any fixed integer a, its ordinary binary representation with digits $\{0, 1\}$ is uniquely determined, but this is not the case for the signed one, with digits in $\{-1, 0, 1\}$. In fact, each integer has infinitely many signed binary representations, which led to the study of algorithms that choose the ones with minimal weight (see e.g. [MO90, KT93, LK97]). The authors of [GH06] showed that typically a number has several signed binary representations with minimal weight, but already in the 1960's Reitwiesner proved in [R60] that the signed representation is unique when adding the constraint $d_i d_{i+1} = 0$. Such representations are usually known as *signed separated binary expansions*, or SSB for short. In [DKL06] it is shown how to obtain SSB expansions through the binary odometer and a three state Markov chain. Furthermore, in [DKL06] the set $K := \{d_1 d_2 \ldots \in \{-1, 0, 1\}^{\mathbb{N}} : \forall i \in \mathbb{N}, d_i d_{i+1} = 0\}$ is introduced as a compactification of \mathbb{Z} . The authors identify the set K, endowed with the left shift σ , with the map $S(x) = 2x \mod \mathbb{Z}$ on the interval $[-\frac{2}{3}, \frac{2}{3}]$, through the conjugation

$$\psi(d_1 d_2 \dots) = \sum_{k=1}^{\infty} \frac{d_k}{2^k}.$$
(5.1)

The dynamical viewpoint allows them to obtain metric properties of the system (K, σ) , such as a σ -invariant measure, the maximal entropy and the frequency of 0 in typical expansions.

In [DK17] this dynamical approach was further developed by considering a family of symmetric doubling maps $\{S_{\alpha} : [-1,1] \rightarrow [-1,1]\}_{\alpha \in [1,2]}$ defined by $S_{\alpha}(x) = 2x - d\alpha$

and

$$d = \begin{cases} -1, & \text{if } x \in [-1, -\frac{1}{2}) \\ 0, & \text{if } x \in [-\frac{1}{2}, \frac{1}{2}], \\ 1, & \text{if } x \in (\frac{1}{2}, 1]. \end{cases}$$

The map S from [DKL06], producing SSB expansions, is then easily identified with the map $S_{\frac{3}{2}}$, through the conjugacy $x \mapsto \frac{3}{2}x$. For each $\alpha \in [1,2]$ iterations of S_{α} produce signed binary expansions of the form $x = \sum_{k=1}^{\infty} \frac{d_k}{2^k}$ with $d_k \in \{-1,0,1\}$ for each number $x \in [-1,1]$. The authors of [DK17] showed that the frequency of 0 in such expansions depends continuously on the parameter α and it takes its maximum value $\frac{2}{3}$, corresponding to the minimum Hamming weight of $\frac{1}{3}$, precisely for α in the subinterval $[\frac{6}{5}, \frac{3}{2}]$. It follows that the SSB expansion of integers produces sequences where typically only $\frac{1}{3}$ of the digits is different from zero. The results from [DK17] are achieved by finding a detailed description of the invariant probability density f_{α} of S_{α} for each value of α and then explicitly computing the frequency of the digit 0 using Birkhoff's Ergodic Theorem. The fact that the family $\{S_{\alpha}\}$ exhibits the dynamical phenomenon of matching was essential for these results.

In this chapter we consider signed binary expansions in the framework of random dynamical systems. The advantage of random systems in this context is that a single random system produces many more number expansions per number than a deterministic map, allowing one to study the properties of many expansions simultaneously. See e.g. [DK03, DdV05, DdV07, DK07, KKV17, DO18] for the use of random systems in the study of different types of number expansions. We will introduce a family of random systems $\{R_{\alpha}\}_{\alpha\in[1,2]}$, called random symmetric doubling maps, such that each element R_{α} produces for typical numbers in the interval [-1,1] infinitely many different signed binary expansions. This is contrary to the map S_{α} , which produces a unique signed binary expansion for each number in [-1,1]. Our main result for the family $\{R_{\alpha}\}_{\alpha\in[1,2]}$ is that the frequency of the digit 0 in typical signed binary expansions produced by any of the maps R_{α} is at most $\frac{1}{2}$, and therefore the Hamming weight is at least $\frac{1}{2}$. This reinforces the result from [DK17] that the maps S_{α} with $\alpha \in \left[\frac{6}{5}, \frac{3}{2}\right]$ perform best in terms of minimal weight.

We obtain this result from Birkhoff's Ergodic Theorem after gathering detailed knowledge on the invariant probability densities of the random maps R_{α} . We first express these densities as infinite sums of indicator functions using the algebraic procedure from [KM18]. To compute the frequency of 0 we need to evaluate the Lebesgue integral of these densities over part of the domain and therefore we convert the infinite sums into finite sums. For this we introduce a random version of the dynamical concept of matching that is available for one-dimensional systems (see e.g. [NN08, DKS09, BCIT13, BSORG13, BCK17, BCMP18, CIT18, CM18, KLMM20]). Our definition of random matching properly extends the one-dimensional notion of matching and we illustrate the concept with examples of random continued fraction maps and random generalised β -transformations. We show that under certain mild conditions, if a random system of piecewise affine maps defined on the same interval has random matching, then any invariant probability density of the system is piecewise constant. The precise formulation of this statement and the conditions are given in the next section. Finally, we use this random matching property to show that for Lebesgue almost all parameters α the invariant density of the random systems R_{α} , producing signed binary expansions, is in fact piecewise constant.

This chapter is outlined as follows. The second section is devoted to random matching for random systems defined on an interval. We first recall some preliminaries on invariant measures for random interval maps. We then define the notion of random matching and state and prove the result about densities of random systems of piecewise affine maps with matching. We also discuss the examples of random continued fraction transformations and random generalised β -transformations. In the third section we introduce and discuss the family $\{R_{\alpha}\}$ of random symmetric doubling maps and the corresponding signed binary expansions. We prove that R_{α} has random matching for Lebesgue almost all $\alpha \in [1, 2]$. We also provide a full description of the *matching intervals*, i.e., intervals of parameters that exhibit comparable matching behaviour, and describe the invariant densities of the maps R_{α} . Finally we prove that typically the frequency of the digit 0 in the signed binary expansions produced by R_{α} does not exceed $\frac{1}{2}$ for any parameter α .

§5.2 Random matching

Matching is a dynamical phenomenon observed in certain families of piecewise smooth interval maps. Recall that, if $T: I \to I$ is such a map on an interval I of real numbers, then we say that T has *matching* if for every discontinuity point c of T or of the derivative T' the orbits of the left and right limits $T^k(c^-) = \lim_{x \to c} T^k(x)$ and $T^k(c^+) = \lim_{x \to c} T^k(x)$ eventually meet, i.e., if for each c there exist positive constants $M = M_c$ and $Q = Q_c$, such that

$$T^{M}(c^{-}) = T^{Q}(c^{+}). (5.2)$$

T is then said to have *strong matching* if, moreover, the orbits of the left and right limits have equal one-sided derivatives at the moment they meet, i.e., if besides (5.2) it also holds that

$$(T^M)'(c^-) = (T^Q)'(c^+).$$
(5.3)

It was proven in [BCMP18, Theorem 1.2] (see also Remark 1.3 in [BCMP18]) that for any piecewise smooth T with strong matching, any invariant probability measure μ that is absolutely continuous with respect to the Lebesgue measure has a piecewise smooth density. For continued fraction transformations (as in [NN08, DKS09, KLMM20] for example) it seems that matching is sufficient to guarantee the existence of a piecewise smooth density (since this is sufficient to construct a natural extension with finitely many pieces). The strong matching condition then enforces some stability in the matching behaviour of certain one-parameter families of continued fraction maps, which becomes visible in the appearance of so called *matching intervals* in the parameter space: If such a family has strong matching for one parameter, then one can find an interval of parameters around it, such that all

the corresponding transformations have matching in the same number of steps and with comparable orbits.

In this section we extend the above definitions of matching and strong matching to random dynamical systems. Let $\Omega \subseteq \mathbb{N}$ be the index set of the available maps, so we have a collection of transformations $\{T_j : I \to I\}_{j \in \Omega}$ defined on the same interval $I \subseteq \mathbb{R}$ at our disposal. Let $\sigma : \Omega^{\mathbb{N}} \to \Omega^{\mathbb{N}}$ be the left shift on one-sided sequences. Recall by Definition 1.2.8 that the random map or pseudo-skew product $R : \Omega^{\mathbb{N}} \times I \to \Omega^{\mathbb{N}} \times I$ is defined by

$$R(\omega, x) = (\sigma(\omega), T_{\omega_1} x).$$

Let $\mathbf{p} = (p_j)_{j \in \Omega}$ be a positive probability vector, representing the probabilities with which we choose the maps T_j . Consider the product measure $m_{\mathbf{p}} \times \mu_{\mathbf{p}}$ from Definition 1.2.9, for $m_{\mathbf{p}}$ the **p**-Bernoulli measure on $\Omega^{\mathbb{N}}$ and $\mu_{\mathbf{p}}$ a probability measure on Ithat is absolutely continuous with resepct to the Lebesgue measure λ , and denote its density by $f_{\mathbf{p}}$. Here we call $\mu_{\mathbf{p}}$ a stationary measure and $f_{\mathbf{p}}$ an invariant density for R.

In the literature there exist various sets of conditions under which the existence of such an invariant measure is guaranteed. See for example [M85, P84, B00, GB03, BG05, I12]. Here we explicitly mention a special case of the conditions by Inoue from [I12] which are simple to state and suit our purposes in the next sections. Assume that the following three conditions hold:

(a1) There is a finite or countable interval partition $\{I_i\}_{i \in \Lambda}$, such that each map T_j is C^1 and monotone on the interior of each interval I_i .

Let C denote the set of all boundary points of the intervals I_i that are in the interior of I. We choose the collection $\{I_i\}$ as small as possible, so that C contains precisely those points that are a critical point of T_j or T'_j for at least one $j \in \Omega$. We call elements $c \in C$ critical points for the corresponding random system R.

- (a2) The random system R is expanding on average, i.e., there exists a constant $0 < \rho < 1$, such that $\sum_{j \in \Omega} \frac{p_j}{|T'_i(x)|} \leq \rho$ holds for each $x \in I \setminus C$.
- (a3) For each $j \in \Omega$ the map

$$x \mapsto \begin{cases} \frac{p_j}{|T'_j(x)|} & \text{if } x \neq c, \\ 0 & \text{otherwise,} \end{cases}$$

is of bounded variation.

It then follows from [I12, Theorem 5.2] that an invariant measure for R of the form $m_{\mathbf{p}} \times \mu_{\mathbf{p}}$ exists. Let \mathcal{R} denote the class of random maps R that satisfy these conditions. We will define random matching for maps in \mathcal{R} , but first we recall some notation on sequences and strings.

For each k > 0 the set $\Omega^k = \{\mathbf{u} = u_1 \cdots u_k : u_i \in \Omega, 1 \le i \le k\}$ is the set of all k-strings of elements in Ω . We let $\Omega^0 = \{\epsilon\}$, with ϵ the empty string. For a finite string \mathbf{u} let $|\mathbf{u}|$ denote its length, i.e., $|\mathbf{u}| = k$ if $\mathbf{u} \in \Omega^k$. Also, for $1 \le n \le k$ we

let $\mathbf{u}_1^n := u_1 \cdots u_n$ and we set $\mathbf{u}_1^0 = \epsilon$. Similarly, for an infinite sequence $\omega \in \Omega^{\mathbb{N}}$ and $n \geq 1$ we use the notation $\omega_1^n := \omega_1 \cdots \omega_n$ with $\omega_1^0 = \epsilon$. Finally, we use square brackets to denote cylinder sets, so

$$[\mathbf{u}] = \{ \omega \in \Omega^{\mathbb{N}} : \omega_1 \cdots \omega_{|\mathbf{u}|} = \mathbf{u} \}.$$
(5.4)

For $\mathbf{u} \in \Omega^k$ and $0 \le n \le k$, let

$$T_{\mathbf{u}} = T_{u_k} \circ T_{u_{k-1}} \circ \cdots \circ T_{u_1}$$
 and $T_{\mathbf{u}}^n = T_{\mathbf{u}_1^n} = T_{u_n} \circ T_{u_{n-1}} \circ \cdots \circ T_{u_1}$

Note that $T_{\mathbf{u}}^0 = T_{\mathbf{u}_1^0} = T_{\epsilon} = id$. Similarly if $\omega \in \Omega^{\mathbb{N}}$, we let $T_{\omega}^n = T_{\omega_1^n} = T_{\omega_n} \circ T_{\omega_{n-1}} \circ \cdots \circ T_{\omega_1}$ for any $n \ge 0$. For $\mathbf{u} \in \Omega^k$ the left and right random orbits of the critical points $c \in C$ are

$$T_{\mathbf{u}}(c^-) = \lim_{x \uparrow c} T_{\mathbf{u}}(x)$$
 and $T_{\mathbf{u}}(c^+) = \lim_{x \downarrow c} T_{\mathbf{u}}(x).$

The one-sided derivatives along \mathbf{u} are given by

$$T'_{\mathbf{u}}(c^{-}) = \lim_{x \uparrow c} \prod_{n=1}^{k} T'_{u_{n}}(T_{\mathbf{u}_{1}^{n-1}}(x)) \quad \text{ and } \quad T'_{\mathbf{u}}(c^{+}) = \lim_{x \downarrow c} \prod_{n=1}^{k} T'_{u_{n}}(T_{\mathbf{u}_{1}^{n-1}}(x)).$$

We use the abbreviation $p_{\mathbf{u}} := p_{u_1} \cdots p_{u_k}$ with $p_{\epsilon} = 1$.

5.2.1 Definition. (Random matching) A random map $R \in \mathcal{R}$ has random matching if for every $c \in C$ there exists an $M = M_c \in \mathbb{N}$ and a set

$$Y = Y_c \subseteq \left\{ T^k_{\omega}(c^-) : \omega \in \Omega^{\mathbb{N}}, \, 1 \le k \le M \right\} \cap \left\{ T^k_{\omega}(c^+) : \omega \in \Omega^{\mathbb{N}}, \, 1 \le k \le M \right\}$$

such that for every $\omega \in \Omega^{\mathbb{N}}$ there exist $k = k_c(\omega), \ell = \ell_c(\omega) \leq M$ with $T_{\omega}^k(c^-), T_{\omega}^{\ell}(c^+) \in Y$.

The main difference with one-dimensional matching as in (5.2) and (5.3) is that in a random system R the critical points have many different random orbits. Definition 5.2.1 states that any random orbit of the left or the right limit of any critical point c passes through the set Y_c at the latest at time M. The indices k, ℓ are introduced to cater for the possibility that these orbits pass through the set Y_c at different moments. Since all points in Y_c are in the orbit of both c^- and c^+ , this implies that all random orbits of the left limit meet with some random orbit of the right limit and vice versa. This corresponds to the statement in (5.2). Note that we do not ask $T^k_{\omega}(c^-) = T^{\ell}_{\omega}(c^+)$.

5.2.2 Definition. (Strong random matching) A random map $R \in \mathcal{R}$ has strong random matching if it has random matching and if for each $c \in C$ and $y \in Y_c$ the following holds. Set

$$\Omega(y)^{-} = \Big\{ \mathbf{u} \in \bigcup_{k=1}^{M} \Omega^{k} : \exists \omega \in \Omega^{\mathbb{N}} \text{ with } \mathbf{u} = \omega_{1} \cdots \omega_{k_{c}(\omega)} \text{ and } T_{\mathbf{u}}(c^{-}) = y \Big\},$$
$$\Omega(y)^{+} = \Big\{ \mathbf{u} \in \bigcup_{k=1}^{M} \Omega^{k} : \exists \omega \in \Omega^{\mathbb{N}} \text{ with } \mathbf{u} = \omega_{1} \cdots \omega_{\ell_{c}(\omega)} \text{ and } T_{\mathbf{u}}(c^{+}) = y \Big\}.$$

Then,

$$\sum_{\mathbf{u}\in\Omega(y)^{-}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c^{-})} = \sum_{\mathbf{u}\in\Omega(y)^{+}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c^{+})}.$$
(5.5)

Definition 5.2.2 guarantees that one can choose the times k, ℓ such that at those times orbits enter the set Y with the same weighted derivative. This is comparable to (5.3). Note that

ι

$$\bigcup_{y \in Y_c} \bigcup_{\mathbf{u} \in \Omega(y)^-} [\mathbf{u}] = \Omega^{\mathbb{N}} = \bigcup_{y \in Y_c} \bigcup_{\mathbf{u} \in \Omega(y)^+} [\mathbf{u}],$$

where $[\mathbf{u}]$ is a cylinder as defined in (5.4), so we have indeed captured all random orbits of c. Note that Definition 5.2.2 depends on the choices of $k_c(\omega)$ and $\ell_c(\omega)$ for each c in Definition 5.2.1.

If Ω consists of one element only, then the random map is actually a deterministic map. In this case Definition 5.2.1 and Definition 5.2.2 reduce to the definitions of onedimensional matching and strong matching given in (5.2) and (5.3), so the random definitions extend the deterministic ones.

In case each map $T_j : I \to I$ is piecewise affine on a finite partition $c_0 < c_1 < \ldots < c_N$ the conditions (a1) and (a3) are automatically satisfied and under some additional assumptions strong random matching has consequences for invariant densities. For this result we consider a subset of the collection of random maps \mathcal{R} . We define the subset $\mathcal{R}_A \subset \mathcal{R}$ to be the set of random systems in \mathcal{R} that satisfy the following additional assumptions:

- (c1) There exists a finite interval partition $\{I_i\}_{1 \le i \le N}$ of $I = [c_0, c_N]$ given by the points $c_0 < c_1 < \ldots < c_N$, such that each map $T_j : I \to I$, $j \in \Omega$, is piecewise affine with respect to this partition. In other words, for each $j \in \Omega$ and $1 \le i \le N$ we can write $T_j|_{(c_{i-1},c_i)}(x) = k_{i,j}x + d_{i,j}$ for some constants $k_{i,j}, d_{i,j}$.
- (c2) For each $1 \leq i \leq N$ there is an $1 \leq n \leq N$, such that

$$\frac{\sum_{j\in\Omega}\frac{p_j}{k_{i,j}}d_{i,j}}{1-\sum_{j\in\Omega}\frac{p_j}{k_{i,j}}} \neq \frac{\sum_{j\in\Omega}\frac{p_j}{k_{n,j}}d_{n,j}}{1-\sum_{j\in\Omega}\frac{p_j}{k_{n,j}}}.$$
(5.6)

(c3) For each $1 \le i \le N$, $\sum_{j \in \Omega} \frac{p_j}{k_{i,j}} \ne 0$.

Using the results from Chapter 3, we will show that for $R \in \mathcal{R}_A$ the following holds.

5.2.3 Theorem. Let $R \in \mathcal{R}_A$. If R has strong random matching, then there exists an invariant probability measure $m_{\mathbf{p}} \times \mu_{\mathbf{p}}$ for R with $\mu_{\mathbf{p}}$ absolutely continuous with respect to Lebesgue and such that its density $f_{\mathbf{p}}$ is piecewise constant. If moreover every map T_j is expanding, i.e., if $|k_{i,j}| > 1$ for each $1 \le i \le N$ and $j \in \Omega$, then any invariant probability density $f_{\mathbf{p}}$ of R is piecewise constant.

Assumptions (c2) and (c3) are used in Chapter 3 to prove that for systems in \mathcal{R}_A there exists an invariant probability density function that can be written as an infinite

sum of indicator functions. We use this fact in the proof below. These conditions, which are not very restrictive, guarantee that the method from Chapter 3 works, but they might not be necessary for the results from Theorem 3.4.1 and Theorem 5.2.3. In fact, the deterministic analog of Theorem 5.2.3, which can be found in [BCMP18, Theorem 1.2], does not have a condition like (5.6). Their proof uses an induced system with a full branched return map instead. One could try to transfer the proof of [BCMP18, Theorem 1.2] to the setting of random interval maps to avoid (c2) and (c3). Then, the recent results from Inoue in [I20] on first return time functions for random systems seem relevant. These results show, however, that an induced system for a random interval map will become position dependent instead of i.i.d., which might make such an extension not so straightforward.

Proof. The set of critical points of R is given by $C = \{c_1, \ldots, c_{N-1}\}$. Any random map $R \in \mathcal{R}_A$ satisfies the conditions of Theorem 3.4.1. Thus, there exists an invariant probability measure $m_{\mathbf{p}} \times \mu_{\mathbf{p}}$ for R with a probability density $f_{\mathbf{p}}$ for $\mu_{\mathbf{p}}$ of the form

$$f_{\mathbf{p}} = \sum_{i=1}^{N-1} \gamma_i \sum_{k \ge 1} \sum_{\mathbf{u} \in \Omega^k} \left(\frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c_i^-)} \mathbf{1}_{[c_0, T_{\mathbf{u}}(c_i^-))} - \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c_i^+)} \mathbf{1}_{[c_0, T_{\mathbf{u}}(c_i^+))} \right), \tag{5.7}$$

for some constants γ_i depending only on the critical points c_i . Fix an *i* and let M, Y be such that R satisfies the conditions of Definition 5.2.1 and Definition 5.2.2 for c_i . Then by (5.5)

$$\sum_{q \in Y} \left(\sum_{\mathbf{u} \in \Omega(y)^{-}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c_i^{-})} \mathbf{1}_{[c_0, T_{\mathbf{u}}(c_i^{-}))} - \sum_{\mathbf{u} \in \Omega(y)^{+}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c_i^{+})} \mathbf{1}_{[c_0, T_{\mathbf{u}}(c_i^{+}))} \right) = 0.$$

For each $1 \leq i \leq N-1$ and each $1 \leq k \leq M$, let

$$\Omega^{i,k}_{-} = \{ \mathbf{u} \in \Omega^k : \exists \, \omega \in \Omega^{\mathbb{N}} \text{ with } \mathbf{u} = \omega_1 \cdots \omega_k \text{ and } k < k_{c_i}(\omega) \}$$

and similarly

$$\Omega^{i,k}_{+} = \{ \mathbf{u} \in \Omega^{k} : \exists \, \omega \in \Omega^{\mathbb{N}} \text{ with } \mathbf{u} = \omega_{1} \cdots \omega_{k} \text{ and } k < \ell_{c_{i}}(\omega) \}.$$

Then $f_{\mathbf{p}}$ can be written as

$$f_{\mathbf{p}} = \sum_{i=1}^{N-1} \gamma_i \sum_{k=1}^{M} \Big(\sum_{\mathbf{u} \in \Omega_{-}^{i,k}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c_i^-)} \mathbf{1}_{[c_0, T_{\mathbf{u}}(c_i^-))} - \sum_{\mathbf{u} \in \Omega_{+}^{i,k}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c_i^+)} \mathbf{1}_{[c_0, T_{\mathbf{u}}(c_i^+))} \Big).$$

Hence $f_{\mathbf{p}}$ is constant on each interval in the finite partition of I specified by the orbit points in the set

$$\bigcup_{i=1}^{N-1}\bigcup_{k=1}^{M}\bigg(\{T_{\mathbf{u}}(c_i^-)\,:\,\mathbf{u}\in\Omega^{i,k}_-\}\cup\{T_{\mathbf{u}}(c_i^+)\,:\,\mathbf{u}\in\Omega^{i,k}_+\}\bigg).$$

This gives the first part of the result.

For the second part, note that under the additional assumption that $|k_{i,j}| > 1$ for all i, j the map R satisfies the conditions of Theorem 3.5.3. As a consequence, any invariant density $f_{\mathbf{p}}$ of R can be written as in (5.7) for some values γ_i . This proves the theorem.

§5.3 Examples

We give two examples of families of random interval maps depending on one parameter. We show that for each of these families there exist parameter intervals such that the systems have strong random matching for every parameter within these intervals. Moreover, within such an interval matching happens in a comparable way, i.e., with the same M and similar sets Y. As in the deterministic case, we call these intervals *matching intervals*. To ease the notation we use the symbol \star to indicate the set of strings obtained by replacing \star with any $j \in \Omega$. E.g., if $\Omega = \{0, 1, 2\}$, then $0 \star = \{00, 01, 02\}$.

§5.3.1 Random signed β -transformations

Let $\beta = \frac{1+\sqrt{5}}{2}$ be the golden mean, so $\beta^2 = \beta + 1$. For each $\alpha \in (1, \beta^2)$, let $\mathbf{p} = (p_0, p_1)$ be a positive probability vector and consider the random system R_{α} given by two generalised β -transformations $T_{\alpha,j} : [-\beta, \beta] \to [-\beta, \beta], j = 0, 1$, defined as follows.



Figure 5.1: The maps $T_{\alpha,0}$ and $T_{\alpha,1}$ for $\alpha \in \left(\frac{3\beta-2}{2}, 4\beta-5\right)$.

Consider the points

$$z_0 = -\beta$$
, $z_1 = -1$, $z_2 = -\frac{1}{\beta}$, $z_3 = \frac{1}{\beta}$, $z_4 = 1$, $z_5 = \beta$,

and let

$$T_{\alpha,0}(x) = \begin{cases} \beta x + \alpha & \text{if } x \in [z_0, z_2), \\ \beta x & \text{if } x \in [z_2, z_4), \\ \beta x - \alpha & \text{if } x \in [z_4, z_5], \end{cases}$$

and

$$T_{\alpha,1}(x) = \begin{cases} \beta x + \alpha & \text{if } x \in [z_0, z_1], \\ \beta x & \text{if } x \in (z_1, z_3], \\ \beta x - \alpha & \text{if } x \in (z_3, z_5]. \end{cases}$$

See Figure (5.1) for the graphs of $T_{\alpha,0}$ and $T_{\alpha,1}$, and Figure (5.2) for some examples of R_{α} , for different values of α .



Figure 5.2: R_{α} for $\alpha = 1$ in (a), $\alpha = \beta$ in (b), and $\alpha = \beta^2$ in (c).

Let

$$J_1 = \left(\frac{3\beta - 2}{2}, 4\beta - 5\right)$$
 and $J_2 = \left(\frac{6 - \beta}{2}, \frac{\beta + 5}{3}\right)$.

We will show that strong random matching happens for any $\alpha \in J_1 \cup J_2$. Note that $C = \{-1, -\frac{1}{\beta}, \frac{1}{\beta}, 1\}$ so that by the symmetry of the maps, to show that R_{α} has random matching, we only need to consider the points $\frac{1}{\beta}$ and 1.



Figure 5.3: R_{α} for $\alpha \in J_1$, J_2 respectively.

We start by considering $\alpha \in J_1$. See Figure (5.3)(a) for the corresponding random system R_{α} . The parameter interval is constructed in such a way that for any $\alpha \in J_1$ the initial parts of the random orbits of the left and right limits to $\frac{1}{\beta}$ and 1 are determined in the following way. For j = 0, 1 and any $\omega \in \{0, 1\}^{\mathbb{N}}$,

$$T_{\alpha,0}(1^{-}) = \beta, \qquad T_{\alpha,\omega}(\beta) = \beta^{2} - \alpha, \qquad T_{\alpha,\omega}^{2}(\beta) = \beta^{2}(\beta - \alpha),$$

$$T_{\alpha,1}(1^{-}) = T_{\alpha,j}(1^{+}) = \beta - \alpha, \quad T_{\alpha,\omega}(\beta - \alpha) = \beta(\beta - \alpha), \quad T_{\alpha,\omega}^{2}(\beta - \alpha) = \beta^{2}(\beta - \alpha).$$

Hence, for $1 \in C$ we can take $M = k_1(\omega) = \ell_1(\omega) = 3$ for each ω , $Y = \{\beta^2(\beta - \alpha)\}$ and one easily checks the conditions of both Definition 5.2.1 and Definition 5.2.2.

For $\frac{1}{\beta}$ the orbits are more complicated. Firstly, $T_{\alpha,j}(\frac{1}{\beta}) = 1 = T_{\alpha,0}(\frac{1}{\beta})$ and $T_{\alpha,1}(\frac{1}{\beta}) = 1 - \alpha$. We saw the orbit of 1 above, so we concentrate on the orbit of $1 - \alpha$. We have $T_{\alpha,j}(1-\alpha) = \beta(1-\alpha) \in (-1, -\frac{1}{\beta})$, so $T_{\alpha,0}(\beta(1-\alpha)) = \beta(\beta-\alpha)$ and

 $T_{\alpha,1}(\beta(1-\alpha)) = \beta^2(1-\alpha)$. The next couple of iterations are depicted in Figure 5.4, where we have used the property that $\beta^2 = \beta + 1$ to compute the orbit points.



Figure 5.4: The first couple of points in the orbit of $1 - \alpha$ under the random generalised β -transformation. We have boxed $\beta^2(\beta - \alpha)$, since this point also appears in all random orbits of 1.

Take $M = k_{\frac{1}{\beta}}(\omega) = \ell_{\frac{1}{\beta}}(\omega) = 7$ for each ω and set $Y = \{\beta^5(\beta - \alpha) - \alpha, \beta^5(\beta - \alpha) - \beta\alpha = \beta^6 - 3\beta^3\alpha\}$. Then,

$$\Omega(\beta^5(\beta - \alpha) - \alpha)^+ = 0 \star \star \star \star 0 \star \cup 1 \star 0 \star \star 0 \star \cup 1 \star 1 \star \star 0 \star$$

and $\Omega(\beta^5(\beta - \alpha) - \alpha)^- = \star \star \star \star \star 0\star$. Hence,

$$\sum_{\mathbf{u}\in\Omega(\beta^5(\beta-\alpha)-\alpha)^+} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(\frac{1}{\beta}^+)} = \frac{p_0^2 + p_1 p_0^2 + p_1^2 p_0}{\beta^7} = \frac{p_0}{\beta^7} = \sum_{\mathbf{u}\in\Omega(\beta^5(\beta-\alpha)-\alpha)^-} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(\frac{1}{\beta}^-)}.$$

A similar computation gives (5.5) for $\beta^5(\beta - \alpha) - \beta\alpha$, so R_{α} has strong random matching.

Note that also in this example the orbits of 1⁺ meet with some of the orbits of 1⁻ earlier, in this case already after one step. Hence, we could also take $Y_1 = \{\beta - \alpha, \beta^2(\beta - \alpha)\}$ and split the random orbits as follows:

$$\Omega(\beta - \alpha)^+ = \{1\} = \Omega(\beta - \alpha)^- \quad \text{and} \quad \Omega(\beta^2(\beta - \alpha))^+ = 0 \star \star = \Omega(\beta^2(\beta - \alpha))^-.$$

Then for some ω the values $k_1(\omega), \ell_1(\omega)$ are lower, but we have to check condition (5.5) for two points instead of one. For the critical point $\frac{1}{\beta}$ we could use $Y = \{\beta - \alpha, \beta^2(\beta - \alpha), \beta^5(\beta - \alpha) - \alpha, \beta^5(\beta - \alpha) - \beta\alpha\}$ or also $Y = \{\beta^2(\beta - \alpha), \beta^5(\beta - \alpha) - \alpha, \beta^5(\beta - \alpha) - \beta\alpha\}$. By the flexibility in the choice of Y given by Definition 5.2.1 one can choose the set Y that is most convenient. Theorem 5.2.3 explains the need for condition (5.5) in Definition 5.2.2.

We now consider $\alpha \in J_2$; see Figure (5.3)(b) for the corresponding random system R_{α} . For j = 0, 1, the initial parts of the random orbits of the left and right limits to $\frac{1}{\beta}$ and 1 are given by

$$T_{\alpha,j}(\frac{1}{\beta}^{-}) = T_{\alpha,0}(\frac{1}{\beta}^{+}) = 1, \quad T_{\alpha,1}(\frac{1}{\beta}^{+}) = 1 - \alpha.$$

and

$$T_{\alpha,j}(1^+) = T_{\alpha,1}(1^-) = \beta - \alpha, \quad T_{\alpha,0}(1^+) = \beta.$$

Moreover, for any $\omega \in \{0,1\}^{\mathbb{N}}$

$$\begin{split} T_{\alpha,\omega}(1) &= \beta - \alpha, \\ T_{\alpha,\omega}(\beta) &= \beta^2 - \alpha, \\ T_{\alpha,\omega}^1(1-\alpha) &= \beta - \alpha(\beta - 1), \\ T_{\alpha,\omega}^1(1-\alpha) &= \beta - \alpha(\beta - 1), \\ \end{split}$$

Note also that $T_{\alpha,j1}(\beta - \alpha) = \beta(\beta^2 - \alpha)$. Since all possible random orbits of $\frac{1}{\beta}$ and 1 pass by the point $\beta - \alpha$ or by one of its iterates, $\beta(\beta^2 - \alpha)$, to study the orbits of these critical points is enough to follow the one of $\beta - \alpha$. This is depicted in Figure 5.5.

Figure 5.5: The first couple of points in the orbit of $\beta - \alpha$ under R_{α} for $\alpha \in J_2$. We have boxed $\beta(\beta^2 - \alpha)$, since this point appears in the orbits of 1, β and $1 - \alpha$.

Take $M = k_{\frac{1}{3}}(\omega) = \ell_{\frac{1}{3}}(\omega) = 8$ for each ω and set

$$Y = \{\beta^4(\beta^3 - 2\alpha) = \beta^3(\beta^4 - 3\alpha) - \alpha, \beta^3(\beta^4 - 3\alpha) = \beta^7 - 2\beta^4\alpha + \alpha\}.$$

Then,

$$\Omega(\beta^4(\beta^3 - 2\alpha))^- = \star \star \star 01 \star \star 1 \cup \star \star \star 00 \star \star 1 \cup \star \star \star 1 \star \star \star 1,$$

and

$$\Omega(\beta^4(\beta^3 - 2\alpha))^+ = 0 \star \star 01 \star \star 1 \cup 0 \star \star 00 \star \star 1 \cup 0 \star \star 1 \star \star \star 1 \cup 1 \star \star \star 0 \star \star 1 \cup 1 \star \star \star 1 \star \star 1.$$

Hence,

$$\sum_{\mathbf{u}\in\Omega(\beta^4(\beta^3-2\alpha))^-} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(\frac{1}{\beta}^-)} = \frac{p_1^2 p_0 + p_1 p_0^2 + p_1^2}{\beta^8} = \frac{p_1}{\beta^8} = \sum_{\mathbf{u}\in\Omega(\beta^4(\beta^3-2\alpha))^+} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(\frac{1}{\beta}^+)}.$$

A similar computation gives (5.5) for $\beta^3(\beta^4 - 3\alpha)$, so R_{α} has strong random matching for any $\alpha \in J_2$.

Note that also in this example the orbits of $\frac{1}{\beta}^{-}$ meet with some of the orbits of $\frac{1}{\beta}^{+}$ earlier, in this case already after one step. Hence, we could also take the points 1, and $\beta(\beta^2 - \alpha)$ } in the set Y and split the random orbits accordingly.

5.3.1 Remark. The random generalised β -transformations R_{α} from Example 5.3.1 satisfy all conditions of Theorem 5.2.3. Hence, for any $\alpha \in J_1 \cup J_2$ any invariant density of the random system R_{α} is piecewise constant.

§5.3.2 Random CF-maps

For $\alpha \in (0,1)$ let $T_{\alpha,0}, T_{\alpha,1} : [\alpha - 1, \alpha] \to [\alpha - 1, \alpha]$ be the Nakada and Ito-Tanaka α -continued fraction transformations, introduced in [N81] and in [TI81] respectively, and given by

$$T_{\alpha,0}(x) = \frac{1}{|x|} - \left\lfloor \frac{1}{|x|} + 1 - \alpha \right\rfloor \quad \text{and} \quad T_{\alpha,1}(x) = \frac{1}{x} - \left\lfloor \frac{1}{x} + 1 - \alpha \right\rfloor,$$

for $x \neq 0$ and $T_{\alpha,0}(0) = 0 = T_{\alpha,1}(0)$. The graphs are shown in Figure 5.6.



Figure 5.6: The Nakada α -continued fraction map $T_{\alpha,0}$ in (a) and the Ito-Tanaka α -continued fraction map $T_{\alpha,1}$ in (b) for $\alpha = \frac{7}{10} \in \left(\frac{5-\sqrt{13}}{2}, \frac{\sqrt{2}}{2}\right)$.

Let R_{α} denote the corresponding pseudo-skew product on $\{0, 1\}^{\mathbb{N}} \times [\alpha - 1, \alpha]$. For $x \in [0, \alpha]$, the two maps coincide and

$$T_{\alpha,0}(x) = T_{\alpha,1}(x) = \frac{1}{x} - n \quad \text{for} \ x \in \left(\frac{1}{\alpha+n}, \frac{1}{\alpha+n-1}\right].$$

For $x \in [\alpha - 1, 0)$, we have

$$T_{\alpha,0}(x) = -\frac{1}{x} - n \quad \text{for } x \in \left[-\frac{1}{\alpha + n - 1}, -\frac{1}{\alpha + n}\right)$$
$$T_{\alpha,1}(x) = \frac{1}{x} + n \quad \text{for } x \in \left[\frac{1}{\alpha - n}, \frac{1}{\alpha - (n + 1)}\right).$$

We first show that for any $\alpha \in \left(\frac{\sqrt{10}-2}{2}, 2-\sqrt{2}\right)$ the map R_{α} has random matching. For this note that the critical points c are all in the set $\left\{\frac{1}{\alpha+n}, -\frac{1}{\alpha+n}, \frac{1}{\alpha-n} : n \in \mathbb{N}\right\}$. For any positive critical point c > 0 and any $j \in \{0, 1\}$, $T_j(c^-), T_j(c^+) \in \{\alpha - 1, \alpha\}$. For c < 0, c is either a critical point for T_0 and a continuity point for T_1 , or a critical point for T_1 and a continuity point for T_0 . Specifically, since $\alpha > \frac{1}{2}$, for $c = -\frac{1}{\alpha+n}$ we have

$$T_0(c^-) = \alpha$$
, $T_0(c^+) = \alpha - 1$, and $T_1(c^-) = T_1(c^+) = 1 - \alpha$,

and for $c = \frac{1}{\alpha - n}$

$$T_1(c^-) = \alpha - 1$$
, $T_1(c^+) = \alpha$, and $T_0(c^-) = T_0(c^+) = 1 - \alpha$.

As a consequence, to show that R_{α} has random matching we only need to consider the orbits of $\alpha - 1$ and α . Due to the choice of endpoints of the parameter interval $\left(\frac{\sqrt{10-2}}{2}, 2-\sqrt{2}\right)$, the first three orbit points of α and $\alpha - 1$ are easily determined. They are given in Figure 5.7. Hence, if we take M = 3 and

$$Y = \left\{\frac{5\alpha - 3}{1 - 2\alpha}, \frac{4 - 7\alpha}{1 - 2\alpha}\right\}, \quad \text{if } c > 0$$

and

$$Y = \left\{\frac{5\alpha - 3}{1 - 2\alpha}, \frac{4 - 7\alpha}{1 - 2\alpha}, 1 - \alpha\right\}, \quad \text{if } c < 0,$$

then R_{α} has random matching according to Definition 5.2.1.



Figure 5.7: The first three elements in the orbits of α and $\alpha - 1$ under the random continued fraction map R_{α} for $\alpha \in \left(\frac{\sqrt{10-2}}{2}, 2-\sqrt{2}\right)$. The digits above the arrows indicate which one of the maps $T_{\alpha,0}$ or $T_{\alpha,1}$ is applied. If there is no digit, then both maps yield the same orbit point. Orbit points in boxes with the same colour are equal.

 R_{α} does not satisfy strong random matching with this choice of Y. To see this, note that $T'_{\alpha,1}(x) = -\frac{1}{x^2}$ for all x where the derivative exists, while $T'_{\alpha,0}(x) = -\frac{1}{x^2}$ if

x > 0 and $T'_{\alpha,0}(x) = \frac{1}{x^2}$ if x < 0. Now take for example $c = \frac{1}{\alpha+n} > 0$ and $y = \frac{4-7\alpha}{1-2\alpha}$. Then $\Omega(y)^- = \star 11 = \{011, 111\}$ and $\Omega(y)^+ = \star \star 1$. For the quantities from (5.5), we obtain

$$\sum_{\mathbf{u}\in\Omega(y)^-}\frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}}(c^-) = -p_1^2c^2(2\alpha-1)^2 \quad \text{and} \quad \sum_{\mathbf{u}\in\Omega(y)^+}\frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}}(c^+) = -p_1c^2(2\alpha-1)^2,$$

which are not equal for any $p_1 \in (0, 1)$.

We now identify a countable number of parameter intervals on which the maps R_{α} have strong matching with the same exponent M = 4, i.e., we identify a countable number of *matching intervals* for the family R_{α} . For $n \geq 4$ let the interval $J_n := (\ell_n, r_n)$ be defined by the left and right endpoints

$$\ell_n = \frac{n+1-\sqrt{n^2-2n+5}}{2}$$
 and $r_n = \sqrt{\frac{n-2}{n}}$, (5.8)

respectively. Set $g := \frac{\sqrt{5}-1}{2}$ for the small golden mean and note that $g < \ell_n < r_n$ for all $n \ge 4$ and that $\lim_{n\to\infty} \ell_n = \lim_{n\to\infty} r_n = 1$. See Figure 5.8 for an illustration of the location of these intervals.



Figure 5.8: The semicircles indicate the locations of the intervals J_n .

The intervals J_n are chosen in such a way that we can determine the first three orbit points of α and $\alpha - 1$. Let $n \geq 4$ and $\alpha \in J_n$. In particular $\alpha > g$ and for j = 0, 1,

$$T_{\alpha,j}(\alpha) = \frac{1-\alpha}{\alpha} > 0$$

The point ℓ_n is chosen so that $\alpha - 1 \in \left(\frac{1}{\alpha - n}, \frac{\alpha + 1}{1 - n(\alpha + 1)}\right) \subseteq \left(-\frac{1}{\alpha + n - 2}, -\frac{1}{\alpha + n - 1}\right)$. Since $\frac{\alpha + 1}{1 - n(\alpha + 1)} < \frac{1}{\alpha - n - 1}$ we get

$$T_{\alpha,1}(\alpha - 1) = \frac{n(\alpha - 1) + 1}{\alpha - 1}$$
 and $T_{\alpha,0}(\alpha - 1) = \frac{\alpha(n - 1) + 2 - n}{1 - \alpha}$.

It also implies $\frac{1-\alpha}{\alpha} \in (\frac{1}{\alpha+n-2}, \frac{1}{\alpha+n-3})$. As a consequence, for l = 0, 1,

$$T_{\alpha,jl}(\alpha) = \frac{\alpha(n-1) + 2 - n}{1 - \alpha} = T_{\alpha,0}(\alpha - 1) > 0.$$

We further divide the interval J_n . For $k \in \{2, 3, ..., n\}$, let

$$i_{n,k} = \frac{-4 + 2n - kn + k + \sqrt{k^2 n^2 - 2k^2 n + k^2 + 4}}{2(n-1)},$$

and note that $J_n \subseteq \bigcup_{k=2}^{n-1}(i_{n,k+1}, i_{n,k}]$. Therefore, for each $\alpha \in J_n$ there exists a $k \in \{2, 3, \ldots, n-1\}$ such that $\alpha \in (i_{n,k+1}, i_{n,k}]$. The last condition is equivalent to

$$\frac{1}{\alpha+k} < \frac{\alpha(n-1)+2-n}{1-\alpha} \le \frac{1}{\alpha+k-1},$$
(5.9)

so that for $\mathbf{u}\in\Omega^3$ it holds that

$$T_{\alpha,\mathbf{u}}(\alpha) = \frac{1 - 2k + kn - \alpha(kn - k + 1)}{\alpha(n-1) + 2 - n}.$$

On the other hand, the choice of r_n guarantees that $T_{\alpha,1}(\alpha-1) = \frac{1+n(\alpha-1)}{n} > \frac{1}{\alpha+1}$. Then for j = 0, 1,

$$T_{\alpha,1j}(\alpha - 1) = \frac{\alpha(n-1) + 2 - n}{-1 - n(\alpha - 1)}.$$

Equation (5.9) holds if and only if

$$\frac{1}{\alpha + k - 1} < \frac{\alpha(n - 1) + 2 - n}{-1 - n(\alpha - 1)} \le \frac{1}{\alpha + k - 2}$$

is satisfied. In this case, for l = 0, 1

$$T_{\alpha,1jl}(\alpha - 1) = \frac{1 - 2k + kn - \alpha(kn - k + 1)}{\alpha(n - 1) + 2 - n}.$$

Figure 5.9 shows all the relevant orbit points of α and $\alpha - 1$.



Figure 5.9: The first few points in the orbits of α and $\alpha - 1$ under the random continued fraction map R_{α} for $\alpha \in J_n \cap (i_{n,k+1}, i_{n,k}]$.

Definition 5.2.1 holds for $\alpha \in J_n \cap (i_{n,k+1}, i_{n,k}]$ with M = 4 and

$$Y = \left\{ \frac{1 - 2k + kn - \alpha(kn - k + 1)}{\alpha(n - 1) + 2 - n} \right\}$$

for any critical point c > 0. For c < 0 we add the point $1 - \alpha$ to Y. Here the values $k_c(\omega)$ and $\ell_c(\omega)$ either equal 1, 3 or 4 according to the number of orbit points in the paths in Figure 5.9. For Definition 5.2.2, for c > 0 and $y \in Y$ we have

 $\Omega(y)^- = \star 0 \star \cup \star 1 \star \star \text{ and } \Omega(y)^+ = \star \star \star, \text{ so that}$

$$\sum_{\mathbf{u}\in\Omega(y)^{-}} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c^{-})} = (-c^{2})p_{0}(\alpha-1)^{2} \cdot \frac{(\alpha(n-1)+2-n)^{2}}{-(\alpha-1)^{2}} + (-c^{2})p_{1}(-(\alpha-1)^{2}) \cdot \frac{(1+n(\alpha-1))^{2}}{-(\alpha-1)^{2}} \cdot \frac{(\alpha(n-1)+2-n)^{2}}{-(1+n(\alpha-1))^{2}} = c^{2}(\alpha(n-1)+2-n)^{2}.$$

and

$$\sum_{\mathbf{u}\in\Omega(y)^+} \frac{p_{\mathbf{u}}}{T'_{\mathbf{u}}(c^+)} = (-c^2)(-\alpha^2)\frac{(1-\alpha)^2}{-\alpha^2} \cdot \frac{(\alpha(n-1)+2-n)^2}{-(1-\alpha)^2} = c^2(\alpha(n-1)+2-n)^2,$$

implying that also condition (5.5) holds. For $c = -1/(\alpha + n)$ we get $\Omega(1 - \alpha)^- = \Omega(1 - \alpha)^+ = \{1\}, \ \Omega(y)^- = 0 \star \star \star \text{ and } \Omega(y)^+ = 00 \star \cup 01 \star \star, \text{ and for } c = 1/(\alpha - n) \text{ we}$ obtain $\Omega(1 - \alpha)^- = \Omega(1 - \alpha)^+ = \{0\}, \ \Omega(y)^- = 10 \star \cup 11 \star \star \text{ and } \Omega(y)^+ = 1 \star \star \star$. In both cases the result follows in a similar fashion. So, the random continued fraction system R_{α} has strong random matching for any \mathbf{p} and any $\alpha \in J_n$.

Note that in this example the orbits of α meet with some of the orbits of $\alpha - 1$ already after two time steps in the point $\frac{\alpha(n-1)+2-n}{1-\alpha}$. Hence,

$$\frac{\alpha(n-1)+2-n}{1-\alpha} \in \Big\{ T^k_{\omega}(c^-) \, : \, \omega \in \Omega^{\mathbb{N}}, \, k \le M \Big\} \cap \Big\{ T^k_{\omega}(c^+) \, : \, \omega \in \Omega^{\mathbb{N}}, \, k \le M \Big\}.$$

Therefore, for a critical point c > 0, we could also take

$$Y = \left\{\frac{\alpha(n-1) + 2 - n}{1 - \alpha}, \frac{1 - 2k + kn - \alpha(kn - k + 1)}{\alpha(n-1) + 2 - n}\right\}$$

and split the random orbits of α for example in the following way:

$$\Omega\left(\frac{\alpha(n-1)+2-n}{1-\alpha}\right)^{+} = \star \star 0 \quad \text{and} \quad \Omega\left(\frac{1-2k+kn-\alpha(kn-k+1)}{\alpha(n-1)+2-n}\right)^{+} = \star \star 1 \star.$$

For the orbits passing through $\alpha - 1$ we have

$$\Omega\left(\frac{\alpha(n-1)+2-n}{1-\alpha}\right)^{-} = \star 0 \quad \text{and} \quad \Omega\left(\frac{1-2k+kn-\alpha(kn-k+1)}{\alpha(n-1)+2-n}\right)^{-} = \star 1 \star \star.$$

One can check that condition (5.5) is satisfied and R_{α} has strong random matching with this choice of Y. Note that in this case many sequences ω have smaller values $k_c(\omega)$ and $\ell_c(\omega)$ than with $Y = \left\{\frac{1-2k+kn-\alpha(kn-k+1)}{\alpha(n-1)+2-n}\right\}$ and that for some $\omega \in \Omega^{\mathbb{N}}$ we do not take $k_c(\omega)$ equal to the first time that the random orbit $T_{\omega}^k(c^-)$ enters Y. For example, for c > 0 and any ω with $\omega_3 = 1$ we have $T_{\omega}^3(c^+) = \frac{\alpha(n-1)+2-n}{1-\alpha} \in Y$, but we take $k_c(\omega) = 4$. The flexibility in the choice of Y and the length of the paths $k_c(\omega)$ and $\ell_c(\omega)$ embedded in Definition 5.2.1 allows one to choose the option that is computationally most convenient.

§5.4 Random transformations and expansions

In the second part of this Chapter we use random matching to study the frequency of the digit 0 in the signed binary expansions produced by a family of random system of piecewise affine maps. In this section we define this family and its relation to binary expansions.

§5.4.1 Random symmetric doubling maps

A signed binary expansion of a number $x \in [-1,1]$ can be obtained by iterating any piecewise affine map $D : [-1,1] \rightarrow [-1,1]$ that is given by D(x) = 2x - d with $d \in \{-1,0,1\}$ on each of its intervals of monotonicity. One can for example take any $a \in [\frac{1}{4}, \frac{1}{2}]$ and then define the symmetric map

$$D_a(x) = \begin{cases} 2x+1, & \text{if } -1 \le x < -a\\ 2x, & \text{if } -a \le x \le a,\\ 2x-1, & \text{if } a < x \le 1. \end{cases}$$

By setting $d_n(x) = d, d \in \{-1, 0, 1\}$, if $D_a^n(x) = 2D_a^{n-1}(x) - d$, one obtains

$$x = \frac{d_1(x)}{2} + \frac{D_a(x)}{2} = \dots = \frac{d_1(x)}{2} + \dots + \frac{d_n(x)}{2^n} + \frac{D_a^n(x)}{2^n} \to \sum_{n \ge 1} \frac{d_n(x)}{2^n},$$

so this gives a signed binary expansion of x. The family of maps $\{D_a\}_{\frac{1}{4} \leq a \leq \frac{1}{2}}$ is the object of study in [DK17]. As can be seen from Figure 5.10(a) the interval [-2a, 2a] is an attractor for the dynamics of D_a . Since this interval depends on a, in [DK17] the authors decided to work instead with the measurably isomorphic family $\{S_\alpha\}_{1\leq \alpha\leq 2}$ given by

$$S_{\alpha}(x) = \begin{cases} 2x + \alpha, & \text{if } -1 \le x < -\frac{1}{2}, \\ 2x, & \text{if } -\frac{1}{2} \le x \le \frac{1}{2}, \\ 2x - \alpha, & \text{if } \frac{1}{2} < x \le 1, \end{cases}$$
(5.10)

see Figure 5.10(b), which transfers the dependence on the parameter from the domain [-1, 1] to the branches of the maps.

While each deterministic map produces for each number in its domain a single signed binary expansion, one can define random dynamical systems that produce for Lebesgue almost all numbers uncountably many different signed binary expansions. The family of random maps $\{R_{\alpha}\}$, which we define next, extends the family of deterministic maps $\{S_{\alpha}\}$. So the dependence on the parameter is visible in the branches of the maps instead of in the domains.

Let $\Omega = \{0,1\}$ and define for $j \in \Omega$ and each parameter $\alpha \in [1,2]$ the maps $T_j = T_{\alpha,j} : [-1,1] \to [-1,1]$ by

$$T_{\alpha,0}(x) = \begin{cases} 2x + \alpha, & \text{if } x \in \left[-1, \frac{1-\alpha}{2}\right], \\ 2x, & \text{if } x \in \left(\frac{1-\alpha}{2}, \frac{1}{2}\right], \\ 2x - \alpha, & \text{if } x \in \left(\frac{1}{2}, 1\right], \end{cases}$$
(5.11)



Figure 5.10: The maps D_c and $S_{\frac{1}{2c}}$ for $c = \frac{7}{20}$. The grey lines indicate the remainder of the maps $x \mapsto 2x + 1$, $x \mapsto 2x$ and $x \mapsto 2x - 1$. The red box in (a) shows the attractor of the map D_c .

and

$$T_{\alpha,1}(x) = \begin{cases} 2x + \alpha, & \text{if } x \in \left[-1, -\frac{1}{2}\right], \\ 2x, & \text{if } x \in \left[-\frac{1}{2}, \frac{\alpha-1}{2}\right], \\ 2x - \alpha, & \text{if } x \in \left[\frac{\alpha-1}{2}, 1\right]. \end{cases}$$
(5.12)

See Figure 5.11 for three examples. The maps $T_{\alpha,0}$ and $T_{\alpha,1}$ differ on the intervals $\left[-\frac{1}{2}, 1-2\alpha\right]$ and $\left[2\alpha-1, \frac{1}{2}\right]$, which are indicated by the grey areas in the pictures. Let $R = R_{\alpha} : \Omega^{\mathbb{N}} \times [-1,1] \to \Omega^{\mathbb{N}} \times [-1,1]$ be the random system obtained from $T_{\alpha,0}$ and $T_{\alpha,1}$, i.e.,

$$R_{\alpha}(\omega, x) = (\sigma(\omega), T_{\alpha, \omega_1}(x)),$$

where σ is the left shift on $\Omega^{\mathbb{N}}$. We call the systems R_{α} random symmetric doubling maps and the subscript α will sometimes be suppressed if it does not lead to confusion.



Figure 5.11: The maps $T_{\alpha,0}$ and $T_{\alpha,1}$ for $\alpha = 1$ in (a), $\alpha = \frac{3}{2}$ in (b), and $\alpha = 2$ in (c). The blue lines correspond to $T_{\alpha,0}$, the pink ones to $T_{\alpha,1}$ and the purple ones to both.

Fix an $\alpha \in [1,2]$. Recall from (5.4) that we use square brackets to denote the cylinder sets in $\Omega^{\mathbb{N}}$. Let $\pi : \Omega^{\mathbb{N}} \times [-1,1] \to [-1,1]$ denote the canonical projection

onto the second coordinate and set

$$s_{n}(\omega, x) = \begin{cases} -1, & \text{if } R^{n-1}(\omega, x) \in \Omega^{\mathbb{N}} \times \left[-1, -\frac{1}{2}\right) \cup [0] \times \left[-\frac{1}{2}, \frac{1-\alpha}{2}\right], \\ 0, & \text{if } R^{n-1}(\omega, x) \in [1] \times \left[-\frac{1}{2}, \frac{1-\alpha}{2}\right] \\ & \cup \Omega^{\mathbb{N}} \times \left(\frac{1-\alpha}{2}, \frac{\alpha-1}{2}\right) \cup [0] \times \left[\frac{\alpha-1}{2}, \frac{1}{2}\right], \\ 1, & \text{if } R^{n-1}(\omega, x) \in [1] \times \left[\frac{\alpha-1}{2}, \frac{1}{2}\right] \cup \Omega^{\mathbb{N}} \times \left(\frac{1}{2}, 1\right]. \end{cases}$$
(5.13)

Then

$$\pi(R^n(\omega, x)) = 2\pi(R^{n-1}(\omega, x)) - s_n(\omega, x)\alpha,$$

so that just as in the deterministic case by iteration we obtain

$$x = \frac{s_1(\omega, x)\alpha}{2} + \dots + \frac{s_n(\omega, x)\alpha}{2^n} + \frac{\pi(R^n(\omega, x))}{2^n} \to \alpha \sum_{n \ge 1} \frac{s_n(\omega, x)}{2^n}$$

In other words, iterations of the random system R give a signed binary expansion for the pair (ω, x) .

Note that for each $x \in [-1, 1]$ there is an $\omega \in \Omega^{\mathbb{N}}$, such that $\pi(R^n_{\alpha}(\omega, x)) = S^n_{\alpha}(x)$, where S_{α} is the map in the family $\{S_{\alpha}\}$ from [DK17]. In particular, the random signed binary expansions produced by the family $\{R_{\alpha}\}$ include, among many others, the SSB expansions. The randomness of the system allows us to choose (up to a certain degree) where and when we want to have a digit 0. Below we investigate the frequency of the digit 0 in typical expansions produced by the maps R. We do so by applying Birkhoff's Ergodic Theorem for invariant measures for R of the form $m \times \mu$ with m a Bernoulli measure and μ absolutely continuous with respect to the Lebesgue measure. For that we need to investigate the density of such measures μ .

§5.4.2 Random matching almost everywhere

For any $\alpha \in (1,2]$ the common partition on which T_0 and T_1 are monotone is given by the points

$$c_0 = -1$$
, $c_1 = -\frac{1}{2}$, $c_2 = \frac{1-\alpha}{2}$, $c_3 = \frac{\alpha-1}{2}$, $c_4 = \frac{1}{2}$, $c_5 = 1$.

Set, in accordance with (a1),

$$I_1 = [c_0, c_1), \quad I_2 = [c_1, c_2], \quad I_3 = (c_2, c_3), \quad I_4 = [c_3, c_4], \quad I_5 = (c_4, c_5],$$

then $C = \{c_1, c_2, c_3, c_4\}$. For $0 , use <math>\mathbf{p} = (p_0, p_1)$ to denote the probability vector with $p_0 = p$ and $p_1 = 1 - p$. Since T_0 and T_1 from (5.11), (5.12) are both piecewise affine with slope 2, we have $\frac{p_j}{|T'_j(x)|} = \frac{p_j}{T'_j(x)} = \frac{p_j}{2}$, j = 0, 1. So the random system R satisfies conditions (a1), (a2), (a3), i.e., $R \in \mathcal{R}$. Due to the symmetry in the map, to verify whether R has strong random matching it is enough to check the conditions of Definitions 5.2.1 and Definitions 5.2.2 for the points $1 = T_0(c_4)^-$ and $1 - \alpha = T_0(c_4)^+$. Before we proceed with a description of the matching behaviour of the family of random systems $\{R_{\alpha}\}$, we first recall the results from [DK17, Propositions 2.1 and 2.3] on matching for the family of deterministic symmetric doubling maps $\{S_{\alpha}\}$, see (5.10). Let

$$M_{\alpha} = \inf\left\{n \ge 0 : \frac{1}{2} < S_{\alpha}^{n}(1) < \alpha - \frac{1}{2}\right\} + 1.$$
(5.14)

Then according to [DK17, Propositions 2.1 and 2.3] for all $\alpha \in [1, 2]$,

$$S^k_{\alpha}(1-\alpha) = S^k_{\alpha}(1) - \alpha \quad \text{for } k < M_{\alpha}$$
(5.15)

and for Lebesgue almost all $\alpha \in [1, 2]$ in fact $M_{\alpha} < \infty$ and

$$S_{\alpha}^{M_{\alpha}+1}\left(\frac{1}{2}^{-}\right) = S_{\alpha}^{M_{\alpha}}(1) = S_{\alpha}^{M_{\alpha}}(1-\alpha) = S_{\alpha}^{M_{\alpha}+1}\left(\frac{1}{2}^{+}\right).$$

In other words, for Lebesgue almost all parameters $\alpha \in [1, 2]$ the map S_{α} has matching with matching exponent $M = M_{\alpha} + 1$ that is determined by the first time the orbit of 1 enters the interval $(\frac{1}{2}, \alpha - \frac{1}{2})$. Moreover, $S_{\alpha}^{M_{\alpha}-1}(1-\alpha) < -\frac{1}{2}$ for all α , $M_{\alpha} = 1$ for $\alpha \in [\frac{3}{2}, 2]$ and $M_{\alpha} > 1$ for $\alpha \in (1, \frac{3}{2})$. Due to the constant slope and the same matching exponent M_{α} of the left and right limits, in this case matching implies strong matching.

5.4.1 Remark. The discrepancy between $M_{\alpha}+1$ here and M_{α} as matching exponent in [DK17] comes from the fact that in [DK17] the orbits are considered as starting from 1 and $1-\alpha$, whereas in (5.2) and (5.3) we followed the convention in [BCMP18] and start at the critical point $c = \frac{1}{2}$ instead.

From this we deduce the following small lemma.

5.4.2 Lemma. For $\alpha \in [1,2]$ and for all $k < M_{\alpha} - 1$, either $S^k(1) \in I_4$ and $S^k(1) - \alpha \in I_1$ or $S^k(1) \in I_5$ and $S^k(1) - \alpha \in I_2$.

Proof. From (5.15) it follows for all $k < M_{\alpha} - 1$ that $S^{k}(1) - \alpha \geq -1$, implying that

$$\alpha - 1 \le S^k(1) \le 1$$
 and $-1 \le S^k(1) - \alpha \le 1 - \alpha$.

The fact that $S^k(1) \in I_4 \cup I_5$ follows since $\frac{\alpha-1}{2} \leq \alpha - 1$. If $S^k(1) \in I_4$, then $S^k(1) - \alpha \leq \frac{1}{2} - \alpha < -\frac{1}{2}$, so $S^k(1) - \alpha \in I_1$. Suppose $S^k(1) \in I_5$. If $S^k(1) - \alpha < -\frac{1}{2}$, this would imply that $S^k(1) \in (\frac{1}{2}, \alpha - \frac{1}{2})$, contradicting the definition of M_α in (5.14). Hence, $S^k(1) - \alpha \in I_2$.

The next result states that a random equivalent of (5.15) holds for $\alpha \in (1, \frac{3}{2})$.

5.4.3 Proposition. For all $\alpha \in [1,2]$, $0 \leq k \leq M_{\alpha}$ and $\mathbf{u} \in \Omega^k$, it holds that $T_{\mathbf{u}}(1), T_{\mathbf{u}}(1-\alpha) \in \{S^k(1), S^k(1) - \alpha\}.$

Proof. First consider $\alpha \in \left[\frac{3}{2}, 2\right]$. Then $M_{\alpha} = 1$ and the result trivially holds. Fix an $\alpha \in \left(1, \frac{3}{2}\right)$. Since T_0 and T_1 agree on I_5 we can find a sequence $\widehat{\omega} \in \Omega^{\mathbb{N}}$ with $\widehat{\omega}_1 = 0$ that gives

$$T_{\widehat{\omega}_1^k}(1) = S^k(1) \quad \text{for all } k \ge 0.$$

Note that $1 \in I_5$ and from $\alpha \in (1, \frac{3}{2})$ we get $1 - \alpha \in I_2$, so

$$T_0(1-\alpha) = T_0(1) = T_1(1) = 2 - \alpha = S(1) \quad \text{and} \quad T_1(1-\alpha) = 2 - 2\alpha = S(1-\alpha).$$
(5.16)

Hence, from the first iterate on, the orbits of 1 and $1 - \alpha$ under the deterministic map S are contained in the orbit of $1 - \alpha$ under the random map R. To prove the statement, we therefore only have to consider $T^n_{\omega}(1-\alpha)$ for any $\omega \in \Omega^{\mathbb{N}}$ and $n \geq 1$. In particular (5.16) implies that

$$T_{\widehat{\omega}_1^k}(1-\alpha) = S^k(1)$$

for all $k \geq 1$. We prove the statement by induction.

The statement obviously holds for k = 0 and by (5.16) also for k = 1. Let $1 \le n < M_{\alpha}$ and suppose the statement holds for all $k \le n$. Then

$$T_{\widehat{\omega}_1^n}(1-\alpha) = S^n(1) \quad \text{and} \quad T_{\omega_1^n}(1-\alpha) \in \{S^n(1), S^n(1)-\alpha\} \quad \text{for all } \omega \in \Omega^{\mathbb{N}}.$$

By Lemma 5.4.2 there are three cases.

1. If $S^n(1) \in I_4$, then $S^{n+1}(1) = 2S^n(1)$ and $S^n(1) - \alpha \in I_1$. So for the random images we get

$$T_0(S^n(1)) = 2S^n(1), \quad T_1(S^n(1)) = 2S^n(1) - \alpha,$$

and

$$T_0(S^n(1) - \alpha) = T_1(S^n(1) - \alpha) = 2S^n(1) - \alpha.$$

2. If $S^n(1) \in I_5$ and $S^n(1) - \alpha \in I_2$, then

$$T_0(S^n(1)) = T_1(S^n(1)) = 2S^n(1) - \alpha = S^{n+1}(1)$$

and

$$T_0(S^n(1) - \alpha) = 2S^n(1) - \alpha, \quad T_1(S^n(1)) = 2S^n(1) - 2\alpha.$$

3. If $S^n(1) \in I_5$ and $S^n(1) \in I_1$ (so $n = M_\alpha - 1$), then

$$T_0(S^n(1)) = T_1(S^n(1)) = 2S^n - \alpha = S^{n+1}(1)$$

and

$$T_0(S^n(1) - \alpha) = T_1(S^n(1) - \alpha) = 2S^n(1) - \alpha = S^{n+1}(1).$$

Hence, for all $\mathbf{u} \in \Omega^n$ and $j = 0, 1, T_{\mathbf{u}j}(1-\alpha) \in \{S^{n+1}(1), S^{n+1}(1) - \alpha\}$, which gives the result.

From this proposition we can deduce that matching is prevalent for the family $\{R_{\alpha}\}$ and we can find the precise matching times. We first prove the following lemma, stating that all the orbit points $S^n(1), S^n(1-\alpha)$ up to the moment of matching are different.

5.4.4 Lemma. For each $k < M_{\alpha}$ the set $\{S^n(1), S^n(1-\alpha) : 0 \le n \le k\}$ has 2(k+1) elements.

Proof. Since $k < M_{\alpha}$ it follows from (5.15) that $S^n(1) \neq S^n(1-\alpha)$ for each n. It also cannot hold that there are $0 \leq n < k < M_{\alpha}$ such that $S^n(1) = S^k(1-\alpha)$ or $S^k(1) = S^n(1-\alpha)$, since this would imply that $|S^k(1) - S^n(1)| = \alpha$ and that would contradict the fact that $S^n(1), S^k(1) \in I_4 \cup I_5$. This leaves the possibility that there are $0 \leq n < k < M_{\alpha}$ such that $S^n(1) = S^k(1)$, i.e., that the orbit of 1 under S is ultimately periodic, or $S^n(1-\alpha) = S^k(1-\alpha)$. Assume $S^n(1) = S^k(1)$ for some n < k. It follows that $S^n(1-\alpha) = S^n(1) - \alpha = S^k(1) - \alpha = S^k(1-\alpha)$, so the orbit of $1 - \alpha$ is also ultimately periodic and by Proposition 5.4.3 all these orbit points lie at distance α of the corresponding orbit points of 1. This contradicts the fact that α is a matching parameter. Hence, the set $\{S^n(1), S^n(1-\alpha) : 0 \leq n \leq k\}$ has 2(k+1) elements.

5.4.5 Theorem. For Lebesgue almost all parameters $\alpha \in [1,2]$ the map R_{α} has strong random matching with $M = M_{\alpha} + 1$, where M_{α} is given by (5.14), and $Y = \{S^{M_{\alpha}}(1)\}$. Moreover, R_{α} does not satisfy the conditions of strong random matching for any K < M.

Proof. First consider $\alpha \in \begin{bmatrix} \frac{3}{2}, 2 \end{bmatrix}$. Then $T_j(1 - \alpha) = 2 - \alpha = T_j(1)$ for j = 0, 1, so random matching occurs for R with M = 2 and $Y = \{2 - \alpha\}$ and both parts of the theorem hold.

Now, fix $\alpha \in [1, \frac{3}{2})$ such that $S = S_{\alpha}$ has matching. Then, $S^{k}(1) \neq S^{k}(1-\alpha)$ for $1 \leq k < M_{\alpha}$ and $S^{M_{\alpha}-1}(1) \in (\frac{1}{2}, \alpha - \frac{1}{2})$, so that $S^{M_{\alpha}}(1) = 2S^{M_{\alpha}-1}(1) - \alpha$. By Proposition 5.4.3 for each $\mathbf{u} \in \Omega^{M_{\alpha}-1}$ either

$$T_{\mathbf{u}}(1-\alpha) = S^{M_{\alpha}-1}(1) > \frac{1}{2}$$

or

$$T_{\mathbf{u}}(1-\alpha) = S^{M_{\alpha}-1}(1) - \alpha < -\frac{1}{2}.$$

In both cases this leads to $T_{\mathbf{u}j}(1-\alpha) = 2S^{M_{\alpha}-1}(1) - \alpha$ for both j = 0, 1. The same statement holds for $T_{\mathbf{u}}(1)$, so that for $c = \frac{1}{2}$ we therefore have

$$T_{1\mathbf{u}j}\left(\frac{1}{2}^{-}\right) = T_{0\mathbf{u}j}\left(\frac{1}{2}^{+}\right) = T_{1\mathbf{u}j}\left(\frac{1}{2}^{+}\right) = T_{\mathbf{u}j}(1-\alpha) = S^{M_{\alpha}}(1)$$

and

$$T_{0\mathbf{u}j}\left(\frac{1}{2}^{-}\right) = T_{\mathbf{u}j}(1) = S^{M_{\alpha}}(1).$$

Hence, we can take $Y_{\frac{1}{2}} = \{S^{M_{\alpha}}(1)\}$. Since this set contains one element only and the maps T_j have the same constant slope, condition (5.5) from Definition 5.2.2 follows immediately. The first part of the theorem now follows since the deterministic maps S_{α} have matching for Lebesgue almost all parameters α . For the critical points $c \neq \frac{1}{2}$ the statement follows by symmetry.

For the second part we assume for $\alpha \in [1, \frac{3}{2})$ that $S = S_{\alpha}$ has matching and we proceed by contradiction. Therefore, assume that R_{α} satisfies the conditions of Definition 5.2.1 and Definition 5.2.2 for $c = \frac{1}{2}$ for some minimal $1 \le K < M = M_{\alpha} + 1$. Suppose that $S^n(1) \in Y_{\frac{1}{2}}$ for some n < K - 1. By Lemma 5.4.4 any **u** for which $T_{\mathbf{u}}(\frac{1}{2}^{\pm}) = S^n(1)$ has length $|\mathbf{u}| = n + 1$. Together with (5.5) and the fact that the maps $T_{\alpha,0}$ and $T_{\alpha,1}$ both have constant slope 2, this implies that

$$\sum_{\mathbf{u}\in\Omega(S^n(1))^-} p_{\mathbf{u}} = \sum_{\mathbf{u}\in\Omega(S^n(1))^+} p_{\mathbf{u}}.$$
(5.17)

For any $\mathbf{u} \in \Omega^{n+1} \setminus \Omega(S^n(1))^-, \mathbf{u}' \in \Omega^{n+1} \setminus \Omega(S^n(1))^+$ we have by Proposition 5.4.3 that $T_{\mathbf{u}}(\frac{1}{2}^-) = T_{\mathbf{u}'}(\frac{1}{2}^+) = S^n(1) - \alpha$. Furthermore,

$$1 = \sum_{\mathbf{u}\in\Omega^{n+1}} p_{\mathbf{u}} = \sum_{\mathbf{u}\in\Omega(S^n(1))^-} p_{\mathbf{u}} + \sum_{\mathbf{u}\in\Omega^{n+1}\setminus\Omega(S^n(1))^-} p_{\mathbf{u}}$$
$$= \sum_{\mathbf{u}\in\Omega(S^n(1))^+} p_{\mathbf{u}} + \sum_{\mathbf{u}\in\Omega^{n+1}\setminus\Omega(S^n(1))^+} p_{\mathbf{u}}.$$

From (5.17) and Proposition 5.4.3 we see that

$$\sum_{\mathbf{u}\in\Omega(S^n(1-\alpha))^-} p_{\mathbf{u}} = \sum_{\mathbf{u}\in\Omega^{n+1}\setminus\Omega(S^n(1))^-} p_{\mathbf{u}} = \sum_{\mathbf{u}\in\Omega^{n+1}\setminus\Omega(S^n(1))^+} p_{\mathbf{u}} = \sum_{\mathbf{u}\in\Omega(S^n(1-\alpha))^+} p_{\mathbf{u}}.$$

This implies that the conditions of Definition 5.2.1 and Definition 5.2.2 hold with $M_{\frac{1}{2}} = n + 1$ and $Y_{\frac{1}{2}} = \{S^n(1), S^n(1-\alpha)\}$, contradicting the minimality of K. In a similar way we can exclude the possibility that $S^n(1-\alpha) \in Y$ for n < K - 1. Since there is an $\widetilde{\omega} \in \Omega^{\mathbb{N}}$ such that for each k < M - 1, $T_{\widetilde{\omega}_1^k}(1-\alpha) = S^k(1-\alpha) = S^k(1) - \alpha$, it must hold that

$$Y_{\frac{1}{2}} = \{S^{K-1}(1), S^{K-1}(1) - \alpha\}$$

To conclude the proof we show that for this set $Y_{\frac{1}{2}}$ condition (5.5) cannot hold. By the constant slope, condition (5.5) can be rephrased as

$$\begin{cases} \sum_{\mathbf{u}\in\Omega(S^{K-1}(1))^{-}} p_{\mathbf{u}} - \sum_{\mathbf{u}\in\Omega(S^{K-1}(1))^{+}} p_{\mathbf{u}} = 0, \\ \sum_{\mathbf{u}\in\Omega(S^{K-1}(1)-\alpha)^{-}} p_{\mathbf{u}} - \sum_{\mathbf{u}\in\Omega(S^{K-1}(1)-\alpha)^{+}} p_{\mathbf{u}} = 0. \end{cases}$$
(5.18)

and by Lemma 5.4.4 any $\mathbf{u} \in \Omega(S^{K-1}(1))^{\pm} \cup \Omega(S^{K-1}(1) - \alpha)^{\pm}$ has length K. Since $K < M_{\alpha} + 1$, so $K - 2 < M_{\alpha} - 1$, Lemma 5.4.2 tells us that there are only two possibilities:

- 1. $S^{K-2}(1) \in I_4$ and $S^{K-2}(1) \alpha \in I_1$;
- 2. $S^{K-2}(1) \in I_5$ and $S^{K-2}(1) \alpha \in I_2$.

If case 1. holds, then $T_0(S^{K-2}(1)) = S^{K-1}(1)$ and

$$T_1(S^{K-2}(1)) = T_0(S^{K-2}(1) - \alpha) = T_1(S^{K-2}(1) - \alpha) = S^{K-1}(1) - \alpha,$$

so that (5.18) becomes

$$\begin{cases} \sum_{\mathbf{u}\in\Omega(S^{K-2}(1))^{-}} p_{\mathbf{u}}p_{0} - \sum_{\mathbf{u}\in\Omega(S^{K-2}(1))^{+}} p_{\mathbf{u}}p_{0} = 0, \\ \sum_{\mathbf{u}\in\Omega(S^{K-2}(1))^{-}} p_{\mathbf{u}}p_{1} + \sum_{\mathbf{u}\in(\Omega(S^{K-2}(1)-\alpha)^{-})} p_{\mathbf{u}} - \sum_{\mathbf{u}\in\Omega(S^{K-2}(1))^{+}} p_{\mathbf{u}}p_{1} - \sum_{\mathbf{u}\in(\Omega(S^{K-2}(1)-\alpha)^{+})} p_{\mathbf{u}} = 0. \end{cases}$$

The last system of equations implies

$$\begin{cases} \sum_{\mathbf{u}\in\Omega(S^{K-2}(1))^{-}} p_{\mathbf{u}} - \sum_{\mathbf{u}\in\Omega(S^{K-2}(1))^{+}} p_{\mathbf{u}} &= 0, \\ \sum_{\mathbf{u}\in(\Omega(S^{K-2}(1)-\alpha)^{-}} p_{\mathbf{u}} - \sum_{\mathbf{u}\in(\Omega(S^{K-2}(1)-\alpha)^{+}} p_{\mathbf{u}} &= 0, \end{cases}$$

which contradicts the minimality of K. For the second case, the same contradiction is obtained in a similar way.

5.4.6 Remark. From the previous result we see that matching occurs for the random systems R_{α} for the same parameters α and at the same time as for the deterministic systems S_{α} . [DK17] contains a complete description of the matching intervals of the maps S_{α} . The interval [1, 2] can be divided into intervals of parameters for which matching of the maps S_{α} occurs after the same number of steps. By the above, these matching intervals also apply to the systems R_{α} .

§5.4.3 A formula for the stationary measure

Let λ be the Lebesgue measure on [-1, 1]. The existence of an invariant measure of the form $m_{\mathbf{p}} \times \mu_{\mathbf{p}}$ with $\mu_{\mathbf{p}} \ll \lambda$ for the random symmetric doubling maps R_{α} is guaranteed by the results of [P84, M85]. Furthermore, since T_0 is expanding and has a unique absolutely continuous invariant measure, it follows from [P84, Corollary 7] that also for R_{α} there is a unique measure $m_{\mathbf{p}} \times \mu_{\mathbf{p}}$ and that R_{α} is ergodic with respect to this measure. To show that $R_{\alpha} \in \mathcal{R}_A$, we check conditions (c1), (c2), (c3). (c1) is immediate and (c3) follows from the constant slope 2 of the maps $T_{\alpha,0}$ and $T_{\alpha,1}$. We check condition (5.6). Note that for any $\alpha \neq 2$,

$$\frac{\sum_{j \in \Omega} \frac{p_j}{k_{3,j}} d_{3,j}}{1 - \sum_{j \in \Omega} \frac{p_j}{k_{3,j}}} = \frac{\sum_{j \in \Omega} \frac{p_j}{2} 0}{1 - \sum_{j \in \Omega} \frac{p_j}{2}} = 0$$

and

$$\frac{\sum_{j\in\Omega}\frac{p_j}{k_{1,j}}d_{1,j}}{1-\sum_{j\in\Omega}\frac{p_j}{k_{1,j}}} = \frac{\sum_{j\in\Omega}\frac{p_j}{2}\alpha}{1-\sum_{j\in\Omega}\frac{p_j}{2}} = 2\alpha \neq 0.$$

Then Theorem 3.5.3 implies that an explicit formula for the density of this measure can be found via the algebraic procedure in Chapter 3 and from Theorem 5.2.3 and Theorem 5.4.5 we know that for Lebesgue almost all parameters α this density is piecewise constant. We will execute the procedure from Chapter 3 and start by introducing the same notation as in Chapter 3. Since Ω consists of two elements only, from now on we will just use p as an index instead of \mathbf{p} whenever appropriate.

Denote by $a_{i,j}$ and $b_{i,j}$ the left and right limits at each critical point $c_i \in C$, i.e., for $1 \leq i \leq 4$ and $j \in \Omega$:

$$a_{i,j} = T_j(c_i^-) = \lim_{x \uparrow c_i} T_j(x),$$
 and $b_{i,j} = T_j(c_i^+) = \lim_{x \downarrow c_i} T_j(x).$

The images of the critical points are then given by

$$\begin{aligned} a_{1,0} &= a_{1,1} = b_{1,0} = \alpha - 1, & b_{1,1} = -1, \\ a_{2,1} &= b_{2,0} = b_{2,1} = 1 - \alpha, & a_{2,0} = 1, \\ a_{3,0} &= a_{3,1} = b_{3,0} = \alpha - 1, & b_{3,1} = -1, \\ a_{4,1} &= b_{4,0} = b_{4,1} = 1 - \alpha, & a_{4,0} = 1. \end{aligned}$$

For $y \in [-1, 1]$ and $1 \le n \le 4$ set

$$\mathrm{KI}_{n}(y) = \sum_{k \ge 1} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{\mathbf{u}}}{2^{k}} \mathbb{1}_{I_{n}}(T_{\mathbf{u}_{1}^{k-1}}(y)).$$
(5.19)

The quantity $\operatorname{KI}_n(y)$ weighs the number of times the random orbits of y enters the interval I_n . The weight depends on the length and probability of each path $\omega \in \Omega^{\mathbb{N}}$ leading the point y to I_n . The fundamental matrix $A = (A_{n,i})$ of R is the 5×4 matrix with entries

$$A_{n,i} = \begin{cases} \sum_{j \in \Omega} p_j (1 + \mathrm{KI}_n(a_{i,j}) - \mathrm{KI}_n(b_{i,j})), & \text{for } n = i, \\ \sum_{j \in \Omega} p_j (\mathrm{KI}_n(a_{i,j}) - \mathrm{KI}_n(b_{i,j}) - 1), & \text{for } n = i + 1, \\ \sum_{j \in \Omega} p_j (\mathrm{KI}_n(a_{i,j}) - \mathrm{KI}_n(b_{i,j})), & \text{else.} \end{cases}$$

Since for R there is a unique invariant probability measure $m_p \times \mu_p$ with $\mu_p \ll \lambda$, Theorem 3.5.3 implies that the null space of the matrix A is one-dimensional. According to Theorem 3.4.1 there is a unique vector $\gamma = (\gamma_1, \gamma_2, \gamma_3, \gamma_4) \in \mathbb{R}^4 \setminus \{\mathbf{0}\}$ with $A\gamma = \mathbf{0}$ and such that the probability density f_p of μ_p has the form (5.7). Using the values of $a_{i,j}$ and $b_{i,j}$ computed above, we can reduce this to

$$f_{p} = (\gamma_{1} + \gamma_{3}) \sum_{k \ge 0} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{1\mathbf{u}}}{2^{k+1}} \left(\mathbf{1}_{[-1,T_{\mathbf{u}}(\alpha-1))} - \mathbf{1}_{[-1,T_{\mathbf{u}}(-1))} \right) + (\gamma_{2} + \gamma_{4}) \sum_{k \ge 0} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{0\mathbf{u}}}{2^{k+1}} \left(\mathbf{1}_{[-1,T_{\mathbf{u}}(1))} - \mathbf{1}_{[-1,T_{\mathbf{u}}(1-\alpha))} \right).$$
(5.20)

By symmetry to determine f_p it is enough to know the random orbits of 1 and $1 - \alpha$ only. From (5.20) we see that the density is piecewise constant when the orbits of 1 and $1 - \alpha$ are finite or when they merge with the same weight. In the former case the

map admits a Markov partition, the latter case happens if R exhibits strong random matching. We focus on the second situation, since we know from Theorem 5.4.5 that this holds for Lebesgue almost all parameters.

Fix an $\alpha \in [1, 2]$ such that R presents strong random matching. Let M be as in Theorem 5.4.5. Then for each i, j, n,

$$\mathrm{KI}_{n}(a_{i,j}) - \mathrm{KI}_{n}(b_{i,j}) = \sum_{k=1}^{M-1} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{\mathbf{u}}}{2^{k}} \big(\mathbb{1}_{I_{n}}(T_{\mathbf{u}_{1}^{k-1}}(a_{i,j})) - \mathbb{1}_{I_{n}}(T_{\mathbf{u}_{1}^{k-1}}(b_{i,j})) \big).$$

From Lemma 5.4.2 and the symmetry of the map we get

$$KI_3(1) - KI_3(1 - \alpha) = 0 = KI_3(-1) - KI_3(\alpha - 1),$$

implying that $A_{3,1} = A_{3,4} = 0$, $A_{3,2} = -1$ and $A_{3,3} = 1$. Hence, any solution vector $\hat{\gamma}$ for $A\hat{\gamma} = \mathbf{0}$ has the form $\hat{\gamma} = (\hat{\gamma}_1, \hat{\gamma}_2, \hat{\gamma}_2, \hat{\gamma}_3)$ and (5.20) becomes

$$f_{p} = (\gamma_{1} + \gamma_{2}) \frac{p_{1}}{2} \sum_{k=0}^{M-2} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{\mathbf{u}}}{2^{k}} \left(\mathbf{1}_{[-1,T_{\mathbf{u}}(\alpha-1))} - \mathbf{1}_{[-1,T_{\mathbf{u}}(-1))} \right) + (\gamma_{2} + \gamma_{3}) \frac{p_{0}}{2} \sum_{k=0}^{M-2} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{\mathbf{u}}}{2^{k}} \left(\mathbf{1}_{[-1,T_{\mathbf{u}}(1))} - \mathbf{1}_{[-1,T_{\mathbf{u}}(1-\alpha))} \right),$$
(5.21)

where $\gamma = (\gamma_1, \gamma_2, \gamma_2, \gamma_3)$ is the unique non-trivial vector in the null space of the fundamental matrix A that makes f_p into a probability density function. In the next section we will derive a number of properties of f_p with the goal of determining the frequency of the digit 0 in the signed binary expansions of $m_p \times \mu_p$ typical points.

§5.4.4 Minimal weight expansions

Recall from (5.13) that the random signed binary expansion of a point (ω, x) has a digit 0 in the *n*-th position precisely if

$$R^{n-1}(\omega, x) \in [1] \times I_2 \cup \Omega^{\mathbb{N}} \times I_3 \cup [0] \times I_4 =: D_0.$$

Since R is ergodic with respect to $m_p \times \mu_p$, it follows from Birkhoff's Ergodic Theorem that the frequency of the digit 0 in $m_p \times \mu_p$ -almost all (ω, x) equals

$$\pi_0(\alpha, p) := \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{1}_{D_0}(R^k(\omega, x)) = (1-p)\mu_p(I_2) + \mu_p(I_3) + p\mu_p(I_4).$$
(5.22)

To give an example, consider $\alpha = 1$, see Figure 5.11(a). It is straightforward to check that the probability density $f_p = (1 - p)1_{[-1,0]} + p1_{[0,1]}$ is invariant. This gives

$$\pi_0(1,p) = p\mu_p\left(\left[0,\frac{1}{2}\right]\right) + (1-p)\mu_p\left(\left[-\frac{1}{2},0\right]\right) = \frac{p^2 + (1-p)^2}{2} \le \frac{1}{2}$$
(5.23)

with equality only for p = 0 or p = 1.

To estimate $\pi_0(\alpha, p)$ for other values of α we use a few lemmata. For $k \geq 1$ set $E_k = \{ \mathbf{u} \in \Omega^k : T_{\mathbf{u}}(1) = S^k(1) \}$ and $F_k = \{ \mathbf{u} \in \Omega^k : T_{\mathbf{u}}(1-\alpha) = S^k(1) \}$. Also, use $(b_n)_{n\geq 1}$ to denote the digits in the signed binary expansion of 1 generated by S, i.e.,

$$b_n = \begin{cases} -1, & \text{if } S^{n-1}(1) < -\frac{1}{2}, \\ 0, & \text{if } -\frac{1}{2} \le S^{n-1}(1) \le \frac{1}{2}, \\ 1, & \text{if } S^{n-1}(1) > \frac{1}{2}. \end{cases}$$

Write $\mathbf{b}_k = b_1 \cdots b_k$ for any $k \ge 1$.

5.4.7 Lemma. For all $1 \le k < M - 1$, $F_k \subseteq E_k$ and $E_k \setminus F_k = \{\mathbf{b}_k\}$.

Proof. First note that the *n*-th signed binary digit of 1 generated by S, n < M - 1, equals 0 if $S^{n-1}(1) \in I_4$ and 1 if $S^{n-1}(1) \in I_5$. We prove the statement by induction. For k = 1 we have $E_1 = \{0, 1\}$ and $F_1 = \{0\}$. Assume the statement holds for some $1 \leq k < M - 2$. If $S^k(1) = T_{\mathbf{b}_k}(1) \in I_4$, then $b_{k+1} = 0$ and we know from the assumptions and since $S^k(1) - \alpha \in I_1$ that

$$T_{\mathbf{b}_k 0}(1) = S^{k+1}(1), \ T_{\mathbf{b}_k 1}(1) = T_{\mathbf{b}_k 0}(1-\alpha) = T_{\mathbf{b}_k 1}(1-\alpha) = S^{k+1}(1) - \alpha$$

Hence, $\mathbf{b}_k 0 \in E_{k+1} \setminus F_{k+1}$ and $\mathbf{b}_k 1 \notin E_{k+1} \cup F_{k+1}$. If $S^k(1) = T_{\mathbf{b}_k}(1) \in I_5$, then $b_{k+1} = 1$ and

$$T_{\mathbf{b}_k 0}(1) = T_{\mathbf{b}_k 1}(1) = T_{\mathbf{b}_k 0}(1-\alpha) = S^{k+1}(1), \ T_{\mathbf{b}_k 1}(1-\alpha) = S^{k+1}(1) - \alpha.$$

So, $\mathbf{b}_k 1 \in E_{k+1} \setminus F_{k+1}$ and $\mathbf{b}_k 0 \in E_{k+1} \cap F_{k+1}$. For any other $\mathbf{u} \in \Omega^k$ it holds that $T_{\mathbf{u}}(1) = T_{\mathbf{u}}(1-\alpha)$, so that either $\mathbf{u}_j \in E_{k+1} \cap F_{k+1}$ or $\mathbf{u}_j \notin E_{k+1} \cup F_{k+1}$, j = 0, 1. This gives the statement.

5.4.8 Lemma. The density f_p is constant and equal to $\frac{1}{\alpha}$ on the interval $[1-\alpha, \alpha-1]$.

Proof. For any $\mathbf{u} \in \Omega^k$, write $\bar{\mathbf{u}} = (1 - u_1) \cdots (1 - u_k)$ and for a subset $E \subseteq \Omega^k$ write $\bar{E} = \{\mathbf{u} \in \Omega^k : \bar{\mathbf{u}} \in E\}$. By Lemma 5.4.7 we have for each k < M,

$$\delta_k := \sum_{\mathbf{u}\in E_k} \frac{p_{\mathbf{u}}}{2^k} - \sum_{\mathbf{u}\in F_k} \frac{p_{\mathbf{u}}}{2^k} = \frac{p_{\mathbf{b}_k}}{2^k}, \quad \bar{\delta}_k := \sum_{\mathbf{u}\in \bar{E}_k^c} \frac{p_{\mathbf{u}}}{2^k} - \sum_{\mathbf{u}\in \bar{F}_k^c} \frac{p_{\mathbf{u}}}{2^k} = \frac{p_{\bar{\mathbf{b}}_k}}{2^k},$$

Recall the formula for the density f_p from (5.21). Using Proposition 5.4.3 we get

$$\begin{split} \frac{p_0}{2} \sum_{k=0}^{M-2} \sum_{\mathbf{u} \in \Omega^k} \frac{p_{\mathbf{u}}}{2^k} \Big(\mathbf{1}_{[-1,T_{\mathbf{u}}(1))} - \mathbf{1}_{[-1,T_{\mathbf{u}}(1-\alpha))} \Big) \\ &= \frac{p_0}{2} \sum_{k=0}^{M-2} \sum_{\substack{\mathbf{u} \in \Omega^k: \\ T_{\mathbf{u}}(1) = S^k(1), \\ T_{\mathbf{u}}(1-\alpha) = S^k(1) - \alpha}} \frac{p_{\mathbf{u}}}{2^k} \mathbf{1}_{[T_{\mathbf{u}}(1),T_{\mathbf{u}}(1-\alpha))} \\ &= \frac{p_0}{2} \sum_{k=0}^{M-2} \sum_{\substack{\mathbf{u} \in \Omega^k: \\ T_{\mathbf{u}}(1) = S^k(1) - \alpha, \\ T_{\mathbf{u}}(1-\alpha) = S^k(1)}} \frac{p_{\mathbf{u}}}{2^k} \mathbf{1}_{[T_{\mathbf{u}}(1),T_{\mathbf{u}}(1-\alpha))} \\ &= \frac{p_0}{2} \sum_{k=0}^{M-2} \delta_k \mathbf{1}_{[S^k(1)-\alpha,S^k(1))}. \end{split}$$

For the other side it holds similarly using the symmetry of the system that

$$\frac{p_1}{2} \sum_{k=0}^{M-2} \sum_{\mathbf{u} \in \Omega^k} \frac{p_{\mathbf{u}}}{2^k} \left(\mathbf{1}_{[-1,T_{\mathbf{u}}(\alpha-1))} - \mathbf{1}_{[-1,T_{\mathbf{u}}(-1))} \right) = \frac{p_1}{2} \sum_{k=0}^{M-2} \bar{\delta}_k \mathbf{1}_{[-S^k(1),\alpha-S^k(1))}.$$

By (5.15) we have for all k < M - 1,

$$S^{k}(1), \alpha - S^{k}(1) \in [\alpha - 1, 1]$$
 and $S^{k}(-1), S^{k}(\alpha - 1) \in [-1, 1 - \alpha],$

so that on $[1 - \alpha, \alpha - 1]$ we obtain

$$f_p \mid_{[1-\alpha,\alpha-1]} (x) = (\gamma_1 + \gamma_2) \frac{p_1}{2} \sum_{k=0}^{M-2} \bar{\delta}_k + (\gamma_2 + \gamma_3) \frac{p_0}{2} \sum_{k=0}^{M-2} \delta_k.$$

Since f_p is a probability density it follows that

$$1 = \int_{[-1,1]} f_p \, d\lambda = (\gamma_1 + \gamma_2) \frac{p_1}{2} \sum_{k=0}^{M-2} \bar{\delta}_k \alpha + (\gamma_2 + \gamma_3) \frac{p_0}{2} \sum_{k=0}^{M-2} \delta_k \alpha.$$
(5.24)

Hence,

$$f_p \mid_{[1-\alpha,\alpha-1]} (x) = \frac{1}{\alpha},$$

which gives the result.

With this information we can compute $\pi_0(\alpha, p)$ for $\alpha \in \left[\frac{3}{2}, 2\right]$. Since in this case $\alpha - 1 \geq \frac{1}{2}$ it follows from Lemma 5.4.8 that

$$\pi_0(\alpha, p) = \frac{\alpha - 1}{\alpha} + \frac{2 - \alpha}{2\alpha} = \frac{1}{2}.$$
 (5.25)

That is, for $\alpha \geq \frac{3}{2}$, and any $0 , the frequency of the digit 0 is equal to <math>\frac{1}{2}$ in the signed binary expansion of $m_p \times \mu_p$ -almost all (ω, x) . For the other values of α we need to do some more work.

5.4.9 Lemma. Let $\gamma = (\gamma_1, \gamma_2, \gamma_2, \gamma_3)$ be the unique vector in the null space of A that makes f_p into a probability density function. Then $\gamma_2 = \frac{1}{\alpha}$.

Proof. Since $S^k(1) \in I_4 \cup I_5$ for all k < M - 1 it follows from the definition of the function KI_n in (5.19) and Proposition 5.4.3 that for $y = 1, 1 - \alpha$,

$$\mathrm{KI}_{4}(y) + \mathrm{KI}_{5}(y) = \sum_{k=0}^{M-2} \sum_{j \in \Omega} \sum_{\mathbf{u} \in \Omega^{k}} \frac{p_{j}}{2} \frac{p_{\mathbf{u}}}{2^{k}} \mathbf{1}_{I_{4} \cup I_{5}}(T_{\mathbf{u}}(y)) = \frac{1}{2} \sum_{k=0}^{M-2} \sum_{\substack{\mathbf{u} \in \Omega^{k}:\\ T_{\mathbf{u}}(y) = S^{k}(1)}} \frac{p_{\mathbf{u}}}{2^{k}},$$

so that

$$\frac{1}{2} \sum_{k=0}^{M-2} \delta_k = \mathrm{KI}_4(1) - \mathrm{KI}_4(1-\alpha) + \mathrm{KI}_5(1) - \mathrm{KI}_5(1-\alpha).$$

A similar statement holds for -1 and $\alpha - 1$. The fourth and fifth line of the linear system $A\gamma = \mathbf{0}$ read

$$p_1(\mathrm{KI}_4(\alpha - 1) - \mathrm{KI}_4(-1))(\gamma_1 + \gamma_2) + p_0(\mathrm{KI}_4(1) - \mathrm{KI}_4(1 - \alpha))(\gamma_2 + \gamma_3) - \gamma_2 + \gamma_3 = 0$$

and

$$p_1(\mathrm{KI}_5(\alpha - 1) - \mathrm{KI}_5(-1))(\gamma_1 + \gamma_2) + p_0(\mathrm{KI}_5(1) - \mathrm{KI}_5(1 - \alpha))(\gamma_2 + \gamma_3) - \gamma_3 = 0,$$

respectively. Adding them up gives

$$\begin{split} \gamma_2 &= p_1(\gamma_1 + \gamma_2)(\mathrm{KI}_4(\alpha - 1) - \mathrm{KI}_4(-1) + \mathrm{KI}_5(\alpha - 1) - \mathrm{KI}_5(-1)) + \\ &\quad p_0(\gamma_2 + \gamma_3)(\mathrm{KI}_4(1) - \mathrm{KI}_4(1 - \alpha) + \mathrm{KI}_5(1) - \mathrm{KI}_5(1 - \alpha)) \\ &= \frac{p_1}{2}(\gamma_1 + \gamma_2)\sum_{k=0}^{M-2} \bar{\delta}_k + \frac{p_0}{2}(\gamma_2 + \gamma_3)\sum_{k=0}^{M-2} \delta_k. \end{split}$$

The result then follows from (5.24).

Combining Lemma 5.4.8 and Lemma 5.4.9 gives the following expression for the density f_p :

$$f_p = \left(\gamma_1 + \frac{1}{\alpha}\right) \frac{p_1}{2} \sum_{k=0}^{M-2} \frac{p_{\bar{\mathbf{b}}_k}}{2^k} \mathbf{1}_{[-S^k(1),\alpha-S^k(1))} + \left(\frac{1}{\alpha} + \gamma_3\right) \frac{p_0}{2} \sum_{k=0}^{M-2} \frac{p_{\mathbf{b}_k}}{2^k} \mathbf{1}_{[S^k(1)-\alpha,S^k(1))},$$
(5.26)

where $\mathbf{b}_k = b_1 \cdots b_k$ denote the first k digits in the signed binary expansion of 1 given by S.

5.4.10 Lemma. Let $\alpha \in (1, \frac{3}{2})$ be a parameter for which the random system R has strong random matching. Then both $\gamma_1, \gamma_3 \geq 0$. As a consequence, $f_p > 0$ and μ_p is equivalent to the Lebesgue measure.

Proof. Let $\gamma = (\gamma_1, \gamma_2, \gamma_2, \gamma_3)$ be the unique vector in the null space of A that makes f_p into a probability density. Set

$$y = \max_{k \in \{1, 2, \dots, M-2\}} \{ S^k(1), \alpha - S^k(1) \}.$$

By Lemma 5.4.4 we can assume that $y \neq 1$. Then

$$\mu_p([y,1]) = m_p \times \mu_p(R^{-1}(\Omega \times [y,1])) = p\mu_p\left(\left[\frac{y-\alpha}{2}, \frac{1-\alpha}{2}\right] \cup \left[\frac{y}{2}, \frac{1}{2}\right]\right).$$

By the definition of y one can see from (5.26) that

$$\mu_p([y,1]) = \frac{p(\gamma_2 + \gamma_3)}{2}(1-y).$$

Furthermore, $T_0(1 - \alpha) = 2 - \alpha$ and $T_1(1 - \alpha) = 2 - 2\alpha$, so in particular $y \ge \max\{2 - \alpha, 2\alpha - 2\}$. It follows that $1 - \alpha \le \frac{y - \alpha}{2} < \frac{1 - \alpha}{2}$. Thus by Lemma 5.4.8 and Lemma 5.4.9, $f_p \mid_{\left[\frac{y - \alpha}{2}, \frac{1 - \alpha}{2}\right]} = \gamma_2$. We proceed by showing that none of the points $S^k(1)$ or $\alpha - S^k(1), 1 \le k \le M - 2$, lie in the interval $\left[\frac{y}{2}, \frac{1}{2}\right]$, which then by (5.26) implies that the density f_p is also constant on the interval $\left[\frac{y}{2}, \frac{1}{2}\right]$. For $k = M - 2 = M_\alpha - 1$, matching for S implies that $\frac{1}{2} < S^{M-2}(1) < \alpha - \frac{1}{2}$ and $\alpha - S^{M-2}(1) > \frac{1}{2}$. Suppose there exists a $k \in \{1, 2, \dots, M - 3\}$ such that $\frac{y}{2} < S^k(1) < \frac{1}{2}$ (or $\frac{y}{2} < \alpha - S^k(1) < \frac{1}{2}$). Then $S^{k+1}(1) > y$ (or $\alpha - S^{k+1}(1) > y$), which gives a contradiction with the definition of y. The same holds for $\alpha - S^k(1)$. Hence, there is a constant $c \ge 0$ such that

$$\frac{p(\gamma_2 + \gamma_3)}{2}(1 - y) = \mu_p([y, 1]) = p\Big(\gamma_2 \frac{(1 - y)}{2} + c\frac{(1 - y)}{2}\Big).$$

So, $0 \le \mu_p(\left[\frac{y}{2}, \frac{1}{2}\right]) = c = \gamma_3$. The proof that $\gamma_1 \ge 0$ goes similarly. The fact that f_p is strictly positive and the equivalence of μ_p and the Lebesgue measure now follow from (5.26).

The following result can be proven in essentially the same way as [DK17, Theorem 4.1]. We include a proof here for convenience.

5.4.11 Lemma (cf. Theorem 4.1 of [DK17]). Fix $0 . The map <math>\alpha \mapsto \pi_0(\alpha, p)$ is continuous on $(1, \frac{3}{2})$.

Proof. In this proof we use $f_{\alpha} = f_{\alpha,p}$ to denote the unique density from (5.26). By (5.22), for the continuity of $\alpha \mapsto \pi_0(\alpha, p)$ it is sufficient to prove L^1 -convergence of the densities f_{α} ; i.e., for any sequence $\{\alpha_k\}_{k\geq 1} \subseteq (1, \frac{3}{2})$ converging to a fixed $\hat{\alpha} \in (1, \frac{3}{2})$, there is convergence $f_{\alpha_k} \to f_{\hat{\alpha}}$ in $L^1(\lambda)$. The proof of this fact goes along the following lines:

- 1. First we show that there is a uniform bound, i.e., independent of k, on the total variation and supremum norm of the densities f_{α_k} . It then follows from Helly's Selection Theorem that there is some subsequence of (f_{α_k}) for which an a.e. and L^1 limit \hat{f} exist.
- 2. We show that $\hat{f} = f_{\hat{\alpha}}$, which by the same proof implies that any subsequence of (f_{α_k}) has a further subsequence converging a.e. to the same limit $f_{\hat{\alpha}}$. Hence, (f_{α_k}) converges to $f_{\hat{\alpha}}$ in measure.
- 3. By the uniform integrability of (f_{α_k}) it then follows from Vitali's Convergence Theorem that the convergence of (f_{α_k}) to $f_{\hat{\alpha}}$ is in L^1 .

Step 1. and 2. use Perron-Frobenius operators. For j = 0, 1 the Perron-Frobenius operator $P_{\alpha,j}$ of $T_{\alpha,j}$ is uniquely defined by the equation

$$\int (P_{\alpha,j}f)g\,d\lambda = \int f(g\circ T_{\alpha,j})\,d\lambda \quad \forall f\in L^1(\lambda), \,g\in L^\infty(\lambda)$$

and the Perron-Frobenius operator P_{α} of R_{α} is then defined by $P_{\alpha}f = pP_{\alpha,0}f + (1-p)P_{\alpha,1}f$. Equivalently, P_{α} is uniquely defined by the equation

$$\int (P_{\alpha}f)g\,d\lambda = p \int f(g \circ T_{\alpha,0})\,d\lambda + (1-p) \int f(g \circ T_{\alpha,1})\,d\lambda \quad \forall f \in L^{1}(\lambda), \, g \in L^{\infty}(\lambda).$$
(5.27)

Since each R_{α} has a unique probability density f_{α} it follows from [P84, Theorem 1] that f_{α} is the L^1 limit of $(\frac{1}{n} \sum_{j=0}^{n-1} P_{\alpha}^j 1)_{n \geq 1}$ and it is the unique probability density that satisfies $P_{\alpha}f_{\alpha} = f_{\alpha}$. From [I12, Theorem 5.2] each f_{α} is a function of bounded variation. We proceed by finding uniform bounds on the total variation and supremum norm of these densities.

Fix $\hat{\alpha} \in (1, \frac{3}{2})$. For the second iterates of the Perron-Frobenius operators we have

$$P_{\alpha}^2 f = \sum_{i,j=0}^{1} p_i p_j P_{\alpha,j}(P_{\alpha,i}f).$$

Since the intervals of monotonicity of any of the maps $T_{\alpha,\mathbf{u}}$ for $\mathbf{u} \in \Omega^2$, only become arbitrarily small for α approaching 1 and $\frac{3}{2}$, we can find a uniform lower bound δ on the length of the intervals of monotonicity of any map $T_{\alpha,\mathbf{u}}$, $\mathbf{u} \in \Omega^2$, for all values α that are close enough to $\hat{\alpha}$. Applying [BG97, Lemma 5.2.1] to $T_{\alpha,j}$, j = 0, 1, and any of the second iterates $T_{\alpha,\mathbf{u}}$, $\mathbf{u} \in \Omega^2$, gives that

$$Var(P_{\alpha,j}f) \le Var(f) + \frac{1}{\delta} \|f\|_1 \quad \text{and} \quad Var(P_{\alpha,\mathbf{u}}f) \le \frac{1}{2} Var(f) + \frac{1}{2\delta} \|f\|_1,$$

where Var denotes the total variation over the interval [-1, 1]. Since these bounds do not depend on α, j, \mathbf{u} , the same estimates hold for P_{α} , so that for any function $f: [-1, 1] \to \mathbb{R}$ of bounded variation and any $n \ge 1$,

$$Var(P_{\alpha}^{n}f) \leq \frac{1}{2^{\lfloor n/2 \rfloor}} Var(f) + \left(2 + \frac{1}{\delta}\right) \|f\|_{1}.$$
(5.28)

Let $\{\alpha_k\}_{k\geq 1}$ with $\alpha_k \to \hat{\alpha}$ be a sequence for which the lower bound δ holds for each k. For each k and n, write $f_{k,n} = \frac{1}{n} \sum_{i=0}^{n-1} P_{\alpha_k} 1$. Since

$$\sup |f_{k,n}| \le Var(f_{k,n}) + \int f_{k,n} \, d\lambda,$$

it follows from (5.28) that there is a uniform constant C > 0 (independent of k, n) such that $Var(f_{k,n})$, $\sup |f_{k,n}| < C$. The same then holds for the limits f_{α_k} . Helly's Selection Theorem then gives the existence of a subsequence $\{k_i\}$ and a function \hat{f} of bounded variation, such that $f_{\alpha_{k_i}} \to \hat{f}$ in $L^1(\lambda)$ and λ -a.e. and with $Var(\hat{f})$, $\sup |\hat{f}| < C$. This finishes 1. By 2. and 3. above, what remains to finish the proof is to show that $P_{\hat{\alpha}}\hat{f} = \hat{f}$. By (5.27) it is enough to show that for any compactly supported C^1 function g: $[-1,1] \to \mathbb{R}$ it holds that

$$\left|\int (P_{\hat{\alpha}}\hat{f})g\,d\lambda - \int \hat{f}g\,d\lambda\right| = 0.$$

Note that

$$\begin{split} \left| \int (P_{\hat{\alpha}}\hat{f})g\,d\lambda - \int \hat{f}g\,d\lambda \right| \leq & p \left| \int \hat{f}(g \circ T_{\hat{\alpha},0})\,d\lambda - \int \hat{f}g\,d\lambda \right| + \\ & (1-p) \left| \int \hat{f}(g \circ T_{\hat{\alpha},1})\,d\lambda - \int \hat{f}g\,d\lambda \right|. \end{split}$$

For j = 0, 1 we can write

$$\begin{split} \left| \int \hat{f}(g \circ T_{\hat{\alpha},j}) \, d\lambda - \int \hat{f}g \, d\lambda \right| &\leq \left| \int \hat{f}(g \circ T_{\hat{\alpha},j}) \, d\lambda - \int f_{\alpha_{k_i}}(g \circ T_{\hat{\alpha},j}) \, d\lambda \right| \\ &+ \left| \int f_{\alpha_{k_i}}(g \circ T_{\hat{\alpha},j}) \, d\lambda - \int f_{\alpha_{k_i}}(g \circ T_{\alpha_{k_i},j}) \, d\lambda \right| \\ &+ \left| \int f_{\alpha_{k_i}}(g \circ T_{\alpha_{k_i},j}) \, d\lambda - \int \hat{f}g \, d\lambda \right|. \end{split}$$

The first and third integral on the right hand side can be bounded by $||g||_{\infty} ||\hat{f} - f_{\alpha_{k_i}}||_1 \to 0$. For the second integral, $||f_{\alpha_{k_i}}||_{\infty} < C$ and $\int |g \circ T_{\hat{\alpha},j} - g \circ T_{\alpha_{k_i},j}| \, d\lambda \to 0$ by the Dominated Convergence Theorem. Hence, $\hat{f} = f_{\hat{\alpha}}$ and $f_{\alpha_k} \to f_{\hat{\alpha}}$ in L^1 . \Box

Figure 5.12 shows a numerical approximation of the graph of the function $(\alpha, p) \mapsto \pi_0(\alpha, p)$. We can now prove that the maximal value of the frequency of the digit 0 is in fact $\frac{1}{2}$.



Figure 5.12: The graph of $(\alpha, p) \mapsto \pi_0(\alpha, p)$.

5.4.12 Theorem. For any $0 and any <math>\alpha \in [1, 2]$ the frequency $\pi_0(\alpha, p)$ is at most $\frac{1}{2}$ for $m_p \times \lambda$ -a.e. $(\omega, x) \in \Omega^{\mathbb{N}} \times [-1, 1]$.

Proof. For $\alpha \in \left[\frac{3}{2}, 2\right]$ the statement follows from (5.25) and for $\alpha = 1$ from (5.23). Let $\alpha \in \left(1, \frac{3}{2}\right)$. The deterministic map $T_{\alpha,0}$ has density $f_0 = \frac{1}{\alpha} \mathbf{1}_{[1-\alpha,1]}$ and $T_{\alpha,1}$ has $f_1 = \frac{1}{\alpha} \mathbf{1}_{[-1,\alpha-1]}$. Hence $\pi_0(\alpha, p) = \frac{1}{2}$ for p = 0, 1. Let $0 and let <math>\alpha$ be a parameter satisfying the conditions of Lemma 5.4.10. We know that f_p is constant and equal to $\frac{1}{\alpha}$ on $[1 - \alpha, \alpha - 1]$. For $x > \alpha - 1$ the density can be written as

$$\begin{split} f_p(x) &= \frac{1}{\alpha} - \left(\frac{(1-p)(\gamma_1 + \gamma_2)}{2} \sum_{k=0}^{M-2} \frac{p_{\bar{\mathbf{b}}_k}}{2^k} \mathbf{1}_{[\alpha-1,x]}(\alpha - S^k(1)) \right. \\ &+ \frac{p(\gamma_2 + \gamma_3)}{2} \sum_{k=0}^{M-2} \frac{p_{\mathbf{b}_k}}{2^k} \mathbf{1}_{[\alpha-1,x]}(S^k(1)) \right) \\ &= \frac{1}{\alpha} - \frac{(1-p)(\gamma_1 + \gamma_2)}{2} - \left(\frac{(1-p)(\gamma_1 + \gamma_2)}{2} \sum_{k=1}^{M-2} \frac{p_{\bar{\mathbf{b}}_k}}{2^k} \mathbf{1}_{[\alpha-1,x]}(\alpha - S^k(1)) \right. \\ &+ \frac{p(\gamma_2 + \gamma_3)}{2} \sum_{k=1}^{M-2} \frac{p_{\mathbf{b}_k}}{2^k} \mathbf{1}_{[\alpha-1,x]}(S^k(1)) \right) \\ &\leq \frac{1}{\alpha} - \frac{(1-p)(\gamma_1 + \gamma_2)}{2}. \end{split}$$

Similarly, for $x < 1 - \alpha$ we get $f_p(x) \le \frac{1}{\alpha} - \frac{p(\gamma_2 + \gamma_3)}{2}$. By (5.22) and Lemma 5.4.10,

$$\pi_{0}(\alpha, p) = \frac{\alpha - 1}{\alpha} + \frac{\alpha - 1}{2\alpha} + p\mu_{\alpha, p}\left(\left[\alpha - 1, \frac{1}{2}\right]\right) + (1 - p)\mu_{\alpha, p}\left(\left[-\frac{1}{2}, 1 - \alpha\right]\right)$$

$$\leq \frac{3(\alpha - 1)}{2\alpha} + \frac{3 - 2\alpha}{2\alpha}\left(1 - \frac{p(1 - p)\alpha}{2}\min\{\gamma_{1} + \gamma_{2}, \gamma_{2} + \gamma_{3}\}\right)$$

$$= \frac{1}{2} - \frac{3 - 2\alpha}{2}\frac{p(1 - p)}{2}\min\{\gamma_{1} + \gamma_{2}, \gamma_{2} + \gamma_{3}\}$$

$$< \frac{1}{2}.$$

Since matching holds for Lebesgue almost all parameters α , the statement now follows from Lemma 5.4.11 and the equivalence of μ_p and the Lebesgue measure.

§5.5 Final remarks

§5.5.1 On the symmetric doubling maps

The numerical approximation of the graph of $(\alpha, p) \mapsto \pi_0(\alpha, p)$ shown in Figure 5.12 seems to suggest some other features of the map that we have not proved. Firstly, it suggests some symmetry. In fact it can be shown that for each fixed α and any $x \in$ [0,1], it holds that $f_p(x) = f_{1-p}(-x)$. For this one needs to consider the fundamental matrix \tilde{A} corresponding to the random system \tilde{R}_{α} obtained by switching the roles of p and 1 - p. Then using the permutation (12)(45), one can relate various of the quantities involved for \tilde{A} to the fundamental matrix A of R_{α} .

Secondly, for any matching parameter α and any $0 the density <math>f_{\alpha,p}$ is a finite combination of indicator functions, whose supports depend on the position of the points in the set $\{S^k(1), \alpha - S^k(1)\}_{k=0}^{M-2}$ and whose coefficients are polynomials in p. So, for such a fixed α and any $x \in [-1, 1]$, the map $p \mapsto f_{\alpha,p}(x)$ is continuous in p.

Thirdly, the graph also suggests that the map presents a minimum at $p = \frac{1}{2}$. Using the above two facts we were only able to show the following:

5.5.1 Proposition. Let $\alpha \in [1, 2]$ be such that R has strong random matching. Then the map $p \mapsto \pi_0(\alpha, p)$ has an extremal value at $p = \frac{1}{2}$.

Proof. By combining (5.22) and the fact that $f_p(x) = f_{1-p}(-x)$ we obtain

$$\pi_0(\alpha, p) = (1 - p)\mu_{1-p}(I_4) + \mu_p(I_3) + p\mu_p(I_4).$$

Computing the derivative with respect to p then gives

$$\partial_p \pi_0(\alpha, p) = -\mu_{1-p}(I_4) - (1-p)\partial_p(\mu_{1-p}(I_4)) + \partial_p(\mu_p(I_3)) + \mu_p(I_4) + \partial_p(\mu_p(I_4)).$$
(5.29)

From Lemma 5.4.8 it follows that $\partial_p(\mu_p(I_3)) = -\partial_p(\mu_{1-p}(I_3))$, implying that

$$\partial_p(\mu_p(I_3)) = 0$$
 at $p = \frac{1}{2}$.

Therefore, by (5.29) $\partial_p \pi_0(\alpha, p) = 0$ at $p = \frac{1}{2}$.

§5.5.2 On random CF-maps

Theorem 5.2.3 states that for random piecewise affine maps of the interval satisfying (c1), (c2) and (c3) strong random matching implies that there exists a piecewise constant invariant density. Condition (5.5) was sufficient for the theorem to work, which was one of the main motivations for Definition 5.2.2.

Theorem 5.2.3 is a random analogue of [BCMP18, Theorem 1.2], except that there the statement has less assumptions. The authors mention in [BCMP18, Remark 1.3] that for piecewise smooth interval maps with strong matching the corresponding invariant probability densities are piecewise smooth. On the other hand, as we noted before, the natural extension construction which for continued fraction transformations is often used to find invariant densities, seems to suggest that matching alone is sufficient to guarantee the existence of a piecewise smooth density. It would be interesting to investigate this further for the random continued fraction transformation.

In a first attempt to investigate to what extent Theorem 5.2.3 can be generalised to piecewise smooth random systems on an interval, we include some numerical simulations. Recall from Example 5.3.2 that the random continued fraction maps R_{α} have strong random matching for α in the intervals J_n with endpoints as in (5.8), see also Figure 5.8. Figure 5.13 shows two simulations of the invariant densities for such



Figure 5.13: Numerical simulations of the invariant probability densities of the random continued fraction maps R_{α} from Example 5.3.2. In (a) we take $\alpha \in J_4$ and $p_0 = 0.3$ and in (b) we have $\alpha \in J_5$ and $p_0 = 0.6$. The dashed lines indicate the positions of the prematching points, i.e., the points in the orbits of α and $\alpha - 1$ before the moment of matching.

systems R_{α} . The densities seem to be piecewise smooth with discontinuities precisely at the orbit points of α and $\alpha - 1$ before matching. This seems to support the claim that strong random matching is sufficient to guarantee the existence of a piecewise smooth invariant density.



Figure 5.14: Numerical simulations of the invariant probability densities of the random continued fraction maps R_{α} from Example 5.3.2 for three values of α between $\frac{1}{2}$ and $2 - \sqrt{2}$. The map in (a) has $\alpha \in (\frac{\sqrt{10}-2}{2}, 2 - \sqrt{2})$, which is the matching interval considered in Example 5.3.2. The orange graph is the graph of the weighted average of the densities of $T_{\alpha,0}$ and $T_{\alpha,1}$ with the appropriate values of p.

In Example 5.3.2 we also considered the maps R_{α} for $\alpha \in \left(\frac{\sqrt{10}-2}{2}, 2-\sqrt{2}\right)$. We showed that R_{α} has random matching with M = 3, but no strong matching at that moment. With a similar approach it can be shown that R_{α} has random matching for various other intervals in $\left[\frac{1}{2}, \frac{\sqrt{5}-1}{2}\right]$. For $\alpha \in \left[\frac{1}{2}, 2-\sqrt{2}\right]$ both deterministic maps $T_{\alpha,0}$ and $T_{\alpha,1}$ have strong matching with $M, Q \leq 2$, as was shown in [N81] and [TI81], and moreover, for both of them the invariant densities are known. In Figure 5.14 we have plotted the weighted average of these densities together with numerical simulations of the densities for various values of $\alpha \in \left[\frac{1}{2}, 2-\sqrt{2}\right]$ and 0 . This makes us wonder whether we need strong random matching to guarantee the existence of a piecewise smooth invariant density for these random systems or whether random matching is sufficient.

§5.6 Appendix

Figure 5.12 has been produced with the following Phyton program. Note that a computer is indeed able to handle the heavy algebraic procedure proposed in Chapter 3 and compute, from the fundamental matrix M, the density function $f_{\mathbf{p}}$ of the stationary measure $\mu_{\mathbf{p}}$.

```
import numpy as np
import array as arr
import sympy as sym
import random as rd
import scipy.optimize as opt
from random import choices
from fractions import Fraction
from decimal import Decimal
from sympy import pprint, Symbol, init printing
from numpy.linalg import matrix rank
from sympy import *
from pylab import *
from scipy import optimize
from matplotlib import pyplot
from sympy. utilities . lambdify import lambdify
sym.init printing()
aNumerator = 1
aDenominator = 3
print ("Rational a, st I a is a maximal quadratic interval
        for alpha-CF: ", Fraction(aNumerator, aDenominator))
\# compute the regular continued fraction expansion of odd length
def continuedFraction(n, d):
    if d = 0: return []
    q = n/d
    r = n - q * d
    return [q] + continuedFraction(d, r)
print ("Regular continued fraction expansion: ",
        continuedFraction(aNumerator, aDenominator))
m = np.sum(continuedFraction(aNumerator, aDenominator))
print ("Matching exponent: ", m)
vectorContinuedFraction = continuedFraction
        (aNumerator, aDenominator) [1:]
lengthVCF= len(vectorContinuedFraction)
if (lengthVCF \% 2) == 0:
    vectorContinuedFraction [lengthVCF - 1] =
        vectorContinuedFraction [lengthVCF-1]-1
    vectorContinuedFraction = np.append
```

```
Chapter 5
```

```
(vectorContinuedFraction,1)
print ("Regular continued fraction expansion with odd length: ",
        vectorContinuedFraction)
\# compute alpha via the valuation function
        of the associated binary vector
#
vectorBinary = np.ones((m,), dtype=int)
partialSum = np.zeros((lengthVCF,), dtype=int)
partialSum[0] = vectorContinuedFraction[0]
for i in range(1, lengthVCF):
    partialSum[i] = vectorContinuedFraction[i]+partialSum[i-1]
for i in range (0, \text{lengthVCF} - 1, 2):
    vectorBinary [partialSum [i]: partialSum [i+1]] = 0
print("Associated binary vector: ", vectorBinary)
alphaInverse = 0
for i in range(1, m+1):
    alphaInverse += vectorBinary[i-1]/2**i
alphaInverseFraction = Fraction(alphaInverse)
alphaVar = Fraction (alphaInverseFraction.denominator,
        alphaInverseFraction.numerator)
print("Alpha variable: ", alphaVar)
\#Matching interval
startInterval = Fraction(2**m+1,2**m*alphaInverseFraction+1)
endInterval = Fraction (2**m-1, 2**m*alphaInverseFraction -1)
print("Matching interval: ", startInterval, endInterval)
\#Random parameter alpha in the matching interval
        (startInterval, endInterval) of matching exponent m
alpha = rd.uniform(startInterval, endInterval)
print (Fraction (alpha), "alpha: ", alpha)
p = Symbol('p')
print("Probability vector: ", p, 1-p)
def t(x):
    if x > 0.5:
        return Fraction (2*x-alpha)
    elif x < -0.5:
        return Fraction (2*x+alpha)
    else:
        return Fraction (2*x)
\# compute the orbits of 1, 1-alpha and their opposites
orbitOne = np.ones((m+1,), dtype=object)
orbitOneWeighted = np.ones((m+1), dtype=object)
orbitOneWeighted [0] = 1-p
```

```
orbitOneNegative = np.ones((m+1,), dtype=object)
orbitOneAlpha = np.ones((m+1,), dtype=object)
orbitOneAlphaNegative = np.ones((m+1,), dtype=object)
for i in range (1,m+1):
    orbitOne[i] = t(orbitOne[i-1])
    if (alpha-1)/2 < orbitOne[i] < 0.5:
        orbitOneWeighted [i]=p
    else:
        orbitOneWeighted [i]=1-p
orbitOne = orbitOne [:m]
orbitOneWeighted = orbitOneWeighted [:m]
for i in range (0,m):
    orbitOneAlpha[i] = Fraction(orbitOne[i]-alpha)
    orbitOneNegative[i] = Fraction(-orbitOne[i])
    orbitOneAlphaNegative[i] = Fraction(-orbitOne[i]+alpha)
\# compute the partition given by points
        in the prematching set and the switch regions
partition = np.append(orbitOneAlphaNegative)
        np.append(orbitOneAlpha,
        np.append(orbitOne, orbitOneNegative)))
partitionSorted = np.sort(partition)
partitionSwitch = np.append(partition,
        [-Fraction(1,2), (alpha-1)/2, -(alpha-1)/2,
        Fraction(1,2)
partitionSortedSwitch = np.sort(partitionSwitch)
functionLOne = np.ones((m,), dtype=object)
functionLOneNegative = np.ones((m,), dtype=object)
for i in range (1,m):
    functionLOne[i] = sym.simplify
        (functionLOne [i-1]*Fraction (1,2)*orbitOneWeighted [i-1])
    functionLOneNegative[i] = sym.simplify
        (functionLOneNegative [i-1]*Fraction (1,2)*)
        (1 - \text{orbitOneWeighted} [i - 1]))
print("L (1)-L_(1-alpha): ", functionLOne)
print("L_(alpha-1)-L_(-1): ", functionLOneNegative)
\# compute the quantities KI n
kiOne = np.zeros((5,), dtype=object)
kiOneWeighted = np.ones((m+1)), dtype=object)
kiOneWeighted [1] = Fraction (1,2)
for i in range (2,m+1):
```

```
kiOneWeighted [i] = sym. simplify
        (kiOneWeighted [i-1]*Fraction (1,2)*orbitOneWeighted [i-2])
for i in range (1,m):
    if orbitOneWeighted [i-1]==p:
        kiOne[3] += kiOneWeighted[i]
        kiOne[0] += -kiOneWeighted[i]
    else:
        kiOne [4] += kiOneWeighted [i]
        kiOne[1] += -kiOneWeighted[i]
kiOne[4] += kiOneWeighted[m]
kiOne[0] += -kiOneWeighted[m]
print ("KIn(1)-KIn(1-alpha): ", sym.simplify(kiOne))
kiOneNegative=np.zeros((5,), dtype=object)
kiOneWeightedNegative=np.ones((m+1,), dtype=object)
kiOneWeightedNegative[1] = Fraction(1,2)
for i in range (2,m+1):
    kiOneWeightedNegative [i] = sym. simplify
        (kiOneWeightedNegative [i-1]*Fraction (1,2)*
        (1 - \text{orbitOneWeighted} [i - 2]))
for i in range (1,m):
    if orbitOneWeighted [i-1]==p:
        kiOneNegative[4] += kiOneWeightedNegative[i]
        kiOneNegative [1] += -kiOneWeightedNegative [i]
    else:
        kiOneNegative[3] += kiOneWeightedNegative[i]
        kiOneNegative [0] += -kiOneWeightedNegative [i]
kiOneNegative[4] += kiOneWeightedNegative[m]
kiOneNegative[0] += -kiOneWeightedNegative[m]
print ("KIn(alpha-1)-KIn(-1): ", sym.simplify(kiOneNegative))
\# compute the fundamental matrix
mFirstRow = np. zeros((3,), dtype=object)
mSecondRow = np.zeros((3,), dtype=object)
mThirdRow = np.zeros((3,), dtype=object)
mFirstRow[0] = -Fraction(1,2)+(1-p)/2*kiOneNegative[1]
mFirstRow [1] = Fraction(1,2)+p/2*kiOne[1]+(1-p)/2*
        kiOneNegative [1]
mFirstRow[2] = p/2*kiOne[1]
mSecondRow[0] = (1-p)/2 * kiOneNegative[3]
mSecondRow [1] = p/2 * kiOne[3] - Fraction(1,2) + (1-p)/2 *
        kiOneNegative [3]
mSecondRow[2] = Fraction(1,2)+p/2*kiOne[3]
```

```
mThirdRow [0] = (1-p)/2 * kiOneNegative [4]
mThirdRow[1] = p/2*kiOne[4]+(1-p)/2*kiOneNegative[4]
mThirdRow [2] = -Fraction(1,2) + p/2 * kiOne[4]
matrixM = np. array ([mFirstRow, mSecondRow, mThirdRow])
fundamentalMatrix = sym. Matrix (matrixM)
vectorGamma = fundamentalMatrix.nullspace()
print ("Solution vector Gamma: ", sym.factor(vectorGamma))
gamma = sym.Matrix(vectorGamma)
x = sym.factor(gamma[0])
y = sym.factor(gamma[1])
z = sym.factor(gamma[2])
densityCoefficient1 = sym.factor((x+y)*(1-p))
densityCoefficient2 = sym.factor((y+z)*p)
functionLOneNegativeWeighted = sym.factor
        (functionLOneNegative*densityCoefficient1)
functionLOneWeighted = sym.factor
        (functionLOne*densityCoefficient2)
\# compute the density
density = np.zeros((len(partitionSortedSwitch),), dtype=object)
for i in range (0,m):
    initial Position = np.where
        (partitionSortedSwitch=orbitOneAlpha[i])[0][0]+1
    finalPosition = np.where
        (partitionSortedSwitch=orbitOne[i])[0][0]
    for k in range (initialPosition, finalPosition+1):
        density [k] += functionLOneWeighted [i]
    initialPositionNegative = np.where
        (partitionSortedSwitch=orbitOneNegative[i])[0][0]+1
    finalPositionNegative = np.where
        (partitionSortedSwitch=orbitOneAlphaNegative[i])[0][0]
    for k in range
        (initialPositionNegative, finalPositionNegative+1):
        density [k] += functionLOneNegativeWeighted [i]
simplifiedDensity=sym.factor(density)
print ("Density function: ", simplifiedDensity)
normalizingConstant = 0
for i in range(1, len(partitionSortedSwitch)):
    normalizingConstant +=
        (partitionSortedSwitch[i]-partitionSortedSwitch[i-1])*
```

```
density[i]
simplifiedNormalizingConstant=sym.factor(1/normalizingConstant)
print ("Normalizing constant: ", simplifiedNormalizingConstant)
normalizedDensity = sym.factor
        (simplifiedNormalizingConstant*density)
print ("Normalized density: ", normalizedDensity)
\# compute the frequency of the zero digit
frequencyZero = 0
measureI4 = 0
measureI2 = 0
measureI3 = 0
startI4 = np.where(partitionSortedSwitch==(alpha - 1)/2)[0][0]+1
endI4 = np.where(partitionSortedSwitch = = 1/2)[0][0]
for k in range (startI4, endI4+1):
    measureI4 +=
        (partitionSortedSwitch[k]-partitionSortedSwitch[k-1])*
        normalizedDensity[k]
measureI4 = measureI4*p
startI2 = np.where(partitionSortedSwitch = -1/2)[0][0]+1
endI2 = np.where(partitionSortedSwitch==-(alpha - 1)/2)[0][0]
for k in range (startI2, endI2+1):
    measureI2 +=
        (partitionSortedSwitch[k]-partitionSortedSwitch[k-1])*
        normalizedDensity [k]
measureI2 = measureI2*(1-p)
startI3 = np.where(partitionSortedSwitch==-(alpha - 1)/2)[0][0]+1
endI3 = np.where(partitionSortedSwitch == (alpha - 1)/2)[0][0]
for k in range (startI3, endI3+1):
    measureI3 +=
        (partitionSortedSwitch[k]-partitionSortedSwitch[k-1])*
        normalizedDensity[k]
frequencyZero = measureI4 + measureI2 + measureI3
print ("Frequency of the 0 digit:", sym.factor(frequencyZero))
frequencyZeroFunction = lambdify((p,)),
        frequencyZero , modules='numpy')
p \text{ scipy} = np. \text{linspace}(0.0, 1.0)
```

```
y_sympy = frequencyZeroFunction(p_scipy)
pyplot.plot(p_scipy, y_sympy, 'b-', label='Freq(p)')
pyplot.xlabel(r'$p$')
pyplot.ylabel(r'$Frequency$')
pyplot.legend(loc='upper right')
pyplot.show()
```

Bibliography

- [A97] J. Aaronson. An introduction to infinite ergodic theory, volume 50 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1997.
- [AGH18] N. Abbasi, M. Gharaei, and A. J. Homburg. Iterated function systems of logistic maps: synchronization and intermittency. *Nonlinearity*, 31(8):3880– 3913, 2018.
- [ANV15] R. Aimino, M. Nicol, and S. Vaienti. Annealed and quenched limit theorems for random expanding dynamical systems. *Probab. Theory Related Fields*, 162(1-2):233–274, 2015.
- [A98] L. Arnold. *Random dynamical systems*. Springer Monographs in Mathematics. Springer-Verlag Berlin Heidelberg, 1998.
- [AS13] P. Arnoux and T. A. Schmidt. Cross sections for geodesic flows and αcontinued fractions. Nonlinearity, 26(3):711–726, 2013.
- [BG05] W. Bahsoun and P. Góra. Position dependent random maps in one and higher dimensions. *Studia Math.*, 166(3):271–286, 2005.
- [B93] M. F. Barnsley. *Fractals everywhere*. Academic Press, 1993.
- [BI09] L. Barreira and G. Iommi. Frequency of digits in the Lüroth expansion. J. Number Theory, 129(6):1479–1490, 2009.
- [BBDK94] J. Barrionuevo, R. M. Burton, K. Dajani, and C. Kraaikamp. Ergodic properties of generalized Lüroth series. *TU Delft Report*, 94-105:1–16, 1994.
- [B13] C. Bektur. Lyapunov exponents for position dependent random maps: formulae and applications. *Stoch. Anal. Appl.*, 31(4):600–608, 2013.
- [BCIT13] C. Bonanno, C. Carminati, S. Isola, and G. Tiozzo. Dynamics of continued fractions and kneading sequences of unimodal maps. *Discrete Contin. Dyn.* Syst., 33(4):1313–1332, 2013.
- [B51] A. D. Booth. A signed binary multiplication technique. The Quarterly Journal of Mechanics and Applied Mathematics, 4(2):236–240, 1951.
- [BSORG13] V. Botella-Soler, J. A. Oteo, J. Ros, and P. Glendinning. Lyapunov exponent and topological entropy plateaus in piecewise linear maps. J. Phys. A, 46(12):125101, 26, 2013.

- [BG92] A. Boyarsky and P. Góra. A dynamical system model for interference effects and the two-slit experiment of quantum physics. *Physics Letters A*, 168(2):103–112, 1992.
- [BG97] A. Boyarsky and P. Góra. Laws of chaos. Probability and its Applications. Birkhäuser Boston, Inc., Boston, MA, 1997. Invariant measures and dynamical systems in one dimension.
- [BL92] A. Boyarsky and Y. S. Lou. Existence of absolutely continuous invariant measures for higher-dimensional random maps. *Dynamics and Stability of Systems*, 7(4):233–244, 1992.
- [BCK17] H. Bruin, C. Carminati, and C. Kalle. Matching for generalised βtransformations. Indag. Math. (N.S.), 28(1):55–73, 2017.
- [BCMP18] H. Bruin, C. Carminati, S. Marmi, and A. Profeti. Matching in a family of piecewise affine maps. *Nonlinearity*, 32(1):172–208, 2018.
- [BK93] R. Burton and G. Keller. Stationary measures for randomly chosen maps. Journal of Theoretical Probability, 6:1–16, 1993.
- [B00] J. Buzzi. Absolutely continuous s.r.b. measures for random lasota-yorke maps. Transactions of the American Mathematical Society, 352(7):3289– 3303, 2000.
- [CWZ13] C. Cao, J. Wu, and Z. Zhang. The efficiency of approximating real numbers by Lüroth expansion. *Czechoslovak Math. J.*, 63(138)(2):497–513, 2013.
- [CIT18] C. Carminati, S. Isola, and G. Tiozzo. Continued fractions with SL(2, Z)-branches: combinatorics and entropy. Trans. Amer. Math. Soc., 370(7):4927–4973, 2018.
- [CMPT10] C. Carminati, S. Marmi, A. Profeti, and G. Tiozzo. The entropy of α -continued fractions: numerical results. *Nonlinearity*, 23(10):2429–2456, 2010.
- [CT12] C. Carminati and G. Tiozzo. A canonical thickening of \mathbb{Q} and the entropy of α -continued fraction transformations. *Ergodic Theory Dynam. Systems*, 32(4):1249–1269, 2012.
- [CT13] C. Carminati and G. Tiozzo. Tuning and plateaux for the entropy of α continued fractions. *Nonlinearity*, 26(4):1049–1070, 2013.
- [CWY14] H. Chen, W. Wang, and M. Yu. Hausdorff dimension of the graph of the error-sum function of α-Lüroth series. Turkish J. Math., 38(5):803–811, 2014.
- [CW14] H. Chen and Z. Wen. The Hausdorff dimension of certain sets in a class of α -Lüroth expansions. *Sci. China Math.*, 57(2):303–313, 2014.

Chapter 5

- [CMO98] H. Cohen, A. Miyaji, and T. Ono. Efficient elliptic curve exponentiation using mixed coordinates. pages 51–65. Springer Berlin Heidelberg, 1998.
- [CM18] D. Cosper and M. Misiurewicz. Entropy locking. Fund. Math., 241(1):83– 96, 2018.
- [DHK09] K. Dajani, Y. Hartono, and C. Kraaikamp. Mixing properties of (α, β) -expansions. Ergodic Theory Dynam. Systems, 29(4):1119–1140, 2009.
- [DHKM12] K. Dajani, D. Hensley, C. Kraaikamp, and V. Masarotto. Arithmetic and ergodic properties of 'flipped' continued fraction algorithms. Acta Arith., 153(1):51–79, 2012.
- [DK07] K. Dajani and C. Kalle. Random β-expansions with deleted digits. Discrete Contin. Dyn. Syst., 18(1):199–217, 2007.
- [DK10] K. Dajani and C. Kalle. A note on the greedy β -transformation with arbitrary digits. In *École de Théorie Ergodique*, volume 20 of *Sémin. Congr.*, pages 83–104. Soc. Math. France, Paris, 2010.
- [DK13] K. Dajani and C. Kalle. Local dimensions for the random β -transformation. New York J. Math., 19:285–303, 2013.
- [DK17] K. Dajani and C. Kalle. Invariant measures, matching and the frequency of 0 for signed binary expansions. arXiv: 1703.06335, 2017.
- [DKM20] K. Dajani, C. Kalle, and M. Maggioni. Matching for random systems with an application to minimal weight expansions. 2020. Submitted to *Nonlinearity*.
- [DK96] K. Dajani and C. Kraaikamp. On approximation by Lüroth series. J. Théor. Nombres Bordeaux, 8(2):331–346, 1996.
- [DK00] K. Dajani and C. Kraaikamp. The mother of all continued fractions. Colloq. Math., 84/85(part 1):109–123, 2000. Dedicated to the memory of Anzelm Iwanik.
- [DK02] K. Dajani and C. Kraaikamp. Ergodic Theory of Numbers. Number v. 29 in Carus Mathematical Monographs. Mathematical Association of America, 2002.
- [DK03] K. Dajani and C. Kraaikamp. Random β-expansions. Ergodic Theory Dynam. Systems, 23(2):461–479, 2003.
- [DKS09] K. Dajani, C. Kraaikamp, and W. Steiner. Metrical theory for α-Rosen fractions. J. Eur. Math. Soc. (JEMS), 11(6):1259–1283, 2009.
- [DKL06] K. Dajani, C. Kraaikamp, and P. Liardet. Ergodic properties of signed binary expansions. Discrete Contin. Dyn. Syst., 15(1):87–119, 2006.

- [DO18] K. Dajani and M. Oomen. Random N-continued fraction expansions. J. Approx. Theory, 227:1–26, 2018.
- [DdV05] K. Dajani and M. de Vries. Measures of maximal entropy for random β -expansions. J. Eur. Math. Soc., 7(1):51–68, 2005.
- [DdV07] K. Dajani and M. de Vries. Invariant densities for random β-expansions. J. Eur. Math. Soc., 9(1):157–176, 2007.
- [EJK90] P. Erdös, I. Joó, and V. Komornik. Characterization of the unique expansions $1 = \sum_{i=1}^{\infty} q^{-n_i}$ and related problems. *Bull. Soc. Math. France*, 118(3):377–390, 1990.
- [EM12] P. Eslami and M. Misiurewicz. Singular limits of absolutely continuous invariant measures for families of transitive maps. J. Difference Equ. Appl., 18(4):739–750, 2012.
- [FLMW10] A. Fan, L. Liao, J. Ma, and B. Wang. Dimension of Besicovitch-Eggleston sets in countable symbolic space. *Nonlinearity*, 23(5):1185–1197, 2010.
- [FB81] N. Friedman and A. Boyarsky. Matrices and eigenfunctions induced by Markov maps. *Linear Algebra Appl.*, 38:141–147, 1981.
- [G59] A. O. Gel'fond. A common property of number systems. Izv. Akad. Nauk SSSR. Ser. Mat., 23:809–814, 1959.
- [G16] R. Giuliano. Convergence results for *r*-iterated means of the denominators of the Lüroth series. *Unif. Distrib. Theory*, 11(2):179–203, 2016.
- [GB03] P. Góra and A. Boyarsky. Absolutely continuous invariant measures for random maps with position dependent probabilities. J. Math. Anal. Appl., 278(1):225–242, 2003.
- [GBL97] P. Góra, A. Boyarsky, and Y. S. Lou. Lyapunov exponents for higherdimensional random maps. J. Appl. Math. Stochastic Anal., 10(3):209–218, 1997.
- [G84] P. Góra. On small stochastic perturbations of mappings of the unit interval. Colloquium Mathematicae, 49(1):73–85, 1984.
- [G09] P. Góra. Invariant densities for piecewise linear maps of the unit interval. Ergodic Theory Dynam. Systems, 29(5):1549–1583, 2009.
- [GBI06] P. Góra, A. Boyarsky, and S. Islam. Invariant densities of random maps have lower bounds on their supports. J. Appl. Math. Stoch. Anal., pages Art. ID 79175, 13, 2006.
- [GH06] P. J. Grabner and C. Heuberger. On the number of optimal base 2 representations of integers. *Des. Codes Cryptogr.*, 40(1):25–39, 2006.

- [GL16] Y. Gui and W. Li. Hausdorff dimensions of sets related to Lüroth expansion. Acta Math. Hungar., 150(2):286–302, 2016.
- [H02] A. Haas. Invariant measures and natural extensions. Canad. Math. Bull., 45(1):97–108, 2002.
- [HK02] Y. Hartono and C. Kraaikamp. On continued fractions with odd partial quotients. *Rev. Roumaine Math. Pures Appl.*, 47(1):43–62 (2003), 2002.
- [HP06] C. Heuberger and H. Prodinger. Analysis of alternative digit sets for nonadjacent representations. *Monatsh. Math.*, 147:219–248, 2006.
- [I12] T. Inoue. Invariant measures for position dependent random maps with continuous random parameters. *Studia Math.*, 208(1):11–29, 2012.
- [I20] T. Inoue. First return maps of random maps and invariant measures. Nonlinearity, 33(1):249–275, 2020.
- [K14] C. Kalle. Isomorphisms between positive and negative β -transformations. Ergodic Theory Dynam. Systems, 34(1):153–170, 2014.
- [KKV17] C. Kalle, T. Kempton, and E. Verbitskiy. The random continued fraction transformation. *Nonlinearity*, 30(3):1182–1203, 2017.
- [KLMM20] C. Kalle, N. Langeveld, M. Maggioni, and S. Munday. Matching for a family of infinite measure continued fraction transformations. 2020.
- [KM18] C. Kalle and M. Maggioni. Invariant densities for random systems of the interval. 2018. To appear in *Ergodic Theory and Dynamical Systems*.
- [KM21] C. Kalle and M. Maggioni. On approximation by random Lüroth expansions. 2021. Submitted to *International Journal of Number Theory*.
- [KS12] C. Kalle and W. Steiner. β -expansions, natural extensions and multiple tilings associated with Pisot units. *Trans. Amer. Math. Soc.*, 364(5):2281–2318, 2012.
- [KKK90] S. Kalpazidou, A. Knopfmacher, and J. Knopfmacher. Lüroth-type alternating series representations for real numbers. Acta Arith., 55(4):311–322, 1990.
- [KKK91] S. Kalpazidou, A. Knopfmacher, and J. Knopfmacher. Metric properties of alternating Lüroth series. *Portugal. Math.*, 48(3):319–325, 1991.
- [KU10] S. Katok and I. Ugarcovici. Structure of attractors for (a, b)-continued fraction transformations. J. Mod. Dyn., 4(4):637–691, 2010.
- [K82] G. Keller. Stochastic stability in some chaotic dynamical systems. Monatsh. Math., 94(4):313–333, 1982.
- [K14] K. Kempton. On the invariant density of the random β -transformation. Acta Math. Hungar., 142(2):403–419, 2014.

- [KMS11] M. Kesseböhmer, S. Munday, and B. Stratmann. Strong renewal theorems and lyapunov spectra for α-farey and α-lüroth systems. Ergodic Theory and Dynamical Systems, 32(3):989–1017, 2011.
- [KMS16] M. Kesseböhmer, S. Munday, and B. Stratmann. Infinite ergodic theory of numbers. De Gruyter Graduate. De Gruyter, Berlin, 2016.
- [K86] Y. Kifer. Ergodic Theory of Random Transformations, volume 10 of Progress in Probability. Birkhäuser Basel, 1986.
- [K69] D. E. Knuth. The Art of Computer Programming. Addison-Wesley Professional, 1st edition, 1969.
- [KY07] E. Kobre and L. S. Young. Extended systems with deterministic local dynamics and random jumps. Communications in Mathematical Physics, 275(3):709–720, 11 2007.
- [K90] C. Kopf. Invariant measures for piecewise linear transformations of the interval. Appl. Math. Comput., 39(2, part II):123–144, 1990.
- [KT93] K. Koyama and Y. Tsuruoka. Speeding up elliptic cryptosystems by using a signed binary window method. pages 345–357. Springer Berlin Heidelberg, 1993.
- [K91] C. Kraaikamp. A new class of continued fraction expansions. Acta Arith., 57(1):1–39, 1991.
- [KSS10] C. Kraaikamp, T. Schmidt, and I. Smeets. Natural extensions for α-rosen continued fractions. J. Math. Soc. Japan, 62(2):649–671, 2010.
- [KSS12] C. Kraaikamp, T. Schmidt, and W. Steiner. Natural extensions and entropy of α-continued fractions. *Nonlinearity*, 25(8):2207–2243, 2012.
- [K67] U. Krengel. Entropy of conservative transformations. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete, 7:161–181, 1967.
- [LK97] C. Laih and W. Kuo. Speeding up the computations of elliptic curves cryptoschemes. Computers and Mathematics with Applications, 33(5):29– 36, 1997.
- [LM94] A. Lasota and M. C. Mackey. Chaos, fractals, and noise, volume 97 of Applied Mathematical Sciences. Springer-Verlag, New York, second edition, 1994. Stochastic aspects of dynamics.
- [LY73] A. Lasota and J.A. Yorke. On the existence of invariant measures for piecewise monotonic transformations. *Trans. Amer. Math. Soc.*, 186:481– 488 (1974), 1973.
- [L81] L. Lewin. *Polylogarithms and associated functions*. North-Holland Publishing Co., New York-Amsterdam, 1981.

- [LCTW18] L. Li, F. Cao, S. Tang, and Y. Wu. Lüroth expansion digits and Maclaurin's inequality. Wuhan Univ. J. Nat. Sci., 23(6):471–474, 2018.
- [LGB⁺13] Z. Li, P. Góra, A. Boyarsky, H. Proppe, and P. Eslami. Family of piecewise expanding maps having singular measure as a limit of ACIMs. *Ergodic Theory Dynam. Systems*, 33(1):158–167, 2013.
- [LY78] T. Y. Li and J. A. Yorke. Ergodic transformations from an interval into itself. Trans. Amer. Math. Soc., 235:183–192, 1978.
- [LM95] D. Lind and B. Marcus. An Introduction to Symbolic Dynamics and Coding. Cambridge University Press, 1995.
- [L83] J. Lüroth. Ueber eine eindeutige Entwickelung von Zahlen in eine unendliche Reihe. Math. Ann., 21(3):411–423, 1883.
- [LM08] L. Luzzi and S. Marmi. On the entropy of Japanese continued fractions. Discrete Contin. Dyn. Syst., 20(3):673–711, 2008.
- [MT13] B. Mance and J. Tseng. Bounded Lüroth expansions: applying Schmidt games where infinite distortion exists. *Acta Arith.*, 158(1):33–47, 2013.
- [MMY97] S. Marmi, P. Moussa, and J. C. Yoccoz. The Brjuno functions and their regularity properties. Comm. Math. Phys., 186(2):265–293, 1997.
- [MO90] F. Morain and J. Olivos. Speeding up the computations on an elliptic curve using addition-subtraction chains. RAIRO - Theoretical Informatics and Applications - Informatique Théorique et Applications, 24(6):531–543, 1990.
- [M85] T. Morita. Random iteration of one-dimensional transformations. Osaka J. Math., 22(3):489–518, 1985.
- [M11] S. Munday. A note on Diophantine fractals for α -Lüroth systems. *Integers*, 11B:Paper No. A10, 14, 2011.
- [NIT77] H. Nakada, S. Ito, and S. Tanaka. On the invariant measure for the transformations associated with some real continued-fractions. *Keio Engineering Reports*, 30(13):159–175, 1977.
- [N81] H. Nakada. Metrical theory for a class of continued fraction transformations and their natural extensions. *Tokyo J. Math.*, 4(2):399–426, 1981.
- [NN08] H. Nakada and R. Natsui. The non-monotonicity of the entropy of α continued fraction transformations. *Nonlinearity*, 21(6):1207–1225, 2008.
- [N11] R. Natsui. On the Legendre constant of α -continued fractions. J. Number Theory, 131(3):487–507, 2011.
- [P60] W. Parry. On the β-expansions of real numbers. Acta Math. Acad. Sci. Hungar., 11:401–416, 1960.

- [P84] S. Pelikan. Invariant densities for random maps of the interval. Trans. Amer. Math. Soc., 281(2):813–825, 1984.
- [P57] O. Perron. Die Lehre von den Kettenbrüchen. Dritte, verbesserte und erweiterte Aufl. Bd. II. Analytisch-funktionentheoretische Kettenbrüche. B. G. Teubner Verlagsgesellschaft, Stuttgart, 1957.
- [R60] G. W. Reitwiesner. Binary arithmetic. volume 1 of Advances in Computers, pages 231–308. Elsevier, 1960.
- [R57] A. Rényi. Representations for real numbers and their ergodic properties. Acta Math. Acad. Sci. Hungar, 8:477–493, 1957.
- [R61] V. A. Rohlin. Exact endomorphisms of a Lebesgue space. Izv. Akad. Nauk SSSR Ser. Mat., 25:499–530, 1961.
- [S68] T. Salát. Zur metrischen Theorie der Lürothschen Entwicklungen der reellen Zahlen. Czechoslovak Math. J., 18 (93):489–522, 1968.
- [S04] B. Schratzberger. S-expansions in dimension two. J. Théor. Nombres Bordeaux, 16(3):705–732, 2004.
- [SF11] L. Shen and K. Fang. The fractional dimensional theory in Lüroth expansion. Czechoslovak Math. J., 61(136)(3):795–807, 2011.
- [S17] L. Shen. Hausdorff dimension of the set concerning with Borel-Bernstein theory in Lüroth expansions. J. Korean Math. Soc., 54(4):1301–1316, 2017.
- [SYZ14] L. Shen, Y. Yu, and Y. Zhou. A note on the largest digits in Lüroth expansion. Int. J. Number Theory, 10(4):1015–1023, 2014.
- [S03] N. Sidorov. Almost every number has a continuum of β -expansions. Amer. Math. Monthly, 110(9):838–842, 2003.
- [S88] C. E. Silva. On μ-recurrent nonsingular endomorphisms. Israel J. Math., 61(1):1–13, 1988.
- [ST91] C. E. Silva and P. Thieullen. The subadditive ergodic theorem and recurrence properties of Markovian transformations. J. Math. Anal. Appl., 154(1):83–99, 1991.
- [SFM17] K. Song, L. Fang, and J. Ma. Level sets of partial maximal digits for Lüroth expansion. Int. J. Number Theory, 13(10):2777–2790, 2017.
- [SX18] Y. Sun and J. Xu. On the maximal run-length function in the Lüroth expansion. *Czechoslovak Math. J.*, 68(143)(1):277–291, 2018.
- [S19] S. Suzuki. Invariant density functions of random β -transformations. Ergodic Theory Dynam. Systems, 39(4):1099–1120, 2019.
- [TI81] S. Tanaka and S. Ito. On a family of continued-fraction transformations and their ergodic properties. *Tokyo J. Math.*, 4(1):153–175, 1981.

- [TW18] X. Tan and X. Wang. On the fast growth rate of the sum of digits of Lüroth expansion. *Math. Appl. (Wuhan)*, 31(2):300–304, 2018.
- [T83] M. Thaler. Transformations on [0, 1] with infinite invariant measures. Israel J. Math., 46(1-2):67–96, 1983.
- [T14] G. Tiozzo. The entropy of Nakada's α-continued fractions: analytical results. Annali della Scuola Normale Superiore di Pisa, 13(4):1009–1037, 2014.
- [V14] J. Vandehey. A simpler normal number construction for simple Lüroth series. J. Integer Seq., 17(6):Article 14.6.1, 18, 2014.
- [ZC16] Z. Zhang and C. Cao. On points containing arithmetic progressions in their Lüroth expansion. Acta Math. Sci. Ser. B (Engl. Ed.), 36(1):257–264, 2016.
- [Z98] R. Zweimüller. Ergodic structure and invariant densities of non-Markovian interval maps with indifferent fixed points. *Nonlinearity*, 11(5):1263–1276, 1998.
- [Z00] R. Zweimüller. Ergodic properties of infinite measure-preserving interval maps with indifferent fixed points. *Ergodic Theory Dynam. Systems*, 20(5):1519–1549, 2000.
- [Z09] R. Zweimüller. Surrey notes on infinite ergodic theory. 2009.