



Universiteit
Leiden
The Netherlands

Integrative taxonomy of araneomorph spiders: Breathing new life into an old science

Rivera Quiroz, F.A.

Citation

Rivera Quiroz, F. A. (2021, April 14). *Integrative taxonomy of araneomorph spiders: Breathing new life into an old science*. Retrieved from <https://hdl.handle.net/1887/3152423>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3152423>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/3152423> holds various files of this Leiden University dissertation.

Author: Rivera Quiroz, F.A.

Title: Integrative taxonomy of araneomorph spiders: Breathing new life into an old science

Issue date: 2021-04-14

Chapter 1

General Introduction

F. ANDRES RIVERA-QUIROZ^{1, 2}

¹ Understanding Evolution group, Naturalis Biodiversity Center, Darwinweg 2, 2333CR Leiden, The Netherlands

² Institute for Biology Leiden (IBL), Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands.

Taxonomy: what it is, what are its problems and the proposed solutions

Taxonomy is the science and practice of the classification and description of things; in biology, it encompasses the description, identification, nomenclature, and classification of organisms based on their inferred relationships [1–3]; this is the general notion of what taxonomy is and what a taxonomist does. However, in the context of biology, taxonomy has a much broader background that involves all sort of biological knowledge, from the formal naming of groups of organisms (e.g. species) to the temporal and spatial distributions of those groups, their morphology, anatomy, behavior, molecular information, among many other data. The science of taxonomy is currently in a state of transition, where content within publications is being digitally mobilized and incorporated into an interconnected knowledge network.

Throughout its more than two centuries of existence, taxonomy has provided a nomenclatural and relational framework to all the biological disciplines setting the foundations for discovering and understanding biodiversity. Nevertheless, this foundational role can often be confounded with a trivial and expendable one, or even be considered as a mere “service provider” [4, 5] to other fields of science. This perception of simplicity in the taxonomic endeavor, together with the realization of the massive number of undescribed—and unknown—taxa [6–12] and the human impact imposed on ecosystems lead to a crisis commonly known as the *taxonomic impediment* [6, 8, 13–15].

This taxonomic impediment (or impediments?) has become symptomatic in many ways, namely: the reduction of the taxonomic workforce and university curricula; the reduction of funding and economic support; and the view of taxonomy as an obsolete science [4, 9, 14–22]. An unintended consequence has also been the reduction in the literature outlets, leaving fewer venues for the publication of fundamental science. This problem has been highlighted by the recent decision to exclude the taxonomic megajournal *Zootaxa* from widely used science evaluation metrics (although this decision was subsequently reversed in the wake of reactions from the taxonomic community) [23–25].

This taxonomic impediment was noted by governments during the late 1990’s and some agencies like the American National Science Foundation (NSF) created programs to enhance taxonomic training and research for future generations. That was the case of the Partnership for Enhancing Expertise in Taxonomy (PEET) [14, 26–28]. Similar projects were created in other countries like a federal directive for training specialists in cladistics in Brazil [19] and the Darwin Initiative in the UK. These programs looked to revive taxonomy, increase the workforce (especially in little known and neglected taxa), and stimulate the production and sharing of information on the internet, making taxonomic knowledge more accessible. A survey of the PEET impact in the creation of



Figure 1.1. “*Linnaeus in the information age*”. a) The tenth edition of Carl Linnaeus’ *Systema Naturae* published in 1758. Modified from Pyle [43]. b) Depicts an anachronistic portrait of Linnaeus—considered the father of Taxonomy—as a *techie*, using modern technology to develop his science. Modified from Godfray [5].

taxonomic careers showed that although PEET trainees greatly impacted scientific production in their respective fields, there were common concerns about funding continuity and the opportunity of finding steady jobs in taxonomy [28]. This same work emphasized the need for multidisciplinary training that, in addition to traditional taxonomic skills, also developed competence in the use of molecular techniques, data management, information dissemination, etc., as well as capabilities for integrating taxonomy with biogeography, ecology, ethology, etc.

It is clear then that taxonomy needs to change to improve taxonomy-based products and keep up with modern, faster data-driven science. Essential elements for 21st century taxonomy include universal access to primary source information [29–31] and the incorporation of new technologies to facilitate better description, analysis, understanding and conservation of biodiversity (Fig. 1.1). In this context, Godfray and Knapp [17] summarized the major changes that taxonomy has gone through in recent decades, transforming from a purely descriptive discipline to a hypothesis-driven science. This transition has been driven by a “philosophical makeover” that incorporated phylogenetic systematic ideas to test relationships between taxa, along with advances in DNA sequencing that have made vast amounts of molecular data available. Similarly, Godfray [5, 30] and Wheeler et al. [15] have articulated the need to build a more robust online taxonomic infrastructure to overcome some common problems in taxonomy, such as

the publication of isolated descriptions, and limitations in data and image sharing and publication. The first decade of this century saw a healthy and necessary debate concerning the significance of taxonomy, its role among the modern biological science, and the epistemological, philosophical and technical reforms needed for taxonomy to stay relevant in the future [5, 8, 9, 15, 17, 18, 20, 21, 29, 30, 32–42]

“Taxonomy as a team sport”

Taxonomy is often understood as the individual efforts of taxonomists working in isolation; however, the current state of taxonomic science and the pace of the loss of biodiversity demand taxonomy become a group effort that greatly takes advantage of the internet and its capabilities to accelerate the rate and quality of its production and distribution [35]. The taxonomy of a group of organisms typically arises from the growing collection of publications in the primary taxonomic literature of that taxon, and does not reside in a single publication or institution. As such, it tends to be scattered among multiple journals, with contributions by various authors; this has made taxonomic literature a (usually) poorly defined integral of the accumulation of literature [30] with a tendency to be unorganized and fragmentary. Godfray [30] suggested the implementation of a unitary model of taxonomy where one group or institution would be in charge of the administration of current taxonomy for a given group. Spider taxonomy, together with a few other examples like ants [44, 45], lepidopterans [46], dipterans [47], orthopterans [48] and fishes [49], are probably the finest working examples of this authoritative “unitary taxonomy” [30, 31, 50] model.

The taxonomy of spiders is probably one of the best curated and polished, being under constant revision since the mid-20th century, and currently being administered by the World Spider Catalog [51]. The indexation and scrutiny of spider taxonomic literature began in the 1940’s by Pierre Bonnet (University of Toulouse, France) and Carl Friedrich Roewer (Bremen, Germany). Afterwards, the Italian arachnologist Paolo M. Brignoli (University of Aquila, Italy) started filling the gaps in the aforementioned works and publishing periodic catalog supplements. This idea was subsequently picked up by the American arachnologist Norman I. Platnick (American Museum of Natural History, New York) who published three catalog supplements between 1989 and 1998 [52–54]. After this, the online version of the World Spider Catalog (WSC) was established and handled by Platnick from 2000 to 2014 [55]. In 2014, the WSC was transformed into a relational database administered by the Natural History Museum Bern (Switzerland) [51]. This catalog provides a fully searchable database that grants access to information on the more than 48,700 currently valid spider species and their synonyms. The WSC website also keeps a fully accessible library with more than 15,400 documents on spider taxonomy that dates back from the current year to Carl Clerck’s species descriptions from 1757, the only accepted spider descriptions previous to the Linnean era (ICZN: Article 3.1) [56, 57].

Improving access, dissemination and usability of information

Economics is the study of the production, distribution, and consumption of goods and services. If the scientific community is working to modernize the production, distribution, and consumption of taxonomic information, then perhaps we should look to economics for guidance. But economics is dominated by the study of physical goods, whose value is largely driven by their scarcity: rare things tend to be more valuable than common things. In contrast to physical goods information becomes more valuable as it becomes more accessible. So to increase the value of taxonomy, we should work to increase accessibility to biodiversity knowledge [58]. Current technologies allow storing and sharing data at an unprecedented pace, which in turn, allows the transformation of old printed literature (e. g. the Biodiversity Heritage Library [59]), and direct publication of new literature in electronic PDF format. This permits world-wide access to an amount of knowledge that was previously reserved to a few great libraries [15], recovering centuries of aggregated data that might otherwise just be forsaken on a library book shelf or at the back of a researcher's drawer. Nevertheless, it is not the best format for exchanging and querying data generating what has been dubbed the “PDF Black Box” [60]. In this context, the Swiss organization Plazi [61] has created software that semiautomatically detects and extracts taxonomic treatments, pieces of text that link specimen data (e.g. collection locality and date, collector, number of specimens, developmental stage, among others) to a taxon name [62–64].

The Plazi initiative [65–67] has greatly benefited from electronic access to taxonomic literature in PDF format and has focused on mining and mobilizing biodiversity and taxonomic data contained in legacy and newly produced taxonomic literature. This can be done in two ways: a retrospective approach that transforms a taxonomic document in PDF format into a semantically categorized document in XML format that allows specimen data to be extracted; and a prospective approach where data is directly produced in XML format as has been implemented by some journals (e. g. Pensoft's ZooKeys and Biodiversity Data Journal, the European Journal of Taxonomy, among others) [58] (Fig. 1.2). These taxonomic treatments are then stored and available in Plazi's Treatment Bank where the specimen data can be directly gathered and analyzed [68]. Also, a main characteristic of this repository is that these specimen records are also contributed to the Global Biodiversity Information Facility [69], where they are aggregated with other sources of data such as collection specimen records and observations networks (among other sources), forming an unparalleled resource for discovering, analyzing and explaining broad biodiversity patterns [63–65, 68].

In this thesis, we explored the application of these principals in **Chapter 2**, focusing on the extraction of data from legacy taxonomic literature and its application for inferring




Figure 1.2.-The “Plazi approach”. Schematic view of the retrospective (green) and prospective (red) data mining from taxonomic literature. Extracted data can then be displayed and analyzed in specialized repositories [61, 70] and databases [61, 69, 71].

phenological patterns that allow optimizing specimen sampling in the field (see also the fieldwork section below). Also, we used taxonomic literature obtained from the WSC to assess the rare phenomenon of genital asymmetry (GA) in spiders (**Chapter 3**, see the *Teutamus politus* section below). This chapter used taxonomic literature as an image repository observing and comparing the incidence and biological background of GA, giving a hypothetical classification based on previous work in spiders and insects GA [72–80] and drawing a preliminary evolutionary scenario for each type of asymmetry we observed [81].

The integrative taxonomy

Integrative Taxonomy was defined by Dayrat [4] as the complementary delimitation of species based on their phylogeography, morphology, genetics, ecology, behavior, etc. Although the term, as minted by Dayrat [4] failed to be the proposed “new paradigm” in the exercise and understanding of the taxonomic science and was even dubbed as potentially misleading and detrimental by Valdecasas et al. [82] due to its *guidelines* on the taxonomic nomenclature, there are some concepts that are worth recovering




Figure 1.3.- The integrative taxonomic pipeline. Schematic workflow of the taxonomic process used in this thesis. It includes the use of the WSC [51] as a taxonomic library; the extraction of data from taxonomic literature [65, 66]; the use of these and other specimen data for biodiversity analyses; the use of molecular data (some of it from Genbank [71]) for building the phylogenies and a thorough documentation with different imaging techniques of the specimens' morphology.

and incorporating, whenever possible, in the common practice of taxonomy. Taxonomy has generally entrusted this species delimitation on morphology due to both practical and historical reasons. Nevertheless, as mentioned before, the current taxonomic crisis requires the inclusion of new technologies and techniques that favor faster and more reliable taxa descriptions; integration means multidisciplinary [4, 83].

This multidisciplinary has become more and more common in some fields by the integration of: morphological data (e.g. photographs, SEM, 3D imaging, CT-scans, geometric morphometrics, among others), molecular data, geographical information (e.g. collection localities and the use of GIS), and other specimen based data (e.g. number and sex of specimens, collection dates, etc.) that allow testing the species hypotheses in different and more robust approaches [4, 82, 84–86]. Pyle [43] visually exemplifies the enormous advances on species documentation by using these modern technologies. The integrative perspective, besides generating pure taxonomic information—hypotheses of classification and relations—, also contributes with molecular data, high resolution images of specimens and their relevant characters, and other kinds of data to global databases [86–92]. This accessible data can be downloaded and tested independently or incorporated to new taxonomic and systematic studies in an easy and flexible way; this can help to expedite the testing of phylogenetic hypotheses and rapidly generate new datasets and inferences with more explanatory power.

In this context, **Chapters 4 and 5** are examples of the implementation of this integrative taxonomy (Fig. 1.3). Both chapters involve the description of new taxa and also, first reports of these families for Thailand. **Chapter 4** describes two new species and one new genus of the family Hahniidae based on molecular and morphological evidence; **Chapter 5** describes three new species of the family Symphytognathidae, employing molecular and morphological evidence; and making use of micro CT-scans and 3D modeling to circumscribe the genus *Crassignatha* and clearly illustrate and discuss the genital characters of the males of this and other related symphytognathid genera. Although 3D modeling had been used before to study spider genitalia [93–95], ours are, to the best of our knowledge, the smallest spider palps that have ever been 3D modeled, being only 0.2 mm in width.

Fieldwork

The process of how we selected the locations where fieldwork was conducted is explained in detail in **Chapter 2**. Here we cover some generalities of the new specimen collection, where and how it was conducted. Southeast Asia (SEA) is one of the most diverse areas on the planet being represented by several conservation hotspots, of which the Indo-Burma and the Sundaland Hotspots are the most extensive and diverse [96, 97] (Fig. 1.4). The Indo-Burma hotspot is largely distributed over continental SEA; this region is characterized by socio-political instability, high population density and a fragmentary nature of its territory. This, in turn, generates a scattered taxonomy of most groups that makes it difficult to describe and assess its biodiversity [96, 98, 99]. Still, studies on vascular plants and terrestrial vertebrates show the Indo-Burma hotspot to be one of the richest and highest in endemism, but also one of the most threatened hotspots [96, 100].

This uniqueness and magnitude of its biodiversity is also reflected, albeit understudied, in invertebrates. The arachnofauna is mostly known from isolated species descriptions, although a few genus, family and broader revisions have been published [101–103]. Especially interesting to us was the case of the liocranid spiders, and from this family, the species *Teutamus politus* Thorell, 1890 (see *T. politus* section below, and **Chapters 2 and 3**) and its close relatives in the so called *Teutamus* group (TG). While planning our fieldwork, we wanted to improve the probabilities of capturing adult specimens of *T. politus* while also sampling other relatives of the TG. Due to the lack of data on these spiders' phenology, we used data from taxonomic literature to select the localities and times with a higher incidence of liocranid spiders based on the number of adults reported in taxonomic literature. Taking into account this information we decided to collect from July to August 2018 in the three provinces with most specimens reported in Thailand. We covered a variety of habitats in each place (Fig. 1.5) ranging from cloud forests at the top of the highest mountain in the country, to temperate pine, oak and *Dipterocarpus* forests down to tropical vegetations. We also sampled in secondary forests and rubber and oil palm plantation to observe the anthropogenic impact on the populations of these spiders.





Figure 1.4.-Map of the Southeast Asian Hotspots. Modified from Mittermeier et al. [96].

Figure 1.5.- Some of the localities sampled in Thailand. a, b) Chiang Mai: a- bamboo forest in the Pha Daeng National Park. b- Cloud forest in Doi Inthanon National Park, the highest mountain in the country. c, d) Phuket and Krabi: c- a patch of *Kerrodoxia elegans* palm tree in Ton Sai. d. Secondary forest near Than Bok Khorani National Park. (Next page, top) ►

Figure 1.6.- Collecting methods. a) Winkler traps hung outside Doi Suthep National Park headquarters. b, c) Pitfall traps at a *Dipterocarpus* sp. forest near Pha Daeng National Park. d) Active search in sifted leaf litter at Doi Suthep National Park. e) Active search in cryptic habitats (i. e. under rocks, logs, bark, among leaf litter, etc) near Than Bok Khorani National Park. (Next page, bottom) ►

Also, we knew based on literature that most liocranid adult specimens were collected using pitfall traps and leaf litter sifting [101, 103–105]. Therefore, we used a combination of collecting methods (Fig. 1.6) that target ground spiders [106–108]. This allowed us to have a better representation of the liocranid species present in each site and also to capture a wide array of ground dwelling arthropods, including several families of spiders. While referring to records in taxonomic literature when planning field work is a common practice, very few have made such records persistently available to



all using FAIR [67, 109] standards in the way we did for the present work. More information about this fieldwork, data extraction and applications can be found in **Chapter 2** of this thesis. Specimens collected during this field trip were used to observe and document the genital asymmetries in *T. politus* (**Chapter 3**), and to describe new species and build molecular phylogenies of the families Hahniidae and Symphytognathidae (**Chapters 4 and 5**).

Teutamus politus

Fig. 1.7

This species is a central part of this thesis, together with other related taxa of the *Teutamus* group, of the Chapters 2 and 3. Very little is known about the biology of this interesting species. It was first described in 1890 as the type species for the genus *Teutamus*. Its description was based on one female specimen from “Pulo Pinang”, nowadays Malaysia [110]. But it wasn’t until 2001 that their name resurfaced among many others in an impressive piece of taxonomic literature on South East Asian ground spiders by the Dutch arachnologist Christa Deebleman-Reinhold [101]. This seminal work encouraged research on this and other related spider groups in the region [103–105, 111–113]. Deebleman-Reinhold [101] described six new species of this genus, observing relevant information about the habitat and distribution of these species. She made the first description of the male of *T. politus* and noted the asymmetric genital opening in the female. Dankittipakul et al. [104] significantly increased the knowledge of the genus *Teutamus* by describing 17 new species. They reported many new specimens of *T. politus* increasing the available information about their geographic and temporal distribution. These studies greatly advanced knowledge about the taxonomy, and geographical distribution of *T. politus* and other related spiders. However, their biology is still a mystery.

Teutamus species are known to inhabit the leaf litter of dark humid undisturbed rainforests, but have also been reported wandering on the forest floor in daytime. Most species were found to have limited distribution ranges, being sampled in just one locality without known overlapping distributions [101]. *T. politus* is the notable exception by having a relatively wider distribution that extends from Malaysia to the south of Thailand [101, 104]. This species became interesting to us due to the clearly asymmetric female genitalia. This character had been noted and illustrated before (Fig. 1.7d) in the female and had even been used as a diagnostic character of the species but correlations to changes in male morphology and other implications like courtship and mating behavior (Fig. 1.1c) had never been studied. More information about this can be found in the third chapter of this thesis. While sampling specimens of this species our attention was




Figure 1.7.- *Teutamus politus*. a, b) Comparison of an ant *Odontomachus* sp. and a female of *T. politus*; exemplifies the phenomenon of ant mimicry in this spider species. Our sampling also showed them to be myrmecophylic, living on top or close to the *Odontomachus* sp. nests. c) Size comparison of male and female during the behavioral experiments. d) Taxonomic illustrations of the genitalia of this species (female epigynum —left—, and male pedipalp —right—). Modified from Deeleman-Reinhold [101]; and Dankittipakul et al. [104].

drawn to the relation they had with ants of the genus *Odontomachus* (Fig 7a). Most of the *T. politus* specimens we caught were close to or on top of these ant nests. *T. politus* bears some morphological similarities to the ants (Figs. 1.7a, b). Although they had been mentioned to be “vaguely ant-mimicking” before [101], this is the first time that this kind of spider-ant relationship has been observed in this genus. Unfortunately we were not able to make more detailed observations of the nature of the relationship. Nevertheless, similar interactions have been well documented in other spiders ranging from *myrmecophily* —living in close relation to ants— to *myrmecomorphy* —morphologically mimicking the ants— [114–118].

The present thesis aims to show the value of generating new taxonomic content in a way that allows for rigorous testing of phylogenetic hypotheses, and boosts the impact and relevance of taxonomic work by incorporating high resolution images, molecules and specimen data. Although taxonomy is a venerable science, it holds large quantities of useful data that can help elucidate broad patterns in biological evolution, biogeography, and ecology, among other disciplines. All of these might eventually help understanding, protecting and preserving our endangered biodiversity. We are sitting on a mine whose precious data patiently awaits to be extracted.

Thesis outline

Legacy literature data extraction and application

Chapter 2: Mining data from legacy taxonomic literature and application for sampling spiders of the *Teutamus* group (Araneae; Liocranidae) in Southeast Asia. *Scientific Reports* 10, 15787. doi:10.1038/s41598-020-72549-8.

The use of taxonomic literature to infer evolutionary patterns

Chapter 3: Imperfect and askew: A review of asymmetric genitalia in araneomorph spiders (Araneae: Araneomorphae). *PLoS One* 15:6, e0220354: 1-26. doi:10.1371/journal.pone.0220354.

New taxa description in an integrative approach

Chapter 4: First records and a new genus of Comb-tailed spiders (Araneae: Hahniidae) from Thailand with comments on the six-eyed species of this family. *European Journal of Taxonomy* 724, 51-69. doi: 10.5852/ejt.2020.724.1157

Chapter 5: First records and three new species of the family Symphytognathidae (Arachnida: Araneae) from Thailand, and the circumscription of the genus *Crassignatha* Wunderlich, 1995. *ZooKeys* 1012, 21–53. doi.org/10.3897/zookeys.1012.57047

References

1. Merriam-Webster. [https://www.merriam-webster.com/dictionary/taxonomy#:~:text=1%3A the study of the,Sentences Learn More about taxonomy \(2020\) Taxonomy](https://www.merriam-webster.com/dictionary/taxonomy#:~:text=1%3A%20the%20study%20of%20the%20sentences%20learn%20more%20about%20taxonomy%20(2020)&context=taxonomy). [12 August 2020]
2. Cambridge Dictionary. <https://dictionary.cambridge.org/dictionary/english/taxonomy> (2020) Taxonomy. [12 August 2020]
3. Wikipedia. <https://en.wikipedia.org/wiki/Taxonomy> (2020) Taxonomy. [12 August 2020]
4. Dayrat B. (2005). Towards integrative taxonomy. *Biol. J. Linn. Soc.* **85**, 407–415 DOI: 10.1111/j.1095-8312.2005.00503.x.
5. Godfray HCJ. (2007). Linnaeus in the information age. *Nature* **446**, 259–260 DOI: 10.1038/446259a.
6. Wilson EO. (1985). The biological diversity crisis: A challenge to science. *Issues Sci. Technol.* **2**, 20–29.
7. May RM. (1988). How Many Species Are There on Earth? *Science*. **241**, 1441–1449 DOI: 10.1126/science.241.4872.1441.
8. Wilson EO. (2003). The encyclopedia of life. *Trends Ecol. Evol.* **18**, 77–80 DOI: 10.1016/S0169-5347(02)00040-X.
9. Wilson EO. (2004). Taxonomy as a fundamental discipline. *Philos. Trans. R. Soc. B Biol. Sci.* **359**, 739–739 DOI: 10.1098/rstb.2003.1440.
10. Curtis TP, Sloan WT, Scannell JW. (2002). Estimating prokaryotic diversity and its limits. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 10494–10499 DOI: 10.1073/pnas.142680199.
11. May RM. (2010). Tropical arthropod species, more or less? *Science*. **329**, 41–42 DOI: 10.1126/science.1191058.
12. Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. (2011). How many species are there on earth and in the ocean? *PLoS Biol.* **9**, e1001127 DOI: 10.1371/journal.pbio.1001127.
13. Cresswell ID, Bridgewater P. (2000). The Global Taxonomy Initiative – Quo vadis? *Biol. Int. – News Mag. Int. Union Biol. Sci.* **38**, 12–21.
14. Giangrande A. (2003). Biodiversity, conservation, and the ‘Taxonomic impediment’. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **13**, 451–459 DOI: 10.1002/aqc.584.
15. Wheeler QD, Raven PH, Wilson EO. (2004). Taxonomy: Impediment or Expedient? *Science*. **303**, 285 DOI: 10.1126/science.303.5656.285.
16. Godfray HCJ, Knapp S. (2004). Taxonomy for the twenty-first century: Introduction. *Philos. Trans. R. Soc. B Biol. Sci.* **359**, 559–569 DOI: 10.1098/rstb.2003.1457.
17. Ebach MC, Holdrege C. (2005). More Taxonomy, Not DNA Barcoding. *Bioscience* **55**, 823 DOI: 10.1641/0006-3568(2005)055[0823:MTNDB]2.0.CO;2.
18. de Carvalho MR. (2005). Revisiting the Taxonomic Impediment. *Science*. **307**, 353b–353b DOI: 10.1126/science.307.5708.353b.
19. de Carvalho MR, Bockmann FA, Amorim DS, Brandão CRF, de Vivo M, de Figueiredo JL, Britski HA, de Pinna MCC, Menezes NA, Marques FPL, Papavero N, Cancellato EM, Crisci J V., McEachran JD, Schelly RC, Lundberg JG, Gill AC, Britz R, Wheeler QD, Stiassny MLJ, Parenti LR, Page LM, Wheeler WC, Faivovich J, Vari RP, Grande L, Humphries CJ, DeSalle R, Ebach MC, Nelson GJ. (2007). Taxonomic impediment or impediment to taxonomy? A commentary on systematics and the cybertaxonomic-automation paradigm. *Evol. Biol.* **34**, 140–143 DOI: 10.1007/s11692-007-9011-6.
20. de Carvalho MR, Bockmann FA, Amorim DS, Brandão CRF. (2008). Systematics must embrace comparative biology and evolution, not speed and automation. *Evol. Biol.* **35**, 150–157 DOI: 10.1007/s11692-008-9018-7.
21. Sangster G, Luksenburg JA. (2015). Declining rates of species described per taxonomist: Slowdown of progress or a side-effect of improved quality in taxonomy? *Syst. Biol.* **64**, 144–151 DOI: 10.1093/sysbio/syu069.
22. Britz R, Hundsdoerfer A, Friz U. (2020). Funding, training, permits- the three big challenges of taxonomy. *Megataxa* **1**, 49–52 DOI: 10.11646/megataxa.1.1.10.
23. Zeppelini D, Dal Molin A, Lamas CJE, Sarmiento C, Rheims CA, Fernandes DRR, Lima EFB, Silva EN, Carvalho-Filho F, Kovac L, Montoya-Lerma J, Moldovan OT, Souza-Dias PGB, Demite PR, Feitosa RM, Boyer SL, Weiner WM, Rodrigues WC. (2020). The dilemma of self-citation in taxonomy. *Nat. Ecol. Evol.* [Correspondence] DOI: 10.1038/s41559-020-01359-y.

24. Retraction Watch. <https://retractionwatch.com/2020/07/28/major-indexing-service-reverses-decision-to-suppress-two-journals-from-closely-followed-metric/#more-120145> (2020) Major indexing service reverses decision to suppress two journals from closely followed metric.
25. Clarivate Analytics. <http://jcr.help.clarivate.com/Content/title-suppressions.htm> (2020) Title Suppressions.
26. NSF. (National Science Foundation, 1994). *Partnership for Enhancing Expertise in Taxonomy (PEET) program announcement*.
27. Rodman JE, Cody JH. (2003). The taxonomic impediment overcome: NSF's partnerships for enhancing expertise in taxonomy (PEET) as a model. *Syst. Biol.* **52**, 428–435 DOI: 10.1080/10635150309326.
28. Agnarsson I, Kuntner M. (2007). Taxonomy in a changing world: Seeking solutions for a science in crisis. *Syst. Biol.* **56**, 531–539 DOI: 10.1080/10635150701424546.
29. Knapp S, Bateman RM, Chalmers NR, Humphries CJ, Rainbow PS, Smith AB, Taylor PD, Vane-Wright RI, Wilkinson M. (2002). Taxonomy needs evolution, not revolution. *Nature* **419**, (6907):559 DOI: 10.1038/419559a.
30. Godfray HCJ. (2002). Challenges for taxonomy. *Nature* **417**, 17–19 DOI: 10.1038/417017a.
31. Scoble MJ. (2004). Unitary or unified taxonomy? *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **359**, 699–710 DOI: 10.1098/rstb.2003.1456.
32. Hebert PDN, Gregory TR. (2005). The promise of DNA barcoding for taxonomy. *Syst. Biol.* **54**, 852–859 DOI: 10.1080/10635150500354886.
33. Miller SE. (2007). DNA barcoding and the renaissance of taxonomy. *Proc. Natl. Acad. Sci.* **104**, 4775–4776 DOI: 10.1073/pnas.0700466104.
34. Godfray HCJ, Mayo SJ, Scoble MJ. (2008). Pragmatism and rigour can coexist in taxonomy. *Evol. Biol.* **35**, 309–311 DOI: 10.1007/s11692-008-9041-8.
35. Knapp S. (CRC Press, 2008). Taxonomy as a team sport. in *The New Taxonomy* (ed. Wheeler, Q. D.) 33–54 DOI: 10.1201/9781420008562.
36. Wheeler QD. (CRC Press, 2008). Introductory: Toward the new taxonomy. in *The New Taxonomy* (ed. Wheeler, Q. D.) DOI: 10.1201/9781420008562.
37. Wheeler QD. (CRC Press, 2008). Taxonomic shock and awe. in *The New Taxonomy* (ed. Wheeler, Q. D.) 211–226 DOI: 10.1201/9781420008562.
38. Clark BR, Godfray HCJ, Kitching IJ, Mayo SJ, Scoble MJ. (2009). Taxonomy as an eScience. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **367**, 953–966 DOI: 10.1098/rsta.2008.0190.
39. Wheeler QD. (2008). *The New Taxonomy*. CRC Press.
40. Zauner H. (2009). Evolving e-taxonomy. *BMC Evol. Biol.* **9**, 141 DOI: 10.1186/1471-2148-9-141.
41. Wheeler QD, Valdecasas AG. (2010). Cybertaxonomy and ecology. *Nat. Educ. Knowl.* **3**, 6.
42. Orr MC, Ferrari RR, Hughes AC, Chen J, Ascher JS, Yan Y-H, Williams PH, Zhou X, Bai M, Rudoy A, Zhang F, Ma K-P, Zhu C-D. (2020). Taxonomy must engage with new technologies and evolve to face future challenges. *Nat. Ecol. Evol.* [Correspondence] DOI: 10.1038/s41559-020-01360-5.
43. Pyle RL. (2016). Towards a global names architecture: The future of indexing scientific names. *Zookeys* **550**, 261–281 DOI: 10.3897/zookeys.550.10009.
44. Agosti D, Johnson NF. <http://antbase.org/> (2005) Antbase. [25 September 2020]
45. Antweb. <https://www.antweb.org/> (2020) AntWeb. Version 8.41. California Academy of Science. [25 September 2020]
46. Beccaloni G, Scoble M, Kitching I, Simonsen T, Robinson G, Pitkin B, Hine A, Lyal C. <https://www.nhm.ac.uk/our-science/data/lepindex/> (2003) The Global Lepidoptera Names Index.
47. Evenhuis N, Pape T. <http://www.diptera.org/> (2020) Systema Dipterorum.
48. Cigliano MM, Braun H. <http://orthoptera.speciesfile.org/HomePage/Orthoptera/HomePage.aspx> (2020) Orthoptera Species File Online. [25 September 2020]
49. Froese R, Pauly D. (2019) FishBase. www.fishbase.org, version [25 September 2020]
50. Godfray HCJ, Clark BR, Kitching IJ, Mayo SJ, Scoble MJ. (2007). The web and the structure of taxonomy. *Syst. Biol.* **56**, 943–55 DOI: 10.1080/10635150701777521.

51. WSC. <http://www.wsc.nmbe.ch/> (2020) World Spider Catalog Version 21.0. *Natural History Museum Bern*, online at <http://wsc.nmbe.ch> DOI: 10.24436/2.
52. Platnick NI. (1989). *Advances in Spider Taxonomy 1981-1987: A Supplement to Brignoli's A Catalogue of the Araneae described between 1940 and 1981*.
53. Platnick NI. (Entomological Society of New York, 1993). *Advances in spider taxonomy 1988-1991, with synonymies and transfers 1940-1980*.
54. Platnick NI. (Entomological Society New York, 1998). *Advances in Spider Taxonomy, 1992-1995 With Redescriptions 1940-1980. Entomological Society New York*.
55. WSC. <https://wsc.nmbe.ch/archive/> (2020) Archive of the World Spider Catalog.
56. ICZN. (The International Trust for Zoological Nomenclature 1999, 2000). *International Code of Zoological Nomenclature*.
57. Platnick NI, Raven RJ. (2013). Spider systematics: Past and future. *Zootaxa* **3683**, 595–600 DOI: 10.11646/zootaxa.3683.5.8.
58. Miller JA, Dikow T, Agosti D, Sautter G, Catapano T, Penev L, Zhang Z-Q, Pentcheff D, Pyle R, Blum S, Parr C, Freeland C, Garnett T, Ford LS, Muller B, Smith L, Strader G, Georgiev T, Bénichou L. (2012). From taxonomic literature to cybertaxonomic content. *BMC Biol.* **10**, 87 DOI: 10.1186/1741-7007-10-87.
59. Gwinn NE, Rinaldo C. (2009). The Biodiversity Heritage Library: sharing biodiversity literature with the world. *IFLA J.* **35**, 25–34 DOI: 10.1177/0340035208102032.
60. Page RDM. (2010). Enhanced display of scientific articles using extended metadata. *J. Web Semant.* **8**, 190–195 DOI: 10.1016/j.websem.2010.03.004.
61. Plazi. (2020). PLAZI Home Page. Available from <http://plazi.org/org> [20th June 2020].
62. Catapano T. (National Center for Biotechnology Information, 2010). NoTaxPub: An Extension of the NLM/NCBI Journal Publishing DTD for Taxonomic Descriptions. in *Journal Article Tag Suite Conference (JATS-Con) Proceedings 2010* DOI: 10.5281/zenodo.3484285.
63. Penev L, Agosti D, Georgiev T, Catapano T, Miller JA, Blagoderov V, Roberts D, Smith VS, Brake I, Rycroft S, Scott B, Johnson NF, Morris RA, Sautter G, Chavan V, Robertson T, Remsen D, Stoev P, Parr C, Knapp S, Kress WJ, Thompson FC, Erwin TL. (2010). Semantic tagging of and semantic enhancements to systematics papers: Zookeys working examples. *Zookeys* **50**, 1–16 DOI: 10.3897/zookeys.50.538.
64. Penev L, Lyal CHC, Weitzman A, Morse DR, King D, Sautter G, Georgiev T, Morris RA, Catapano T, Agosti D. (2011). XML schemas and mark-up practices of taxonomic literature. *Zookeys* **150**, 89–116 DOI: 10.3897/zookeys.150.2213.
65. Agosti D, Egloff W. (2009). Taxonomic information exchange and copyright: the Plazi approach. *BMC Res. Notes* **2**, 53 DOI: 10.1186/1756-0500-2-53.
66. Agosti D, Catapano T, Sautter G, Egloff W. (2019). The Plazi Workflow: The PDF prison break for biodiversity data. *Biodivers. Inf. Sci. Stand.* **3**, e37046 DOI: 10.3897/biss.3.37046.
67. Chester C, Agosti D, Sautter G, Catapano T, Martens K, Gérard I, Bénichou L. (2019). EJT editorial standard for the semantic enhancement of specimen data in taxonomy literature. *Eur. J. Taxon.* **586**, DOI: 10.5852/ejt.2019.586.
68. Miller JA, Agosti D, Penev L, Sautter G, Georgiev T, Catapano T, Patterson D, King D, Pereira S, Vos RA, Sierra S. (2015). Integrating and visualizing primary data from prospective and legacy taxonomic literature. *Biodivers. data J.* **3**, e5063 DOI: 10.3897/BDJ.3.e5063.
69. GBIF. (2019) Global Biodiversity Information Facility Home Page. Available from: <http://www.gbif.org> [4th April 2019].
70. Zenodo. <https://zenodo.org/> (2020) Zenodo - Research. Shared.
71. Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I. (2019). GenBank. *Nucleic Acids Res.* DOI: 10.1093/nar/gky989.
72. Huber BA. (2004). Evidence for functional segregation in the directionally asymmetric male genitalia of the spider *Metagonia mariguitarensis* (Gonzalez-Sponga) (Pholcidae: Araneae). *J. Zool.* **262**, 317–326 DOI: 10.1017/S0952836903004709.
73. Agnarsson I. (2006). Asymmetric female genitalia and other remarkable morphology in a new genus of cobweb spiders (Theridiidae, Araneae) from Madagascar. *Biol. J. Linn. Soc.* **87**, 211–232

DOI: 10.1111/j.1095-8312.2006.00569.x.

74. Huber BA. (2006). Cryptic female exaggeration: The asymmetric female internal genitalia of *Kaliana yuruani* (Araneae: Pholcidae). *J. Morphol.* **267**, 705–712 DOI: 10.1002/jmor.10431.
75. Huber BA, Sinclair BJ, Schmitt M. (2007). The evolution of asymmetric genitalia in spiders and insects. *Biol. Rev.* **82**, 647–698 DOI: 10.1111/j.1469-185X.2007.00029.x.
76. Holwell GI, Herberstein ME. (2010). Chirally dimorphic male genitalia in praying mantids (Ciulfina: Liturgusidae). *J. Morphol.* **271**, 1176–1184 DOI: 10.1002/jmor.10861.
77. Burger M. (2010). Complex female genitalia indicate sperm dumping in armored goblin spiders (Arachnida, Araneae, Oonopidae). *Zoology* **113**, 19–32 DOI: 10.1016/j.zool.2009.04.002.
78. Schilthuizen M. (2013). Something gone awry: unsolved mysteries in the evolution of asymmetric animal genitalia. *Anim. Biol.* **63**, 1–20 DOI: 10.1163/15707563-00002398.
79. Huber BA, Nuñeza OM. (2015). Evolution of genital asymmetry, exaggerated eye stalks, and extreme palpal elongation in Panjange spiders (Araneae: Pholcidae). *Eur. J. Taxon.* **169**, DOI: 10.5852/ejt.2015.169.
80. Holwell GI, Kazakova O, Evans F, O’Hanlon JC, Barry KL. (2015). The functional significance of chiral genitalia: Patterns of asymmetry, functional morphology and mating success in the praying mantis *Ciulfina baldersoni*. *PLoS One* **10**, DOI: 10.1371/journal.pone.0128755.
81. Rivera-Quiroz FA, Schilthuizen M, Petcharad B, Miller JA. (2020). Imperfect and askew: A review of asymmetric genitalia in araneomorph spiders (Araneae: Araneomorphae). *PLoS One* **15**, e0220354 DOI: 10.1371/journal.pone.0220354.
82. Valdecasas AG, Williams D, Wheeler QD. (2008). ‘Integrative taxonomy’ then and now: A response to Dayrat (2005). *Biol. J. Linn. Soc.* **96**, 211–216 DOI: 10.1111/j.1095-8312.2007.00919.x.
83. Wake MH. (2003). What is ‘integrative biology’? in *Integrative and Comparative Biology* DOI: 10.1093/icb/43.2.239.
84. Padial JM, Miralles A, De la Riva I, Vences M. (2010). The integrative future of taxonomy. *Front. Zool.* **7**, 16 DOI: 10.1186/1742-9994-7-16.

85. Schlick-Steiner BC, Steiner FM, Seifert B, Stauffer C, Christian E, Crozier RH. (2010). Integrative Taxonomy: A Multisource Approach to Exploring Biodiversity. *Annu. Rev. Entomol.* **55**, 421–438 DOI: 10.1146/annurev-ento-112408-085432.
86. Xu X, Liu F, Chen J, Li D, Kuntner M. (2015). Integrative taxonomy of the primitively segmented spider genus *Ganthela* (Araneae: Mesothelae: Liphistiidae): DNA barcoding gap agrees with morphology. *Zool. J. Linn. Soc.* **175**, 288–306 DOI: 10.1111/zoj.12280.
87. Lopardo L, Giribet G, Hormiga G. (2011). Morphology to the rescue: Molecular data and the signal of morphological characters in combined phylogenetic analyses—a case study from mysmenid spiders (Araneae, Mysmenidae), with comments on the evolution of web architecture. *Cladistics* **27**, 278–330 DOI: 10.1111/j.1096-0031.2010.00332.x.
88. Lopardo L, Hormiga G. (2015). Out of the twilight zone: Phylogeny and evolutionary morphology of the orb-weaving spider family Mysmenidae, with a focus on spinneret spigot morphology in symphytognathoids (Araneae, Araneioidea). *Zool. J. Linn. Soc.* **173**, 527–786 DOI: 10.1111/zoj.12199.
89. Ramírez MJ. (2014). The morphology and phylogeny of dionychan spiders (Araneae, Araneomorphae). *Bull. Am. Museum Nat. Hist.* **390**, 1–374 DOI: doi.org/10.1206/821.1.
90. Cao X, Liu J, Chen J, Zheng G, Kuntner M, Agnarsson I. (2016). Rapid dissemination of taxonomic discoveries based on DNA barcoding and morphology. *Sci. Rep.* **6**, 37066 DOI: 10.1038/srep37066.
91. Wheeler WC, Coddington JA, Crowley LM, Dimitrov D, Goloboff PA, Griswold CE, Hormiga G, Prendini L, Ramírez MJ, Sierwald P, Almeida-Silva L, Alvarez-Padilla F, Arnedo MA, Benavides Silva LR, Benjamin SP, Bond JE, Grismado CJ, Hasan E, Hedin M, Izquierdo MA, Labarque FM, Ledford J, Lopardo L, Maddison WP, Miller JA, Piacentini LN, Platnick NI, Polotow D, Silva-Dávila D, Scharff N, Szűts U, Ubick D, Vink CJ, Wood HM, Zhang J. (2017). The spider tree of life: phylogeny of Araneae based on target-gene analyses from an extensive taxon sampling. *Cladistics* **33**, 574–616 DOI: 10.1111/cla.12182.
92. Feng C, Miller JA, Lin Y, Shu Y. (2019). Further study of two chinese cave spiders (Araneae, mysmenidae), with description of a

new genus. *Zookeys* **870**, 77–100 DOI: 10.3897/zookeys.870.35971.

93. Lipke E, Hammel JU, Michalik P. (2015). First evidence of neurons in the male copulatory organ of a spider (Arachnida, Araneae). *Biol. Lett.* **11**, DOI: 10.1098/rsbl.2015.0465.

94. Sentenská L, Müller CHG, Pekár S, Uhl G. (2017). Neurons and a sensory organ in the pedipalps of male spiders reveal that it is not a numb structure. *Sci. Rep.* **7**, 12209 DOI: 10.1038/s41598-017-12555-5.

95. Dederichs TM, Müller CHG, Sentenská L, Lipke E, Uhl G, Michalik P. (2019). The innervation of the male copulatory organ of spiders (Araneae) - A comparative analysis. *Front. Zool.* **16**, DOI: 10.1186/s12983-019-0337-6.

96. Mittermeier RA, Robles-Gil P, Hoffmann M, Pilgrim J, Brooks TM, Goettsch-Mittermeier C, Lamoreux JF, Fonseca GAB da. (2004). *Hotspots Revisited. Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions*.

97. Hrdina A, Romportl D. (2017). Evaluating Global Biodiversity Hotspots-Very Rich and even More Endangered. *J. Landsc. Ecol.* **10**, 107–115 DOI: 10.1515/jlecol-2017-0013.

98. Hanson T, Brooks TM, Da Fonseca GAB, Hoffmann M, Lamoreux JF, MacHlis G, Mittermeier CG, Mittermeier RA, Pilgrim JD. (2009). Warfare in biodiversity hotspots. *Conserv. Biol.* **3**, 578–587 DOI: 10.1111/j.1523-1739.2009.01166.x.

99. CEPF. <https://www.cepf.net/our-work/biodiversity-hotspots/indo-burma/species> (2020) Critical Ecosystem Partnership Fund: protecting biodiversity by empowering people.

100. Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C. (2011). Global Biodiversity Conservation: The Critical Role of Hotspots. in *Biodiversity Hotspots* DOI: 10.1007/978-3-642-20992-5_1.

101. Deeleman-Reinhold C. (Leiden: Brill, 2001). *Forest spiders of South East Asia: with a revision of the sac and ground spiders (Araneae: Clubionidae, Corinnidae, Liocranidae, Gnaphosidae, Prodidomidae and Trochanterriidae)*.

102. Jäger P, Dankittipakul P. (2010). Clubionidae from Laos and Thailand (Arachnida: Araneae). *Zootaxa* **2730**, 23–43 DOI: 10.11646/zootaxa.2730.1.2.

103. Dankittipakul P, Deeleman-Reinhold C. (2013). Delimitation of the spider genus

Sesieutes Simon, 1897, with descriptions of five new species from South East Asia (Araneae: Corinnidae). *J. Nat. Hist.* **47**, 167–195 DOI: 10.1080/00222933.2012.742165.

104. Dankittipakul P, Tavano M, Singtripop T. (2012). Seventeen new species of the spider genus *Teutamus* Thorell, 1890 from Southeast Asia (Araneae: Liocranidae). *J. Nat. Hist.* **46**, 1689–1730 DOI: 10.1080/00222933.2012.681314.

105. Dankittipakul P, Tavano M, Singtripop T. (2011). Neotype designation for *Sphingius thecatus* Thorell 1890 synonymies new records and descriptions of six new species from Southeast Asia (Araneae Liocranidae). *Zootaxa* 1–20.

106. Colwell RK, Coddington JA. (1994). Estimating Terrestrial Biodiversity through Extrapolation. *Philos. Trans. R. Soc. B Biol. Sci.* **345**, 101–118 DOI: 10.1098/rstb.1994.0091.

107. Scharff N, Coddington J a., Griswold CE, Hormiga G, Bjørn P de P. (2003). When to quit? estimating spider species richness in a northern European deciduous forest. *J. Arachnol.* **31**, 246–273 DOI: 10.1636/0161-8202(2003)031[0246:WTQESS]2.0.CO;2.

108. Coddington JA, Agnarsson I, Miller JA, Kuntner M, Hormiga G. (2009). Undersampling bias: The null hypothesis for singleton species in tropical arthropod surveys. *J. Anim. Ecol.* **78**, 573–584 DOI: 10.1111/j.1365-2656.2009.01525.x.

109. Wilkinson MD, Dumontier M, Aalbersberg IJ, Appleton G, Axton M, Baak A, Blomberg N, Mons B, et al. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* **3**, DOI: 10.1038/sdata.2016.18.

110. Thorell T. (1890). Arachnidi di Pinang raccolti nel 1889 dai Signori L. Loria e L. Fea. *Ann. del Mus. Civ. di Stor. Nat. di Genova* **30**, 269–383.

111. Ono H. (2009). Three new spiders of the family Clubionidae, Liocranidae and Gnaphosidae (Arachnida, Araneae) from Vietnam. *Bull. Natl. Museum Nat. Sci. Tokyo* **35**, 1–8.

112. Dankittipakul P, Deeleman-Reinhold C. (2012). A new spider species of the genus *Sudharmia* from Sumatra, Indonesia (Araneae, Liocranidae). *Dongwuxue Yanjiu* **33**, 187–190 DOI: 10.3724/SP.J.1141.2012.02187.

113. Dankittipakul P, Tavano M, Singtripop T. (2013). Revision of the spider genus *Jacaena* Thorell, 1897, with descriptions of four new species from

Chapter • 1

Thailand (Araneae: Corinnidae). *J. Nat. Hist.* **47**, 1539–1567 DOI: 10.1080/00222933.2012.763059.

114. Emerton JH. (1911). New spiders from New England. *Trans. Connect. Acad. Arts Sci.* **16**, 383–407.

115. Penniman AJ. (1978). Taxonomic and natural history notes on *Phrurolithus fratellus* Gertsch (Araneae: Clubionidae). *J. Arachnol.* **6**, 125–132 DOI: 10.2307/3705134.

116. Cushing PE. (1997). Myrmecomorphy and Myrmecophily in Spiders: A Review. *Florida Entomol.* **8**, 165–193 DOI: 10.2307/3495552.

117. Cushing PE. (2012). Spider-ant associations: An updated review of myrmecomorphy, myrmecophily, and myrmecophagy in spiders. *Psyche (London)* **2012**, DOI: 10.1155/2012/151989.

118. Ramírez MJ, Grismado CJ, Ubick D, Ovtsharenko V, Cushing PE, Platnick NI, Wheeler WC, Prendini L, Crowley LM, Horner N V. (2019). Myrmecicultoridae, a New Family of Myrmecophilic Spiders from the Chihuahuan Desert (Araneae: Entelegynae). *Am. Museum Novit.* **3930**, 1–24 DOI: 10.1206/3930.1.