



Universiteit
Leiden
The Netherlands

Integrative taxonomy of araneomorph spiders: Breathing new life into an old science

Rivera Quiroz, F.A.

Citation

Rivera Quiroz, F. A. (2021, April 14). *Integrative taxonomy of araneomorph spiders: Breathing new life into an old science*. Retrieved from <https://hdl.handle.net/1887/3152423>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3152423>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/3152423> holds various files of this Leiden University dissertation.

Author: Rivera Quiroz, F.A.

Title: Integrative taxonomy of araneomorph spiders: Breathing new life into an old science

Issue date: 2021-04-14

Chapter 6

Epilogue

F. ANDRES RIVERA-QUIROZ*^{1, 2}

¹ Understanding Evolution group, Naturalis Biodiversity Center, Darwinweg 2, 2333CR Leiden, The Netherlands

² Institute for Biology Leiden (IBL), Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands.

General discussion

The present thesis highlights the relevance and usefulness of taxonomic literature (old and new) and exemplifies the benefits of adopting an integral approach to the description of new taxa. As mentioned in **Chapter 1**, taxonomy as a science has accumulated data and knowledge for more than 250 years. The quality and usefulness of the facts recorded in taxonomic literature have greatly improved from the early (purely) descriptive texts to the modern works that are rich in detailed and integrated data. These developments have improved the rigor of phylogenetic inference, documented distribution patterns of taxa through time and space, and revealed broad evolutionary patterns and other interesting phenomena. My work illustrates some applications of legacy data contained in literature (**Chapters 2 and 3**), and also explores an integrative perspective that involves new taxonomic descriptions and generation of phylogenetic hypotheses integrating molecular and morphological data (**Chapters 4 and 5**).

Taxonomy has provided a system under which groups of natural entities can be catalogued and biological data can be aggregated by the use of a taxonomic name (e.g. genus or species). Nevertheless, slow pace of identification, description, and categorization, plus the huge number of unknown species and human impact on natural habitats and biodiversity have made traditional taxonomy obsolete and partially unreliable. As a response to this, several authors have made patent the urgent need for taxonomy to accelerate its description and knowledge accumulation process by the change of some practices and use of new technologies [1–9]. One of the main drivers of this change involves the access to primary taxonomic literature and specimen information [1, 6]. My thesis heavily relies on the utilization of some of these “e-taxonomic” products, like the BHL [10], WSC [11], Plazi [12], and GBIF [13]; and also helps to test and collaborates on the improvement of the program Golden Gate Imagine, the software used for data mining in **Chapter 2**. Electronic access to literature is one of the primary tools in my dissertation, being fundamental for acquiring primary taxonomic data used in **Chapter 2** to analyze the distribution of the *Teutamus* group; observing the patterns and formulating an evolutionary hypotheses for the origin of genital asymmetry in spiders (**Chapter 3**); and gathering taxonomic information that helped us identify and eventually describe new species and test the phylogenies of the the spider families Hahniidae (**Chapter 4**) and the Symphytognathidae (**Chapter 5**).

The approach used in my thesis for mining and gathering specimen data from taxonomic literature—used in **Chapter 2**—proved to be a powerful tool for analyzing species distributions. Here, I applied this knowledge to plan our fieldwork targeting adult specimens of one particular group of ground dwelling spiders; however, there are many other possible uses that remain to be explored in the future. Some examples of these applications include species estimations, catalog building and taxonomic inconsistencies identification, inclusion of underrepresented taxa in global datasets, species distribution patterns detection, among other applications. This potent tool has been gaining traction among taxonomic journals and publishers (like Pensoft, Zootaxa, Zookeys, among

others) in its prospective approach [14, 15]; and is also being used in projects similar to the one presented here that have extracted data and analyzed the legacy taxonomic literature of groups as diverse as on damselflies [16] and *Tyrannosaurus rex* [17, 18].

Besides the aforementioned specimen data, taxonomic literature has also linked other types of information like illustrations, photographs, and molecular data to taxonomic names. In this way, I think of taxonomic literature as a massive (yet largely unstructured) data repository. This repository has accumulated biological data in the form of text called treatments [19–21]. The quality and usefulness of some of these data greatly varies depending on the time, author and even taxonomic group, and has —without any doubt — improved with utilization of modern techniques and technologies [22]. Spider taxonomy — probably one of the best curated bodies of taxonomic literature (see Chapter 1) — has accumulated and made available thousands of taxonomic documents that represent a huge collection of species, facts, and images. Similar to other arthropod groups, spider species delimitation greatly relies on the comparative morphology of the genital characters; therefore, it does not come as a surprise that sexual characters are some of the best studied and understood, having detailed images and descriptions that might allow for the detection of interesting phenomena and broad evolutionary patterns. One example of this is the evolution of asymmetry in spider genitalia [23, 24]. For **Chapter 3**, I conducted a casual but taxonomically broad study of the evolution of this character in spiders, assigning the identified cases to traditional categories of asymmetry [23–26] and suggested the evolutionary patterns and causes behind the development of this morphological character. Moreover, I attempted to test some of these hypotheses by examining the behavioral implications of the evolution of genital asymmetry in the species *Teutamus politus*; although I collected a fair number of live specimens during our fieldwork in an attempt to observe courtship and mating, the behavioral experiments were unsuccessful. Nevertheless, ours was the most comprehensive study of asymmetry in spiders encountering dozens of species in several families that show some kind of asymmetry and had been overlooked in the scarce previous reviews on the topic.

The sampling methods I used during our fieldwork — although intended for the collection of *Teutamus* group spiders — also captured much more material. Some of these specimens were used in this thesis for **Chapters 3, 4 and 5**; but considerably more material remains to be studied. Our sampling collected thousands of specimens from several arthropod groups including insects, millipedes, mites, Opiliones, among many others. Together with my supervisor Jeremy Miller and our collaborator Booppa Petcharad, I collected more than 4,600 spider specimens of which 1,454 were adults representing 35 spider families. Only the families Liocranidae, Hahniidae and Symphytognathidae (the latter two both new records for Thailand) were identified to species level. Still, from this relatively small selection of taxa, I described a total of five new species following the integrative approach I mentioned in the introduction and including morphological and molecular data. All of the specimens collected for the present project are now deposited in the collection of Naturalis Biodiversity Center, where they can be archived and eventually contribute to other taxonomic, systematic, ecological and evolutionary studies.

In this context, **Chapters 4 and 5** focused on the generation of new taxonomic content using the integrative approach [27–33]. In these two chapters I described five new species and made the first reports of two families for Thailand. Both families — as in the aforementioned *Teutamus* group — are ground dwelling, mostly inhabiting leaf litter. They are relatively small to tiny, ranging from less than 1 cm down to less than 0.5 mm! In fact, one of the newly described species, *Anapistula choojaiae* n. sp., could be considered among the smallest spiders ever discovered together with a couple other species in the genus *Patu* [34, 35]. Both chapters are examples of the integration of new taxonomic descriptions with high resolution photographs, molecular data and character evolution, similar to what has been done previously in other spider studies [27, 29, 31, 33]. In the case of the Hahniidae (**Chapter 4**) by reviewing what is known about the eye size reduction and eventual eye loss within this family; and for the Symphytognathidae (**Chapter 5**) by using 3D modeling to accurately document and compare the genital morphology allowing us to better circumscribe the genus *Crassignatha*, and transferring there some species previously misplaced in the genus *Patu*.

I consider that this thesis demonstrates the use of many of the new taxonomic *e-tools* like specimen information databases (e.g. GBIF) and literature repositories, and data mining and management resources (e.g. BHL, WSC, PLAZI). In this way, the present work illustrates the re-use and re-analysis of specimen data and morphological images contained in literature, and also features the use of an integrative taxonomic approach for new descriptions that allow for species documentation, as well as the inference of evolutionary hypotheses. I feel that the common use of these tools will, without any doubt, help overcome the taxonomic impediment while collaborating on the effort of describing and understanding our (greatly endangered) biodiversity.

References

1. Godfray HCJ. (2002). Challenges for taxonomy. *Nature* **417**, 17–19 DOI: 10.1038/417017a.
2. Godfray HCJ. (2007). Linnaeus in the information age. *Nature* **446**, 259–260 DOI: 10.1038/446259a.
3. Godfray HCJ, Clark BR, Kitching IJ, Mayo SJ, Scoble MJ. (2007). The web and the structure of taxonomy. *Syst. Biol.* **56**, 943–55 DOI: 10.1080/10635150701777521.
4. de Carvalho MR, Bockmann FA, Amorim DS, Brandão CRF, de Vivo M, de Figueiredo JL, Britski HA, de Pinna MCC, Menezes NA, Marques FPL, Papavero N, Cancellato EM, Crisci J V., McEachran JD, Schelly RC, Lundberg JG, Gill AC, Britz R, Wheeler QD, Stiassny MLJ, Parenti LR, Page LM, Wheeler WC, Faivovich J, Vari RP, Grande L, Humphries CJ, DeSalle R, Ebach MC, Nelson GJ. (2007). Taxonomic impediment or impediment to taxonomy? A commentary on systematics and the cybertaxonomic-automation paradigm. *Evol. Biol.* **34**, 140–143 DOI: 10.1007/s11692-007-9011-6.
5. Wheeler QD. (2008). *The New Taxonomy*. CRC Press.
6. Wheeler QD, Raven PH, Wilson EO. (2004). Taxonomy: Impediment or Expedient? *Science*. **303**, 285 DOI: 10.1126/science.303.5656.285.
7. Godfray HCJ, Knapp S. (2004). Taxonomy for the twenty-first century: Introduction. *Philos. Trans. R. Soc. B Biol. Sci.* **359**, 559–569 DOI: 10.1098/rstb.2003.1457.
8. Clark BR, Godfray HCJ, Kitching IJ, Mayo

- SJ, Scoble MJ. (2009). Taxonomy as an eScience. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **367**, 953–966 DOI: 10.1098/rsta.2008.0190.
9. Orr MC, Ferrari RR, Hughes AC, Chen J, Ascher JS, Yan Y-H, Williams PH, Zhou X, Bai M, Rudoy A, Zhang F, Ma K-P, Zhu C-D. (2020). Taxonomy must engage with new technologies and evolve to face future challenges. *Nat. Ecol. Evol.* [Correspondence] DOI: 10.1038/s41559-020-01360-5.
10. Gwinn NE, Rinaldo C. (2009). The Biodiversity Heritage Library: sharing biodiversity literature with the world. *IFLA J.* **35**, 25–34 DOI: 10.1177/0340035208102032.
11. WSC. <http://www.wsc.nmbe.ch/> (2020) World Spider Catalog Version 21.0. *Natural History Museum Bern*, online at <http://wsc.nmbe.ch> DOI: 10.24436/2.
12. Plazi. (2020). PLAZI Home Page. Available from <http://plazi.org/org> [20th June 2020].
13. GBIF. (2019) Global Biodiversity Information Facility Home Page. Available from: <http://www.gbif.org> [4th April 2019].
14. Chester C, Agosti D, Sautter G, Catapano T, Martens K, Gérard I, Bénichou L. (2019). EJT editorial standard for the semantic enhancement of specimen data in taxonomy literature. *Eur. J. Taxon.* **586**, DOI: 10.5852/ejt.2019.586.
15. Agosti D, Catapano T, Sautter G, Egloff W. (2019). The Plazi Workflow: The PDF prison break for biodiversity data. *Biodivers. Inf. Sci. Stand.* **3**, e37046 DOI: 10.3897/biss.3.37046.
16. Plazi.org. <https://tinyurl.com/y6d94ega> (2020) Taxonomic search Lestes.
17. Miller J, Braumuller Y, Kishor P, Shorthouse D, Dimitrova M, Sautter G, Agosti D. (2019). Mobilizing Data from Taxonomic Literature for an Iconic Species (Dinosauria, Theropoda, Tyrannosaurus rex). *Biodivers. Inf. Sci. Stand.* DOI: 10.3897/biss.3.37078.
18. Plazi.org. <https://tinyurl.com/y3f3kq7u> (2020) Taxonomic search Tyrannosaurus.
19. Catapano T. (National Center for Biotechnology Information, 2010). NoTaxPub: An Extension of the NLM/NCBI Journal Publishing DTD for Taxonomic Descriptions. in *Journal Article Tag Suite Conference (JATS-Con) Proceedings 2010* DOI: 10.5281/zenodo.3484285.
20. Penev L, Agosti D, Georgiev T, Catapano T, Miller JA, Blagoderov V, Roberts D, Smith VS, Brake I, Rycroft S, Scott B, Johnson NF, Morris RA, Sautter G, Chavan V, Robertson T, Remsen D, Stoev P, Parr C, Knapp S, Kress WJ, Thompson FC, Erwin TL. (2010). Semantic tagging of and semantic enhancements to systematics papers: Zookeys working examples. *Zookeys* **50**, 1–16 DOI: 10.3897/zookeys.50.538.
21. Penev L, Lyal CHC, Weitzman A, Morse DR, King D, Sautter G, Georgiev T, Morris RA, Catapano T, Agosti D. (2011). XML schemas and mark-up practices of taxonomic literature. *Zookeys* **150**, 89–116 DOI: 10.3897/zookeys.150.2213.
22. Pyle RL. (2016). Towards a global names architecture: The future of indexing scientific names. *Zookeys* **550**, 261–281 DOI: 10.3897/zookeys.550.10009.
23. Huber BA, Sinclair BJ, Schmitt M. (2007). The evolution of asymmetric genitalia in spiders and insects. *Biol. Rev.* **82**, 647–698 DOI: 10.1111/j.1469-185X.2007.00029.x.
24. Schilthuizen M. (2013). Something gone awry: unsolved mysteries in the evolution of asymmetric animal genitalia. *Anim. Biol.* **63**, 1–20 DOI: 10.1163/15707563-00002398.
25. Palmer AR. (Elsevier, 2005). Antisymmetry. in *Variation* (eds. Hallgrinsson, B. & Hall, B.) 359–397 DOI: 10.1016/B978-012088777-4/50018-1.
26. Huber BA. (2010). Mating positions and the evolution of asymmetric insect genitalia. *Genetica* **138**, 19–25 DOI: 10.1007/s10709-008-9339-6.
27. Lopardo L, Giribet G, Hormiga G. (2011). Morphology to the rescue: Molecular data and the signal of morphological characters in combined phylogenetic analyses—a case study from mysmenid spiders (Araneae, Mysmenidae), with comments on the evolution of web architecture. *Cladistics* **27**, 278–330 DOI: 10.1111/j.1096-0031.2010.00332.x.
28. Xu X, Liu F, Chen J, Li D, Kuntner M. (2015). Integrative taxonomy of the primitively segmented spider genus *Ganthea* (Araneae: Mesothelae: Liphistiidae): DNA barcoding gap agrees with morphology. *Zool. J. Linn. Soc.* **175**, 288–306 DOI: 10.1111/zoj.12280.
29. Lopardo L, Hormiga G. (2015). Out of the twilight zone: Phylogeny and evolutionary morphology of the orb-weaving spider family

- Mysmenidae, with a focus on spinneret spigot morphology in symphytognathoids (Araneae, Araneioidea). *Zool. J. Linn. Soc.* **173**, 527–786 DOI: 10.1111/zoj.12199.
30. Ramírez MJ. (2014). The morphology and phylogeny of dionychan spiders (Araneae, Araneomorphae). *Bull. Am. Museum Nat. Hist.* **390**, 1–374 DOI: doi.org/10.1206/821.1.
31. Cao X, Liu J, Chen J, Zheng G, Kuntner M, Agnarsson I. (2016). Rapid dissemination of taxonomic discoveries based on DNA barcoding and morphology. *Sci. Rep.* **6**, 37066 DOI: 10.1038/srep37066.
32. Wheeler WC, Coddington JA, Crowley LM, Dimitrov D, Goloboff PA, Griswold CE, Hormiga G, Prendini L, Ramírez MJ, Sierwald P, Almeida-Silva L, Alvarez-Padilla F, Arnedo MA, Benavides Silva LR, Benjamin SP, Bond JE, Grismado CJ, Hasan E, Hedin M, Izquierdo MA, Labarque FM, Ledford J, Lopardo L, Maddison WP, Miller JA, Piacentini LN, Platnick NI, Polotow D, Silva-Dávila D, Scharff N, Szűts T, Ubick D, Vink CJ, Wood HM, Zhang J. (2017). The spider tree of life: phylogeny of Araneae based on target-gene analyses from an extensive taxon sampling. *Cladistics* **33**, 574–616 DOI: 10.1111/cla.12182.
33. Feng C, Miller JA, Lin Y, Shu Y. (2019). Further study of two chinese cave spiders (Araneae, mysmenidae), with description of a new genus. *Zookeys* **870**, 77–100 DOI: 10.3897/zookeys.870.35971.
34. Forster RR, Platnick NI. (1977). A review of the spider family Symphytognathidae (Arachnida, Araneae). *Am. museum Novit.*
35. Mammola S, Michalik P, Hebets EA, Isaia M. (2017). Record breaking achievements by spiders and the scientists who study them. *PeerJ* **5**, e3972 DOI: 10.7717/peerj.3972.