



Universiteit
Leiden
The Netherlands

Investigations on the role of impaired lysosomes of macrophages in disease

Lienden, M.J.C. van der

Citation

Lienden, M. J. C. van der. (2021, March 18). *Investigations on the role of impaired lysosomes of macrophages in disease*. Retrieved from <https://hdl.handle.net/1887/3152425>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3152425>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3152425> holds various files of this Leiden University dissertation.

Author: Lienden, M.J.C. van der

Title: Investigations on the role of impaired lysosomes of macrophages in disease

Issue Date: 2021-03-18

Chapter 3

Localization of active endogenous and exogenous GBA by correlative light-electron microscopy in human fibroblasts

Manuscript published as:

Van Meel, E., Bos, E., **van der Lienden, M. J. C.**, Overkleeft, H. S., van Kasteren, S. I., Koster, A. J. & Aerts, J. M. F. G. Localization of active endogenous and exogenous β -glucocerebrosidase by correlative light-electron microscopy in human fibroblasts. *Traffic* **20**, 346–356 (2019).

Abstract

β -Glucocerebrosidase (GBA) is the enzyme that degrades glucosylceramide in lysosomes. Defects in GBA that result in overall loss of enzymatic activity give rise to the lysosomal storage disorder Gaucher disease, which is characterized by the accumulation of glucosylceramide in tissue macrophages. Gaucher disease is currently treated by infusion of mannose receptor-targeted recombinant GBA. The recombinant GBA is thought to reach the lysosomes of macrophages, based on the impressive clinical response that is observed in Gaucher patients (type 1) receiving this enzyme replacement therapy. In this study, we used cyclophellitol-derived activity-based probes with a fluorescent reporter that irreversibly bind to the catalytic pocket of GBA, to visualize the active enzymes in a correlative microscopy approach. The uptake of pre-labeled recombinant enzyme was monitored by fluorescence and electron microscopy in human fibroblasts that stably expressed the mannose receptor. The endogenous active enzyme was simultaneously visualized by *in situ* labeling with the activity-based probe containing an orthogonal fluorophore. This method revealed the efficient delivery of recombinant GBA to lysosomal target compartments that contained endogenous active enzyme.

Introduction

The lysosomal acid β -glucosidase, β -glucocerebrosidase (GBA, EC 3.2.1.45), is essential to the catabolism of glycosphingolipids in lysosomes as it removes the β -D-glucopyranose group from glucosylceramide. The enzyme encoded by the *GBA* gene is synthesized as a 497 amino acid protein in the endoplasmic reticulum. While the majority of the lysosomal enzymes are modified with the mannose 6-phosphate recognition marker to mediate their transport from the biosynthetic pathway to the lysosomes, GBA does not contain phosphomannosyl residues.^{1,3} Instead, GBA is routed via a mannose 6-phosphate independent targeting pathway, by binding to the lysosomal membrane protein, LIMP II.⁴ LIMP II interacts with newly synthesized GBA at the site of the endoplasmic reticulum. After passage through the Golgi complex, the LIMP II-GBA complex is directed to endosomes and lysosomes, likely through a dileucine-based sorting motif in its C-terminal cytosolic tail.^{5,6} The importance of LIMP II is evident as individuals with LIMP II deficiency develop progressive myoclonic epilepsy with glomerulosclerosis and neurological manifestations, named action myoclonus-renal failure syndrome (AMRF).^{7,8} In AMRF, mutations in the *SCARB2* gene encoding LIMP II cause the failure of normally synthesized GBA to interact with LIMP II. As a result, GBA is secreted from the cells, leading to reduced lysosomal levels of the enzyme in various cell types.^{4,7,9,10}

Mutations in *GBA* result in a prominent loss of GBA in lysosomes and cause the autosomal recessive lysosomal storage disorder Gaucher disease.¹¹ Gaucher disease is characterized by the accumulation of the substrate glucosylceramide in tissue macrophages.¹² The clinical manifestations of the disease are remarkably variable, but usually include enlargement of the liver and spleen, infiltration of the bone marrow by storage macrophages, thrombocytopenia, anemia and bone disease, and can include neurological symptoms. Currently approved treatments are substrate reduction therapy aiming to reduce substrate buildup in macrophages and enzyme replacement therapy (ERT), in which human recombinant GBA (hrGBA) is administered intravenously. To mediate uptake by macrophages, hrGBA is modified to expose its mannose moieties, which bind to the mannose receptor (Man-R) present at the surface of these cells. Endocytosis of the Man-R delivers hrGBA to the endo-lysosomal system.

The development of activity-based probes (ABPs) that covalently and irreversibly tag GBA with high sensitivity and almost complete selectivity has allowed the ultra-sensitive visualization of active GBA molecules *in vitro* and *in vivo* in cells and organisms.¹³ These cyclophellitol-derived ABPs react with the catalytic nucleophile Glu340 of GBA to form an enzyme-substrate complex, linked through an ester bond that is stable under native physiological conditions. When applied to the cell culture medium, the ABPs rapidly enter the cells, a process that appears independent of endocytosis¹³, and bind to the intracellular pools of GBA. The presence of a fluorescent reporter allows the detection by light microscopy.¹³ As fluorescence microscopy does not provide information about the underlying ultrastructure, in this study a correlative light and electron microscopy (CLEM) approach was employed to allow the more detailed localization of active GBA. Moreover, the cellular fate of hrGBA following binding to the Man-R was investigated. The uptake of ABP-prelabeled hrGBA was monitored while simultaneously visualizing the endogenous enzyme by using ABPs with different fluorescent reporters. Light microscopy showed co-localization of endocytosed and endogenous GBA, but also distinct punctae

for either endocytosed or endogenous enzyme. By CLEM, the endogenous GBA was localized to lysosomes, where it showed substantial overlap with endocytosed hrGBA. Of note, some lysosomes did not appear to be reached by hrGBA during the course of the experiment. In addition, the endocytosed enzyme was detected in earlier compartments of the endo-lysosomal system that showed no detectable endogenous GBA. This method will be valuable in determining the efficiency of ERT for Gaucher disease and potentially other lysosomal storage diseases.

Results

GBA was readily detected in normal human dermal fibroblasts (NHDFs) by fluorescence microscopy upon *in situ* labeling with the ABP, MDW941 (**Figure 1**), consistent with previous observations by our lab.¹³ Under these conditions, approximately 50% of total GBA was labeled, as determined by GBA activity assays. Pre-incubation with conduritol B epoxide (CBE) for 16h prior to the *in situ* labeling with MDW941, which should block the active site of all available GBA molecules¹⁴, resulted in no detectable signal (**Supplemental figure 1**), neither was any signal detected upon DMSO or CBE incubation in the absence of probe (**Supplemental figure 1**). These data indicate that the fluorescent signal does not represent unbound, endocytosed probe and is specific to active GBA. Importantly, *in situ* labeled GBA showed clear overlap with total GBA as detected by confocal immunofluorescence microscopy after staining with antibodies to GBA (**Figure 1**, upper panels, **Supplemental figure 2**). Significant overlap was also detected with the late endosomal/lysosomal marker LAMP-1, although not all LAMP-1-containing compartments appeared positive for GBA (**Figure 1**, lower panels, **Supplemental figure 2**). Loading of the late compartments of the endo-lysosomal system with the endocytic tracer dextran-FITC (3h pulse, 6h chase), resulted in substantial co-localization with MDW941, although not all MDW941 positive punctae contained dextran (**Supplemental figure 3**).




Figure 1: Active GBA accumulates in late endosomes and lysosomes. *In situ* labeling of active GBA in NHDFs with 5 nM MDW941 for 2h shows a punctate staining pattern (red) by confocal fluorescence microscopy. Significant overlap is observed with total GBA (see arrows in insets upper panels, green) and the late endosomal/lysosomal marker LAMP-1 (see arrows in insets lower panels, green) upon immunostaining with specific antibodies. The arrowheads show LAMP-1 positive compartments that contain no detectable levels of MDW941. Nuclei were stained with DAPI (blue). Scale bars, 25 μ m.

To identify the endo-lysosomal compartments that accumulated active GBA, a correlative microscopy approach was applied. First, GBA was *in situ* labeled as in the previous experiments, after which the cells were fixed, gelatin embedded and prepared for electron microscopy according to the Tokuyasu technique.¹⁵ The ultrathin cryosections on grids were first mounted on glass slides with 50% glycerol to allow their imaging by confocal microscopy. This revealed the presence of numerous fluorescent punctae per cell profile (**Figure 2A**). Subsequently, the sections were contrasted and embedded with uranyl acetate/methylcellulose and imaged with an electron microscope. Overlay of the fluorescent images and electron micrographs showed that most of the fluorescent signal localized to membrane-bound compartments (**Figure 2B, C**), while the CBE-pre-treated and DMSO-treated sections showed no fluorescent signal. The membrane-bound compartments mainly represented lysosomes, as identified by the presence of the characteristic internal membrane sheets and their electron dense appearance (**Figure 2D, E**). The earlier compartments of the endo-lysosomal system, the endosomes, characterized by their more electron lucent appearance and intraluminal vesicles and the absence of internal membrane sheets, occasionally showed low levels of active GBA, but in most cases the signal was below the detection limit (**Figure 2D, E**). These data indicate that the majority of the active GBA molecules in fibroblasts reside in the lysosomes.




Figure 2: Correlative microscopy shows the accumulation of active GBA in lysosomes of NHDFs. (A) Confocal fluorescence image of an ultrathin cryosection of NHDFs in situ labeled with MDW941 for 2h at 5 nM. The red punctae represent active GBA (MDW941), while the blue dots represent fluorescent electron dense beads used for correlation of the images. Scale bar, 10 μ m. Merged fluorescence and electron microscopy images of boxed areas 1 and 2 are shown in (B) and (C), respectively. Scale bars, 3 μ m. (D, E). Magnifications of boxed areas in (B) and (C), respectively. MDW941 (red), which labels active GBA, is present in lysosomes. E, endosome; L, lysosome; M, mitochondrion; PM, plasma membrane; *, bead used for correlation. The white arrowheads indicate lysosomes positive for MDW941, recognized by the internal membrane sheets, while the black arrowheads indicate endosomes negative for the ABP. Scale bars, D, 250 nm; E, 500 nm.

To allow the efficient internalization of hrGBA by human fibroblasts, cell lines were generated that stably express the mannose receptor (Man-R) with a C-terminal V5 tag. In this way, the uptake of hrGBA by the modified fibroblasts mimics the mode of uptake by tissue macrophages, which express Man-R at their surface. Fibroblasts that were obtained from a patient homozygous for the nonsense mutation p.W178X⁷ contained no detectable levels of LIMP II by Western blotting (Figure 3A). As a result, the intracellular level of

GBA was significantly reduced. In control fibroblasts, three bands were detected for GBA by Western blotting (**Figure 3A**), i.e. the complex glycosylated form, the ER form and the late endosomal/lysosomal form.¹⁶ In the LIMP II^{-/-} cells low levels of the ER form were detected (**Figure 3A**) and the intracellular GBA activity was reduced to approximately $5 \pm 0.9\%$ of the level in control fibroblasts (**Figure 3B**). These findings are consistent with the absence of the lysosomal targeting receptor LIMP II, which results in the increased secretion of GBA by these cells. Lentiviral transduction of both NHDFs and LIMP II^{-/-} fibroblasts with *MRC1* resulted in the expression of the Man-R as determined by anti-V5 immunoblotting (**Figure 3A**) and immunofluorescence microscopy (**Supplemental figure 4**), with the LIMP II^{-/-} cells containing slightly higher expression levels of Man-R than the NHDFs.




Figure 3: Man-R expressing fibroblasts endocytose increased amounts of MDW933-labeled hrGBA. A. NHDFs and LIMP II^{-/-} fibroblast lysates with or without Man-R with a V5-tag were subjected to SDS-PAGE and immunoblotting. No detectable levels of LIMP II are present in the LIMP

*II^{-/-} cells, while the levels of GBA are significantly reduced (lane 3 and 4). The Man-R is expressed in the control and LIMP II^{-/-} fibroblast lines (lane 2 and 4), as detected with an antibody to the V5 tag. GAPDH shows equal protein loading. B. GBA activity in the various cell lines as determined with 4-methylumbelliferyl-coupled specific substrate. The values are averages of 3 independent experiments $\pm$ standard deviations, with the activity in NHDFs set to 100%. The activity in LIMP II^{-/-} fibroblast lines is ~5% of NHDFs, while the Man-R expressing NHDFs show a somewhat increased activity. * $p < 0.01$. C. Representative wet slab gel of at least 3 independent experiments shows the increased uptake of hrGBA by Man-R expressing NHDFs and LIMP II^{-/-} fibroblasts, as compared to the Man-R negative fibroblasts. The addition of mannan reduces the hrGBA uptake to the same levels in all four cell lines, likely representing the uptake by non-specific, fluid-phase endocytosis. 1% of total MDW933-hrGBA that was added to the culture medium (input) was loaded for comparison. β -tubulin shows equal protein loading.*

Subsequently, hrGBA was *in vitro* labeled with MDW933 (an ABP functionalized with a green fluorescent BODIPY dye ¹³) and, after the removal of unreacted ABP (**Supplemental figure 5**), added to the culture medium to follow its uptake in the various cell lines. A 6h-incubation with MDW933-hrGBA resulted in the enhanced internalization of hrGBA by fibroblasts that expressed the Man-R, both NHDFs and LIMP II^{-/-}, as compared to those that did not express the receptor (**Figure 3C**). Somewhat higher levels of hrGBA were observed in the LIMP II^{-/-} cells as compared to the NHDFs, which is likely due to the higher levels of Man-R present in the former, resulting in increased uptake. Competition with mannan, to block the Man-R, reduced the uptake to the same level in all cell lines. This level likely reflects the uptake by fluid-phase endocytosis. The effect of mannan that was observed in NHDFs and LIMP II^{-/-} fibroblasts that did not overexpress the Man-R might be caused by the block of another lectin responsible for hrGBA internalization, as these cells are unlikely to express the Man-R endogenously.

To visualize the delivery of hrGBA to the intracellular target compartments and find out whether it reaches the lysosomes that contain active GBA, MDW933-labeled hrGBA was applied to the culture medium of the four different cell lines, while endogenous GBA was labeled simultaneously with MDW941. Analysis by SDS-PAGE showed labeling of the endogenous enzyme with the red probe in NHDFs, identified as two bands representing the complex glycosylated and lysosomal forms of GBA (**Figure 4**). In LIMP II^{-/-} fibroblasts the GBA levels were in most instances too low to be detected by this method (**Figure 4**). Importantly, uptake of MDW933-labeled hrGBA did not result in significant cross-labeling of the endogenous enzyme due to unreacted MDW933, as only a single band was labeled with MDW933, representing hrGBA. In addition, the signal for MDW941 had a similar intensity in the lanes with or without MDW933-hrGBA. Likewise, no cross-labeling of internalized hrGBA with MDW941 appeared to occur.




Figure 4. Simultaneous uptake of ABP-labeled hrGBA and labeling of the active endogenous enzyme. The four different fibroblast cell lines were allowed to take up 2 $\mu\text{g/mL}$ MDW933-hrGBA for 6h, while endogenous GBA was in situ labeled during the final 2h with MDW941. A representative wet slab gel shows endocytosed hrGBA (upper panel, scanned green channel MDW933) and labeled endogenous GBA (middle panel, scanned red channel MDW941). The lower panel shows the overlay of the two channels. While endogenous GBA is readily detected in NHDFs, the signal is too low in LIMP II^{-/-} fibroblasts. β -tubulin shows equal protein loading.

Subsequently, the same experimental set-up was used in confocal fluorescence microscopy. The Man-R expressing cells represented a heterogeneous population, as they expressed varying levels of receptor. While the signal for MDW933-labeled hrGBA was below the detection level in all cells without Man-R, a punctate staining pattern was observed in NHDFs that expressed sufficient levels of Man-R (**Figure 5**, upper panels). These punctae significantly overlapped with endogenous GBA as visualized with MDW941, suggesting the delivery of hrGBA to lysosomes. However, some punctae were mainly positive for either hrGBA or endogenous GBA (see insets **Figure 5**, **Supplemental figure 2**). A similar result was observed in NHDFs that were immunostained for LAMP-1 after the 6h-uptake with hrGBA (**Supplemental figure 6**). There was a significant overlap of hrGBA and LAMP-1, in addition to LAMP-1-positive compartments that did not contain detectable levels of hrGBA. Importantly, no signal for MDW933-hrGBA could be visualized after uptake in the presence of mannan, which further suggests that no cross-labeling of endogenous GBA occurred with MDW933. As expected, endogenous GBA failed to be detected in the LIMP II^{-/-} fibroblasts (**Figure 5**, middle panels). However,

uptake of MDW933-labeled hrGBA resulted in a punctate pattern that showed partial overlap with LAMP-1 (**Figure 5**, lower panels, see insets, **Supplemental figure 2**). These data suggest that hrGBA is delivered to late endosomes/lysosomes in the absence of LIMP II.




Figure 5: Endocytosed hrGBA partially co-localizes with active GBA and LAMP-1. Confocal fluorescence microscopy of NHDFs and LIMP II^{-/-} fibroblasts expressing the Man-R after a 6h uptake of 2 μg/mL MDW933-hrGBA (green) and 2h in situ labeling of endogenous GBA (red) with 5 nM MDW941 (upper and middle panels) or immunofluorescent staining of LAMP-1 (red, lower panels; see also Supplemental figure 4 for LAMP-1 staining in NHDFs). Substantial co-localization was observed in NHDFs between endocytosed hrGBA and endogenous GBA (see arrows in inset upper panels). In LIMP II^{-/-} fibroblasts, the endocytosed enzyme reached a portion of the LAMP-1 positive compartments (arrows in inset lower panels). The arrowheads indicate compartments positive for either endocytosed hrGBA or endogenous GBA/LAMP-1. Scale bars, 10 μm.

With CLEM the nature of the various fluorescent punctae could be identified. As the uptake of labeled hrGBA was continuous for 6h, we anticipated that the hrGBA-containing punctae would represent both early and late compartments of the endo-lysosomal system. Consistent with this prediction, hrGBA was present in endosomes, as well as in lysosomes

(Figure 6A-D). Significant overlap was observed between endocytosed and endogenous GBA in lysosomes (Figure 6E). Interestingly, some lysosomes appeared to be negative for endocytosed hrGBA (Figure 6E). In agreement with the previous data, no endogenous GBA was detected by CLEM in the LIMP II^{-/-} fibroblasts. However, endocytosed hrGBA was delivered to the endosomes and lysosomes (Figure 7), suggesting that the absence of LIMP II does not have a major effect on the lysosomal delivery of GBA upon endocytosis.




Figure 6: hrGBA is delivered to GBA-containing lysosomes in NHDFs as determined by CLEM. (A-E). Overlay of electron micrograph and confocal fluorescence image of an ultrathin cryosection of NHDFs expressing the Man-R. The cells were allowed to take up MDW933-labeled hrGBA (green) for 6h, while endogenous GBA was labeled by adding MDW941 to the culture medium at a concentration of 5 nM for 2h (red). Nuclei were stained with DAPI (blue). Scale bar in A, 1 μ m. (B-D). Magnification of boxed area 1 from A. E, endosome; L, lysosome. The arrow indicates an endosome that contains endocytosed hrGBA (green). The lysosomes contain both endocytosed hrGBA and endogenous enzyme (red). (E). Magnification of boxed area 2 from A showing lysosomes with both endocytosed hrGBA (green) and endogenous GBA (red). One lysosome (arrow) contains mainly the endogenous enzyme. M, mitochondrion; PM, plasma membrane. Scale bar, 500 nm.

Discussion

In this study we have presented a correlative microscopy technique that combines the use of ABPs, for the detection of active lysosomal enzyme by fluorescence microscopy, with electron microscopy. This allows the visualization of the positive structures at high resolution. The cyclophellitol-derived ABPs that irreversibly bind to the active site of the lysosomal β -glucosidase GBA, readily penetrate through the cellular membranes, which provides the great benefit that they can be applied to living cells.¹³ In our approach, *in situ*

labeling of human fibroblasts with ABP, equipped with a fluorescent tag, was followed by fixation and sample preparation according to the Tokuyasu technique.^{15,17} The ultrathin cryosections were directly imaged for their fluorescent signal by confocal microscopy, without the need of on-section labeling reactions that could increase background signals or influence the sensitivity of detection. The subsequent imaging by electron microscopy provides a relatively straightforward technique to localize the active enzyme to distinct subcellular compartments.




Figure 7: hrGBA is delivered to lysosomes in LIMP II^{-/-} fibroblasts as determined by CLEM. (A, B). Overlay of electron micrograph and confocal fluorescent image of an ultrathin cryosection showing LIMP II^{-/-} fibroblasts + Man-R, after a 6h-uptake of MDW933-labeled hrGBA (green) and in situ labeling of endogenous GBA (red, not detected). Nuclei were stained with DAPI (blue). Both endosomes (E) and lysosomes (L) are positive for hrGBA, however, not all of them are reached (arrows). M, mitochondrion; *, bead used for correlation. Scale bars, 500 nm.

This correlative approach to localize GBA resulted in the detection of active enzyme mainly in lysosomes. By confocal fluorescence microscopy, the ABP co-localized well with our antibody to GBA and was blocked by pre-incubation with the inhibitor CBE, showing the specificity of GBA labeling. In addition, the ABP co-localized significantly with the lysosomal membrane protein LAMP-1, which is present in late endosomes and lysosomes.^{18,19} By CLEM we were able to determine that the highest levels of active GBA were present in lysosomes, based on the morphological characteristics of these compartments, while endosomes appeared negative. Thus the LAMP-1-containing, ABP-negative compartments most likely represent late endosomes with GBA levels below the detection limit.

Subsequently, we aimed to follow the uptake of recombinant GBA and relate it to the localization of the endogenous enzyme using two-color CLEM. To this end, hrGBA was *in vitro* labeled with ABP containing a green fluorophore, while the endogenous pool of GBA was *in situ* labeled with a red ABP. By confocal fluorescence microscopy we observed a significant overlap of green and red fluorescent signals, while there were also punctae positive for either fluorophore. Correlative microscopy showed that the overlapping green/red signals localized to lysosomes, while the green dots represented early and late endosomes containing hrGBA and the red dots lysosomes that had not been reached by endocytosed hrGBA. The observation that many, but not all lysosomes were reached by endocytosed hrGBA could be a technicality due to the imaging of sections instead of intact cells, or other factors such as the duration of uptake. Alternatively, an interesting possibility could be that there are different populations of lysosomes, some more and others less accessible by endocytosis.

Altogether, the imaging of GBA with ABPs in this correlative microscopy approach resulted in the localization of the enzyme to specific subcellular compartments, without the need for additional labeling.^{20,21} This method could be applied to model organisms as well, as these ABPs can be delivered to various tissues upon intravenous or intracerebroventricular injection in rodents.^{13,22} Moreover, correction of lysosomal enzyme deficiency by ERT is presently pursued for several other lysosomal storage disorders, such as Fabry disease and Pompe disease, which are caused by deficiency of lysosomal α -galactosidase A (GLA) and lysosomal α -glucosidase (GAA), respectively.²³⁻²⁵ As ABPs have already been designed for GLA and GAA^{26,27}, this method might be applied to analyze the targeting of these expensive drugs. Finally, mutations in *GBA* have recently been recognized as a risk factor for multiple myeloma and carriers of one mutant *GBA* allele are more likely to develop Parkinsonism and Lewy body dementia.^{28,29} Extending our knowledge of this enzyme could be instrumental to the understanding of these, and possibly other, pathologies.

Material and Methods

Cell lines

Normal human dermal fibroblasts (NHDFs) were obtained from Lonza and LIMP II^{-/-} fibroblasts from a patient homozygous for the mutation c.533G>A, resulting in an early stop codon at W178.⁷ All cell lines were maintained in Dulbecco's Modified Eagle's Medium with 4.5 g/L glucose, sodium pyruvate and sodium bicarbonate (D6546, Sigma), supplemented with 10% (v/v) fetal bovine serum, 100,000 units/L penicillin, 100 mg/L streptomycin and 2 mM glutamax (Thermo Fisher Scientific Inc.) at 37°C in a 5% CO₂ humidified incubator.

NHDFs or LIMP II^{-/-} fibroblasts stably expressing the mannose receptor (Man-R) were generated by lentiviral transduction. Human spleen mRNA was used to amplify the human *MRC1* (*MRC1*, NM_002438.3) coding sequence by PCR using the following oligonucleotides: sense 5'-GGG GAC AAG TTT GTA CAA AAA AGC AGG CTT CGG TAC CAC CAT GAG GCT ACC CCT GC -3' and antisense 5'-GGG GAC CAC TTT GTA CAA GAA AGC TGG GTC GAT GAC CGA GTG TTC ATT CTG -3'. The fragment was cloned into pDNOR-221, and subsequently sub-cloned into pLenti6.3/TO/V5-DEST using the Gateway system (Invitrogen), generating a C-terminal V5 fusion protein. Due to toxicity of the construct in *E. coli* CopyCutter™ EPI400™ chemically competent *E. coli* cells were used according to the manufacturer's protocol (Epicentre). The full sequence was verified by DNA sequencing.

To produce lentiviral particles, HEK293T cells in 6-well plates were transfected at ~70% confluency with the envelope and packaging plasmids pMD2.G, pRSV, pMDL-RRE and pLenti6.3-MRC1 at a 1:1 ratio, 3 µg DNA in total, and 9 µg polyethylenimine (PEI 25K, Polysciences Inc.). The culture medium was replaced 16h after transfection with culture medium containing 20 mM HEPES and collected twice after a 24h incubation period. The culture medium containing viral particles was passed over a 0.45 µm filter and applied to the control and LIMP II^{-/-} fibroblast lines for 8-24h. Transduced cells were selected for by incubation with 2.5 µg/mL blasticidin (Sigma) containing culture medium for several weeks.

In situ labeling of GBA and fluorescence microscopy

Cells were grown to ~70% confluency on glass coverslips in 12-well plates. GBA was *in situ* labeled by adding MDW941 ('red', BODIPY 573/612) to the culture medium at a concentration of 5 nM. The cells were incubated for 2h in a humidified incubator at 37°C with 5% CO₂. Pre-incubation with CBE (Enzo Life Sciences Inc.) was performed at a concentration of 0.3 mM for 16h. Unlabeled cells were incubated with similar percentages (v/v) of DMSO. Uptake with FITC-CM-dextran (MW 70,000, Sigma) was performed at a concentration of 1 mg/mL for 3h (pulse). Subsequently, the cells were washed 5 times with PBS and incubated in culture medium for 6h (chase). *In situ* labeling with 5 nM MDW941 was performed during the last 2h of the chase.

At the end of the incubations, the cells were washed 3 times with PBS and fixed with 4% (w/v) formaldehyde (Sigma)/PBS for 30 minutes at RT. After the fixation, the cells were washed with PBS and distilled H₂O, after which they were either stained for immunofluorescence microscopy or mounted directly on a microscope slide with ProLong Diamond antifade reagent containing DAPI (Molecular Probes). Immunofluorescence

staining was performed as described.³⁰ Antibodies used were mouse anti-GBA monoclonal (8E4, generated in the Aerts lab) at dilution 1:500, rabbit anti-LAMP-1 (Abcam) at dilution 1:400, mouse anti-V5 (Invitrogen) at dilution 1:250 and Alexa Fluor conjugated IgGs (H+L) donkey anti-mouse Alexa 488 and donkey anti-rabbit Alexa 488 (Invitrogen). The cells were imaged under a Leica TCS SP8 confocal microscope with a 63x/1.40 NA HC Plan Apo CS2 oil immersion objective and equipped with a hybrid detector (HyD).

In vitro labeling of hrGBA with MDW933

hrGBA (Cerezyme) was a kind gift from Genzyme (Boston, MA). 5 μ L hrGBA at a concentration of 1 μ g/ μ L was incubated for 16h at 37°C with 40 μ L of 150 mM McIlvaine buffer pH=5.2, 0.2% (w/v) sodium taurocholate (Calbiochem), 0.1% (v/v) Triton X-100 and 2 μ M MDW933 ('green', BODIPY 493/503), while protected from the light. After the incubation, an aliquot was kept to confirm that the residual hrGBA activity was reduced >99% (see below). The remainder was passed over a protein desalting spin column, MWCO 7K (Thermo Scientific) according to the manufacturer's protocol, to remove any unreacted probe (see also **Supplemental figure 3**).

GBA activity assay

The enzymatic activity of hrGBA preparations or GBA in cell lysates was measured by incubating 10-20 μ L sample with 100 μ L substrate mix, which consisted of 3.75 mM 4-methylumbelliferyl β -D-glucopyranoside (Glycosynth Limited), 0.2% (w/v) sodium taurocholate (Calbiochem), 0.1% (w/v) BSA, 0.1% (v/v) Triton X-100 in 150 mM McIlvaine buffer pH=5.2, in a 96-well plate at 37°C. The reactions were stopped with 200 μ L 1 M glycine-NaOH pH=10.3 and read in an LS55 fluorescence spectrometer (PerkinElmer) at λ_{ex} 366 nm, λ_{em} 445 nm.

Uptake of MDW933-labeled hrGBA and in situ labeling of endogenous GBA

All uptake experiments were performed with 2 μ g/mL MDW933-labeled hrGBA for 6h. When indicated, MDW941 was added during the last 2h of the incubation at a concentration of 5 nM. To block uptake of hrGBA by the Man-R, mannan from *Saccharomyces cerevisiae* (Sigma) was added simultaneously at 3 mg/mL. At the end of the incubations, the cells were washed 3 times with PBS and either lysed (for the GBA activity assay/SDS-PAGE) or fixed for microscopy.

SDS-PAGE and Western blotting

Cells were lysed by the addition of ice-cold lysis buffer, i.e. 25 mM potassium phosphate buffer pH=6.5, 0.1% Triton X-100 and a protease inhibitor cocktail (cOmplete, Roche Diagnostics GmbH). Total protein concentration in the lysates was determined by the BCA protein assay (Pierce BCA protein assay kit, Thermo Fisher Scientific Inc.) in a 96-well plate format, according to the manufacturer's protocol. The samples were read in an EMax Plus microplate reader (Molecular Devices). Equal protein amounts (5 μ g) were resolved on 10% polyacrylamide gels and transferred onto 0.2 μ m nitrocellulose membranes (Amersham, GE Healthcare). Antibodies used were rabbit anti-LIMP II (Novus Biologicals), mouse anti-GBA (8E4, generated in the Aerts lab), rabbit anti-GAPDH (Cell Signaling), mouse anti-V5 (Invitrogen) and rabbit anti- β -tubulin (Cell

Signaling) and secondary antibodies used were donkey anti-rabbit Alexa Fluor 647 IgG (H+L) (Invitrogen), goat anti-mouse IgG (H+L)-HRP conjugate and goat anti-rabbit IgG (H+L)-HRP conjugate (BioRad). HRP-conjugated antibodies were visualized by chemiluminescence (Pierce ECL Plus Western blotting substrate, Thermo Fisher Scientific Inc.). The wet slab gels and immunoblots were scanned on a Typhoon FLA 9500 scanner (GE Healthcare).

Correlative microscopy

NHDFs were grown in 100 mm culture dishes to ~70% confluency. GBA was *in situ* labeled with 5 nM MDW941 for 2h at 37°C, either with or without pre-incubation with 0.3 mM CBE for 16h as a control for the specificity of the signal. To check for autofluorescence, one dish was incubated for 2h with a similar percentage (v/v) of DMSO. The uptake experiments were performed with NHDFs + Man-R and LIMP II^{-/-} fibroblasts + Man-R, with 2 µg/mL MDW933-labeled hrGBA for 6h. During the last 2h endogenous GBA was *in situ* labeled with 5 nM MDW941. As a control for the Man-R-specific uptake of MDW933-hrGBA, the uptake in NHDFs + Man-R, followed by the *in situ* labeling was performed in the presence of 3 mg/mL mannan.

At the end of the incubations, the culture medium was removed and the cells were washed 3 times quickly with culture medium at 37°C. Subsequently, the cells were fixed with 2% (w/v) formaldehyde + 0.2% glutaraldehyde (Electron Microscopy Sciences) in 0.1 M PHEM buffer for 2h at RT, while protected from the light. Embedding of the samples and cryosectioning was performed as described.¹⁵ For correlation purposes, fluospheresTM carboxylate-modified microspheres blue (350/440) with a diameter of 100 nm (Thermo Fisher Scientific Inc.) were added to the grids, which were coated with a formvar film, either before or after collection of the sections by incubating the grids for 2 minutes on the beads (1:2000 dilution of 2% solids). Subsequently, the sections were incubated on 2% gelatin for 30 minutes at 37°C and on 15 mM glycine/PBS for 10 minutes at RT. The nuclei were stained with 50 ng/mL DAPI (Sigma)/PBS for 5 minutes. After a quick wash with distilled H₂O, the ultrathin cryosections on titanium grids (Agar Scientific) were mounted on microscope glass slides in 50% glycerol/H₂O and imaged under a Leica TCS SP8 confocal microscope with a 63x/1.40 NA HC Plan Apo CS2 oil immersion objective. Immediately after imaging, the glycerol was removed from the grids by washing in distilled H₂O. The sections were contrasted with 1.8% methylcellulose/0.6% uranyl acetate/H₂O at pH=4.8 and analyzed in a FEI Tecnai 12 BioTwin electron microscope at 120 kV. Correlation of the images was performed using Adobe Photoshop CC 2015.5.

Labeling specificity in correlative microscopy

The specificity of the green and red fluorescent signals for hrGBA and endogenous GBA, respectively, was determined in the following way. First of all, the *in vitro* labeling reaction was optimized to label over 99% of hrGBA, which was confirmed by an enzymatic activity assay to result in near absence of hrGBA activity. In this way, cross-labeling of hrGBA with red ABP was prevented during the *in situ* labeling. Consistent herewith, no red ABP signal was detected in fibroblasts that lack GBA, due to a mutation in the receptor LIMP II, upon endocytosis of labeled hrGBA. Secondly, any unreacted green ABP was removed from the preparation before addition to the culture medium. SDS-PAGE analysis showed substantial reduction of unreacted ABP. Any remaining unreacted probe was not sufficient

to result in detectable cross-labeling of endogenous GBA as determined by SDS-PAGE of the cell lysates. Finally, the uptake of ABP-labeled hrGBA in the presence of mannan did not result in a detectable green fluorescent signal in the cells by microscopy, indicating that the endogenous enzyme was not labeled with green probe.


References

1. Aerts, J. M., Schram, A. W., Strijland, A., van Weely, S., Jonsson, L. M., Tager, J. M., Sorrell, S. H., Ginns, E. I., Barranger, J. A., Murray, G. J. Glucocerebrosidase, a lysosomal enzyme that does not undergo oligosaccharide phosphorylation. *Biochim Biophys. Acta.*, **964**, 303-308 (1988).
2. Rijnboutt, S., Aerts, H. M., Geuze, H. J., Tager, J. M., Strous, G. J. Mannose 6-phosphate-independent membrane association of cathepsin D, glucocerebrosidase, and sphingolipid-activating protein in HepG2 cells. *J. Biol. Chem.*, **266**, 4862-4868 (1991).
3. Rijnboutt, S., Kal, A. J., Geuze, H. J., Aerts, H., Strous, G. J. Mannose 6-phosphate-independent targeting of cathepsin D to lysosomes in HepG2 cells. *J. Biol. Chem.*, **266**, 23586-23592 (1991).
4. Reczek, D., Schwake, M., Schroder, J., Hughes, H., Blanz, J., Jin, X., Brondyk, W., Van Patten, S., Edmunds, T., Saftig, P. LIMP-2 is a receptor for lysosomal mannose-6-phosphate-independent targeting of beta-glucocerebrosidase. *Cell*, **131**, 770-783 (2007).
5. Ogata, S., Fukuda, M. Lysosomal targeting of Limp II membrane glycoprotein requires a novel Leu-Ile motif at a particular position in its cytoplasmic tail. *J. Biol. Chem.*, **269**, 5210-5217 (1994).
6. Sandoval, I. V., Arredondo, J. J., Alcalde, J., Gonzalez Noriega, A., Vandekerckhove, J., Jimenez, M. A., Rico, M. The residues Leu(Ile)₄₇₅-Ile(Leu, Val, Ala)₄₇₆, contained in the extended carboxyl cytoplasmic tail, are critical for targeting of the resident lysosomal membrane protein LIMP II to lysosomes. *J. Biol. Chem.*, **269**, 6622-6631 (1994).
7. Balreira, A., Gaspar, P., Caiola, D., Chaves, J., Beirao, I., Lima, J. L., Azevedo, J. E., Miranda, M. C. A nonsense mutation in the LIMP-2 gene associated with progressive myoclonic epilepsy and nephrotic syndrome. *Hum. Mol. Genet.*, **17**, 2238-2243 (2008).
8. Berkovic, S. F., Dibbens, L. M., Oshlack, A., Silver, J. D., Katerelos, M., Vears, D. F., Lullmann-Rauch, R., Blanz, J., Zhang, K. W., Stankovich, J., Kalnins, R. M., Dowling, J. P., Andermann, E., Andermann, F., Faldini, E., D'Hooge, R., Vadlamudi, L., Macdonell, R. A., Hodgson, B. L., Bayly, M. A., Savige, J., Mulley, J. C., Smyth, G. K., Power, D. A., Saftig, P., Bahlo, M. Array-based gene discovery with three unrelated subjects shows SCARB2/LIMP-2 deficiency causes myoclonus epilepsy and glomerulosclerosis. *Am. J. Hum. Genet.*, **82**, 673-684 (2008).
9. Gaspar, P., Kallemeijn, W. W., Strijland, A., Scheij, S., Van Eijk, M., Aten, J., Overkleeft, H. S., Balreira, A., Zunke, F., Schwake, M., Sa Miranda, C., Aerts, J. M. Action myoclonus-renal failure syndrome: diagnostic applications of activity-based probes and lipid analysis. *J. Lipid Res.*, **55**, 138-145 (2014).
10. Malini, E., Zampieri, S., Deganuto, M., Romanello, M., Sechi, A., Bembi, B., Dardis, A. Role of LIMP-2 in the intracellular trafficking of beta-glucosidase in different human cellular models. *FASEB J.*, **29**, 3839-3852 (2015).
11. Brady, R. O., Kanfer, J. N., Bradley, R. M., Shapiro, D. Demonstration of a deficiency of glucocerebrosidase-cleaving enzyme in Gaucher's disease. *J. Clin. Invest.*, **45**, 1112-1115 (1966).
12. Grabowski, G. A., Petsko, G. A., Kolodny, E. H. Gaucher disease. In: D. Valle, A. L. Beaudet, B. Vogelstein, K. W. Kinzler, S. E. Antonarakis, A. Ballabio, K. M. Gibson, and G. Mitchell (Eds.), *The Online Metabolic and Molecular Bases of Inherited Disease*. New York: McGraw-Hill (2014).
13. Witte, M. D., Kallemeijn, W. W., Aten, J., Li, K. Y., Strijland, A., Donker-Koopman, W. E., van den Nieuwendijk, A. M., Bleijlevens, B., Kramer, G., Florea, B. I., Hooibrink, B., Hollak, C. E., Ottenhoff, R., Boot, R. G., van der Marel, G. A., Overkleeft, H. S., Aerts, J.


- M. Ultrasensitive in situ visualization of active glucocerebrosidase molecules. *Nat. Chem. Biol.*, **6**, 907-913 (2010).
14. Daniels, L. B., Glew, R. H., Radin, N. S., Vunnam, R. R. A revised fluorometric assay for Gaucher's disease using conduritol-beta-epoxide with liver as the source of Beta-glucosidase. *Clin. Chim Acta.*, **106**, 155-163 (1980).
15. Slot, J. W., Geuze, H. J. Cryosectioning and immunolabeling. *Nat Protoc.*, **2**, 2480-2491 (2007).
16. Van Weely, S., Aerts, J. M., Van Leeuwen, M. B., Heikoop, J. C., Donker-Koopman, W. E., Barranger, J. A., Tager, J. M., Schram, A. W. Function of oligosaccharide modification in glucocerebrosidase, a membrane-associated lysosomal hydrolase. *Eur. J. Biochem.*, **191**, 669-677 (1990).
17. Tokuyasu, K. T. A technique for ultracryotomy of cell suspensions and tissues. *J. Cell. Biol.*, **57**, 551-565 (1973).
18. Barriocanal, J. G., Bonifacino, J. S., Yuan, L., Sandoval, I. V. Biosynthesis, glycosylation, movement through the Golgi system, and transport to lysosomes by an N-linked carbohydrate-independent mechanism of three lysosomal integral membrane proteins. *J. Biol. Chem.*, **261**, 16755-16763 (1986).
19. Marsh, M., Schmid, S., Kern, H., Harms, E., Male, P., Mellman, I., Helenius, A. Rapid analytical and preparative isolation of functional endosomes by free flow electrophoresis. *J. Cell. Biol.*, **104**, 875-886 (1987).
20. Ngo, J. T., Adams, S. R., Deerinck, T. J., Boassa, D., Rodriguez-Rivera, F., Palida, S. F., Bertozzi, C. R., Ellisman, M. H., Tsien, R. Y. Click-EM for imaging metabolically tagged nonprotein biomolecules. *Nat. Chem. Biol.*, **12**, 459-465 (2016).
21. van Elsland, D. M., Bos, E., de Boer, W., Overkleeft, H. S., Koster, A. J., van Kasteren, S. I. Detection of bioorthogonal groups by correlative light and electron microscopy allows imaging of degraded bacteria in phagocytes. *Chem Sci*, **7**, 752-758 (2016).
22. Herrera Moro Chao, D., Kallemeijn, W. W., Marques, A. R., Orre, M., Ottenhoff, R., van Roomen, C., Foppen, E., Renner, M. C., Moeton, M., van Eijk, M., Boot, R. G., Kamphuis, W., Hol, E. M., Aten, J., Overkleeft, H. S., Kalsbeek, A., Aerts, J. M. Visualization of Active Glucocerebrosidase in Rodent Brain with High Spatial Resolution following In Situ Labeling with Fluorescent Activity Based Probes. *PLoS One*, **10**, e0138107 (2015).
23. Eng, C. M., Guffon, N., Wilcox, W. R., Germain, D. P., Lee, P., Waldek, S., Caplan, L., Linthorst, G. E., Desnick, R. J., International Collaborative Fabry Disease Study Group Safety and efficacy of recombinant human alpha-galactosidase A replacement therapy in Fabry's disease. *N. Engl. J. Med.*, **345**, 9-16 (2001).
24. Schiffmann, R., Kopp, J. B., Austin, H. A., 3rd, Sabnis, S., Moore, D. F., Weibel, T., Balow, J. E., Brady, R. O. Enzyme replacement therapy in Fabry disease: a randomized controlled trial. *JAMA*, **285**, 2743-2749 (2001).
25. Van den Hout, J. M., Reuser, A. J., de Klerk, J. B., Arts, W. F., Smeitink, J. A., Van der Ploeg, A. T. Enzyme therapy for pompe disease with recombinant human alpha-glucosidase from rabbit milk. *J. Inherit Metab. Dis.*, **24**, 266-274 (2001).
26. Jiang, J., Kuo, C. L., Wu, L., Franke, C., Kallemeijn, W. W., Florea, B. I., van Meel, E., van der Marel, G. A., Codee, J. D., Boot, R. G., Davies, G. J., Overkleeft, H. S., Aerts, J. M. Detection of Active Mammalian GH31 alpha-Glucosidases in Health and Disease Using In-Class, Broad-Spectrum Activity-Based Probes. *ACS Cent. Sci.*, **2**, 351-358 (2016).
27. Willems, L. I., Beenakker, T. J., Murray, B., Scheij, S., Kallemeijn, W. W., Boot, R. G., Verhoek, M., Donker-Koopman, W. E., Ferraz, M. J., van Rijssel, E. R., Florea, B. I., Codee,

- J. D., van der Marel, G. A., Aerts, J. M., Overkleeft, H. S. Potent and selective activity-based probes for GH27 human retaining alpha-galactosidases. *J. Am. Chem. Soc.*, **136**, 11622-11625 (2014).
28. Aflaki, E., Westbroek, W., Sidransky, E. The Complicated Relationship between Gaucher Disease and Parkinsonism: Insights from a Rare Disease. *Neuron*, **93**, 737-746 (2017).
29. Nair, S., Branagan, A. R., Liu, J., Boddupalli, C. S., Mistry, P. K., Dhodapkar, M. V. Clonal Immunoglobulin against Lysolipids in the Origin of Myeloma. *N. Engl. J. Med.*, **374**, 555-561 (2016).
30. Qian, Y., Flanagan-Steet, H., van Meel, E., Steet, R., Kornfeld, S. A. The DMAP interaction domain of UDP-GlcNAc:lysosomal enzyme N-acetylglucosamine-1-phosphotransferase is a substrate recognition module. *Proc. Natl. Acad. Sci. USA.*, **110**, 10246-10251 (2013).


Supplemental Figures




Supplemental figure 1: In situ labeling of NHDFs with MDW941 specifically visualizes active GBA. While in situ labeling of active GBA with MDW941 shows a punctate staining pattern in NHDFs (left upper panel), pre-incubation with CBE followed by in situ labeling with MDW941 results in loss of the fluorescent signal (right upper panel). Incubation with DMSO or CBE shows no detectable autofluorescence in lysosomes with the same imaging settings (lower panels). Nuclei were stained with DAPI. Scale bars, 10 μm.




Supplemental figure 2: 2D intensity histograms of confocal fluorescence microscopy images. The 2D intensity histograms of the merged fluorescence microscopy images from figure 1 (A) and figure 5 (B) are shown. The histograms were generated with the ImageJ plugin 'Coloc 2', using Costes threshold regression and 30 Costes randomizations.




Supplemental figure 3: MDW941 reaches lysosomes marked with the endocytic tracer dextran. Confocal fluorescence images of NHDFs after uptake of FITC-dextran (green) and in situ labeling of endogenous GBA with 5 nM MDW941 (red) for 2h. The 3h-uptake with 1 mg/mL dextran-FITC was followed by a 6h-chase to mark lysosomes. The arrows show examples of co-localization of dextran with MDW941. Not all MDW941 positive compartments were reached by dextran. Nuclei were stained with DAPI (blue). The 2D intensity histogram of the merged image is shown. Scale bar, 10 μ m.



Supplemental figure 4: Expression of the Man-R is detected by immunofluorescence microscopy. NHDFs and LIMP II^{-/-} fibroblasts + or - Man-R were stained with anti-V5 antibodies (green) to detect the presence of the Man-R with a C-terminal V5 tag. As the cells were permeabilized, the signal is detected in intracellular compartments by confocal fluorescence microscopy. Nuclei were stained with DAPI. Note that no signal is detected in the Man-R negative cells. Scale bars, 25 μ m.



Supplemental figure 5. In vitro labeling of hrGBA with MDW933 and removal of unbound ABP. Representative wet slab gel showing equivalent amounts (lanes '2' contain the double amounts of lanes '1') of MDW933 used in the in vitro reaction (lane 1 and 2), MDW933-labeled hrGBA at the end of the reaction with residual free ABP (lane 3 and 4) and MDW933-labeled GBA after passage over a protein desalting column to remove unreacted MDW933 (lane 5 and 6). Note that MDW933 runs ahead of the dye front (indicated with the arrowhead) and is greatly reduced in lanes 5 and 6.



Supplemental figure 6: Endocytosed hrGBA partially co-localizes with the late endosomal/lysosomal marker LAMP-1 in NHDFs. Confocal fluorescence images of NHDFs expressing the Man-R, after a 6h uptake of MDW933-hrGBA (green) and immunostaining of LAMP-1 (red). The arrows in the insets show co-localization of endocytosed GBA and LAMP-1, while the arrowheads show punctae that are positive for either MDW933-hrGBA or LAMP-1. Nuclei were stained with DAPI (blue). The 2D intensity histogram of the merged image is shown. Scale bar, 10 μm .