



Universiteit  
Leiden  
The Netherlands

## Targeting glycolysis in endothelial cells to prevent intraplaque neovascularization and atherogenesis in mice

Perrotta, P.

### Citation

Perrotta, P. (2021, March 24). *Targeting glycolysis in endothelial cells to prevent intraplaque neovascularization and atherogenesis in mice*. Retrieved from <https://hdl.handle.net/1887/3152433>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3152433>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3152433> holds various files of this Leiden University dissertation.

**Author:** Perrotta, P.

**Title:** Targeting glycolysis in endothelial cells to prevent intraplaque neovascularization and atherogenesis in mice

**Issue Date:** 2021-03-24

# Chapter 3

---

## *Animal models of atherosclerosis*

---

Emini Veseli B, **Perrotta P**, De Meyer GRA, Roth L, Van der Donckt C, Martinet W, De Meyer GRY.

*Eur J Pharmacol.* 2017;816:3-13

## Abstract

An ideal animal model of atherosclerosis resembles human anatomy and pathophysiology and has the potential to be used in medical and pharmaceutical research to obtain results that can be extrapolated to human medicine. Moreover, it must be easy to acquire, can be maintained at a reasonable cost, is easy to handle and shares the topography of the lesions with humans. In general, animal models of atherosclerosis are based on accelerated plaque formation due to a cholesterol-rich/Western-type diet, manipulation of genes involved in the cholesterol metabolism, and the introduction of additional risk factors for atherosclerosis. Mouse and rabbit models have been mostly used, followed by pigs and non-human primates. Each of these models has its advantages and limitations. The mouse has become the predominant species to study experimental atherosclerosis because of its rapid reproduction, ease of genetic manipulation and its ability to monitor atherogenesis in a reasonable time frame. Both Apolipoprotein E deficient (ApoE<sup>-/-</sup>) and LDL-receptor (LDLr<sup>-/-</sup>) knockout mice have been frequently used, but also ApoE/LDLr double-knockout, ApoE3-Leiden and PCSK9-AAV mice are valuable tools in atherosclerosis research. However, a great challenge was the development of a model in which intraplaque microvessels, haemorrhages, spontaneous atherosclerotic plaque ruptures, myocardial infarction and sudden death occur consistently. These features are present in ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup> mice, which can be used as a validated model in pre-clinical studies to evaluate novel plaque-stabilizing drugs.

**Keywords:** animal models, atherosclerosis, plaque rupture, ApoE, LDL receptor, PCSK9

## Introduction

Atherosclerosis is a progressive inflammatory disease characterized by accumulation of lipids in the arterial vessel wall, which starts early in life. Disease progression leads to build-up of atherosclerotic plaques that cause narrowing of the arterial lumen. Atherosclerotic plaques often remain stable for years, but can rapidly become unstable, rupture and trigger thrombus formation. Accordingly, in addition to restriction of the vessel lumen, the presence of atherosclerotic plaques is linked to an increased risk of acute cardiovascular events such as myocardial infarction (MI) and stroke. The use of animal models of atherosclerosis is an essential tool to improve the understanding of the molecular mechanisms behind atherosclerotic plaque formation and progression, as well as the occurrence of plaque rupture and its associated cardiovascular events. Moreover, animal models allow to assess novel pharmacological treatments that can prevent or slow down the onset of atherosclerosis. In general, animal models for atherosclerosis are based on accelerated plaque formation due to: (1) a cholesterol-rich/Western-type diet, (2) manipulation of genes involved in the cholesterol metabolism, and (3) the introduction of additional risk factors for atherosclerosis, such as diabetes.

In this review, we will discuss the animal models that have contributed to the understanding of atherosclerosis and its clinical consequences, and that allow significant improvement in treatment.

Numerous studies have shown that high plasma levels of low-density lipoprotein (LDL) represent one of the most prominent risk factors of atherosclerosis. Indeed, LDL tends to accumulate in the sub-endothelial space of the arterial wall and progressively undergoes oxidative modifications to form oxidized LDL (oxLDL). This induces an inflammatory response characterized by overexpression of chemotactic proteins such as monocyte chemoattractant protein-1 (MCP-1), and adhesion molecules (vascular cell adhesion molecule-1 (VCAM-1), E-selectin and P-selectin) by endothelial cells.<sup>1,2</sup> Adhesion molecules promote the infiltration of blood-carried monocytes into the inflamed arterial wall. After differentiation into macrophages, these cells engulf oxLDL, transform into foam cells and contribute to plaque development by secreting multiple mediators of the inflammatory process in the vessel wall.<sup>3</sup> The inflammatory response also promotes recruitment of circulating

monocytes and T-cells that stimulate the migration of vascular smooth muscle cells (SMCs) from the tunica media into the sub-endothelial space where they exhibit abnormally high proliferation and secrete extracellular matrix proteins that also contribute to atheroma growth.<sup>1</sup> Advanced human plaques are characterized by a large necrotic core, many lipid laden and activated macrophages, few SMCs, intraplaque neovascularisation and haemorrhages, and a thin fibrous cap that separates the plaque from the blood stream. Rupture of the fibrous cap of such high-risk vulnerable plaques leads to luminal thrombosis, arterial occlusion or embolism in distant vascular beds, resulting in MI, stroke or sudden death.<sup>4</sup> In humans, atherosclerotic plaques can usually be found in the aorta, coronary arteries and in the carotid and cerebral arteries.<sup>5</sup>

In 1908, Ignatowski investigated for the first time plaque formation in the aortic wall of rabbits that were fed a cholesterol-rich diet.<sup>6</sup> Since then many other animal species such as mice, birds, pigs and non-human primates have been used as an experimental model of atherosclerosis.<sup>1, 7</sup>

An ideal animal model resembles human anatomy and pathophysiology, and has the potential to be used in medical and pharmaceutical research to obtain results that can be extrapolated to human medicine. Moreover, it is important that animals used as models are easy to acquire, can be maintained at a reasonable cost, are easy to handle and have well-defined genetic characteristics. A valuable animal model for atherosclerosis research not only shares the crucial aspects of the disease process with humans but also the topography of the lesions. In addition, the animals preferably develop lesions in a spontaneous manner after consumption of a diet similar as in humans.<sup>8</sup> Although several animals develop atherosclerotic plaques after a cholesterol-rich diet, the topography of the lesions is not always similar as compared to humans. Furthermore, it is important to note that in the majority of atherosclerosis models, animals do not spontaneously develop the complications seen in humans such as plaque rupture, MI, stroke and sudden death.

### **Mouse models of atherosclerosis**

Over the past decades, the mouse has become the predominant species to study experimental atherosclerosis because of its rapid reproduction, ease of genetic

manipulation and its ability to monitor atherogenesis in a reasonable time frame.<sup>8-11</sup> However, mice are relatively resistant to the development of atherosclerosis due to their significantly different lipid profile as compared to humans. Therefore, genetic manipulation of their lipid metabolism is mandatory.<sup>8, 12</sup> In mice, most of the cholesterol is transported in high density lipoprotein (HDL) like particles. Accordingly, mice contain only low concentrations of the atherogenic LDL and very low density lipoprotein (VLDL). Mice deficient in the receptor clearing these LDL particles (LDLR<sup>-/-</sup> mice) develop significantly higher plasma levels of cholesterol. Apolipoprotein E (ApoE) is a glycoprotein synthesized mainly in the liver and the brain and functions as a ligand for receptors that clear chylomicrons and VLDL remnants.<sup>12</sup> Deficiency in this glycoprotein (ApoE<sup>-/-</sup>) leads to increased plasma levels of total cholesterol, mostly in the VLDL and chylomicron fractions,<sup>13</sup> which are quadrupled by a high-fat or Western-type diet.<sup>14</sup> Both mouse models have extensively been used to study the mechanisms underlying the initiation and progression of atherosclerosis. Atherosclerotic lesions in mice develop in regions of the vasculature subjected to low and/or oscillatory shear stress.<sup>9, 10</sup> Predilection sites in the mouse are the aortic root, lesser curvature of the aortic arch and branch points of the brachiocephalic, left carotid and subclavian arteries. However, on a high-cholesterol diet, ApoE<sup>-/-</sup> mice develop plaques more rapidly and with a more advanced phenotype as compared to LDLR<sup>-/-</sup> mice<sup>15</sup>, making the ApoE<sup>-/-</sup> model widely used in experimental atherosclerosis studies.

The Apolipoprotein (Apo) E3-Leiden mutation is associated with a genetic form of hyperlipidaemia. Therefore, ApoE3-Leiden transgenic mice can also be used as a model for atherosclerosis, but in comparison with ApoE<sup>-/-</sup> and LDLR<sup>-/-</sup> mice, they show rather low levels of total plasma cholesterol and triglycerides when fed a normal diet. Nevertheless, these mice are highly responsive to fat-, sugar-, and cholesterol-containing diets resulting in strongly elevated lipoprotein profiles.<sup>16</sup> Regardless of lesion development, varying from fatty streaks to mild, moderate, and severe plaques, ApoE3-Leiden mice lack the critical events such as plaque rupture, thrombus formation, and/or haemorrhage, which are of major importance in human atherosclerosis.<sup>17, 18</sup>

High plasma levels of lipoprotein (a) [Lp(a)], which is a complex of LDL and a large glycoprotein called Apolipoprotein (a) [Apo(a)], is an independent risk factor for the

development of atherosclerosis in humans.<sup>19, 20</sup> Virtually all species other than primates lack Apo(a), hampering the use of convenient animal models to study its role in atherosclerotic plaque development. Therefore, transgenic mice that express human Apo(a) are used. When fed a Western-type diet, these mice show the presence of macrophage-like cells in combination with the development of fatty-streak lesions at the base of the aorta.<sup>21</sup> In humans, plasma Apo(a) is almost entirely covalently bound to LDL, whereas in mice, Apo(a) circulates as non-lipoprotein associated Apo(a).<sup>21</sup> Therefore, Apo(a) transgenic mice can be used to identify the role of Apo(a) in atherogenesis, independent of human LDL.

The most commonly used mouse models of atherosclerosis are described in detail below and in Figure 1.

#### *Apolipoprotein E deficient (ApoE<sup>-/-</sup>) mice*

ApoE is a glycoprotein with a molecular size of approximately 34 kDa. It is synthesized mainly in the liver and brain, and is a structural component of all lipoprotein particles except low-density lipoproteins. It serves as a ligand for cell-surface lipoprotein receptors whose function is to clear chylomicrons and VLDL remnants. It is also synthesized by monocytes and macrophages.<sup>22</sup> Other functions include cholesterol homeostasis, local redistribution of cholesterol within tissues, immunoregulation and dietary absorption and biliary excretion of cholesterol.<sup>23, 24</sup>

In 1992, the first line of ApoE<sup>-/-</sup> mice was developed almost contemporaneously in two laboratories.<sup>13, 14</sup> The deletion of the ApoE gene was done in mouse embryonic stem cells by homologous recombination. ApoE<sup>-/-</sup> mice were healthy, had a similar body weight as wild-type mice, and were born at the expected frequency.<sup>14, 25</sup> However, their lipoprotein profile disclosed significant differences with the wild-type mates. The ability of ApoE<sup>-/-</sup> mice to clear plasma lipoproteins is severely impaired resulting in plasma cholesterol levels of 400-600 mg/dl when fed a normal diet, whereas wild-type mice have levels of 75-110 mg/dl.<sup>26, 27</sup> This drastic change is due to an increase in VLDL-sized particles. The development of significant hypercholesterolaemia, even when fed a normal diet, suggests that in the absence of an environmental stimulus, deficiency of ApoE is sufficient to cause massive changes in lipoprotein metabolism. Furthermore, the lack of ApoE boosts the



sensitivity to dietary fat and cholesterol. After several weeks of feeding a Western-type diet (consisting of 21% fat and 0.15% cholesterol, which is similar to the everyday diet of Western countries), plasma cholesterol levels double in wild-type mice, whereas in ApoE-deficient mice a fourfold increase in total plasma cholesterol is observed.<sup>14</sup> Extensive atherosclerosis is seen in mice on both types of diet by 2 to 3 months of age.<sup>28</sup> On the other hand, heterozygous ApoE-deficient mice do not show an increase in plasma cholesterol levels even when fed a Western-type diet, presuming that a 50% decrease in ApoE is not sufficient to increase plasma lipids. Of note, plasma cholesterol levels in mice are not affected by age or sex of the animal.<sup>26</sup>

The entire spectrum of atherosclerotic lesions is present in ApoE<sup>-/-</sup> mice.<sup>25</sup> Monocyte attachment to endothelial cells is noticed from 6 weeks of age, and after 8 weeks foam cell lesion development is detectable. After 15 to 20 weeks, intermediate lesions are present containing mostly SMCs as well as fibrous plaques consisting of SMCs, extracellular matrix and a necrotic core covered with a fibrous cap.<sup>26</sup> In more advanced lesions, fibro-fatty nodules are a nidus for calcification and plaques become more calcified with time.<sup>29</sup> When fed a Western-type diet, the time course for lesion formation is tremendously accelerated.<sup>25</sup> Compared to mice fed a low-fat diet, lesions are 3-4 times larger within the same period of time. This response implies a diet-dependent mechanism, i.e. increased fat leads to increased plasma cholesterol, which in turn leads to increased atherosclerosis, which resembles the diet-dependency of atherosclerotic heart disease observed in humans.<sup>14</sup>

ApoE<sup>-/-</sup> mice tend to develop atherosclerotic plaques at vascular branch points, with predilection for the aortic root, the lesser curvature of the aortic arch, the principal branches of the aorta as well as the pulmonary and carotid arteries.<sup>26</sup> Sequential events of plaque formation in ApoE<sup>-/-</sup> mice are considerably similar to those in well-established larger animal models of atherosclerosis and in humans.<sup>26</sup> Although this mouse model is used by many research groups, it has some limitations. For instance, ApoE is a multifunctional protein that has an impact on inflammation, oxidation, reverse cholesterol transport by macrophages, and smooth muscle proliferation and migration. These functions might affect atherosclerotic plaque development in ApoE<sup>-/-</sup> mice, independent of plasma lipid levels.<sup>30</sup> Furthermore, not

LDL, which is characteristic of human atherosclerosis, but VLDL is the most abundant lipoprotein in ApoE<sup>-/-</sup> mice.<sup>14</sup> However, the major limitation of the 'classical' mouse models of atherosclerosis is the rarity of plaque rupture and thrombosis,<sup>31,32</sup> whereas these events are fairly common in humans and can lead to MI and stroke.<sup>25</sup> It has been suggested that this might be due to the tiny diameter of the mouse vessels; as the vessel diameter decreases, the surface tension increases exponentially, impeding the likelihood of plaque rupture.<sup>25</sup> However, also other explanations have to be taken into account as discussed below in the section 'mouse models of atherosclerotic plaque rupture'.

A method for the induction of accelerated atherogenesis and plaque rupture is the placement of a perivascular collar or cuff, mainly in ApoE<sup>-/-</sup> mice. In their study, Sasaki et al., claim that cuff placement around the left carotid artery results in an animal model of plaque rupture. By using the ligation technique to induce neo-intimal hyperplasia, they observed lipid- and collagen-rich lesions accompanied with intraplaque haemorrhage and plaque rupture. Furthermore, a decrease in collagen content, and formation of fibrinogen-positive thrombi were detected, analogous to plaque rupture in humans.<sup>33</sup> Along with this observation, perivascular carotid collar placement also reproduces the induction of rapid and site-controlled atherosclerosis<sup>34</sup>, while maintaining the structural integrity of the endothelium. Formed plaques are located primarily in the area proximal to the collar. The advantages of this model over the conventional animal models of mechanically induced atherosclerosis include the closer resemblance to human plaque morphology and endothelial expression pattern.<sup>34</sup>

#### *LDL receptor-deficient (LDLR<sup>-/-</sup>) mice*

The LDL receptor is a membrane receptor with a molecular weight of 160 kDa, which mediates the endocytosis of cholesterol-rich LDL and thus maintains the plasma level of LDL. It also facilitates the cellular uptake of apolipoprotein B- and E-containing lipoproteins. LDL receptor deficiency along with mutations in the gene encoding for the LDL receptor count for the phenotypic events described in familial hypercholesterolaemia.<sup>35,36</sup> Mice with a targeted inactivation of the LDL receptor were created in 1993.<sup>37,38</sup> Compared with wild-type, LDLR<sup>-/-</sup> mice display modestly

elevated plasma cholesterol levels and develop no or only mild atherosclerosis when fed a normal diet.<sup>38</sup> In terms of lipoprotein particles, the increase is higher among IDL and LDL sized particles, whereas HDL and triglycerides remain unaffected.<sup>37, 38</sup> It is worth to note that this is different from ApoE<sup>-/-</sup> mice, in which cholesterol is primarily accumulated in large lipoprotein particles such as chylomicron remnants, VLDL and IDL particles (*vide supra*).<sup>14, 39</sup> The response to high-fat/high cholesterol Western-type diets shows a remarkable change in lipoprotein profile of these mice with a high probability for atherosclerotic lesion development.

The plaques that develop in LDLr<sup>-/-</sup> mice are generally the same as those seen in ApoE<sup>-/-</sup> mice.<sup>40</sup> A Western-type diet induces larger and more advanced lesions with a collagen-rich fibrous cap, a necrotic core containing cholesterol clefts and cellular enrichment adjacent to the lumen.<sup>41</sup> The plaque development occurs in a time-dependent manner, initially in the proximal aorta, and spreading toward the distal aorta. Similar to humans, the locations where the blood flow is disturbed are more prone to atherosclerotic lesions.<sup>40</sup> By making LDLr<sup>-/-</sup> and ApoE<sup>-/-</sup> mice homozygous for the ApoB-100 allele, total plasma cholesterol levels of approximately 300 mg/dl were obtained on a normal diet. LDLr<sup>-/-</sup> ApoB<sup>100/100</sup> mice developed more atherosclerotic lesions than the ApoE<sup>-/-</sup> ApoB<sup>100/100</sup> mice, even with a normal diet.<sup>42,</sup>

43

The LDLr<sup>-/-</sup> mouse model has some advantages in comparison with ApoE<sup>-/-</sup> mice. Firstly, plasma cholesterol is mostly carried by LDL particles, which generates a more human-like lipid profile. Secondly, the absence of the LDL receptor does not have an impact on inflammation as compared to ApoE deficiency. Thus, atherosclerotic plaque development in this mouse model is based on elevated plasma lipid levels and not caused by other functions linked to the LDL receptor.<sup>8</sup> Thirdly, the LDLr<sup>-/-</sup> mouse model shares the characteristics observed in human familial hypercholesterolaemia, which is caused by the absence of functional LDL receptors.<sup>44, 45</sup>

#### *ApoE/LDL receptor double-knockout mice*

Introduced shortly after ApoE<sup>-/-</sup> and LDLr<sup>-/-</sup> mice, ApoE/LDL receptor double knock out mice represent a model that develops more severe hyperlipidaemia and

atherosclerosis than the former ones.<sup>46</sup> It is an animal model with spontaneous atherosclerotic plaque development and it has been reported that even on regular chow diet, the progression of atherosclerosis is usually more marked in ApoE/LDL receptor double knock out mice than in mice deficient for ApoE alone.<sup>47</sup> There is no significant difference in the lipoprotein profile of the double knockouts compared to ApoE<sup>-/-</sup> mice, they both have high levels of VLDL and LDL<sup>48</sup>, except the marked elevations in B48 and B100 apolipoproteins.<sup>49</sup> This mouse model is considered suitable to study the anti-atherosclerotic effects of possible treatments, without the need of an atherogenic diet.<sup>25</sup>

### *ApoE3-Leiden mice*

Although ApoE<sup>-/-</sup> mice and LDLr<sup>-/-</sup> mice are the two most frequently used mouse models for atherosclerosis, also ApoE3-Leiden mice are utilized in many studies. Apolipoprotein (Apo)E3-Leiden is associated with a genetic form of hyperlipidaemia and is particularly expressed in a Dutch family. Transgenic mice have been generated using a genomic 27-kilobase DNA construct (containing the ApoE gene, ApoC1 gene and all regulatory elements) isolated from the APOE3-Leiden proband, to study the effect of the ApoE3-Leiden mutation *in vivo*.<sup>17, 50</sup>

Remarkably, although these mice are less susceptible for atherosclerosis than the ApoE deficient mice, they also show dramatically elevated total plasma cholesterol and triglyceride levels when fed a Western-type diet. This is mainly attributed to an increase in VLDL/LDL particles, which demonstrates that ApoE3-Leiden mice have a human-like lipoprotein profile.<sup>51</sup> Another advantage is that ApoE3-Leiden mice have the ability to synthesize functional ApoE. This offers the possibility to study the effect of elevated plasma lipid levels without disturbing inflammatory processes, which is an important limitation of the 'classical' ApoE<sup>-/-</sup> mouse model.<sup>52</sup>

The ApoE3-Leiden mice develop atherosclerotic lesions in the aorta and large vessels when fed a Western-type diet. Lesions are also observed in the proximal coronary arteries, the aortic root, the aortic arch and its main branch points, the thoracic aorta, the abdominal aorta, the renal artery branch points, the abdominal aorta bifurcation, and the iliac artery bifurcations. It is interesting to note that this strain develops early foam cell lesions on normal chow diet. However, more complex

and advanced lesions are observed after 1, 3 and 6 months of Western-type diet feeding.<sup>17</sup>

ApoE3-Leiden mice are used as a model to elucidate factors involved in the metabolism of ApoE and the aetiology of familial dyslipidaemia in particular. Furthermore, ApoE3-Leiden mice are utilized to study complications of venous bypass grafting, a clinical procedure that bypasses an atherosclerotic obstruction in an artery. Similar to humans, the grafted vein in this model undergoes remodelling, which is a consequence of exposure to higher blood pressure and shear stress but also vessel injury due to surgery. This process results in the formation of intimal hyperplasia and accelerated atherosclerosis, which may lead to obstruction of the graft.<sup>17, 53, 54</sup>

Because similarities were found between lesions in vein grafts and native atherosclerosis, a murine model of vein graft disease has been established.<sup>55</sup> In this model, the thoracic caval vein of a donor mouse is grafted in the carotid artery of a receiver mouse. This procedure has been used in mice that are susceptible to atherosclerosis (ApoE<sup>-/-</sup> or ApoE3-Leiden mice) so that vein grafts with accelerated atherosclerosis could be studied.<sup>56-58</sup> It has been shown that vein grafts in these transgenic mice are morphologically similar to rupture-prone plaques in humans. The lesions in this model have the typical characteristics of late stage atherosclerosis, including the presence of foam cells, a large necrotic core, intraplaque neovascularization, calcification and cholesterol clefts.<sup>59</sup>

### *PCSK9-AAV mice*

Besides the abovementioned models, a new line of mouse model without germline genetic engineering is emerging in the research field of atherosclerosis. The so called pro-protein convertase subtilisin/kexin type 9 (PCSK9) - adeno associated virus (AAV) mice were described independently by two research groups in 2014 as a rapid, versatile and cost-effective animal model for atherosclerosis.<sup>60, 61</sup> PCSK9, a newly identified human subtilase, is a serine protease with plasma concentrations of  $\approx 100$  to 200 ng/mL and it is highly expressed in the liver.<sup>62, 63</sup> Several studies have shown that PCSK9 reduces hepatic uptake of LDL by increasing the endosomal and lysosomal degradation of LDL receptors.<sup>64</sup> In brief, after protein

maturation and secretion, circulating PCSK9 binds the LDL receptors on the cell surface and is subsequently co-internalized together with the receptor. This distracts the normal recycling process of the receptor to the plasma membrane and promotes degradation in the lysosome.<sup>63</sup>

Recombinant AAV vectors support long-term transgene expression in many animal models<sup>65-67</sup> and humans.<sup>68</sup> Following single intravenous injection with human D374Y<sup>61</sup> or murine D377Y<sup>60</sup> gain-of-function mutant PCSK9, mice were stably expressing PCSK9<sup>DY</sup> mRNA in the liver. AAV viral infection does not elicit any adverse effects in the animals and no signs of liver damage or immunologic response were observed following infection. At 30 days after injection, total serum cholesterol in PCSK9<sup>DY</sup>-AAV transgenes was doubled compared to control mice. These differences remained the same even after 1 year post-infection, confirming a chronic effect of a single AAV injection.<sup>61</sup> Western-type diet exacerbated hyperlipidaemia in PCSK9<sup>DY</sup>-AAV mice, leading to plasma cholesterol levels of up to 1165 mg/dl, while chow diet fed mice barely reached 316 mg/dl. The lipoprotein profile of Western-type diet fed PCSK9<sup>DY</sup>-AAV mice showed an equal distribution between VLDL and LDL particles.<sup>61</sup> PCSK9<sup>DY</sup> transgenic mice develop atherosclerosis in a dose-dependent manner. Hyperlipidaemia provokes the build-up of lesions throughout the vasculature resembling those of LDLR<sup>-/-</sup> mice, which is exacerbated by HFD feeding.<sup>60, 61</sup> Aortic root lesions show advanced plaque development with foam cells, smooth muscle cells, macrophage infiltration and fibrous tissue, but importantly, lesions progress to the fibro-atheromatous stage.<sup>60, 61</sup> and within the time frame of 15-20 weeks, vascular calcification occurs.<sup>69</sup> When Roche-Molina et al. combined PCSK9<sup>DY</sup> expression and ApoE deficiency, they revealed an expected synergistic effect: the lesions doubled in size with no significant differences in lipoprotein profile as compared to single mutants on the same diet.<sup>61</sup>

Overall, the induction of hyperlipidaemia and atherosclerosis in animals with different genetic backgrounds, the robust stability after single administration of mutant human PCSK9 and the fact that there are no major biosafety concerns in using AAVs as vectors, makes the PCSK9-AAV model a valuable tool in atherosclerosis research.<sup>61</sup>

## Mouse models of atherosclerotic plaque rupture

Despite major advances in cardio- and cerebrovascular research, plaque rupture remains the leading cause of acute events.<sup>70</sup> Therefore, the need for the development of plaque-stabilizing therapies is high. Several research groups have tried to develop suitable models of plaque rupture for the last 15 years but in these models rupture occurs only sporadically, after a long period of time, or depends on mechanical injury.<sup>11,71,72</sup> Moreover, the reproducibility is low and events as seen in humans are rarely observed.

As discussed earlier, atherosclerotic plaques in mice develop in specific sites such as the aortic root, the lesser curvature of the aortic arch and the branch points of the brachiocephalic, the left carotid and the subclavian arteries. However, mice show only minor plaque development in the coronary and carotid arteries, which are the main sites of atherosclerotic plaque development in humans.<sup>8-10</sup> To induce plaque rupture in mice, several approaches based on surgical (such as arterial ligation or the positioning of a cuff around an artery) or genetic manipulation have been proposed (Table 1). Although these models have been useful in understanding the concepts of plaque rupture, none of them exhibit the full combination of the characteristics seen in human vulnerable/ruptured plaques. Moreover, plaque rupture with a superimposed occlusive thrombus, the most common complication of human atherosclerosis, is rarely observed.<sup>73</sup> Consequently, clinical events such as MI or ischemic stroke are almost never seen in these models.<sup>9,70</sup> Furthermore, most of these models do not show 'spontaneous' plaque ruptures. When spontaneous ruptures are observed, they only occur sporadically and after a long period of time. However, recently a model of consistent, spontaneous atherosclerotic plaque ruptures in mice has been described, as discussed below.

### *Apolipoprotein E-deficient Fibrillin-1 mutant (ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup>) mice*

The extracellular matrix is a complex network of predominantly elastin and collagen, which is essential to provide structural, adhesive and biochemical signalling support to the vessel wall. In elastic arteries, elastin is the most abundant protein. The elastic fibres comprise the elastin core, which is surrounded by a mantle of fibrillin-rich

microfibrils. <sup>74</sup>The elastic-fibre-associated microfibrils have as the main structural component fibrillin-1, a large glycoprotein of about 350 kDa, whose major role is in binding and sequestering growth factors, such as transforming growth factor- $\beta$  (TGF- $\beta$ ), as well as providing the scaffold for the deposition and the cross-linking of elastin.

<sup>75, 76</sup>

Recently, we reported the effect of an impaired elastin structure of the vessel wall on the progression of atherosclerosis by cross-breeding ApoE<sup>-/-</sup> mice with mice containing a heterozygous mutation (C1039G<sup>+/-</sup>) in the fibrillin-1 (Fbn1) gene. <sup>76</sup> Mutations in the Fbn1 gene lead to the Marfan syndrome, a genetic disorder characterized by fragmentation of elastic fibres. <sup>77</sup>This results in increased arterial stiffening, elevated pulse pressure and progressive aortic dilatation. <sup>76, 78, 79</sup> Moreover, the mutation leads to the development of highly unstable plaques in ApoE<sup>-/-</sup> mice, resulting in spontaneous plaque rupture with end-points including MI and sudden death. <sup>76, 80</sup> Importantly, these events do not – or only very occasionally – occur in ApoE<sup>-/-</sup> mice on a Western-type diet or in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice fed a normal diet. <sup>80, 81</sup> These findings underscore the importance of elastin fragmentation in combination with a Western-type diet as prerequisites for atherosclerotic plaque rupture in mice.

ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice have significantly larger plaques with a highly unstable phenotype, characterized by a large necrotic core (occupying about 30% of total plaque area), and a strongly diminished collagen content. Accelerated atherogenesis in these mice is likely the result of enhanced vascular inflammation, leading to increased monocyte attraction, oxidation and accumulation of lipids. <sup>82</sup> Inducible nitric oxide synthase (iNOS), a marker for activated macrophages and inflammation, is significantly more expressed in plaques of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on either Western-type diet or normal diet as compared to ApoE<sup>-/-</sup> mice on Western-type diet. Accordingly, inflammatory cytokines tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 $\beta$  (IL-1 $\beta$ ) and interleukin-6 (IL-6) are highly increased. In addition, a higher infiltration of T-cells and their activation marker interferon- $\gamma$  (IFN- $\gamma$ ) is present, the latter playing an important role in collagen turn-over by inhibiting SMCs to synthesise collagen, required to repair and maintain fibrous cap integrity. <sup>83, 84</sup> Moreover, matrix metalloproteinase (MMP)-2, -9, -12 and -13 expression or activity



is increased in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice. MMP-2 and MMP-9 are implicated in both atherosclerosis and angiogenesis. <sup>85</sup> For example, ApoE<sup>-/-</sup> mice lacking MMP-2 develop smaller and more stable plaques, whereas macrophages overexpressing active MMP-9 promote neovascularisation, intraplaque haemorrhage <sup>86, 87</sup> and features of plaque rupture in ApoE<sup>-/-</sup> mice. <sup>87</sup> In the latter case, those features were attributed to elastin degradation, underscoring its role in plaque destabilisation and rupture. MMP-12 and MMP-13 additionally contribute to elastin and (type I) collagen degradation, respectively. Taken together, in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet enhanced collagen/extracellular matrix breakdown together with decreased synthesis and repair are likely responsible for weakening of the fibrous cap and rendering it more rupture-prone. <sup>84, 85</sup>













Extensive neovascularisation and intraplaque haemorrhages consistently occur in the brachiocephalic and common carotid arteries of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet. These features are rarely seen in murine atherosclerosis models but are known to highly affect plaque progression and vulnerability in humans. <sup>88</sup> In ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a Western-type diet, intraplaque neo-vessels, likely arising from adventitial vasa vasorum, clearly sprout out of the media. <sup>89, 90</sup> Neo-vessels are not only present at the base of the plaque but are also frequently observed in its centre, similar to human pathology. <sup>88, 90</sup> Angiogenesis requires extracellular matrix degradation by proteases, including MMPs, to enable endothelial cell migration into the surrounding tissue. <sup>85</sup> In addition, degradation of the extracellular matrix induces release of sequestered angiogenic factors such as vascular endothelial growth factor (VEGF) and TGF- $\beta$  <sup>85, 86</sup>, also observed in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet. The extent of neovascularisation in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice correlates with the degree of elastin fragmentation in the vessel wall. However, degradation of the extracellular matrix alone is not sufficient to induce neovascularisation in atherosclerotic plaques, because microvessels are not present in plaques of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a normal diet. This observation indicates that an additional factor is needed to trigger plaque neovascularisation. Hypoxia, a well-known angiogenesis trigger <sup>91</sup>, is strongly increased in plaques of brachiocephalic and carotid arteries in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet. By contrast, hypoxia in the ascending aorta is minor, which likely explains the

absence of neo-vessels at that site. Thus, the highly permeable arterial wall, due to degradation of the extracellular matrix, combined with intraplaque hypoxia seems required for neo-vessel formation in atherosclerotic plaques of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet. Importantly, those neo-vessels are highly leaky. Moreover, the presence of intraplaque erythrocytes near neo-vessels at the base of the plaque points to intraplaque haemorrhages, substantiating ruptured neo-vessels as source of intraplaque bleeding.<sup>86, 88, 92</sup> Erythrocytes are important sources of free cholesterol, thereby increasing necrotic core size. Hence, neovascularisation, besides supplying plaques with leukocytes and lipoproteins, can promote focal plaque expansion when microvessels rupture or become thrombotic.<sup>88, 91, 92</sup> Taken together, these observations in this mouse model are in line with current concepts of human vulnerable plaques.

In addition to enhanced plaque vulnerability, plaque rupture is consistently present in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a Western-type diet, but only very rarely in ApoE<sup>-/-</sup> mice on a Western-type diet. Moreover, fibrin-rich mural thrombi are present in brachiocephalic, carotid and coronary arteries and ascending aortas. Both intrinsic (i.e. a highly unstable plaque phenotype) and extrinsic factors (i.e. forces acting on the plaque) are elementary for plaque rupture.<sup>93</sup> In general, rupture occurs when the mechanical stress applied on the fibrous cap exceeds its tensile strength. The latter is mainly determined by the collagen content of the plaque, which is significantly decreased in plaques of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet.<sup>84, 93</sup> Elevated pulse pressure (as a consequence of arterial stiffening)<sup>76</sup> leads to repetitive plaque deformation, increasing the tensile stress on the cap.<sup>79, 94</sup> When applied chronically, this can lead to plaque fatigue, making it prone to rupture.<sup>93, 94</sup> Moreover, due to the progressive aortic dilatation and outward remodelling (as a result of the large plaques), the collagen and elastin fibres of the cap are stretched and become more rigid, increasing the susceptibility to mechanical stress. Aortic dilatation is highly pronounced in the ascending aorta of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a Western-type diet, suggesting that this mechanism is responsible for rupture of unstable plaques at this site. In brachiocephalic and carotid arteries, intraplaque neovascularisation and haemorrhage are frequently present, further increasing plaque size and vulnerability to rupture.

In addition, sudden death is observed in ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a Western-type diet, mainly between 16 and 23 weeks, with 50% mortality after 20 weeks. Moreover, ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a Western-type diet that died suddenly show a significantly higher frequency of coronary stenosis compared to survivors, suggesting that the presence of coronary artery plaque plays an important role in cardiac death. The majority of ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on Western-type diet show infarcted areas, which compromise cardiac function even more. Although it is not known whether the increased infarcted area is the result of plaque rupture or due to pronounced plaque formation and coronary artery stenosis, these findings are remarkable because coronary artery plaque and spontaneous MIs almost never develop in ApoE<sup>-/-</sup> mice on a Western-type diet. Also in humans, differences in fibrillin-1 genotype have shown to greatly affect plaque progression and severity of coronary artery disease, underscoring the pathophysiological relevance of fibrillin-1 mutations in cardiovascular disease.<sup>79</sup>

Thus, elastin fragmentation in combination with a Western-type diet leads to plaque destabilisation and rupture in ApoE<sup>-/-</sup> mice. ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice show many features of human end-stage atherosclerosis, such as an enlarged necrotic core, a thin fibrous cap with an important loss of collagen fibres, outward remodelling and the presence of intraplaque microvessels and haemorrhage, resulting in plaque rupture, MI and sudden death. Therefore, ApoE<sup>-/-</sup>Fbn1<sup>C1039G<sup>+/-</sup></sup> mice on a Western-type diet offer the opportunity to investigate the role of key factors involved in plaque destabilisation, including intraplaque neovascularisation, which will provide more insight into the mechanisms of plaque disruption and potential targets for therapeutic interventions.<sup>80, 95-97</sup>

|   | Model  | Lipid profile   | Plaque distribution and characteristics (20 weeks WD)   | Advantages & limitations  |
|---|--|---|---|---|
| ApoE <sup>-/-</sup>                           | Disruption of the ApoE gene<br>   | <b>Plasma cholesterol:</b><br>400-600 mg/dl on ND<br>>1000 mg/dl on WD<br><br><b>Lipoproteins:</b><br>↓ VLDL<br>↓ LDL<br>↓ HDL              | <br><b>Fibrous plaques:</b><br>Smooth muscle cells<br>Extracellular matrix<br>Inflammatory cells<br>Necrotic core  | <ul style="list-style-type: none"> <li>⊕ Develops atherosclerosis on ND</li> <li>⊖ No human-like lipid profile</li> <li>⊖ ApoE plays a role in inflammation → influence plaque development</li> <li>⊖ No spontaneous plaque rupture, thrombosis and complications</li> </ul>                                    |
| LDLr <sup>-/-</sup>                           | Disruption of the LDL receptor gene<br>   | <b>Plasma cholesterol:</b><br>200-300 mg/dl on ND<br>>1000 mg/dl on WD<br><br><b>Lipoproteins:</b><br>↓ VLDL<br>↓ LDL<br>= HDL              | <br><b>Fibrous plaques:</b><br>Smooth muscle cells<br>Extracellular matrix<br>Inflammatory cells<br>Necrotic core  | <ul style="list-style-type: none"> <li>⊕ Human-like lipid profile (LDL)</li> <li>⊕ Functional ApoE → no impact on inflammation</li> <li>⊖ Complex lesion development requires a WD</li> <li>⊖ No spontaneous plaque rupture, thrombosis and complications</li> </ul>  |
| ApoE <sup>-/-</sup> LDLr <sup>-/-</sup>       | Disruption of the ApoE and the LDL receptor gene<br>                                    | <b>Plasma cholesterol:</b><br>400-600 mg/dl on ND<br>>1000 mg/dl on WD<br><br><b>Lipoproteins:</b><br>↓ VLDL<br>↓ LDL<br>↓ HDL              | <br><b>Fibrous plaques:</b><br>Smooth muscle cells<br>Extracellular matrix<br>Inflammatory cells<br>Necrotic core  | <ul style="list-style-type: none"> <li>⊕ Develops atherosclerosis on ND</li> <li>⊖ No spontaneous plaque rupture, thrombosis and complications</li> </ul>   |
| ApoE3-Leiden                                  | ApoE3-Leiden mutation via DNA construct (ApoE, ApoC1) from the ApoE3-Leiden proband<br> | <b>Plasma cholesterol:</b><br>100-200 mg/dl on ND<br>>1000 mg/dl on WD<br><br><b>Lipoproteins:</b><br>↓ VLDL<br>↓ LDL<br>↓ HDL (only on WD) | <br><b>Fibrous plaques:</b><br>Smooth muscle cells<br>Extracellular matrix<br>Inflammatory cells<br>Necrotic core  | <ul style="list-style-type: none"> <li>⊕ Functional ApoE → no impact on inflammation</li> <li>⊖ Complex lesion development requires a WD</li> <li>⊖ No spontaneous plaque rupture, thrombosis and complications</li> </ul>  |
| PCSK9-AAV                                     | Single Adeno-Associated Virus-mediated gene transfer of mutant PCSK9<br>              | <b>Plasma cholesterol:</b><br>300 mg/dl on ND<br>>1000 mg/dl on WD<br><br><b>Lipoproteins:</b><br>↓ VLDL (only on WD)<br>↓ LDL<br>= HDL     | <br><b>Fibrous plaques:</b><br>Smooth muscle cells<br>Extracellular matrix<br>Inflammatory cells<br>Necrotic core  | <ul style="list-style-type: none"> <li>⊕ No genetic modification needed</li> <li>⊖ Lesion development requires a WD</li> <li>⊖ No spontaneous plaque rupture, thrombosis and complications</li> </ul>   |
| ApoE <sup>-/-</sup> Fbn1 <sup>C1039G+/-</sup> | ApoE <sup>-/-</sup> mice X mice with a mutation in the Fbn1 gene (C1039G)<br>         | <b>Plasma cholesterol:</b><br>400-600 mg/dl on ND<br>>1000 mg/dl on WD<br><br><b>Lipoproteins:</b><br>↓ VLDL<br>↓ LDL<br>↓ HDL              | <br><b>Rupture-prone plaques:</b><br>↓ Smooth muscle cells<br>↓ Extracellular matrix<br>↑ Inflammatory cells<br>↑ Necrotic core<br>Neovascularisation<br>Haemorrhage | <ul style="list-style-type: none"> <li>⊕ Accelerated plaque development</li> <li>⊕ Spontaneous plaque rupture with complications (MI, stroke) and sudden death</li> <li>⊕ Intra-plaque neovascularisation and haemorrhage present</li> <li>⊖ Male mice on WD die prematurely due to aneurysm rupture</li> </ul> |

**Figure 1. Overview of current mouse models of atherosclerosis.** This figure describes the different models with their total plasma cholesterol levels on normal (ND) and Western-type diet (WD), lipoprotein profile, plaque characteristics, advantages and limitations. The distribution of the plaques in the thoracic aorta and the complexity is shown for mice fed a WD for 20 weeks. The composition of the most complex lesions at that time point is displayed (usually found in the aortic root or brachiocephalic artery).

## Rabbits

The rabbit has been one of the most frequently used animals in atherosclerosis research because of their easy handling and relatively inexpensive maintenance.<sup>8</sup> However, there has been a reduced trend of using this animal model since 2000, probably due to the availability of ApoE and LDL receptor knock-out mice.<sup>98</sup> Multiple approaches and models have been used to study atherosclerosis and its complication in rabbits, including genetically hypercholesterolaemic rabbits such as Watanabe heritable hyperlipidaemic rabbits (WHHL)<sup>99</sup>, New Zealand White rabbits fed a cholesterol-rich diet<sup>100</sup>, and very recently ApoE<sup>-/-</sup> rabbits.<sup>101</sup>

Rabbits have a lipoprotein metabolism that is similar to humans (except for their hepatic lipase deficiency) and show significant differences with mice. Unlike mice, in which HDL is the predominant plasma lipoprotein, rabbits transport significant amounts of cholesterol via ApoB-containing particles (VLDL and LDL).<sup>102</sup> Consequently, rabbits have been useful to point out the role of elevated plasma cholesterol as a critical factor in the initiation of atherosclerosis. Limitations of rabbit models include a highly abnormal diet required for the development of hypercholesterolaemia, massive inflammation and hepatic toxicity due to the long term high cholesterol feeding.<sup>1</sup>

### *Watanabe heritable hyperlipidaemic (WHHL) rabbits*

WHHL rabbits are a mutant strain that shows spontaneous hypercholesterolaemia and atherosclerosis due to a defect in the LDL receptor.<sup>99</sup> Homozygous WHHL rabbits fed a normal diet are hypercholesterolaemic from birth with LDL as the predominant lipoprotein. They exhibit various types of atherosclerotic lesions ranging from early fatty streaks to advanced lesions in the aorta, coronary arteries and cerebral artery.<sup>100, 103</sup> These rabbits also show an increased risk of MI. The WHHL rabbit was one of the first rabbit models in which the effect of statins to suppress plaque destabilization and to reduce thrombogenicity was investigated. High-fructose and high fat–diet fed WHHL rabbits develop early insulin resistance and glucose tolerance and show aortic lesions with a lipid core and calcifications.

This model has allowed researchers to investigate the effect of insulin resistance on atherosclerosis lesion formation.<sup>104</sup>

#### *New Zealand White (NZW) rabbits*

NZW rabbits are commonly used to study atherosclerosis. NZW rabbits that are fed a normal diet have low plasma cholesterol levels (mostly < 50 mg/dl) and consequently do not develop spontaneous atherosclerosis. However, supplementing the diet with 0.3-0.5% cholesterol increases the plasma cholesterol level up to 1000 mg/dl. The plaque composition is determined by the level of dietary cholesterol and the duration of cholesterol feeding. One protocol uses adult rabbits that are fed a cholesterol-rich diet (1.0-1.5% cholesterol) for a short period of time (about 8 weeks). By using such a diet, rabbits develop severe hypercholesterolaemia with plasma cholesterol levels between 1500 and 3000 mg/dl, which are never seen in humans, resulting in atherosclerotic plaques primarily composed of macrophage-derived foam cells. For this reason, the most used protocol lasts 20 to 26 weeks and consists of a diet containing 0.3% cholesterol. On average, cholesterol levels rise to about 800 mg/dl. This protocol develops atherosclerotic plaques in the aortic arch and thoracic aorta, rather than in the abdominal aorta (less pronounced plaque formation)<sup>100, 102</sup>, whereas in humans, plaques are commonly found in the abdominal aorta. Coronary atherosclerosis is also observed in cholesterol-fed rabbits but is usually restricted to the left coronary arterial trunks. Depending on the length of the cholesterol feeding, also plaque calcification occurs. However, there is no evidence for spontaneous plaque rupture.

#### *Apolipoprotein E knockout (ApoE<sup>-/-</sup>) rabbits*

Recently, ApoE<sup>-/-</sup> rabbits have been reported as a model to study the relationship between atherosclerosis and human hyperlipidaemia.<sup>101</sup> ApoE<sup>-/-</sup> rabbits can be generated using genome editing enzymes such as zinc finger nucleases, transcription activator-like effector nucleases (TALENs) or RNA-guided CRISPR-associated protein 9 (Cas9) endonucleases. Because the rabbit lipoprotein profile is similar to humans<sup>105</sup>, the ApoE<sup>-/-</sup> rabbit represents an attractive alternative to the

ApoE<sup>-/-</sup> mouse. Even on a normal diet, ApoE<sup>-/-</sup> rabbits show mild hyperlipidaemia with plasma total cholesterol levels around 200 mg/dl. However, when fed a cholesterol-rich diet (0.3% cholesterol and 3% soybean oil for 2 weeks) their plasma total cholesterol levels increase to about 1000 mg/dl (vs. about 170 mg/dl in cholesterol-fed wild-type rabbits). ApoE<sup>-/-</sup> rabbits develop more pronounced aortic atherosclerosis than wild-type rabbits when fed with a cholesterol diet for 10 weeks.<sup>101</sup> Because both ApoE and the LDL receptor play an important role in mediating cholesterol metabolism, ApoE<sup>-/-</sup> rabbits together with LDL receptor-deficient WHHL rabbits may be valuable models for the study of human hyperlipidaemia: ApoE<sup>-/-</sup> rabbits show elevation of remnant lipoproteins, whereas WHHL rabbits have high levels of LDL accompanied by low HDL.<sup>101</sup>

### **Large animal models (pigs and non-human primates)**

Although small animal models have provided insight into the mechanisms that drive atherosclerosis, additional strategies are required to translate these findings into improved prevention and treatment of symptomatic atherosclerosis in humans. Efficient large animal models of atherosclerosis may be useful to deal with these challenges. Indeed, the translation of the knowledge obtained from studies in mice to the development of drugs for human atherosclerosis can benefit from a bridging tool such as porcine models of atherosclerosis. Not only the effects of pharmacological treatments on atherosclerosis can be studied in such models but also clinical imaging end-points can be evaluated as guiding tool for subsequent phase II clinical trials.<sup>106</sup> Gene-editing tools for large animals have made it possible to create gene-modified minipigs that develop atherosclerosis with many similarities to humans in terms of predilection for lesion sites and histopathology. For instance, minipigs with liver-specific expression of human D374Y-PCSK9 show severe hypercholesterolaemia and development of progressive atherosclerotic lesions.<sup>107</sup> Together with existing porcine models of atherosclerosis that are based on spontaneous mutations or severe diabetes, such models may provide new approaches for translational research in atherosclerosis<sup>106</sup>. Non-human primates show hypercholesterolaemia when fed a high fat/high cholesterol diet and develop coronary fibro-fatty atherosclerotic plaques, similar to humans.<sup>8</sup> Yet, working with

monkeys is expensive, highly regulated, and requires very specialized laboratory animal science skills. Therefore, these models are not frequently used. A few years ago, however, also knockout non-human primates have been created, which may reinforce the interest in large animal models with accelerated atherosclerosis.<sup>106, 108</sup>

In conclusion, many efforts have been made to develop animal models that resemble human atherosclerosis as good as possible. However, each of the current animal models has its advantages and limitations, as summarized in Table 2. A great challenge was the development of an animal model of spontaneous (i.e. without mechanical interventions) plaque rupture with human-like endpoints such as MI, stroke and sudden death. These features are present in ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup> mice, making it a promising model to evaluate potential plaque stabilizing therapies.

**Table 1.** Mouse models of atherosclerotic plaque rupture

| Strain  | mechanism                                 | Duration (weeks) | Plaque disruption | Luminal thrombus         | Intraplaque neo-vessels | IPH    | Outward remodelling | 'Human-like' complications     | Comments  | Ref. |
|---|---|------------------|-------------------|--------------------------|-------------------------|--------|---------------------|--------------------------------|---|------|
| ApoE <sup>-/-</sup>                             | 'ageing'                                  | 60               | 12%               | 3%                       | N.D.                    | N.D.   | N.D.                | Coronary thrombosis            | Long term experiment<br>Low rate of plaque rupture and thrombosis | 109  |
| ApoE <sup>-/-</sup>                             | Mixed C57BL/6-129SvJ background           | 30-65            | 52%               | 73%                      | N.D.                    | N.D.   | N.D.                | MI ('some cases') sudden death | Long term experiment mixed background                             | 110  |
| ApoE <sup>-/-</sup>                             | Collar: Ad-p53 in SMC + phenylephrine     | 15-17            | 40%               | 5%                       | N.D.                    | 35%    | N.D.                | N.D.                           | Plaque rupture not spontaneous<br>complicated manipulation        | 111  |
| ApoE <sup>-/-</sup>                             | Active MMP-9 overexpression in Mφ         | 41               | 40%               | Fibrin deposition (100%) | N.D.                    | 90%    | N.D.                | Sudden death (20%)             | Long term experiment, complicated manipulation                    | 87   |
| ApoE <sup>-/-</sup>                             | Cuff (+ ligation)                         | 13-14            | 29-63%            | 17-42%                   | N.D.                    | 31-47% | N.D.                | N.D.                           | Plaque rupture not spontaneous                                    | 33   |
| ApoE <sup>-/-</sup> ; TM <sup>Pro</sup>         | Collar, genetic hypercoagulability        | 16-17            | +                 | +                        | N.D.                    | +      | +                   | N.D.                           | Plaque rupture not spontaneous                                    | 112  |
| ApoE <sup>-/-</sup>                             | Partial ligation carotid + renal arteries | 16               | +                 | 50%                      | N.D.                    | 80%    | N.D.                | N.D.                           | Plaque rupture not spontaneous                                    | 113  |
| ApoE <sup>-/-</sup>                             | uPA overexpression in Mφ                  | 43-48            | 78%               | Fibrin deposition (67%)  | N.D.                    | 61%    | N.D.                | N.D.                           | Long term experiment, complicated manipulation                    | 114  |
| ApoE <sup>-/-</sup>                             | Tandem stenosis                           | 14-22            | 32%               | +                        | +                       | 50%    | +                   | N.D.                           | Plaque rupture not spontaneous                                    | 72   |
| ApoE <sup>-/-</sup> ; Fbn1 <sup>C1039G+/-</sup> | Elastin fragmentation                     | 20-35            | 50-70%            | carotid                  | +                       | 90%    | +                   | MI, stroke, sudden death       | Spontaneous plaque rupture  | 80   |

IPH, intraplaque haemorrhage; Ref., reference; +, present; MI, myocardial infarction N.D., not determined; Ad, adenovirus; Mφ, macrophages; uPA, Urokinase-type plasminogen active



**Table 2.** Most important advantages and limitations of commonly used models of atherosclerosis

|                           | <b>Advantages</b>   | <b>Limitations</b>  |
|---------------------------|---|---|
| <b>Mice</b>               | <ul style="list-style-type: none"> <li>• Relatively cheap</li> <li>• Efficient</li> <li>• Easy crossbreeding</li> <li>• Low cost for pharmacological intervention studies</li> </ul>  | <ul style="list-style-type: none"> <li>• Differences with human plaques:               <ul style="list-style-type: none"> <li>– Less coronary plaque formation</li> <li>– No intraplaque neovascularisation and haemorrhage</li> <li>– Rare plaque rupture and thrombosis</li> </ul> </li> </ul> <p>However, these limitations are overcome in ApoE<sup>-/-</sup>Fbn1<sup>C1039G+/-</sup> mice.</p>       |
| <b>Rabbits</b>            | <ul style="list-style-type: none"> <li>• Allows translation research such as catheterisation of the aorta (coronary arteries are too small)</li> <li>• Relatively cheap</li> <li>• Easy to breed and handle</li> </ul>  | <ul style="list-style-type: none"> <li>• Differences in the localisation of the plaques as compared to humans:               <ul style="list-style-type: none"> <li>– Plaques are most pronounced in the aortic arch and descending thoracic aorta (in contrast to the abdominal aorta in humans)</li> </ul> </li> </ul>  |
| <b>Pigs</b>               | <ul style="list-style-type: none"> <li>• Similarities with human plaques:               <ul style="list-style-type: none"> <li>– Localisation: coronary arteries, abdominal aorta, ileo-femoral</li> <li>– Neovascularisation towards the plaque</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li>• Plaque development mostly ends in the foam cell stage (this can be overcome with the selection of natural mutants, mechanical damage, introduction of other risk factors, genetic engineering with minipigs)</li> <li>• Thrombosis due to plaque rupture is rare (in contrast to humans)</li> <li>• Relatively expensive and more difficult to handle</li> </ul> |
| <b>Non-human primates</b> | <ul style="list-style-type: none"> <li>• Very similar plaque formation as compared to humans (both micro- and macroscopic)</li> <li>• Plaque formation in the coronary arteries</li> </ul>  | <ul style="list-style-type: none"> <li>• Non-human primates are expensive and highly regulated</li> <li>• Specialized training is needed</li> </ul>   |

## References

1. Fuster JJ, Castillo AI, Zaragoza C, Ibáñez B, Andrés V. Animal models of atherosclerosis. *Progress in Molecular Biology and Translational Science*. 2012;1-23
2. Tabas I, García-Cardena G, Owens GK. Recent insights into the cellular biology of atherosclerosis. *J. Cell Biol.* 2015;209:13-22
3. Sakakura K, Nakano M, Otsuka F, Ladich E, Kolodgie FD, Virmani R. Pathophysiology of atherosclerosis plaque progression. *Heart Lung Circ.* 2013;22:399-411
4. Berliner JA, Navab M, Fogelman AM, Frank JS, Demer LL, Edwards PA, Watson AD, Lusis AJ. Atherosclerosis: Basic mechanisms : Oxidation, inflammation, and genetics. *Circulation*. 1995;91:2488-2496
5. Lusis AJ. Atherosclerosis. *Nature*. 2000;407:233-241
6. Konstantinov IE, Jankovic GM. Alexander i. Ignatowski: A pioneer in the study of atherosclerosis. *Tex. Heart Inst. J.* 2013;40:246-249
7. Kapourchali FR, Surendiran G, Chen L, Uitz E, Bahadori B, Moghadasian MH. Animal models of atherosclerosis. *World J. Clin. Cases*. 2014;2:126-132
8. Getz GS, Reardon CA. Animal models of atherosclerosis. *Arterioscler. Thromb. Vasc. Biol.* 2012;32:1104-1115
9. Bond AR, Jackson CL. The fat-fed apolipoprotein e knockout mouse brachiocephalic artery in the study of atherosclerotic plaque rupture. *J. Biomed. Biotechnol.* 2011;2011:1-10
10. VanderLaan PA, Reardon CA, Getz GS. Site specificity of atherosclerosis: Site-selective responses to atherosclerotic modulators. *Arterioscler. Thromb. Vasc. Biol.* 2004;24:12-22
11. Schwartz SM, Galis ZS, Rosenfeld ME, Falk E. Plaque rupture in humans and mice. *Arterioscler. Thromb. Vasc. Biol.* 2007;27:705-713
12. Meir KS, Leitersdorf E. Atherosclerosis in the apolipoprotein e-deficient mouse: A decade of progress. *Arterioscler. Thromb. Vasc. Biol.* 2004;24:1006-1014

13. Piedrahita JA, Zhang SH, Hageman JR, Oliver PM, Maeda N. Generation of mice carrying a mutant apolipoprotein e gene inactivated by gene targeting in embryonic stem cells. *PNAS*. 1992;89:4471-4475
14. Plump AS, Smith JD, Hayek T, Aalto-Setälä K, Walsh A, Verstuyft JG, Rubin EM, Breslow JL. Severe hypercholesterolemia and atherosclerosis in apolipoprotein e-deficient mice created by homologous recombination in es cells. *Cell*. 1992;71:343-353
15. Silvestre-Roig C, de Winther MP, Weber C, Daemen MJ, Lutgens E, Soehnlein O. Atherosclerotic plaque destabilization. *Circ. Res*. 2014;114:214-226
16. Zadelaar S, Kleemann R, Verschuren L, de Vries-Van der Weij J, van der Hoorn J, Princen HM, Kooistra T. Mouse models for atherosclerosis and pharmaceutical modifiers. *Arterioscler. Thromb. Vasc. Biol*. 2007;27:1706-1721
17. Lutgens E, Daemen M, Kockx M, Doevendans P, Hofker M, Havekes L, Wellens H, de Muinck ED. Atherosclerosis in apoe\*3-leiden transgenic mice : From proliferative to atheromatous stage. *Circulation*. 1999;99:276-283
18. Ross R. Cell biology of atherosclerosis. *Annu Rev Physiol*. 1995;57:791-804
19. Breslow JL. Transgenic mouse models of lipoprotein metabolism and atherosclerosis. *Proc. Natl. Acad. Sci. U. S. A*. 1993;90:8314-8318
20. Gencer B, Kronenberg F, Stroes ES, Mach F. Lipoprotein(a): The revenant. *Eur. Heart J*. 2017
21. Lawn RM, Wade DP, Hammer RE, Chiesa G, Verstuyft JG, Rubin EM. Atherogenesis in transgenic mice expressing human apolipoprotein(a). *Nature*. 1992;360:670-672
22. Curtiss LK, Boisvert WA. Apolipoprotein e and atherosclerosis. *Curr. Opin. Lipidol*. 2000;11:243-251
23. Sehayek E, Shefer S, Nguyen LB, Ono JG, Merkel M, Breslow JL. Apolipoprotein e regulates dietary cholesterol absorption and biliary cholesterol excretion: Studies in c57bl/6 apolipoprotein e knockout mice. *PNAS*. 2000;97:3433-3437
24. Mahley R. Apolipoprotein e: Cholesterol transport protein with expanding role in cell biology. *Science*. 1988;240:622-630

25. Jawien J, Nastalek P, Korbut R. Mouse models of experimental atherosclerosis. *J. Physiol. Pharmacol.* 2004;55:503-517
26. Nakashima Y, Plump AS, Raines EW, Breslow JL, Ross R. Apoe-deficient mice develop lesions of all phases of atherosclerosis throughout the arterial tree. *Arterioscler. Thromb. Vasc. Biol.* 1994;14:133-140
27. Plump AS, Breslow JL. Apolipoprotein e and the apolipoprotein e-deficient mouse. *Annu. Rev. Nutr.* 1995;15:495-518
28. Reddick RL, Zhang SH, Maeda N. Atherosclerosis in mice lacking apo e. Evaluation of lesional development and progression [published erratum appears in *arterioscler thromb* 1994 may;14(5):839]. *Arterioscler. Thromb. Vasc. Biol.* 1994;14:141-147
29. Rattazzi M, Bennett BJ, Bea F, Kirk EA, Ricks JL, Speer M, Schwartz SM, Giachelli CM, Rosenfeld ME. Calcification of advanced atherosclerotic lesions in the innominate arteries of apoe-deficient mice: Potential role of chondrocyte-like cells. *Arterioscler. Thromb. Vasc. Biol.* 2005;25:1420-1425
30. Getz GS, Reardon CA. Apoprotein e as a lipid transport and signaling protein in the blood, liver, and artery wall. *J. Lipid Res.* 2009;50 Suppl:S156-161
31. Smith JD, Breslow JL. The emergence of mouse models of atherosclerosis and their relevance to clinical research. *J. Intern. Med.* 1997;242:99-109
32. Plump AS, Lum PY. Genomics and cardiovascular drug development. *J. Am. Coll. Cardiol.* 2009;53:1089-1100
33. Sasaki T, Kuzuya M, Nakamura K, Cheng XW, Shibata T, Sato K, Iguchi A. A simple method of plaque rupture induction in apolipoprotein e-deficient mice. *Arterioscler. Thromb. Vasc. Biol.* 2006;26:1304-1309
34. von der Thusen JH, van Berkel TJ, Biessen EA. Induction of rapid atherogenesis by perivascular carotid collar placement in apolipoprotein e-deficient and low-density lipoprotein receptor-deficient mice. *Circulation.* 2001;103:1164-1170
35. Marais AD. Familial hypercholesterolaemia. *Clin. Biochem. Rev.* 2004;25:49-68
36. Defesche JC. Low-density lipoprotein receptor--its structure, function, and mutations. *Semin. Vasc. Med.* 2004;4:5-11

37. Ishibashi S, Brown MS, Goldstein JL, Gerard RD, Hammer RE, Herz J. Hypercholesterolemia in low density lipoprotein receptor knockout mice and its reversal by adenovirus-mediated gene delivery. *J. Clin. Invest.* 1993;92:883-893
38. Ishibashi S, Goldstein JL, Brown MS, Herz J, Burns DK. Massive xanthomatosis and atherosclerosis in cholesterol-fed low density lipoprotein receptor-negative mice. *J. Clin. Invest.* 1994;93:1885-1893
39. Zhang SH, Reddick RL, Piedrahita JA, Maeda N. Spontaneous hypercholesterolemia and arterial lesions in mice lacking apolipoprotein e. *Science.* 1992;258:468-471
40. Knowles JW, Maeda N. Genetic modifiers of atherosclerosis in mice. *Arterioscler. Thromb. Vasc. Biol.* 2000;20:2336-2345
41. Hartvigsen K, Binder CJ, Hansen LF, Rafia A, Juliano J, Horkko S, Steinberg D, Palinski W, Witztum JL, Li AC. A diet-induced hypercholesterolemic murine model to study atherogenesis without obesity and metabolic syndrome. *Arterioscler. Thromb. Vasc. Biol.* 2007;27:878-885
42. Véniant MM, Sullivan MA, Kim SK, Ambroziak P, Chu A, Wilson MD, Hellerstein MK, Rudel LL, Walzem RL, Young SG. Defining the atherogenicity of large and small lipoproteins containing apolipoprotein b100. *J. Clin. Invest.* 2000;106:1501-1510
43. Veniant MM, Withycombe S, Young SG. Lipoprotein size and atherosclerosis susceptibility in apoe<sup>-/-</sup> and ldlr<sup>-/-</sup> mice. *Arterioscler. Thromb. Vasc. Biol.* 2001;21:1567-1570
44. Hobbs HH, Russell DW, Brown MS, Goldstein JL. The ldl receptor locus in familial hypercholesterolemia: Mutational analysis of a membrane protein. *Annu. Rev. Genet.* 1990;24:133-170
45. Lee YT, Lin HY, Chan YW, Li KH, To OT, Yan BP, Liu T, Li G, Wong WT, Keung W, Tse G. Mouse models of atherosclerosis: A historical perspective and recent advances. *Lipids Health Dis.* 2017;16:12
46. Bonthu S, Heistad DD, Chappell DA, Lamping KG, Faraci FM. Atherosclerosis, vascular remodeling, and impairment of endothelium-dependent relaxation in genetically altered hyperlipidemic mice. *Arterioscler. Thromb. Vasc. Biol.* 1997;17:2333-2340

47. Witting PK, Pettersson K, Ostlund-Lindqvist AM, Westerlund C, Eriksson AW, Stocker R. Inhibition by a coantioxidant of aortic lipoprotein lipid peroxidation and atherosclerosis in apolipoprotein e and low density lipoprotein receptor gene double knockout mice. *FASEB J.* 1999;13:667-675
48. Caligiuri G, Levy B, Pernow J, Thoren P, Hansson GK. Myocardial infarction mediated by endothelin receptor signaling in hypercholesterolemic mice. *Proc. Natl. Acad. Sci. U. S. A.* 1999;96:6920-6924
49. Ishibashi S, Herz J, Maeda N, Goldstein JL, Brown MS. The two-receptor model of lipoprotein clearance: Tests of the hypothesis in "knockout" mice lacking the low density lipoprotein receptor, apolipoprotein e, or both proteins. *Proc. Natl. Acad. Sci. U. S. A.* 1994;91:4431-4435
50. van den Maagdenberg AM, Hofker MH, Krimpenfort PJ, de Bruijn I, van Vlijmen B, van der Boom H, Havekes LM, Frants RR. Transgenic mice carrying the apolipoprotein e3-leiden gene exhibit hyperlipoproteinemia. *J Biol Chem.* 1993;268:10540-10545
51. van Vlijmen BJ, van den Maagdenberg AM, Gijbels MJ, van der Boom H, HogenEsch H, Frants RR, Hofker MH, Havekes LM. Diet-induced hyperlipoproteinemia and atherosclerosis in apolipoprotein e3-leiden transgenic mice. *J Clin Invest.* 1994;93:1403-1410
52. Gijbels MJ, van der Cammen M, van der Laan LJ, Emeis JJ, Havekes LM, Hofker MH, Kraal G. Progression and regression of atherosclerosis in apoe3-leiden transgenic mice: An immunohistochemical study. *Atherosclerosis.* 1999;143:15-25
53. de Vries MR, Simons KH, Jukema JW, Braun J, Quax PHA. Vein graft failure: From pathophysiology to clinical outcomes. *Nat. Rev. Cardiol.* 2016;13:451-470
54. Karper JC, de Vries MR, van den Brand BT, Hoefler IE, Fischer JW, Jukema JW, Niessen HWM, Quax PHA. Toll-like receptor 4 is involved in human and mouse vein graft remodeling, and local gene silencing reduces vein graft disease in hypercholesterolemic apoe\*3leiden mice. *Arterioscler. Thromb. Vasc. Biol.* 2011;31:1033-1040
55. Zou Y, Dietrich H, Hu Y, Metzler B, Wick G, Xu Q. Mouse model of venous bypass graft arteriosclerosis. *Am. J. Pathol.* 1998;153:1301-1310

56. Dietrich H, Hu Y, Zou Y, Huemer U, Metzler B, Li C, Mayr M, Xu Q. Rapid development of vein graft atheroma in apoe-deficient mice. *Am. J. Pathol.* 2000;157:659-669
57. Lardenoye JHP, De Vries MR, Grimbergen JM, Havekes LM, Knaapen MW, Kockx MM, van Hinsbergh VW, van Bockel JH, Quax PH. Inhibition of accelerated atherosclerosis in vein grafts by placement of external stent in apoe\*3-leiden transgenic mice. *Arterioscler. Thromb. Vasc. Biol.* 2002;22:1433-1438
58. Lardenoye JHP, de Vries MR, Löwik CW, Xu Q, Dhore CR, Cleutjens JP, van Hinsbergh VW, van Bockel JH, Quax PH. Accelerated atherosclerosis and calcification in vein grafts: A study in apoe\*3 leiden transgenic mice. *Circ. Res.* 2002;91:577-584
59. Lardenoye JH, Delsing DJ, de Vries MR, Deckers MM, Princen HM, Havekes LM, van Hinsbergh VW, van Bockel JH, Quax PH. Accelerated atherosclerosis by placement of a perivascular cuff and a cholesterol-rich diet in apoe\*3leiden transgenic mice. *Circ. Res.* 2000;87:248-253
60. Bjorklund MM, Hollensen AK, Hagensen MK, Dagnaes-Hansen F, Christoffersen C, Mikkelsen JG, Bentzon JF. Induction of atherosclerosis in mice and hamsters without germline genetic engineering. *Circ. Res.* 2014;114:1684-1689
61. Roche-Molina M, Sanz-Rosa D, Cruz FM, Garcia-Prieto J, Lopez S, Abia R, Muriana FJ, Fuster V, Ibanez B, Bernal JA. Induction of sustained hypercholesterolemia by single adeno-associated virus-mediated gene transfer of mutant hpcsk9. *Arterioscler. Thromb. Vasc. Biol.* 2015;35:50-59
62. Denis M, Marcinkiewicz J, Zaid A, Gauthier D, Poirier S, Lazure C, Seidah NG, Prat A. Gene inactivation of proprotein convertase subtilisin/kexin type 9 reduces atherosclerosis in mice. *Circulation.* 2012;125:894-901
63. Akram ON, Bernier A, Petrides F, Wong G, Lambert G. Beyond ldl cholesterol, a new role for pcsk9. *Arterioscler. Thromb. Vasc. Biol.* 2010;30:1279-1281
64. Li J, Tumanut C, Gavigan JA, Huang WJ, Hampton EN, Tumanut R, Suen KF, Trauger JW, Spraggon G, Lesley SA, Liao G, Yowe D, Harris JL.

- Secreted pcsk9 promotes ldl receptor degradation independently of proteolytic activity. *Biochem. J.* 2007;406:203-207
65. Cerrone M, Noorman M, Lin X, Chkourko H, Liang FX, van der Nagel R, Hund T, Birchmeier W, Mohler P, van Veen TA, van Rijen HV, Delmar M. Sodium current deficit and arrhythmogenesis in a murine model of plakophilin-2 haploinsufficiency. *Cardiovasc. Res.* 2012;95:460-468
66. Kaspar BK, Roth DM, Lai NC, Drumm JD, Erickson DA, McKirnan MD, Hammond HK. Myocardial gene transfer and long-term expression following intracoronary delivery of adeno-associated virus. *J. Gene Med.* 2005;7:316-324
67. Suhy DA, Kao SC, Mao T, Whiteley L, Denise H, Souberbielle B, Burdick AD, Hayes K, Wright JF, Lavender H, Roelvink P, Kolykhalov A, Brady K, Moschos SA, Hauck B, Zelenia O, Zhou S, Scribner C, High KA, Renison SH, Corbau R. Safe, long-term hepatic expression of anti-hcv shRNA in a nonhuman primate model. *Mol. Ther.* 2012;20:1737-1749
68. Zsebo K, Yaroshinsky A, Rudy JJ, Wagner K, Greenberg B, Jessup M, Hajjar RJ. Long-term effects of aav1/serca2a gene transfer in patients with severe heart failure: Analysis of recurrent cardiovascular events and mortality. *Circ. Res.* 2014;114:101-108
69. Goettsch C, Hutcheson JD, Hagita S, Rogers MA, Creager MD, Pham T, Choi J, Mlynarchik AK, Pieper B, Kjolby M, Aikawa M, Aikawa E. A single injection of gain-of-function mutant pcsk9 adeno-associated virus vector induces cardiovascular calcification in mice with no genetic modification. *Atherosclerosis.* 2016;251:109-118
70. Ylä-Herttuala S, Bentzon JF, Daemen M, Falk E, Garcia-Garcia HM, Herrmann J, Hofer I, Jukema JW, Krams R, Kwak BR, Marx N, Naruszewicz M, Newby A, Pasterkamp G, Serruys PWJC, Waltenberger J, Weber C, Tokgözoğlu L. Stabilisation of atherosclerotic plaques. *Thromb. Haemost.* 2011;106:1-19
71. Ni M, Chen WQ, Zhang Y. Animal models and potential mechanisms of plaque destabilisation and disruption. *Heart.* 2009;95:1393-1398
72. Chen YC, Bui AV, Diesch J, Manasseh R, Hausding C, Rivera J, Haviv I, Agrotis A, Htun NM, Jowett J, Hagemeyer CE, Hannan RD, Bobik A, Peter



- K. A novel mouse model of atherosclerotic plaque instability for drug testing and mechanistic/therapeutic discoveries using gene and microRNA expression profiling. *Circ. Res.* 2013;113:252-265
73. Bentzon JF, Falk E. Atherosclerotic lesions in mouse and man: Is it the same disease? *Curr. Opin. Lipidol.* 2010;21:434-440
74. Kielty CM, Sherratt MJ, Shuttleworth CA. Elastic fibres. *J. Cell Sci.* 2002;115:2817-2828
75. Judge DP, Dietz HC. Marfan's syndrome. *Lancet.* 2005;366:1965-1976
76. Van Herck JL, De Meyer GRY, Martinet W, Van Hove CE, Foubert K, Theunis MH, Apers S, Bult H, Vrints CJ, Herman AG. Impaired fibrillin-1 function promotes features of plaque instability in apolipoprotein e-deficient mice. *Circulation.* 2009;120:2478-2487
77. Judge DP, Biery NJ, Keene DR, Geubtner J, Myers L, Huso DL, Sakai LY, Dietz HC. Evidence for a critical contribution of haploinsufficiency in the complex pathogenesis of marfan syndrome. *J. Clin. Invest.* 2004;114:172-181
78. Mariko B, Pezet M, Escoubet B, Bouillot S, Andrieu J-P, Starcher B, Quaglino D, Jacob M-P, Huber P, Ramirez F, Faury G. Fibrillin-1 genetic deficiency leads to pathological ageing of arteries in mice. *J. Pathol.* 2011;224:33-44
79. Medley TL, Cole TJ, Gatzka CD, Wang WY, Dart AM, Kingwell BA. Fibrillin-1 genotype is associated with aortic stiffness and disease severity in patients with coronary artery disease. *Circulation.* 2002;105:810-815
80. Van der Donckt C, Van Herck JL, Schrijvers DM, Vanhoutte G, Verhoye M, Blockx I, Van Der Linden A, Bauters D, Lijnen HR, Sluimer JC, Roth L, Van Hove CE, Franssen P, Knaapen MW, Hervent AS, De Keulenaer GW, Bult H, Martinet W, Herman AG, De Meyer GRY. Elastin fragmentation in atherosclerotic mice leads to intraplaque neovascularization, plaque rupture, myocardial infarction, stroke, and sudden death. *Eur. Heart J.* 2015;36:1049-1058
81. Van der Donckt C, Roth L, Vanhoutte G, Blockx I, Bink DI, Ritz K, Pintelon I, Timmermans JP, Bauters D, Martinet W, Daemen MJ, Verhoye M, De Meyer GRY. Fibrillin-1 impairment enhances blood-brain barrier

- permeability and xanthoma formation in brains of apolipoprotein e-deficient mice. *Neuroscience*. 2015;295:11-22
82. Fulop T, Larbi A, Fortun A, Robert L, Khalil A. Elastin peptides induced oxidation of ldl by phagocytic cells. *Pathol. Biol. (Paris)*. 2005;53:416-423
83. Koenig W, Khuseyinova N. Biomarkers of atherosclerotic plaque instability and rupture. *Arterioscler. Thromb. Vasc. Biol*. 2007;27:15-26
84. Libby P. Mechanisms of acute coronary syndromes and their implications for therapy. *N. Engl. J. Med*. 2013;368:2004-2013
85. Raffetto JD, Khalil RA. Matrix metalloproteinases and their inhibitors in vascular remodeling and vascular disease. *Biochem. Pharmacol*. 2008;75:346-359
86. de Nooijer R, Verkleij CJ, von der Thüsen JH, Jukema JW, van der Wall EE, van Berkel TJ, Baker AH, Biessen EA. Lesional overexpression of matrix metalloproteinase-9 promotes intraplaque hemorrhage in advanced lesions but not at earlier stages of atherogenesis. *Arterioscler. Thromb. Vasc. Biol*. 2005;26:340-346
87. Gough PJ, Gomez IG, Wille PT, Raines EW. Macrophage expression of active mmp-9 induces acute plaque disruption in apoe-deficient mice. *J. Clin. Invest*. 2006;116:59-69
88. Virmani R, Kolodgie FD, Burke AP, Finn AV, Gold HK, Tulenko TN, Wrenn SP, Narula J. Atherosclerotic plaque progression and vulnerability to rupture: Angiogenesis as a source of intraplaque hemorrhage. *Arterioscler. Thromb. Vasc. Biol*. 2005;25:2054-2061
89. Moulton KS, Olsen BR, Sonn S, Fukai N, Zurakowski D, Zeng X. Loss of collagen xviii enhances neovascularization and vascular permeability in atherosclerosis. *Circulation*. 2004;110:1330-1336
90. Rademakers T, Douma K, Hackeng TM, Post MJ, Sluimer JC, Daemen MJAP, Biessen EAL, Heeneman S, van Zandvoort MAMJ. Plaque-associated vasa vasorum in aged apolipoprotein e-deficient mice exhibit proatherogenic functional features in vivo. *Arterioscler. Thromb. Vasc. Biol*. 2013;33:249-256
91. Sluimer JC, Kolodgie FD, Bijnens APJJ, Maxfield K, Pacheco E, Kutys B, Duimel H, Frederik PM, van Hinsbergh VWM, Virmani R, Daemen MJAP.

- Thin-walled microvessels in human coronary atherosclerotic plaques show incomplete endothelial junctions. *J. Am. Coll. Cardiol.* 2009;53:1517-1527
92. Kockx MM, Cromheeke KM, Knaapen MW, Bosmans JM, De Meyer GRY, Herman AG, Bult H. Phagocytosis and macrophage activation associated with hemorrhagic microvessels in human atherosclerosis. *Arterioscler. Thromb. Vasc. Biol.* 2003;23:440-446
93. Slager CJ, Wentzel JJ, Gijssen FJH, Thury A, van der Wal AC, Schaar JA, Serruys PW. The role of shear stress in the destabilization of vulnerable plaques and related therapeutic implications. *Nat. Clin. Pract. Cardiovasc. Med.* 2005;2:456-464
94. Huang Y, Teng Z, Sadat U, He J, Graves MJ, Gillard JH. In vivo mri-based simulation of fatigue process: A possible trigger for human carotid atherosclerotic plaque rupture. *BioMed. Eng. OnLine.* 2013;12:36
95. Roth L, Rombouts M, Schrijvers DM, Martinet W, De Meyer GRY. Cholesterol-independent effects of atorvastatin prevent cardiovascular morbidity and mortality in a mouse model of atherosclerotic plaque rupture. *Vascul. Pharmacol.* 2016;80:50-58
96. Roth L, Rombouts M, Schrijvers DM, Lemmens K, De Keulenaer GW, Martinet W, De Meyer GRY. Chronic intermittent mental stress promotes atherosclerotic plaque vulnerability, myocardial infarction and sudden death in mice. *Atherosclerosis.* 2015;242:288-294
97. Roth L, Van Dam D, Van der Donckt C, Schrijvers DM, Lemmens K, Van Brussel I, De Deyn PP, Martinet W, De Meyer GRY. Impaired gait pattern as a sensitive tool to assess hypoxic brain damage in a novel mouse model of atherosclerotic plaque rupture. *Physiol. Behav.* 2015;139:397-402
98. Fan J, Kitajima S, Watanabe T, Xu J, Zhang J, Liu E, Chen YE. Rabbit models for the study of human atherosclerosis: From pathophysiological mechanisms to translational medicine. *Pharmacol. Ther.* 2015;146:104-119
99. Watanabe Y. Serial inbreeding of rabbits with hereditary hyperlipidemia (whhl-rabbit) \*1 incidence and development of atherosclerosis and xanthoma. *Atherosclerosis.* 1980;36:261-268

100. Baumgartner C, Brandl J, Münch G, Ungerer M. Rabbit models to study atherosclerosis and its complications – transgenic vascular protein expression in vivo. *Prog. Biophys. Mol. Biol.* 2016;121:131-141
101. Niimi M, Yang D, Kitajima S, Ning B, Wang C, Li S, Liu E, Zhang J, Eugene Chen Y, Fan J. Apoe knockout rabbits: A novel model for the study of human hyperlipidemia. *Atherosclerosis.* 2016;245:187-193
102. Fan J, Watanabe T. Cholesterol-fed and transgenic rabbit models for the study of atherosclerosis. *J. Atheroscler. Thromb.* 2000;7:26-32
103. Atkinson JB, Hoover RL, Berry KK, Swift LL. Cholesterol-fed heterozygous watanabe heritable hyperlipidemic rabbits: A new model for atherosclerosis. *Atherosclerosis.* 1989;78:123-136
104. Ning B, Wang X, Yu Y, Waqar AB, Yu Q, Koike T, Shiomi M, Liu E, Wang Y, Fan J. High-fructose and high-fat diet-induced insulin resistance enhances atherosclerosis in watanabe heritable hyperlipidemic rabbits. *Nutr. Metab. (Lond.).* 2015;12
105. Brousseau ME, Hoeg JM. Transgenic rabbits as models for atherosclerosis research. *J. Lipid Res.* 1999;40:365-375
106. Shim J, Al-Mashhadi RH, Sorensen CB, Bentzon JF. Large animal models of atherosclerosis--new tools for persistent problems in cardiovascular medicine. *J. Pathol.* 2016;238:257-266
107. Al-Mashhadi RH, Sorensen CB, Kragh PM, Christoffersen C, Mortensen MB, Tolbod LP, Thim T, Du Y, Li J, Liu Y, Moldt B, Schmidt M, Vajta G, Larsen T, Purup S, Bolund L, Nielsen LB, Callesen H, Falk E, Mikkelsen JG, Bentzon JF. Familial hypercholesterolemia and atherosclerosis in cloned minipigs created by DNA transposition of a human pcsk9 gain-of-function mutant. *Sci. Transl. Med.* 2013;5:166ra161
108. Niu Y, Shen B, Cui Y, Chen Y, Wang J, Wang L, Kang Y, Zhao X, Si W, Li W, Xiang Andy P, Zhou J, Guo X, Bi Y, Si C, Hu B, Dong G, Wang H, Zhou Z, Li T, Tan T, Pu X, Wang F, Ji S, Zhou Q, Huang X, Ji W, Sha J. Generation of gene-modified cynomolgus monkey via cas9/rna-mediated gene targeting in one-cell embryos. *Cell.* 2014;156:836-843

109. Calara F, Silvestre M, Casanada F, Yuan N, Napoli C, Palinski W. Spontaneous plaque rupture and secondary thrombosis in apolipoprotein e-deficient and ldl receptor-deficient mice. *J. Pathol.* 2001;195:257-263
110. Williams H. Characteristics of intact and ruptured atherosclerotic plaques in brachiocephalic arteries of apolipoprotein e knockout mice. *Arterioscler. Thromb. Vasc. Biol.* 2002;22:788-792
111. von der Thusen JH, van Vlijmen BJM, Hoeben RC, Kockx MM, Havekes LM, van Berkel TJC, Biessen EAL. Induction of atherosclerotic plaque rupture in apolipoprotein e-/- mice after adenovirus-mediated transfer of p53. *Circulation.* 2002;105:2064-2070
112. Borissoff JI, Otten JJT, Heeneman S, Leenders P, van Oerle R, Soehnlein O, Loubele STBG, Hamulyák K, Hackeng TM, Daemen MJAP, Degen JL, Weiler H, Esmon CT, van Ryn J, Biessen EAL, Spronk HMH, ten Cate H. Genetic and pharmacological modifications of thrombin formation in apolipoprotein e-deficient mice determine atherosclerosis severity and atherothrombosis onset in a neutrophil-dependent manner. *PLoS One.* 2013;8:e55784
113. Jin Sx, Shen Lh, Nie P, Yuan W, Hu Lh, Li Dd, Chen Xj, Zhang Xk, He B. Endogenous renovascular hypertension combined with low shear stress induces plaque rupture in apolipoprotein e-deficient mice. *Arterioscler. Thromb. Vasc. Biol.* 2012;32:2372-2379
114. Hu JH, Du L, Chu T, Otsuka G, Dronadula N, Jaffe M, Gill SE, Parks WC, Dichek DA. Overexpression of urokinase by plaque macrophages causes histological features of plaque rupture and increases vascular matrix metalloproteinase activity in aged apolipoprotein e-null mice. *Circulation.* 2010;121:1637-1644

