



Universiteit
Leiden
The Netherlands

From molecules to monitoring: integrating genetic tools into freshwater quality assessments

Beentjes, K.K.

Citation

Beentjes, K. K. (2021, April 8). *From molecules to monitoring: integrating genetic tools into freshwater quality assessments*. Retrieved from <https://hdl.handle.net/1887/3158798>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3158798>

Note: To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <https://hdl.handle.net/1887/3158798> holds various files of this Leiden University dissertation.

Author: Beentjes, K.K.

Title: From molecules to monitoring: integrating genetic tools into freshwater quality assessments

Issue Date: 2021-04-08

REFERENCES

Afgan E, Baker D, Batut B, Van Den Beek M, Bouvier D, Ech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D (2018) The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Research* 46:W537-W544. doi: 10.1093/nar/gky379.

Agersnap S, Larsen WB, Knudsen SW, Strand D, Thomsen PF, Hesselsøe M, Mortensen PB, Vrålstad T, Møller PR (2017) Monitoring of noble, signal and narrow-clawed crayfish using environmental DNA from freshwater samples. *PLoS ONE* 12:e0179261. doi: 10.1371/journal.pone.0179261.

Alberdi A, Aizpurua O, Gilbert MTP, Bohmann K (2018) Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution* 9:134-147. doi: 10.1111/2041-210X.12849.

Alcamo JM, Vörösmarty CJ, Naiman RJ, Lettenmaier DP, Pahl-Wostl C (2008) A grand challenge for freshwater research: understanding the global water system. *Environmental Research Letters* 3:010202. doi: 10.1088/1748-9326/3/1/010202.

Alp M, Keller I, Westram AM, Robinson CT (2012) How river structure and biological traits influence gene flow: a population genetic study of two stream invertebrates with differing dispersal abilities. *Freshwater Biology* 57:969-981. doi: 10.1111/j.1365-2427.2012.02758.x.

Amend AS, Seifert KA, Bruns TD (2010) Quantifying microbial communities with 454 pyrosequencing: Does read abundance count? *Molecular Ecology* 19:5555-5565. doi: 10.1111/j.1365-294X.2010.04898.x.

Andruszkiewicz EA, Starks HA, Chavez FP, Sassoubre LM, Block BA, Boehm AB (2017) Biomonitoring of marine vertebrates in Monterey Bay using eDNA metabarcoding. *PLoS ONE* 12:e0176343. doi: 10.1371/journal.pone.0176343.

Andújar C, Arribas P, Gray C, Bruce C, Woodward G, Yu DW, Vogler AP (2018a) Metabarcoding of freshwater invertebrates to detect the effects of a pesticide spill. *Molecular Ecology* 27:146-166. doi: 10.1111/mec.14410.

Andújar C, Arribas P, Yu DW, Vogler AP, Emerson BC (2018b) Why the COI barcode should be the community DNA metabarcode for the Metazoa. *Molecular Ecology* 27:3968-3975. doi: 10.1111/mec.14844.

References

- Apothéloz-Perret-Gentil L, Cordonier A, Straub F, Iseli J, Esling P, Pawlowski J (2017) Taxonomy-free molecular diatom index for high-throughput eDNA biomonitoring. *Molecular Ecology Resources* 17:1231-1242. doi: 10.1111/1755-0998.12668.
- Aylagas E, Borja Á, Rodríguez-Ezpeleta N (2014) Environmental status assessment using DNA metabarcoding: Towards a genetics based marine biotic index (gAMBI). *PLoS ONE* 9:e90529. doi: 10.1371/journal.pone.0090529.
- Aylagas E, Borja Á, Irigoien X, Rodríguez-Ezpeleta N (2016) Benchmarking DNA metabarcoding for biodiversity-based monitoring and assessment. *Frontiers in Marine Science* 3:96. doi: 10.3389/fmars.2016.00096.
- Aylagas E, Borja Á, Muxika I, Rodríguez-Ezpeleta N (2018) Adapting metabarcoding-based benthic biomonitoring into routine marine ecological status assessment networks. *Ecological Indicators* 95:194-202. doi: 10.1016/j.ecolind.2018.07.044.
- Bagley M, Pilgrim E, Knapp M, Yoder C, Santo Domingo J, Banerji A (2019) High-throughput environmental DNA analysis informs a biological assessment of an urban stream. *Ecological Indicators* 104:378-389. doi: 10.1016/j.ecolind.2019.04.088.
- Baird DJ, Hajibabaei M (2012) Biomonitoring 2.0: a new paradigm in ecosystem assessment made possible by next-generation DNA sequencing. *Molecular Ecology* 21:2039-2044. doi: 10.1111/j.1365-294X.2012.05519.x.
- Barbour MT, Gerritsen J, Snyder B, Stribling J (1999) *Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish*. Washington, DC: U.S. Environmental Protection Agency.
- Barmentlo SH, Schrama M, De Snoo GR, Van Bodegom PM, Nieuwenhuijzen A, Vijver MG (in prep) Experimental evidence for neonicotinoid driven decline in aquatic emerging insects.
- Barmentlo SH, Schrama M, Van Bodegom PM, De Snoo GR, Musters CJM, Vijver MG (2019) Neonicotinoids and fertilizers jointly structure naturally assembled freshwater macroinvertebrate communities. *Science of the Total Environment* 691:36-44. doi: 10.1016/j.scitotenv.2019.07.110.
- Barnes MA, Turner CR, Jerde CL, Renshaw MA, Chadderton WL, Lodge DM (2014) Environmental conditions influence eDNA persistence in aquatic systems. *Environmental Science & Technology* 48:1819-27. doi: 10.1021/es404734p.
- Barnes MA, Turner CR (2015) The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics* 17:1-17. doi: 10.1007/s10592-015-0775-4.
- Barnosky AD, Matzke N, Tomiya S, Wogan GOU, Swartz B, Quental TB, Marshall C, McGuire JL, Lindsey EL, Maguire KC, Mersey B, Ferrer EA (2011) Has the Earth's sixth mass extinction already arrived? *Nature* 471:51-7. doi: 10.1038/nature09678.

- Barsoum N, Bruce C, Forster J, Ji Y-Q, Yu DW (2019) The devil is in the detail: Metabarcoding of arthropods provides a sensitive measure of biodiversity response to forest stand composition compared with surrogate measures of biodiversity. *Ecological Indicators* 101:313-323. doi: 10.1016/J.ECOLIND.2019.01.023.
- Baselga A, Fujisawa T, Crampton-Platt A, Bergsten J, Foster PG, Monaghan MT, Vogler AP (2013) Whole-community DNA barcoding reveals a spatio-temporal continuum of biodiversity at species and genetic levels. *Nature Communications* 4:1892. doi: 10.1038/ncomms2881.
- Beentjes KK, Speksnijder AGCL, Van der Hoorn BB, Van Tol J (2015) DNA barcoding program at Naturalis Biodiversity Center, the Netherlands. *Genome* 58:193. doi: 10.1139/gen-2015-0087.
- Beermann AJ, Zizka VMA, Elbrecht V, Baranov V, Leese F (2018) DNA metabarcoding reveals the complex and hidden responses of chironomids to multiple stressors. *Environmental Sciences Europe* 30:26. doi: 10.1186/s12302-018-0157-x.
- Beermann AJ, Werner M-T, Elbrecht V, Zizka VMA, Leese F (2020) DNA metabarcoding improves the detection of multiple stressor responses of stream invertebrates to increased salinity, fine sediment deposition and reduced flow velocity. *Science of The Total Environment*:141969. doi: 10.1016/j.scitotenv.2020.141969.
- Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Wheeler DL (2005) GenBank. *Nucleic Acids Research* 33:D34-8. doi: 10.1093/nar/gki063.
- Bergsten J, Bilton DT, Fujisawa T, Elliott M, Monaghan MT, Balke M, Hendrich L, Geijer J, Herrmann J, Foster GN, Ribera I, Nilsson AN, Barraclough TG, Vogler AP (2012) The effect of geographical scale of sampling on DNA barcoding. *Systematic Biology* 61:851-869. doi: 10.1093/sysbio/sys037.
- Berney C, Fahrni J, Pawlowski J (2004) How many novel eukaryotic “kingdoms”? Pitfalls and limitations of environmental DNA surveys. *BMC Biology* 2:13. doi: 10.1186/1741-7007-2-13.
- Biggs J, Ewald N, Valentini A, Gaboriaud C, Dejean T, Griffiths RA, Foster J, Wilkinson JW, Arnell A, Brotherton P, Williams P, Dunn F (2015) Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biological Conservation* 183:19-28. doi: 10.1016/j.biocon.2014.11.029.
- Birk S, Bonne W, Borja A, Brucet S, Courrat A, Poikane S, Solimini A, Van De Bund W, Zampoukas N, Hering D (2012) Three hundred ways to assess Europe’s surface waters: An almost complete overview of biological methods to implement the Water Framework Directive. *Ecological Indicators* 18:31-41. doi: 10.1016/j.ecolind.2011.10.009.
- Bista I, Carvalho GR, Walsh K, Seymour M, Hajibabaei M, Lallias D, Christmas M, Creer S (2017) Annual time-series analysis of aqueous eDNA reveals ecologically relevant dynamics

References

of lake ecosystem biodiversity. *Nature Communications* 8:14087. doi: 10.1038/ncomms14087.

Bista I, Carvalho GR, Tang M, Walsh K, Zhou X, Hajibabaei M, Shokralla S, Seymour M, Bradley D, Liu S, Christmas M, Creer S (2018) Performance of amplicon and shotgun sequencing for accurate biomass estimation in invertebrate community samples. *Molecular Ecology Resources*. doi: 10.1111/1755-0998.12888.

Brown EA, Chain FJJ, Crease TJ, MacIsaac HJ, Cristescu ME (2015) Divergence thresholds and divergent biodiversity estimates: can metabarcoding reliably describe zooplankton communities? *Ecology and Evolution* 5:2234-2251. doi: 10.1002/ece3.1485.

Buchner D, Beermann AJ, Laini A, Rolauuffs P, Vitecek S, Hering D, Leese F (2019) Analysis of 13,312 benthic invertebrate samples from German streams reveals minor deviations in ecological status class between abundance and presence/absence data. *PLoS ONE* 14:e0226547. doi: 10.1371/journal.pone.0226547.

Buhay JE (2009) "COI-like" sequences are becoming problematic in molecular systematic and DNA barcoding studies. *Journal of Crustacean Biology* 29:96-110. doi: 10.1651/08-3020.1.

Bush A, Compson Z, Monk W, Porter T, Steeves R, Emilson E, Gagne N, Hajibabaei M, Roy M, Baird D (2019) Studying ecosystems with DNA metabarcoding: lessons from aquatic biomonitoring. *bioRxiv*:578591. doi: 10.1101/578591.

Butchart SHM, Walpole M, Collen B, Van Strien A, Scharlemann JPW, Almond REA, Baillie JEM, Bomhard B, Brown C, Bruno J, Carpenter KE, Carr GM, Chanson J, Chenery AM, Csirke J, Davidson NC, Dentener F, Foster M, Galli A, Galloway JN, Genovesi P, Gregory RD, Hockings M, Kapos V, Lamarque J-F, Leverington F, Loh J, McGeoch MA, McRae L, Minasyan A, Hernández Morcillo M, Oldfield TEE, Pauly D, Quader S, Revenga C, Sauer JR, Skolnik B, Spear D, Stanwell-Smith D, Stuart SN, Symes A, Tierney M, Tyrrell TD, Vié J-C, Watson R (2010) Global biodiversity: indicators of recent declines. *Science* 328:1164-8. doi: 10.1126/science.1187512.

Buxton AS, Groombridge JJ, Zakaria NB, Griffiths RA (2017) Seasonal variation in environmental DNA in relation to population size and environmental factors. *Scientific Reports* 7:46294. doi: 10.1038/srep46294.

Buxton AS, Groombridge JJ, Griffiths RA (2018) Seasonal variation in environmental DNA detection in sediment and water samples. *PLoS ONE* 13:e0191737. doi: 10.1371/journal.pone.0191737.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. doi: 10.1186/1471-2105-10-421.

Carew ME, Pettigrove VJ, Metzeling L, Hoffmann AA (2013) Environmental monitoring using next generation sequencing: rapid identification of macroinvertebrate bioindicator

species. *Frontiers in Zoology* 10:45. doi: 10.1186/1742-9994-10-45.

Carvalho L, Poikane S, Lyche Solheim A, Phillips G, Borics G, Catalan J, De Hoyos C, Drakare S, Dudley BJ, Järvinen M, Laplace-Tretyure C, Maileht K, McDonald C, Mischke U, Moe J, Morabito G, Nöges P, Nöges T, Ott I, Pasztaleniec A, Skjelbred B, Thackeray SJ (2013) Strength and uncertainty of phytoplankton metrics for assessing eutrophication impacts in lakes. *Hydrobiologia* 704:127-140. doi: 10.1007/s10750-012-1344-1.

Chagnon M, Kreutzweiser D, Mitchell EAD, Morrissey CA, Noome DA, Van Der Sluijs JP (2015) Risks of large-scale use of systemic insecticides to ecosystem functioning and services. *Environmental Science and Pollution Research* 22:119-134. doi: 10.1007/s11356-014-3277-x.

Chain FJJ, Brown EA, MacIsaac HJ, Cristescu ME (2016) Metabarcoding reveals strong spatial structure and temporal turnover of zooplankton communities among marine and freshwater ports. *Diversity and Distributions* 22:493-504. doi: 10.1111/ddi.12427.

Chamberlain S (2017) BOLD: Interface to Bold Systems 'API'. R package version 0.5.0.

Chará-Serna AM, Epele LB, Morrissey CA, Richardson JS (2019) Nutrients and sediment modify the impacts of a neonicotinoid insecticide on freshwater community structure and ecosystem functioning. *Science of the Total Environment* 692:1291-1303. doi: 10.1016/j.scitotenv.2019.06.301.

Chariton AA, Ho KT, Proestou D, Bik H, Simpson SL, Portis LM, Cantwell MG, Baguley JG, Burgess RM, Pelletier MM, Perron M, Gunsch C, Matthews RA (2014) A molecular-based approach for examining responses of eukaryotes in microcosms to contaminant-spiked estuarine sediments. *Environmental Toxicology and Chemistry* 33:359-369. doi: 10.1002/etc.2450.

Chonova T, Kurmayer R, Rimet F, Labanowski J, Vasselon V, Keck F, Illmer P, Bouchez A (2019) Benthic diatom communities in an alpine river impacted by waste water treatment effluents as revealed using DNA metabarcoding. *Frontiers in Microbiology* 10:653. doi: 10.3389/fmicb.2019.00653.

Civade R, Dejean T, Valentini A, Roset N, Raymond JC, Bonin A, Taberlet P, Pont D (2016) Spatial representativeness of environmental DNA metabarcoding signal for fish biodiversity assessment in a natural freshwater system. *PLoS ONE* 11:e0157366. doi: 10.1371/journal.pone.0157366.

Clare EL, Chain FJJ, Littlefair JE, Cristescu ME, Deiner K (2016) The effects of parameter choice on defining molecular operational taxonomic units and resulting ecological analyses of metabarcoding data. *Genome* 59:981-990. doi: 10.1139/gen-2015-0184.

Clarke RT, Hering D (2006) Errors and uncertainty in bioassessment methods - major results and conclusions from the STAR project and their application using STARBUGS. *Hydrobiologia* 566:433-439. doi: 10.1007/s10750-006-0079-2.

References

- Collen B, Whitton F, Dyer EE, Baillie JEM, Cumberlidge N, Darwall WRT, Pollock C, Richman NI, Soulsby A-M, Böhm M (2014) Global patterns of freshwater species diversity, threat and endemism. *Global Ecology and Biogeography* 23:40-51. doi: 10.1111/geb.12096.
- Cordier T, Esling P, Lejzerowicz F, Visco J, Ouadahi A, Martins C, Cedhagen T, Pawlowski J (2017) Predicting the ecological quality status of marine environments from eDNA metabarcoding data using supervised machine learning. *Environmental Science and Technology* 51:9118-9126. doi: 10.1021/acs.est.7b01518.
- Cordier T, Forster D, Dufresne Y, Martins CI, Stoeck T, Pawlowski J (2018) Supervised machine learning outperforms taxonomy-based environmental DNA metabarcoding applied to biomonitoring. *Molecular Ecology Resources*. doi: 10.1111/1755-0998.12926.
- Cordier T, Frontalini F, Cermakova K, Apothéoz-Perret-Gentil L, Treglia M, Scantamburlo E, Bonamin V, Pawlowski J (2019) Multi-marker eDNA metabarcoding survey to assess the environmental impact of three offshore gas platforms in the North Adriatic Sea (Italy). *Marine Environmental Research* 146:24-34. doi: 10.1016/j.marenvres.2018.12.009.
- Corse E, Meglécz E, Archambaud G, Ardisson M, Martin JF, Tougard C, Chappaz R, Dubut V (2017) A from-benchtop-to-desktop workflow for validating HTS data and for taxonomic identification in diet metabarcoding studies. *Molecular Ecology Resources* 17:e146-e159. doi: 10.1111/1755-0998.12703.
- Corvalan C, Hales S, McMichael A (2005) *Ecosystems and Human Well-being: Health Synthesis*. Washington, DC: Island Press.
- Costanza R (1992) Toward an operational definition of ecosystem health. In: Costanza R, Norton BG, Haskell BD eds. *Ecosystem Health: New Goals for Environmental Management*. Washington, DC: Island Press, 239-256.
- Côté IM, Darling ES, Brown CJ (2016) Interactions among ecosystem stressors and their importance in conservation. *Proceedings of the Royal Society B: Biological Sciences* 283:1-9. doi: 10.1098/rspb.2015.2592.
- Creedy TJ, Ng WS, Vogler AP (2019) Toward accurate species-level metabarcoding of arthropod communities from the tropical forest canopy. *Ecology and Evolution* 9:3105-3116. doi: 10.1002/ece3.4839.
- Cristescu ME (2019) Can environmental RNA revolutionize biodiversity science? *Trends in Ecology and Evolution* 34:694-697. doi: 10.1016/j.tree.2019.05.003.
- Curry CJ, Gibson JF, Shokralla S, Hajibabaei M, Baird DJ (2018) Identifying North American freshwater invertebrates using DNA barcodes: are existing COI sequence libraries fit for purpose? *Freshwater Science* 37:178-189. doi: 10.1086/696613.
- Darling JA, Mahon AR (2011) From molecules to management: adopting DNA-based

- methods for monitoring biological invasions in aquatic environments. *Environmental Research* 111:978-88. doi: 10.1016/j.envres.2011.02.001.
- Davis JM, Rosemond AD, Eggert SL, Cross WF, Wallace JB (2010) Long-term nutrient enrichment decouples predator and prey production. *Proceedings of the National Academy of Sciences* 107:121-126. doi: 10.1073/pnas.0908497107.
- Deagle BE, Thomas AC, Shaffer AK, Trites AW, Jarman SN (2013) Quantifying sequence proportions in a DNA-based diet study using Ion Torrent amplicon sequencing: which counts count? *Molecular Ecology Resources* 13:620-33. doi: 10.1111/1755-0998.12103.
- Deagle BE, Jarman SN, Coissac E, Pompanon F, Taberlet P (2014) DNA metabarcoding and the cytochrome c oxidase subunit I marker: not a perfect match. *Biology Letters* 10:20140562. doi: 10.1098/rsbl.2014.0562.
- Deagle BE, Thomas AC, McInnes JC, Clarke LJ, Vesterinen EJ, Clare EL, Kartzinel TR, Eveson JP (2019) Counting with DNA in metabarcoding studies: How should we convert sequence reads to dietary data? *Molecular Ecology* 28:391-406. doi: 10.1111/mec.14734.
- Deiner K, Altermatt F (2014) Transport distance of invertebrate environmental DNA in a natural river. *PLoS ONE* 9:e88786. doi: 10.1371/journal.pone.0088786.
- Deiner K, Knapp RA, Boiano DM, May B (2013) Increased accuracy of species lists developed for alpine lakes using morphology and cytochrome oxidase I for identification of specimens. *Molecular Ecology Resources* 13:820-831. doi: 10.1111/1755-0998.12130.
- Dejean T, Valentini A, Duparc A, Pellier-Cuit S, Pompanon F, Taberlet P, Miaud C (2011) Persistence of environmental DNA in freshwater ecosystems. *PLoS ONE* 6:e23398. doi: 10.1371/journal.pone.0023398.
- Dejean T, Valentini A, Miquel C, Taberlet P, Bellemain E, Miaud C (2012) Improved detection of an alien invasive species through environmental DNA barcoding: the example of the American Bullfrog *Lithobates catesbeianus*. *Journal of Applied Ecology* 49:953-959. doi: 10.1111/j.1365-2664.2012.02171.x.
- DeSalle R, Egan MG, Siddall M (2005) The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions of the Royal Society B: Biological Sciences* 360:1905-16. doi: 10.1098/rstb.2005.1722.
- Dickie IA, Boyer S, Buckley HL, Duncan RP, Gardner PP, Hogg ID, Holdaway RJ, Lear G, Makiola A, Morales SE, Powell JR, Weaver L (2018) Towards robust and repeatable sampling methods in eDNA-based studies. *Molecular Ecology Resources* 18:940-952. doi: 10.1111/1755-0998.12907.
- Van Dijk TC, Van Staalduinen MA, Van der Sluijs JP (2013) Macro-invertebrate decline in surface water polluted with imidacloprid. *PLoS ONE* 8:e62374. doi: 10.1371/journal.pone.0062374.

References

- Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B (2014) Defaunation in the Anthropocene. *Science* 345:401-406. doi: 10.1126/science.1251817.
- Djurhuus A, Port J, Closek CJ, Yamahara KM, Romero-Maraccini O, Walz KR, Goldsmith DB, Michisaki R, Breitbart M, Boehm AB, Chavez FP (2017) Evaluation of filtration and DNA extraction methods for environmental DNA biodiversity assessments across multiple trophic levels. *Frontiers in Marine Science* 4:314. doi: 10.3389/fmars.2017.00314.
- Doi H, Uchii K, Takahara T, Matsuhashi S, Yamanaka H, Minamoto T (2015) Use of droplet digital PCR for estimation of fish abundance and biomass in environmental DNA surveys. *PLoS ONE* 10:e0122763. doi: 10.1371/journal.pone.0122763.
- Donohue I, Donohue LA, Ainín BN, Irvine K (2009) Assessment of eutrophication pressure on lakes using littoral invertebrates. *Hydrobiologia* 633:105-122. doi: 10.1007/s10750-009-9868-8.
- Drew LW (2011) Are we losing the science of taxonomy? *BioScience* 61:942–946. doi: 10.1525/bio.2011.61.12.4.
- Dudgeon D, Arthington AH, Gessner MO, Kawabata Z-I, Knowler DJ, Lévêque C, Naiman RJ, Prieur-Richard A-H, Soto D, Stiassny MLJ, Sullivan CA (2006) Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews of the Cambridge Philosophical Society* 81:163-82. doi: 10.1017/S1464793105006950.
- Edgar RC (2016) UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing. Cold Spring Harbor Laboratory. doi: 10.1101/081257.
- Eichmiller JJ, Miller LM, Sorensen PW (2016) Optimizing techniques to capture and extract environmental DNA for detection and quantification of fish. *Molecular Ecology Resources* 16:56-68. doi: 10.1111/1755-0998.12421.
- Elbrecht V, Taberlet P, Dejean T, Valentini A, Usseglio-Polatera P, Beisel J-N, Coissac E, Boyer F, Leese F (2016) Testing the potential of a ribosomal 16S marker for DNA metabarcoding of insects. *PeerJ* 4:e1966. doi: 10.7717/peerj.1966.
- Elbrecht V, Vamos EE, Meissner K, Aroviita J, Leese F (2017a) Assessing strengths and weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine stream monitoring. *Methods in Ecology and Evolution* 8:1265-1275. doi: 10.1111/2041-210X.12789.
- Elbrecht V, Peinert B, Leese F (2017b) Sorting things out: Assessing effects of unequal specimen biomass on DNA metabarcoding. *Ecology and Evolution* 7:6918-6926. doi: 10.1002/ece3.3192.
- Elbrecht V, Vamos EE, Steinke D, Leese F (2018a) Estimating intraspecific genetic diversity from community DNA metabarcoding data. *PeerJ* 6:e4644. doi: 10.7717/peerj.4644.

- Elbrecht V, Hebert PDN, Steinke D (2018b) Slippage of degenerate primers can cause variation in amplicon length. *Scientific Reports* 8:10999. doi: 10.1038/s41598-018-29364-z.
- Elbrecht V, Bourlat SJ, Lindner A, Noll NW, Sorg M, Zizka VMA, Alexander M (2020) Pooling size sorted malaise trap fractions to maximise taxon recovery with metabarcoding. *bioRxiv*. doi: 10.1101/2020.06.09.118950.
- Elbrecht V, Leese F (2015) Can DNA-based ecosystem assessments quantify species abundance? Testing primer bias and biomass - sequence relationships with an innovative metabarcoding protocol. *PLoS ONE* 10:e0130324. doi: 10.1371/journal.pone.0130324.
- Elbrecht V, Leese F (2017) Validation and development of COI metabarcoding primers for freshwater macroinvertebrate bioassessment. *Frontiers in Environmental Science* 5:11. doi: 10.3389/fenvs.2017.00011.
- Erdozain M, Thompson DG, Porter TM, Kidd KA, Kreutzweiser DP, Sibley PK, Swystun T, Chartrand D, Hajibabaei M (2019) Metabarcoding of storage ethanol vs. conventional morphometric identification in relation to the use of stream macroinvertebrates as ecological indicators in forest management. *Ecological Indicators* 101:173-184. doi: 10.1016/j.ecolind.2019.01.014.
- European Union (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy. *Official Journal of the European Parliament* L327:1-82. doi: 10.1039/ap9842100196.
- Evans NT, Olds BP, Renshaw MA, Turner CR, Li Y, Jerde CL, Mahon AR, Pfrender ME, Lamberti GA, Lodge DM (2016) Quantification of mesocosm fish and amphibian species diversity via environmental DNA metabarcoding. *Molecular Ecology Resources* 16:29-41. doi: 10.1111/1755-0998.12433.
- Evans NT, Shirey PD, Wieringa JG, Mahon AR, Lamberti GA (2017a) Comparative cost and effort of fish distribution detection via environmental DNA analysis and electrofishing. *Fisheries* 42:90-99. doi: 10.1080/03632415.2017.1276329.
- Evans NT, Li Y, Renshaw MA, Olds BP, Deiner K, Turner CR, Jerde CL, Lodge DM, Lamberti GA, Pfrender ME (2017b) Fish community assessment with eDNA metabarcoding: effects of sampling design and bioinformatic filtering. *Canadian Journal of Fisheries and Aquatic Sciences* 74:1362-1374. doi: 10.1139/cjfas-2016-0306.
- Evers CHM, Van den Broek AJM, Buskens R, Van Leerdam A, Knobens RAE, Van Herpen FCJ (2012) Omschrijving MEP en maatlaten voor sloten en kanalen voor de KRW 2015-2021. STOWA rapport 2012-34.
- Ficetola GF, Miaud C, Pompanon F, Taberlet P (2008) Species detection using environmental DNA from water samples. *Biology Letters* 4:423-5. doi: 10.1098/rsbl.2008.0118.

References

- Ficetola GF, Taberlet P, Coissac E (2016) How to limit false positives in environmental DNA and metabarcoding? *Molecular Ecology Resources* 16:604-607. doi: 10.1111/1755-0998.12508.
- Fujii K, Doi H, Matsuoka S, Nagano M, Sato H, Yamanaka H (2019) Environmental DNA metabarcoding for fish community analysis in backwater lakes: A comparison of capture methods. *PLoS ONE* 14:e0210357. doi: 10.1371/journal.pone.0210357.
- Garlapati D, Charankumar B, Ramu K, Madeswaran P, Ramana Murthy M V. (2019) A review on the applications and recent advances in environmental DNA (eDNA) metagenomics. *Reviews in Environmental Science and Biotechnology* 18:389-411. doi: 10.1007/s11157-019-09501-4.
- Geiger MF, Moriniere J, Hausmann A, Haszprunar G, Wägele W, Hebert PDN, Rulik B (2016) Testing the Global Malaise Trap Program - How well does the current barcode reference library identify flying insects in Germany? *Biodiversity Data Journal* 4. doi: 10.3897/BDJ.4.e10671.
- Gibson J, Shokralla S, Porter TM, King I, Van Konynenburg S, Janzen DH, Hallwachs W, Hajibabaei M (2014) Simultaneous assessment of the macrobiome and microbiome in a bulk sample of tropical arthropods through DNA metasytematics. *Proceedings of the National Academy of Sciences* 111:8007-12. doi: 10.1073/pnas.1406468111.
- Gibson JF, Shokralla S, Curry C, Baird DJ, Monk WA, King I, Hajibabaei M (2015) Large-scale biomonitoring of remote and threatened ecosystems via high-throughput sequencing. *PLoS ONE* 10:e0138432. doi: 10.1371/journal.pone.0138432.
- Gleick PH (1993) *Water in Crisis: A Guide to the World's Fresh Water Resources*. New York: Oxford University Press, Inc.
- Goldberg CS, Pilliod DS, Arkle RS, Waits LP (2011) Molecular detection of vertebrates in stream water: a demonstration using Rocky Mountain Tailed Frogs and Idaho Giant Salamanders. *PLoS ONE* 6:e22746. doi: 10.1371/journal.pone.0022746.
- Goldberg CS, Turner CR, Deiner K, Klymus KE, Thomsen PF, Murphy MA, Spear SF, McKee A, Oyler-McCance SJ, Cornman RS, Laramie MB, Mahon AR, Lance RF, Pilliod DS, Strickler KM, Waits LP, Fremier AK, Takahara T, Herder JE, Taberlet P (2016) Critical considerations for the application of environmental DNA methods to detect aquatic species. *Methods in Ecology and Evolution*. doi: 10.1111/2041-210X.12595.
- Goldberg CS, Strickler KM, Fremier AK (2018) Degradation and dispersion limit environmental DNA detection of rare amphibians in wetlands: Increasing efficacy of sampling designs. *Science of the Total Environment* 633:695-703. doi: 10.1016/j.scitotenv.2018.02.295.
- Gosselain V, Descy JP, Everbecq E (1994) The phytoplankton community of the River Meuse, Belgium: seasonal dynamics (year 1992) and the possible incidence of zooplankton grazing. *Hydrobiologia* 289:179-191. doi: 10.1007/BF00007419.

- Grey EK, Bernatchez L, Cassey P, Deiner K, Deveney M, Howland KL, Lacoursière-Roussel A, Leong SCY, Li Y, Olds B, Pfrender ME, Prowse TAA, Renshaw MA, Lodge DM (2018) Effects of sampling effort on biodiversity patterns estimated from environmental DNA metabarcoding surveys. *Scientific Reports* 8:8843. doi: 10.1038/s41598-018-27048-2.
- Guardiola M, Wangenstein OS, Taberlet P, Coissac E, Uriz MJ, Turon X (2016) Spatio-temporal monitoring of deep-sea communities using metabarcoding of sediment DNA and RNA. *PeerJ* 4:e2807. doi: 10.7717/peerj.2807.
- Haase P, Murray-Bligh J, Lohse S, Pauls S, Sundermann A, Gunn R, Clarke R (2006) Assessing the impact of errors in sorting and identifying macroinvertebrate samples. *Hydrobiologia* 566:505-521. doi: 10.1007/s10750-006-0075-6.
- Haase P, Pauls SU, Schindehütte K, Sundermann A (2010) First audit of macroinvertebrate samples from an EU Water Framework Directive monitoring program: human error greatly lowers precision of assessment results. *Journal of the North American Benthological Society* 29:1279-1291.
- Hajibabaei M, Shokralla S, Zhou X, Singer GAC, Baird DJ (2011) Environmental barcoding: a next-generation sequencing approach for biomonitoring applications using river benthos. *PLoS ONE* 6:e17497. doi: 10.1371/journal.pone.0017497.
- Hajibabaei M, Spall JL, Shokralla S, Van Konynenburg S (2012) Assessing biodiversity of a freshwater benthic macroinvertebrate community through non-destructive environmental barcoding of DNA from preservative ethanol. *BMC Ecology* 12:28. doi: 10.1186/1472-6785-12-28.
- Van der Hammen H (1992) *De macrofauna van Noord-Holland*. K.U. Nijmegen.
- Hanashiro FTT, Mukherjee S, Souffreau C, Engelen J, Brans KI, Busschaert P, De Meester L (2019) Freshwater bacterioplankton metacommunity structure along urbanization gradients in Belgium. *Frontiers in Microbiology* 10:743. doi: 10.3389/fmicb.2019.00743.
- Hänfling B, Handley LL, Read DS, Hahn C, Li J, Nichols P, Blackman RC, Oliver A, Winfield IJ (2016) Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Molecular Ecology* 25:3101-3119. doi: 10.1111/mec.13660.
- Harper LR, Lawson Handley L, Hahn C, Boonham N, Rees HC, Gough KC, Lewis E, Adams IP, Brotherton P, Phillips S, Hänfling B (2018) Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested newt (*Triturus cristatus*). *Ecology and Evolution* 8:6330-6341. doi: 10.1002/ece3.4013.
- Harper LR, Buxton AS, Rees HC, Bruce K, Brys R, Halfmaerten D, Read DS, Watson H V, Sayer CD, Jones EP, Priestley V, Mächler E, Múrria C, Garcés-Pastor S, Medupin C, Burgess K, Benson G, Boonham N, Griffiths RA, Lawson Handley L, Hänfling B (2019a) Prospects and challenges of environmental DNA (eDNA) monitoring in freshwater ponds. *Hydrobiologia*

References

826:25-41. doi: 10.1007/s10750-018-3750-5.

Harper LR, Handley LL, Carpenter AI, Murray-Dickson G, Muri C Di, Macgregor CJ, Logan TW, Law A, Breithaupt T, Read DS, McDevitt A, Hänfling B (2019b) Environmental DNA (eDNA) metabarcoding of pond water as a tool to survey conservation and management priority mammals. *bioRxiv*:546218. doi: 10.1101/546218.

Hebert PDN, Cywinska A, Ball SL, DeWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences* 270:313-21. doi: 10.1098/rspb.2002.2218.

Hebert PDN, Ratnasingham S, Zakharov E V., Telfer AC, Levesque-Beaudin V, Milton MA, Pedersen S, Jannetta P, Dewaard JR (2016) Counting animal species with DNA barcodes: Canadian insects. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371:20150333. doi: 10.1098/rstb.2015.0333.

Heisler J, Glibert PM, Burkholder JM, Anderson DM, Cochlan W, Dennison WC, Dortch Q, Gobler CJ, Heil CA, Humphries E, Lewitus A, Magnien R, Marshall HG, Sellner K, Stockwell DA, Stoecker DK, Suddleson M (2008) Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae* 8:3-13. doi: 10.1016/j.hal.2008.08.006.

Hering D, Borja A, Jones JI, Pont D, Boets P, Bouchez A, Bruce K, Drakare S, Hänfling B, Kahlert M, Leese F, Meissner K, Mergen P, Reyjol Y, Segurado P, Vogler A, Kelly M (2018) Implementation options for DNA-based identification into ecological status assessment under the European Water Framework Directive. *Water Research* 138:192-205. doi: 10.1016/j.watres.2018.03.003.

Hinlo R, Gleeson D, Lintermans M, Furlan E (2017) Methods to maximise recovery of environmental DNA from water samples. *PLoS ONE* 12:e0179251. doi: 10.1371/journal.pone.0179251.

Hoborn D (2020) BIOSCAN: DNA barcoding to accelerate taxonomy and biogeography for conservation and sustainability. *Genome*:1-4. doi: 10.1139/gen-2020-0009.

Hopkins GW, Freckleton RP (2002) Declines in the numbers of amateur and professional taxonomists: implications for conservation. *Animal Conservation* 5:245-249. doi: 10.1017/S1367943002002299.

Hubbell SP (2001) *The unified neutral theory of biodiversity and biogeography*. Princeton University Press.

Huemer P, Mutanen M, Sefc KM, Hebert PDN (2014) Testing DNA barcode performance in 1000 species of European Lepidoptera: Large geographic distances have small genetic impacts. *PLoS ONE* 9:e115774. doi: 10.1371/journal.pone.0115774.

Huson DH, Auch AF, Qi J, Schuster SC (2007) MEGAN analysis of metagenomic data.

Genome Research 17:377-86. doi: 10.1101/gr.5969107.

Jackson RB, Carpenter SR, Dahm CN, McKnight DM, Naiman RJ, Postel SL, Running SW (2001) Water in a changing world. *Ecological Applications* 11:1027-1045. doi: 10.1890/1051-0761(2001)011[1027:WIACW]2.0.CO;2.

Jackson JK, Battle JM, White BP, Pilgrim EM, Stein ED, Miller PE, Sweeney BW (2014) Cryptic biodiversity in streams: a comparison of macroinvertebrate communities based on morphological and DNA barcode identifications. *Freshwater Science* 33:312-324. doi: 10.1086/675225.

Jane SF, Wilcox TM, Mckelvey KS, Young MK, Schwartz MK, Lowe WH, Letcher BH, Whiteley AR (2015) Distance, flow and PCR inhibition: eDNA dynamics in two headwater streams. *Molecular Ecology Resources* 15:216-227. doi: 10.1111/1755-0998.12285.

Jerde CL, Mahon AR, Chadderton WL, Lodge DM (2011) "Sight-unseen" detection of rare aquatic species using environmental DNA. *Conservation Letters* 4:150-157. doi: 10.1111/j.1755-263X.2010.00158.x.

Jeunen GJ, Knapp M, Spencer HG, Taylor HR, Lamare MD, Stat M, Bunce M, Gemmell NJ (2019) Species-level biodiversity assessment using marine environmental DNA metabarcoding requires protocol optimization and standardization. *Ecology and Evolution* 9:1323-1335. doi: 10.1002/ece3.4843.

Jo T, Murakami H, Yamamoto S, Masuda R, Minamoto T (2019) Effect of water temperature and fish biomass on environmental DNA shedding, degradation, and size distribution. *Ecology and Evolution* 9:1135-1146. doi: 10.1002/ece3.4802.

Jones FC (2008) Taxonomic sufficiency: The influence of taxonomic resolution on freshwater bioassessments using benthic macroinvertebrates. *Environmental Reviews* 16:45-69. doi: 10.1139/A07-010.

Joshi NA, Fass JN (2011) Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files (Version 1.33). Available at <https://github.com/najoshi/sickle>.

Karr JR (1999) Defining and measuring river health. *Freshwater Biology* 41:221-234. doi: 10.1046/j.1365-2427.1999.00427.x.

Keeley N, Wood SA, Pochon X (2018) Development and preliminary validation of a multi-trophic metabarcoding biotic index for monitoring benthic organic enrichment. *Ecological Indicators* 85:1044-1057. doi: 10.1016/j.ecolind.2017.11.014.

Kelly RP, Port JA, Yamahara KM, Martone RG, Lowell N, Thomsen PF, Mach ME, Bennett M, Prahler E, Caldwell MR, Crowder LB (2014) Environmental monitoring. Harnessing DNA to improve environmental management. *Science* 344:1455-6. doi: 10.1126/science.1251156.

References

- Kelly RP, Closek CJ, O'Donnell JL, Kralj JE, Shelton AO, Samhouri JF (2017) Genetic and manual survey methods yield different and complementary views of an ecosystem. *Frontiers in Marine Science* 3:283. doi: 10.3389/FMARS.2016.00283.
- Kenney MA, Sutton-Grier AE, Smith RF, Gresens SE (2009) Benthic macroinvertebrates as indicators of water quality: The intersection of science and policy. *Terrestrial Arthropod Reviews* 2:99-128. doi: 10.1163/187498209X12525675906077.
- Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO (2013) Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research* 41:e1. doi: 10.1093/nar/gks808.
- Klymus KE, Richter CA, Chapman DC, Paukert C (2015) Quantification of eDNA shedding rates from invasive bighead carp *Hypophthalmichthys nobilis* and silver carp *Hypophthalmichthys molitrix*. *Biological Conservation* 183:77-84. doi: 10.1016/j.biocon.2014.11.020.
- Klymus KE, Marshall NT, Stepien CA (2017) Environmental DNA (eDNA) metabarcoding assays to detect invasive invertebrate species in the Great Lakes. *PLoS ONE* 12:e0177643. doi: 10.1371/journal.pone.0177643.
- Krol L, Van der Hoorn B, Gorsich EE, Trimbos K, Van Bodegom PM, Schrama M (2019) How does eDNA compare to traditional trapping? Detecting mosquito communities in South-African freshwater ponds. *Frontiers in Ecology and Evolution* 7:260. doi: 10.3389/fevo.2019.00260.
- Kvist S (2013) Barcoding in the dark?: A critical view of the sufficiency of zoological DNA barcoding databases and a plea for broader integration of taxonomic knowledge. *Molecular Phylogenetics and Evolution* 69:39-45. doi: 10.1016/j.ympev.2013.05.012.
- Lacoursière-Roussel A, Côté G, Leclerc V, Bernatchez L (2015) Quantifying relative fish abundance with eDNA: a promising tool for fisheries management. *Journal of Applied Ecology*. doi: 10.1111/1365-2664.12598.
- Lacoursière-Roussel A, Howland K, Normandeau E, Grey EK, Archambault P, Deiner K, Lodge DM, Hernandez C, Leduc N, Bernatchez L (2018) eDNA metabarcoding as a new surveillance approach for coastal Arctic biodiversity. *Ecology and Evolution* 8. doi: 10.1002/ece3.4213.
- Langer-Jaesrich M, Köhler HR, Gerhardt A (2010) Assessing toxicity of the insecticide thiacloprid on *Chironomus riparius* (Insecta: Diptera) using multiple end points. *Archives of Environmental Contamination and Toxicology* 58:963-972. doi: 10.1007/s00244-009-9420-x.
- Lanzén A, Lekang K, Jonassen I, Thompson EM, Troedsson C (2017) DNA extraction replicates improve diversity and compositional dissimilarity in metabarcoding of eukaryotes in marine sediments. *PLoS ONE* 12:e0179443. doi: 10.1371/journal.pone.0179443.

- Laroche O, Wood SA, Tremblay LA, Ellis JI, Lear G, Pochon X (2018) A cross-taxa study using environmental DNA/RNA metabarcoding to measure biological impacts of offshore oil and gas drilling and production operations. *Marine Pollution Bulletin* 127:97-107. doi: 10.1016/j.marpolbul.2017.11.042.
- Larsson AJM, Stanley G, Sinha R, Weissman IL, Sandberg R (2018) Computational correction of index switching in multiplexed sequencing libraries. *Nature Methods* 15:305-307. doi: 10.1038/nmeth.4666.
- Lawson Handley L, Read DS, Winfield IJ, Kimbell H, Johnson H, Li J, Hahn C, Blackman R, Wilcox R, Donnelly R, Szitenberg A, Hänfling B (2019) Temporal and spatial variation in distribution of fish environmental DNA in England's largest lake. *Environmental DNA* 1:26-39. doi: 10.1002/edn3.5.
- Leese F, Altermatt F, Bouchez A, Ekrem T, Hering D, Meissner K, Mergen P, Pawlowski J, Piggott J, Rimet F, Steinke D, Taberlet P, Weigand A, Abarenkov K, Beja P, Bervoets L, Björnsdóttir S, Boets P, Boggero A, Bones A, Borja Á, Bruce K, Bursić V, Carlsson J, Čiampor F, Čiamporová-Zatovičová Z, Coissac E, Costa F, Costache M, Creer S, Csabai Z, Deiner K, DelValls Á, Drakare S, Duarte S, Eleršek T, Fazi S, Fišer C, Flot J-F, Fonseca V, Fontaneto D, Grabowski M, Graf W, Guðbrandsson J, Hellström M, Hershkovitz Y, Hollingsworth P, Japoshvili B, Jones J, Kahlert M, Kalamujic Stroil B, Kasapidis P, Kelly M, Kelly-Quinn M, Keskin E, Kõljalg U, Ljubešić Z, Maček I, Mächler E, Mahon A, Marečková M, Mejdandzic M, Mircheva G, Montagna M, Moritz C, Mulk V, Naumoski A, Navodaru I, Padišák J, Pálsson S, Panksep K, Penev L, Petrusek A, Pfannkuchen M, Primmer C, Rinkevich B, Rotter A, Schmidt-Kloiber A, Segurado P, Speksnijder A, Stoev P, Strand M, Šulčius S, Sundberg P, Traugott M, Tsigenopoulos C, Turon X, Valentini A, Van der Hoorn B, Várбірó G, Vasquez Hadjilyra M, Viguri J, Vitonyté I, Vogler A, Vrålstad T, Wägele W, Wenne R, Winding A, Woodward G, Zegura B, Zimmermann J (2016) DNAqua-Net: Developing new genetic tools for bioassessment and monitoring of aquatic ecosystems in Europe. *Research Ideas and Outcomes* 2:e11321. doi: 10.3897/rio.2.e11321.
- Leray M, Yang JY, Meyer CP, Mills SC, Agudelo N, Ranwez V, Boehm JT, Machida RJ (2013) A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in Zoology* 10:34. doi: 10.1186/1742-9994-10-34.
- Levi T, Allen JM, Bell D, Joyce J, Russell JR, Tallmon DA, Vulstek SC, Yang C, Yu DW (2019) Environmental DNA for the enumeration and management of Pacific salmon. *Molecular Ecology Resources* 19:597-608. doi: 10.1111/1755-0998.12987.
- Li J, Lawson Handley LJ, Read DS, Hänfling B (2018a) The effect of filtration method on the efficiency of environmental DNA capture and quantification via metabarcoding. *Molecular Ecology Resources* 18:1102-1114. doi: 10.1111/1755-0998.12899.
- Li F, Peng Y, Fang W, Altermatt F, Xie Y, Yang J, Zhang X (2018b) Application of environmental DNA metabarcoding for predicting anthropogenic pollution in rivers. *Environmental Science*

References

and Technology 52:11708-11719. doi: 10.1021/acs.est.8b03869.

Lim NKM, Tay YC, Srivathsan A, Tan JWT, Kwik JTB, Baloglu B, Meier R, Yeo DCJ (2016) Next-generation freshwater bioassessment: eDNA metabarcoding with a conserved metazoan primer reveals species-rich and reservoir-specific communities. *Royal Society Open Science* 3:160635. doi: 10.1098/rsos.160635.

Lobo J, Shokralla S, Costa MH, Hajibabaei M, Costa FO (2017) DNA metabarcoding for high-throughput monitoring of estuarine macrobenthic communities. *Scientific Reports* 7:15618. doi: 10.1038/s41598-017-15823-6.

Lodge DM, Turner CR, Jerde CL, Barnes MA, Chadderton L, Egan SP, Feder JL, Mahon AR, Pfrender ME (2012) Conservation in a cup of water: estimating biodiversity and population abundance from environmental DNA. *Molecular Ecology* 21:2555-8. doi: 10.1111/j.1365-294X.2012.05600.x.

Lugg WH, Griffiths J, Van Rooyen AR, Weeks AR, Tingley R (2018) Optimal survey designs for environmental DNA sampling. *Methods in Ecology and Evolution* 9:1049-1059. doi: 10.1111/2041-210X.12951.

Macher JN, Salis RK, Blakemore KS, Tollrian R, Matthaei CD, Leese F (2016) Multiple-stressor effects on stream invertebrates: DNA barcoding reveals contrasting responses of cryptic mayfly species. *Ecological Indicators* 61:159-169. doi: 10.1016/j.ecolind.2015.08.024.

Macher JN, Vivancos A, Piggott JJ, Centeno FC, Matthaei CD, Leese F (2018) Comparison of environmental DNA and bulk-sample metabarcoding using highly degenerate cytochrome c oxidase I primers. *Molecular Ecology Resources* 18:1456-1468. doi: 10.1111/1755-0998.12940.

Mächler E, Deiner K, Steinmann P, Altermatt F (2014) Utility of environmental DNA for monitoring rare and indicator macroinvertebrate species. *Freshwater Science* 33:1174-1183.

Mächler E, Deiner K, Spahn F, Altermatt F (2016) Fishing in the water: effect of sampled water volume on environmental DNA-based detection of macroinvertebrates. *Environmental Science and Technology* 50:305-312. doi: 10.1021/acs.est.5b04188.

Magoč T, Salzberg SL (2011) FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27:2957-2963. doi: 10.1093/bioinformatics/btr507.

Majaneva M, Diserud OH, Eagle SHC, Boström E, Hajibabaei M, Ekrem T (2018) Environmental DNA filtration techniques affect recovered biodiversity. *Scientific Reports* 8:4682. doi: 10.1038/s41598-018-23052-8.

Marshall JC, Steward AL, Harch BD (2006) Taxonomic resolution and quantification of freshwater macroinvertebrate samples from an Australian dryland river: the benefits and costs of using species abundance data. *Hydrobiologia* 572:171-194. doi: 10.1007/s10750-005-9007-0.

- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17:10. doi: 10.14806/ej.17.1.200.
- Matson PA, Parton WJ, Power AG, Swift MJ (1997) Agricultural intensification and ecosystem properties. *Science* 277:504-509. doi: 10.1126/science.277.5325.504.
- Matsuhashi S, Doi H, Fujiwara A, Watanabe S, Minamoto T (2016) Evaluation of the environmental DNA method for estimating distribution and biomass of submerged aquatic plants. *PLoS ONE* 11:e0156217. doi: 10.1371/journal.pone.0156217.
- Merkes CM, McCalla SG, Jensen NR, Gaikowski MP, Amberg JJ (2014) Persistence of DNA in carcasses, slime and avian feces may affect interpretation of environmental DNA data. *PLoS ONE* 9:e113346. doi: 10.1371/journal.pone.0113346.
- Meusnier I, Singer GAC, Landry J-F, Hickey DA, Hebert PDN, Hajibabaei M (2008) A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics* 9:214. doi: 10.1186/1471-2164-9-214.
- Miya M, Sato Y, Fukunaga T, Sado T, Poulsen JY, Sato K, Minamoto T, Yamamoto S, Yamanaka H, Araki H, Kondoh M, Iwasaki W (2015) MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species. *Royal Society Open Science* 2:150088. doi: 10.1098/rsos.150088.
- Van der Molen D, Pot R, Evers C, Van Herpen F, Van Nieuwerburgh L (2016) Referenties en maatlatten voor natuurlijke watertypen voor de KRW 2015-2021. STOWA rapport 2012-31.
- Montagud D, Soria JM, Soria-Perpiñà X, Alfonso T, Vicente E (2018) A comparative study of four indexes based on zooplankton as trophic state indicators in reservoirs. *Limnetica* 38. doi: 10.23818/limn.38.06.
- Morrissey CA, Mineau P, Devries JH, Sanchez-Bayo F, Liess M, Cavallaro MC, Liber K (2015) Neonicotinoid contamination of global surface waters and associated risk to aquatic invertebrates: A review. *Environment International* 74:291-303. doi: 10.1016/j.envint.2014.10.024.
- Moushomi R, Wilgar G, Carvalho G, Creer S, Seymour M (2019) Environmental DNA size sorting and degradation experiment indicates the state of *Daphnia magna* mitochondrial and nuclear eDNA is subcellular. *Scientific Reports* 9:12500. doi: 10.1038/s41598-019-48984-7.
- Moyer GR, Díaz-Ferguson E, Hill JE, Shea C (2014) Assessing environmental DNA detection in controlled lentic systems. *PLoS ONE* 9:e103767. doi: 10.1371/journal.pone.0103767.
- Nicholson A, McIsaac D, MacDonald C, Gec P, Mason BE, Rein W, Wrobel J, De Boer M, Milián-García Y, Hanner RH (2020) An analysis of metadata reporting in freshwater environmental DNA research calls for the development of best practice guidelines. *Environmental DNA*. doi: 10.1002/edn3.81.

References

- O'Donnell JL, Kelly RP, Shelton AO, Samhoury JF, Lowell NC, Williams GD (2017) Spatial distribution of environmental DNA in a nearshore marine habitat. *PeerJ* 5:e3044. doi: 10.7717/peerj.3044.
- Oksanen J, Kindt R, Legendre P, O'Hara R (2007) *VEGAN: Community Ecology Package*. R package version 2.5-5, <https://www.rdocumentation.org/packages/vegan>.
- Olds BP, Jerde CL, Renshaw MA, Li Y, Evans NT, Turner CR, Deiner K, Mahon AR, Brueseke MA, Shirey PD, Pfrender ME, Lodge DM, Lamberti GA (2016) Estimating species richness using environmental DNA. *Ecology and Evolution*. doi: 10.1002/ece3.2186.
- Pauls SU, Alp M, Bálint M, Bernabò P, Čiampor F, Čiamporová-Zaťovičová Z, Finn DS, Kohout J, Leese F, Lencioni V, Paz-Vinas I, Monaghan MT (2014) Integrating molecular tools into freshwater ecology: developments and opportunities. *Freshwater Biology* 59:1559-1576. doi: 10.1111/fwb.12381.
- Pawlowski J, Kelly-Quinn M, Altermatt F, Apothéloz-Perret-Gentil L, Beja P, Boggero A, Borja A, Bouchez A, Cordier T, Domaizon I, Feio MJ, Filipe AF, Fornaroli R, Graf W, Herder J, Van der Hoorn B, Iwan Jones J, Sagova-Mareckova M, Moritz C, Barquín J, Piggott JJ, Pinna M, Rimet F, Rinkevich B, Sousa-Santos C, Specchia V, Trobajo R, Vasselon V, Vitecek S, Zimmerman J, Weigand A, Leese F, Kahlert M (2018) The future of biotic indices in the ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems. *Science of the Total Environment* 637-638:1295-1310. doi: 10.1016/j.scitotenv.2018.05.002.
- Pawluczyk M, Weiss J, Links MG, Egaña Aranguren M, Wilkinson MD, Egea-Cortines M, Egaña Aranguren M, Wilkinson MD, Egea-Cortines M (2015) Quantitative evaluation of bias in PCR amplification and next-generation sequencing derived from metabarcoding samples. *Analytical and Bioanalytical Chemistry* 407:1841-1848. doi: 10.1007/s00216-014-8435-y.
- Pereira AS, Dâmaso-Rodrigues ML, Amorim A, Daam MA, Cerejeira MJ (2018) Aquatic community structure in Mediterranean edge-of-field waterbodies as explained by environmental factors and the presence of pesticide mixtures. *Ecotoxicology* 27:661-674. doi: 10.1007/s10646-018-1944-2.
- Piggott JJ, Townsend CR, Matthaei CD (2015) Reconceptualizing synergism and antagonism among multiple stressors. *Ecology and Evolution* 5:1538-1547. doi: 10.1002/ece3.1465.
- Pilliod DS, Goldberg CS, Arkle RS, Waits LP, Richardson J (2013) Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples. *Canadian Journal of Fisheries and Aquatic Sciences* 70:1123-1130. doi: 10.1139/cjfas-2013-0047.
- Pilliod DS, Goldberg CS, Arkle RS, Waits LP (2014) Factors influencing detection of eDNA from a stream-dwelling amphibian. *Molecular Ecology Resources* 14:109-116. doi: 10.1111/1755-0998.12159.

- Pimm SL, Jenkins CN, Abell R, Brooks TM, Gittleman JL, Joppa LN, Raven PH, Roberts CM, Sexton JO (2014) The biodiversity of species and their rates of extinction, distribution, and protection. *Science* 344:1246752. doi: 10.1126/science.1246752.
- Piñol J, Mir G, Gomez-Polo P, Agustí N (2015) Universal and blocking primer mismatches limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of arthropods. *Molecular Ecology Resources* 15:819-30. doi: 10.1111/1755-0998.12355.
- Pisa LW, Amaral-Rogers V, Belzunces LP, Bonmatin JM, Downs CA, Goulson D, Kreutzweiser DP, Krupke C, Liess M, Mcfield M, Morrissey CA, Noome DA, Settele J, Simon-Delso N, Stark JD, Van Der Sluijs JP, Van Dyck H, Wiemers M (2014) Effects of neonicotinoids and fipronil on non-target invertebrates. *Environmental Science and Pollution Research* 22:68-102. doi: 10.1007/s11356-014-3471-x.
- Pochon X, Wood SA, Keeley NB, Lejzerowicz F, Esling P, Drew J, Pawlowski J (2015) Accurate assessment of the impact of salmon farming on benthic sediment enrichment using foraminiferal metabarcoding. *Marine Pollution Bulletin* 100:370-382. doi: 10.1016/j.marpolbul.2015.08.022.
- Pochon X, Zaiko A, Fletcher LM, Laroche O, Wood SA (2017) Wanted dead or alive? Using metabarcoding of environmental DNA and RNA to distinguish living assemblages for biosecurity applications. *PLoS ONE* 12:e0187636. doi: 10.1371/journal.pone.0187636.
- Pompanon F, Deagle BE, Symondson WOC, Brown DS, Jarman SN, Taberlet P (2012) Who is eating what: diet assessment using next generation sequencing. *Molecular Ecology* 21:1931-50. doi: 10.1111/j.1365-294X.2011.05403.x.
- Pont D, Rocle M, Valentini A, Civade R, Jean P, Maire A, Roset N, Schabuss M, Zornig H, Dejean T (2018) Environmental DNA reveals quantitative patterns of fish biodiversity in large rivers despite its downstream transportation. *Scientific Reports* 8:10361. doi: 10.1038/s41598-018-28424-8.
- Porter TM, Hajibabaei M (2018) Automated high throughput animal CO1 metabarcode classification. *Scientific Reports* 8:219675. doi: 10.1038/s41598-018-22505-4.
- Pot R (2015) QBWAT, programma voor beoordeling van de biologische waterkwaliteit volgens de Nederlandse maatlaten voor de Kaderrichtlijn Water, versie 5.33.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO (2013) The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research* 41:D590-D596. doi: 10.1093/nar/gks1219.
- Raby M, Nowierski M, Perlov D, Zhao X, Hao C, Poirier DG, Sibley PK (2018) Acute toxicity of 6 neonicotinoid insecticides to freshwater invertebrates. *Environmental Toxicology and Chemistry* 37:1430-1445. doi: 10.1002/etc.4088.

References

- Rapport DJ (1989) What constitutes ecosystem health? *Perspectives in Biology and Medicine* 33:120-132. doi: 10.1353/pbm.1990.0004.
- Ratnasingham S, Hebert PDN (2007) BOLD: the Barcode of Life Data System (www.barcodinglife.org). *Molecular Ecology Notes* 7:355-364. doi: 10.1111/j.1471-8286.2006.01678.x.
- Rees HC, Baker CA, Gardner DS, Maddison BC, Gough KC (2017) The detection of great crested newts year round via environmental DNA analysis. *BMC Research Notes* 10:327. doi: 10.1186/s13104-017-2657-y.
- Regier HA (1993) The notion of natural and cultural integrity. In: Woodley S, Kay J eds. *Ecological Integrity and the Management of Ecosystems*. Delray Beach, FL: St Lucie Press, 3-18.
- Renshaw MA, Olds BP, Jerde CL, Mcveigh MM, Lodge DM (2015) The room temperature preservation of filtered environmental DNA samples and assimilation into a phenol-chloroform-isoamyl alcohol DNA extraction. *Molecular Ecology Resources* 15:168-176. doi: 10.1111/1755-0998.12281.
- Rockström J, Steffen W, Noone K, Persson A, Chapin FSI, Lambin E, Lenton TM, Scheffer M, Folke C, Schellnhuber HJ, Nykvist B, Wit CA de, Hughes T, Van der Leeuw S, Rodhe H, Sörlin S, Snyder PK, Costanza R, Svedin U, Falkenmark M, Karlberg L, Corell RW, Fabry VJ, Hansen J, Walker B, Liverman D, Richardson K, Crutzen P, Foley J (2009) Planetary boundaries: exploring the safe operating space for humanity. *Ecology and Society* 14:32.
- Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584. doi: 10.7717/peerj.2584.
- Roussel J-M, Paillisson J-M, Tréguier A, Petit E (2015) The downside of eDNA as a survey tool in water bodies. *Journal of Applied Ecology* 52:n/a-n/a. doi: 10.1111/1365-2664.12428.
- RStudio (2015) RStudio: Integrated development environment for R (Version 0.99.902).
- Sánchez-Bayo F, Goka K, Hayasaka D (2016) Contamination of the aquatic environment with neonicotinoids and its implication for ecosystems. *Frontiers in Environmental Science* 4:71. doi: 10.3389/fenvs.2016.00071.
- Sansom BJ, Sassoubre LM (2017) Environmental DNA (eDNA) shedding and decay rates to model freshwater mussel eDNA transport in a river. *Environmental Science and Technology* 51:14244-14253. doi: 10.1021/acs.est.7b05199.
- Schmidt-Kloiber A, Hering D (2015) www.freshwaterecology.info - An online tool that unifies, standardises and codifies more than 20,000 European freshwater organisms and their ecological preferences. *Ecological Indicators* 53:271-282. doi: 10.1016/j.ecolind.2015.02.007.
- Schmidt BR, Kéry M, Ursenbacher S, Hyman OJ, Collins JP (2013) Site occupancy models in

- the analysis of environmental DNA presence/absence surveys: A case study of an emerging amphibian pathogen. *Methods in Ecology and Evolution* 4:646-653. doi: 10.1111/2041-210X.12052.
- Schmieder R, Edwards R (2011) Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863-864. doi: 10.1093/bioinformatics/btr026.
- Schneider J, Valentini A, Dejean T, Montarsi F, Taberlet P, Glaizot O, Fumagalli L (2016) Detection of invasive mosquito vectors using environmental DNA (eDNA) from water samples. *PLoS ONE* 11:e0162493. doi: 10.1371/journal.pone.0162493.
- Schrama M, Barmantlo SH, Hunting ER, Van Logtestijn RSP, Vijver MG, Van Bodegom PM (2017) Pressure-induced shifts in trophic linkages in a simplified aquatic food web. *Frontiers in Environmental Science* 5:75. doi: 10.3389/fenvs.2017.00075.
- Schreiner VC, Szöcs E, Bhowmik AK, Vijver MG, Schäfer RB (2016) Pesticide mixtures in streams of several European countries and the USA. *Science of the Total Environment* 573:680-689. doi: 10.1016/j.scitotenv.2016.08.163.
- Schwarzenbach RP, Escher BI, Fenner K, Hofstetter TB, Johnson CA, Von Gunten U, Wehrli B (2006) The challenge of micropollutants in aquatic systems. *Science* 313:1072-1077. doi: 10.1126/science.1127291.
- Scrimgeour GJ, Wicklum D (1996) Aquatic ecosystem health and integrity: problems and potential solutions. *Journal of the North American Benthological Society* 15:254. doi: 10.2307/1467951.
- Seymour JR, Amin SA, Raina JB, Stocker R (2017) Zooming in on the phycosphere: The ecological interface for phytoplankton-bacteria relationships. *Nature Microbiology* 2:17065. doi: 10.1038/nmicrobiol.2017.65.
- Shaw JLA, Clarke LJ, Wedderburn SD, Barnes TC, Weyrich LS, Cooper A (2016) Comparison of environmental DNA metabarcoding and conventional fish survey methods in a river system. *Biological Conservation* 197:131-138. doi: 10.1016/j.biocon.2016.03.010.
- Shokralla S, Singer GAC, Hajibabaei M (2010) Direct PCR amplification and sequencing of specimens' DNA from preservative ethanol. *BioTechniques* 48:233-4. doi: 10.2144/000113362.
- Sigsgaard EE, Carl H, Møller PR, Thomsen PF (2015) Monitoring the near-extinct European weather loach in Denmark based on environmental DNA from water samples. *Biological Conservation* 183:46-52. doi: 10.1016/j.biocon.2014.11.023.
- Sigsgaard EE, Nielsen IB, Carl H, Krag MA, Knudsen SW, Xing Y, Holm-Hansen TH, Møller PR, Thomsen PF (2017) Seawater environmental DNA reflects seasonality of a coastal fish community. *Marine Biology* 164:128. doi: 10.1007/s00227-017-3147-4.

References

- Sigsgaard EE, Jensen MR, Winkelmann IE, Møller PR, Hansen MM, Thomsen PF (2020) Population-level inferences from environmental DNA - Current status and future perspectives. *Evolutionary Applications* 13:245-262. doi: 10.1111/eva.12882.
- Smart AS, Tingley R, Weeks AR, Van Rooyen AR, McCarthy MA (2015) Environmental DNA sampling is more sensitive than a traditional survey technique for detecting an aquatic invader. *Ecological Applications* 25:1944-1952. doi: 10.1890/14-1751.1.
- Smart AS, Weeks AR, Van Rooyen AR, Moore A, McCarthy MA, Tingley R (2016) Assessing the cost-efficiency of environmental DNA sampling. *Methods in Ecology and Evolution* 7:1291-1298. doi: 10.1111/2041-210X.12598.
- Song H, Buhay JE, Whiting MF, Crandall KA (2008) Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proceedings of the National Academy of Sciences* 105:13486-13491. doi: 10.1073/pnas.0803076105.
- De Souza LS, Godwin JC, Renshaw MA, Larson E (2016) Environmental DNA (eDNA) detection probability is influenced by seasonal activity of organisms. *PLoS ONE* 11:e0165273. doi: 10.1371/journal.pone.0165273.
- Spear MJ, Embke HS, Krysan PJ, Vander Zanden MJ (2020) Application of eDNA as a tool for assessing fish population abundance. *Environmental DNA:edn*3.94. doi: 10.1002/edn3.94.
- Spear SF, Groves JD, Williams LA, Waits LP (2015) Using environmental DNA methods to improve detectability in a hellbender (*Cryptobranchus alleganiensis*) monitoring program. *Biological Conservation* 183:38-45. doi: 10.1016/j.biocon.2014.11.016.
- Spens J, Evans AR, Halfmaerten D, Knudsen SW, Sengupta ME, Mak SST, Sigsgaard EE, Hellström M (2017) Comparison of capture and storage methods for aqueous microbial eDNA using an optimized extraction protocol: advantage of enclosed filter. *Methods in Ecology and Evolution* 8:635-645. doi: 10.1111/2041-210X.12683.
- Šporka F, Vlek HE, Bulánková E, Krno I (2006) Influence of seasonal variation on bioassessment of streams using macroinvertebrates. *Hydrobiologia* 566:543-555. doi: 10.1007/s10750-006-0073-8.
- Stat M, John J, DiBattista JD, Newman SJ, Bunce M, Harvey ES (2019) Combined use of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity. *Conservation Biology* 33:196-205. doi: 10.1111/cobi.13183.
- Stein ED, Martinez MC, Stiles S, Miller PE, Zakharov E V (2014) Is DNA barcoding actually cheaper and faster than traditional morphological methods: results from a survey of freshwater bioassessment efforts in the United States? *PLoS ONE* 9:e95525. doi: 10.1371/journal.pone.0095525.

- Stewart K, Ma H, Zheng J, Zhao J (2017) Using environmental DNA to assess population-wide spatiotemporal reserve use. *Conservation Biology* 31:1173-1182. doi: 10.1111/cobi.12910.
- Stewart KA (2019) Understanding the effects of biotic and abiotic factors on sources of aquatic environmental DNA. *Biodiversity and Conservation* 28:983-1001. doi: 10.1007/s10531-019-01709-8.
- Stoeck T, Frühe L, Forster D, Cordier T, Martins CIM, Pawlowski J (2018) Environmental DNA metabarcoding of benthic bacterial communities indicates the benthic footprint of salmon aquaculture. *Marine Pollution Bulletin* 127:139-149. doi: 10.1016/j.marpolbul.2017.11.065.
- Stoeckle MY, Soboleva L, Charlop-Powers Z (2017) Aquatic environmental DNA detects seasonal fish abundance and habitat preference in an urban estuary. *PLoS ONE* 12:e0175186. doi: 10.1371/journal.pone.0175186.
- STOWA (2014) *Handboek Hydrobiologie. Biologisch onderzoek voor de ecologische beoordeling van Nederlandse zoete en brakke oppervlaktewateren. Deels aangepaste versie.*
- Strayer DL (2006) Challenges for freshwater invertebrate conservation. *Journal of the North American Benthological Society* 25:271-287. doi:10.1899/0887-3593(2006)25[271:CFFIC]2.0.CO;2.
- Strayer DL, Dudgeon D (2010) Freshwater biodiversity conservation: recent progress and future challenges.
- Stribling JB, Pavlik KL, Holdsworth SM, Leppo EW (2008) Data quality, performance, and uncertainty in taxonomic identification for biological assessments. *Journal of the North American Benthological Society* 27:906-919. doi: 10.1899/07-175.1.
- Strickler KM, Fremier AK, Goldberg CS (2015) Quantifying effects of UV-B, temperature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation* 183:85-92. doi: 10.1016/j.biocon.2014.11.038.
- Suikkanen S, Pulina S, Engström-Öst J, Lehtiniemi M, Lehtinen S, Brutemark A (2013) Climate change and eutrophication induced shifts in northern summer plankton communities. *PLoS ONE* 8:e66475. doi: 10.1371/journal.pone.0066475.
- Sumon KA, Ritika AK, Peeters ETHM, Rashid H, Bosma RH, Rahman MS, Fatema MK, Van den Brink PJ (2018) Effects of imidacloprid on the ecology of sub-tropical freshwater microcosms. *Environmental Pollution* 236:432-441. doi: 10.1016/j.envpol.2018.01.102.
- Suter GW (1993) A critique of ecosystem health concepts and indexes. *Environmental Toxicology and Chemistry* 12:1533-1539. doi: 10.1002/etc.5620120903.
- Sweeney BW, Battle JM, Jackson JK, Dapkey T (2011) Can DNA barcodes of stream macroinvertebrates improve descriptions of community structure and water quality? *Journal of the North American Benthological Society* 30:195-216. doi: 10.1899/10-016.1.

References

- Sworobowicz L, Grabowski M, Mamos T, Burzyński A, Kilikowska A, Sell J, Wysocka A (2015) Revisiting the phylogeography of *Asellus aquaticus* in Europe: insights into cryptic diversity and spatiotemporal diversification. *Freshwater Biology* 60:1824-1840. doi: 10.1111/fwb.12613.
- Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012a) Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* 21:2045-50. doi: 10.1111/j.1365-294X.2012.05470.x.
- Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH (2012b) Environmental DNA. *Molecular Ecology* 21:1789-93. doi: 10.1111/j.1365-294X.2012.05542.x.
- Takahara T, Minamoto T, Doi H (2013) Using environmental DNA to estimate the distribution of an invasive fish species in ponds. *PLoS ONE* 8:e56584. doi: 10.1371/journal.pone.0056584.
- Thomas AC, Nguyen PL, Howard J, Goldberg CS (2019) A self-preserving, partially biodegradable eDNA filter. *Methods in Ecology and Evolution*:2041-210X.13212. doi: 10.1111/2041-210X.13212.
- Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E (2012a) Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS ONE* 7:e41732. doi: 10.1371/journal.pone.0041732.
- Thomsen PF, Kielgast J, Iversen LL, Wiuf C, Rasmussen M, Gilbert MTP, Orlando L, Willerslev E (2012b) Monitoring endangered freshwater biodiversity using environmental DNA. *Molecular Ecology* 21:2565-73. doi: 10.1111/j.1365-294X.2011.05418.x.
- Thomsen PF, Willerslev E (2015) Environmental DNA - An emerging tool in conservation for monitoring past and present biodiversity. *Biological Conservation* 183:4-18. doi: 10.1016/j.biocon.2014.11.019.
- Tréguier A, Paillisson J-M, Dejean T, Valentini A, Schlaepfer MA, Roussel J-M (2014) Environmental DNA surveillance for invertebrate species: advantages and technical limitations to detect invasive crayfish *Procambarus clarkii* in freshwater ponds. *Journal of Applied Ecology* 51:871-879. doi: 10.1111/1365-2664.12262.
- Turner CR, Barnes MA, Xu CCY, Jones SE, Jerde CL, Lodge DM (2014a) Particle size distribution and optimal capture of aqueous microbial eDNA. *Methods in Ecology and Evolution* 5:676-684. doi: 10.1111/2041-210X.12206.
- Turner CR, Miller DJ, Coyne KJ, Corush J (2014b) Improved methods for capture, extraction, and quantitative assay of environmental DNA from Asian bigheaded carp (*Hypophthalmichthys* spp.). *PLoS ONE* 9:e114329. doi: 10.1371/journal.pone.0114329.
- Turner CR, Uy KL, Everhart RC (2015) Fish environmental DNA is more concentrated in aquatic sediments than surface water. *Biological Conservation* 183:93-102. doi: 10.1016/j.

biocon.2014.11.017.

Ushio M, Murakami H, Masuda R, Sado T, Miya M, Sakurai S, Yamanaka H, Minamoto T, Kondoh M (2018) Quantitative monitoring of multispecies fish environmental DNA using high-throughput sequencing. *Metabarcoding and Metagenomics* 2:1-15. doi: 10.1101/113472.

Uthicke S, Lamare M, Doyle JR (2018) eDNA detection of corallivorous seastar (*Acanthaster cf. solaris*) outbreaks on the Great Barrier Reef using digital droplet PCR. *Coral Reefs* 37:1229-1239. doi: 10.1007/s00338-018-1734-6.

Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, Bellemain E, Besnard A, Coissac E, Boyer F, Gaboriaud C, Jean P, Poulet N, Roset N, Copp GH, Geniez P, Pont D, Argillier C, Baudoin J-M, Peroux T, Crivelli AJ, Olivier A, Acqueberge M, Le Brun M, Møller PR, Willerslev E, Dejean T (2015) Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Molecular Ecology* 25:929-942. doi: 10.1111/mec.13428.

Visco JA, Apothéloz-Perret-Gentil L, Cordonier A, Esling P, Pillet L, Pawlowski J (2015) Environmental monitoring: inferring the diatom index from next-generation sequencing data. *Environmental Science and Technology* 49:7597-7605. doi: 10.1021/es506158m.

Wangensteen OS, Palacín C, Guardiola M, Turon X (2018) DNA metabarcoding of littoral hard-bottom communities: high diversity and database gaps revealed by two molecular markers. *PeerJ* 6:e4705. doi: 10.7717/peerj.4705.

Weigand H, Beermann AJ, Čiampor F, Costa FO, Csabai Z, Duarte S, Geiger MF, Grabowski M, Rimet F, Rulik B, Strand M, Szucsich N, Weigand AM, Willassen E, Wyler SA, Bouchez A, Borja A, Čiamporová-Zaťovičová Z, Ferreira S, Dijkstra K, Eisendle U, Freyhof J, Gadawski P, Graf W, Haegerbaeumer A, Van der Hoorn BB, Japoshvili B, Keresztes L, Keskin E, Leese F, Macher J, Mamos T, Paz G, Pešić V, Pfannkuchen DM, Pfannkuchen MA, Price BW, Rinkevich B, Teixeira MAL, Várbbíró G, Ekrem T (2019) DNA barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-analysis and recommendations for future work. *bioRxiv*:576553. doi: 10.1101/576553.

Weigand AM, Macher JN (2018) A DNA metabarcoding protocol for hyporheic freshwater meiofauna: Evaluating highly degenerate COI primers and replication strategy. *Metabarcoding and Metagenomics* 2:e26869. doi: 10.3897/mbmg.2.26869.

Wetzel RG (2001) *Limnology: Lake and River Ecosystems*. San Diego, CA: Academic Press.

Wilcox TM, McKelvey KS, Young MK, Jane SE, Lowe WH, Whiteley AR, Schwartz MK (2013) Robust detection of rare species using environmental DNA: the importance of primer specificity. *PLoS ONE* 8:e59520. doi: 10.1371/journal.pone.0059520.

Wilcox TM, McKelvey KS, Young MK, Lowe WH, Schwartz MK (2015) Environmental DNA particle size distribution from Brook Trout (*Salvelinus fontinalis*). *Conservation Genetics Resources* 7:639-641. doi: 10.1007/s12686-015-0465-z.

References

- Willerslev E, Hansen AJ, Binladen J, Brand TB, Gilbert MTP, Shapiro B, Bunce M, Wiuf C, Gilichinsky DA, Cooper A (2003) Diverse plant and animal genetic records from holocene and pleistocene sediments. *Science* 300:791-795. doi: 10.1126/science.1084114.
- Williams N, Sweetman J (2019) Effects of neonicotinoids on the emergence and composition of chironomids in the Prairie Pothole Region. *Environmental Science and Pollution Research* 26:3862-3868. doi: 10.1007/s11356-018-3683-6.
- Wright-Stow AE, Winterbourn MJ (2003) How well do New Zealand's stream-monitoring indicators, the macroinvertebrate community index and its quantitative variant, correspond? *New Zealand Journal of Marine and Freshwater Research* 37:461-470. doi: 10.1080/00288330.2003.9517180.
- Wu Z, Cai Y, Liu X, Xu CP, Chen Y, Zhang L (2013) Temporal and spatial variability of phytoplankton in Lake Poyang: The largest freshwater lake in China. *Journal of Great Lakes Research* 39:476-483. doi: 10.1016/j.jglr.2013.06.008.
- WWF (2014) The Living Planet Report 2014. doi: 10.1016/j.edurev.2013.01.001.
- Yamamuro M, Komuro T, Kamiya H, Kato T, Hasegawa H, Kameda Y (2019) Neonicotinoids disrupt aquatic food webs and decrease fishery yields. *Science* 366:620-623. doi: 10.1126/science.aax3442.
- Zimmermann J, Jahn R, Gemeinholzer B (2011) Barcoding diatoms: Evaluation of the V4 subregion on the 18S rRNA gene, including new primers and protocols. *Organisms Diversity and Evolution* 11:173-192. doi: 10.1007/s13127-011-0050-6.
- Zimmermann J, Glöckner G, Jahn R, Enke N, Gemeinholzer B (2015) Metabarcoding vs. morphological identification to assess diatom diversity in environmental studies. *Molecular Ecology Resources* 15:526-42. doi: 10.1111/1755-0998.12336.
- Zinger L, Bonin A, Alsos IG, Bálint M, Bik H, Boyer F, Chariton AA, Creer S, Coissac E, Deagle BE, De Barba M, Dickie IA, Dumbrell AJ, Ficetola GF, Fierer N, Fumagalli L, Gilbert MTP, Jarman S, Jumpponen A, Kausarud H, Orlando L, Pansu J, Pawlowski J, Tedersoo L, Thomsen PF, Willerslev E, Taberlet P (2019) DNA metabarcoding - Need for robust experimental designs to draw sound ecological conclusions. *Molecular Ecology* 28:1857-1862. doi: 10.1111/mec.15060.
- Zizka VMA, Geiger MF, Leese F (2020) DNA metabarcoding of stream invertebrates reveals spatio-temporal variation but consistent status class assessments in a natural and urban river. *Ecological Indicators* 115. doi: 10.1016/j.ecolind.2020.106383.