



Universiteit  
Leiden  
The Netherlands

## Laser-generated toroidal helium plasmas

Kooij, V.L.

### Citation

Kooij, V. L. (2021, April 28). *Laser-generated toroidal helium plasmas*. *Casimir PhD Series*. Retrieved from <https://hdl.handle.net/1887/3161377>

Version: Publisher's Version

License: [Licence agreement concerning inclusion of doctoral thesis in the Institutional Repository of the University of Leiden](#)

Downloaded from: <https://hdl.handle.net/1887/3161377>

**Note:** To cite this publication please use the final published version (if applicable).

Cover Page



Universiteit Leiden



The handle <http://hdl.handle.net/1887/3161377> holds various files of this Leiden University dissertation.

**Author:** Kooij, V.L.

**Title:** Laser-generated toroidal helium plasmas

**Issue date:** 2021-04-28

## *Photographs of selected experimental set-ups*

*To give a more vivid impression of the research presented in this dissertation, we present photographs of selected experimental set-ups used for this research. References are given to the research in this dissertation to which these photos relate.*



Photo 1: The pulsed high power optical set-up used to generate the transient toroidal helium plasmas studied in this work. On the left, the Quanta-Ray GCR-3 Q-switched Nd:YAG laser is visible, capable of generating high power 10 ns laser pulses with a wavelength of 1064 nm and with energies up to 275 mJ. On the right, the Continuum NY61-10 laser is visible, similarly capable of generating laser pulses with energies up to 300 mJ. At the top of the photo, a circular aperture is visible in the black aluminium plate, through which the high power laser pulses are guided on their way into the plasma reactor. The dimensions of the breadboard on which the optical set-up has been built is 600 x 450 mm in size. This photo is referenced on page 15 and in figure 2.2.




Photo 2: The sub-microsecond temporal resolution, microwave interferometric set-up, used to measure the complex transmission coefficient of 57 GHz microwave radiation traversing a transient toroidal helium plasma. The dimensions of the breadboard on which the interferometer has been built is approximately 230 x 150 mm in size. The copper pyramidal horn antennas, between which the toroidal plasma will be generated, are visible to the right of the breadboard. The central component of the interferometer is the black four-port  $90^\circ$  3 dB directional coupler visible at the bottom of the breadboard. The microwave radiation is guided through rectangular TE<sub>10</sub> mode wave-guide with dimensions of 3.76 x 1.88 mm. During measurements, the whole interferometric set-up will be mounted inside the plasma reactor. This photo is referenced on page 69 and in figure 4.2.




Photo 3: The tailor-made 2.465 GHz iris coupled TE<sub>101</sub> mode  $\lambda_g/2$  rectangular microwave cavity during integration into the plasma reactor. The microwave cavity will be positioned in such a way that the anti-node of the electric field of the TE<sub>101</sub> mode coincides with the focal point of the lens creating the laser-induced breakdown plasmas from which the toroidal plasmas emerge. The brass stubs just visible to the left of the plasma reactor form the tailor-made three stub tuner. The aluminium beyond cut-off view ports used to let the laser pulses safely enter and exit the microwave cavity, and to image the toroidal plasma from the side, are visible on that part of the microwave cavity, that is already inside the plasma reactor. The microwave cavity goes straight through the left flange of the plasma reactor, and is vacuum sealed to ensure that experiments can be performed in a helium atmosphere. The guide wavelength for the rectangular wave-guide with internal dimensions of 90 x 40 mm is approximately 165 mm. This photo is referenced on page 88 and in figure 5.2.





# Bibliography

- Abel N. H. (1826). Auflösung einer mechanischen aufgabe. *Journal für die reine und angewandte Mathematik* 1, 153–157. DOI: [10/dtp6wf](https://doi.org/10/dtp6wf) (cited on p. 26).
- Abramowitz M. and Stegun I. (1972). *Handbook of mathematical functions*. Dover Publications (cited on p. 66).
- Akhmetov D. G. (2009). *Vortex rings*. Springer-Verlag Berlin Heidelberg (cited on p. 22).
- Alberti A., Munafò A., Koll M., Nishihara M., Pantano C., Freund J. B., Elliott G. S. and Panesi M. (2020). Laser-induced non-equilibrium plasma kernel dynamics. *Journal of Physics D: Applied Physics* 53, 025201. DOI: [10/dnkg](https://doi.org/10/dnkg) (cited on pp. 46, 48, 57).
- Alberti A., Munafò A., Pantano C., Freund J. B. and Panesi M. (2019a). Modelling of air breakdown by single-mode and multi-mode lasers. *AIAA Scitech 2019 Forum*, 1250. DOI: [10/dbtn](https://doi.org/10/dbtn) (cited on pp. 22, 46, 48, 57, 58).
- Alberti A., Munafò A., Pantano C., Freund J. B. and Panesi M. (2019b). Radiation supported plasma waves in non-equilibrium laser discharges. *2019 Working Group Radiation of High Temperature Gas, March 26th* (cited on pp. 46, 48, 57).
- Arfken G. B. and Weber H. J. (1995). *Mathematical methods for physicists*. 4th edition. Academic Press (cited on pp. 26, 27).
- Ashcroft N. W. and Mermin N. D. (1976). *Solid state physics*. Saunders College Publishing (cited on pp. 63, 65).
- Atkinson R. and Lissberger P. H. (1992). Sign conventions in magneto-optical calculations and measurements. *Applied Optics* 31, 6076–6081. DOI: [10/fgwh6m](https://doi.org/10/fgwh6m) (cited on p. 71).
- Bak M. S., Im S. and Cappelli M. A. (2014). Successive laser-induced breakdowns in atmospheric pressure air and premixed ethane-air mixtures. *Combustion and Flame* 161, 1744–1751. DOI: [10/f56s6n](https://doi.org/10/f56s6n) (cited on pp. 10, 14, 22, 45, 53).

- Bak M. S., Wermer L. and Im S. (2015). Schlieren imaging investigation of successive laser-induced breakdowns in atmospheric-pressure air. *Journal of Physics D: Applied Physics* 48, 485203. DOI: [10/dbg5](#) (cited on pp. [10](#), [14](#), [22](#), [23](#), [45](#), [53](#)).
- Bellan P. M. (2006). *Fundamentals of plasma physics*. Cambridge University Press (cited on p. [66](#)).
- Besant W. H. (1859). *A treatise on hydrostatics and hydrodynamics*. Deighton, Bell, and Co. (cited on p. [33](#)).
- Bethe H. A., Fuchs K., Hirschfelder J. O., Magee J. L., Peierls R. E. and Von Neumann J. (1958). *Blast wave*. Los Alamos Scientific Laboratory New Mexico (cited on p. [53](#)).
- Boivin R. F., Kline J. L. and Scime E. E. (2001). Electron temperature measurement by a helium line intensity ratio method in helicon plasmas. *Physics of Plasmas* 8, 5303. DOI: [10/b4mcnd](#) (cited on p. [99](#)).
- Bracewell R. N. (1956). Strip integration in radio astronomy. *Australian Journal of Physics* 9, 198–217. DOI: [10/fvr5xr](#) (cited on p. [27](#)).
- Bradley D., Sheppard C. G. W., Suardjaja I. M. and Woolley R. (2004). Fundamentals of high-energy spark ignition with lasers. *Combustion and Flame* 138, 55–77. DOI: [10/fqfq3b](#) (cited on pp. [10](#), [14](#), [22](#), [45](#), [54](#)).
- Broekaert J. A. C. and Siemens V. (2004). Recent trends in atomic spectrometry with microwave-induced plasmas. *Spectrochimica Acta Part B: Atomic Spectroscopy* 59, 1823–1839. DOI: [10/dz45kv](#) (cited on p. [88](#)).
- Candel S. M. (1981). An algorithm for the fourier-bessel transform. *Computer Physics Communications* 23, 343–353. DOI: [10/cdp6t5](#) (cited on p. [27](#)).
- Carbone E. A. D., Schregel C. G. and Czarnetzki U. (2016). Ignition and afterglow dynamics of a high pressure nanosecond pulsed helium micro-discharge: ii. rydberg molecules kinetics. *Plasma Sources Science and Technology* 25, 054004. DOI: [10/dmqd](#) (cited on pp. [21](#), [45](#)).
- Chanin L. M., Phelps A. V. and Biondi M. A. (1962). Measurements of the attachment of low-energy electrons to oxygen molecules. *Physical Review* 128, 219. DOI: [10/fjf4r7](#) (cited on pp. [42](#), [44](#), [45](#)).
- Chen F. F. (1974). *Introduction to plasma physics*. Plenum Press (cited on pp. [9](#), [10](#), [61](#), [63](#), [66](#), [67](#), [77](#), [81](#)).
- Chen Y., Lewis J. W. L. and Parigger C. (2000). Spatial and temporal profiles of pulsed laser-induced air plasma emissions. *Journal of Quantitative Spectroscopy and Radiative Transfer* 67, 91–103. DOI: [10/b95r8g](#) (cited on pp. [23](#), [33](#), [48](#)).
- Chorin A. J. and Marsden J. E. (1993). *A mathematical introduction to fluid mechanics*. 3rd edition. Springer-Verlag New York (cited on pp. [61–63](#)).



- Cioffi D. F. (1990). Supernova remnants as probes of the interstellar medium. *Physical Processes in Hot Cosmic Plasmas*, 1–16. DOI: [10/dvt2](https://doi.org/10/dvt2) (cited on pp. 53, 54).
- Collin R. E. (2001). *Foundations for microwave engineering*. 2nd edition. Wiley-IEEE Press (cited on p. 71).
- Cooper J. (1966). Plasma spectroscopy. *Reports on Progress in Physics* 29, 35–130. DOI: [10/fw3p5n](https://doi.org/10/fw3p5n) (cited on pp. 21, 26, 27, 29).
- Cristoforetti G., Legnaioli S., Pardini L., Palleschi V., Salvetti A. and Tognoni E. (2006). Spectroscopic and shadowgraphic analysis of laser induced plasmas in the orthogonal double pulse pre-ablation configuration. *Spectrochimica Acta Part B: Atomic Spectroscopy* 61, 340–350. DOI: [10/bcvs2j](https://doi.org/10/bcvs2j) (cited on pp. 54, 57).
- de Klerk A. J. J. M., van der Veen R. I., Dalhuisen J. W. and Bouwmeester D. (2017). Knotted optical vortices in exact solutions to maxwell's equations. *Physical Review A* 95, 053820. DOI: [10/fbfc](https://doi.org/10/fbfc) (cited on p. 10).
- Dribinski V., Ossadtchi A., Mandelshtam V. A. and Reisler H. (2002). Reconstruction of abel-transformable images: the gaussian basis-set expansion abel transform method. *Review of Scientific Instruments* 73, 2634–2642. DOI: [10/d7pm4v](https://doi.org/10/d7pm4v) (cited on pp. 26, 27).
- Drude P. (1900). Zur elektronentheorie der metalle. *Annalen der Physik* 306, 566. DOI: [10/cmck9z](https://doi.org/10/cmck9z) (cited on p. 63).
- Dumitrache C., VanOsdol R., Limbach C. M. and Yalin A. P. (2017). Control of early flame kernel growth by multi-wavelength laser pulses for enhanced ignition. *Scientific Reports* 7, 10239. DOI: [10/gbwm93](https://doi.org/10/gbwm93) (cited on pp. 10, 14, 22, 45).
- Fridman A. (2008). *Plasma chemistry*. Cambridge University Press (cited on p. 91).
- Gharib M., Mendoza S., Rosenfeld M., Beizai M. and Pereira F. J. A. (2017). Toroidal plasmoid generation via extreme hydrodynamic shear. *Proceedings of the National Academy of Sciences* 114, 12657–12662. DOI: [10/gcn523](https://doi.org/10/gcn523) (cited on pp. 10, 14).
- Ghosh S. and Mahesh K. (2008). Numerical simulation of the fluid dynamic effects of laser energy deposition in air. *Journal of Fluid Mechanics* 605, 329–354. DOI: [10/fkqmf7](https://doi.org/10/fkqmf7) (cited on pp. 22, 23, 39, 45, 48).
- Glasstone S. and Dolan P. J. (1977). *The effects of nuclear weapons*. 3rd edition. Department of Defense Washington D.C. (cited on p. 57).
- Goedbloed J. P. and Poedts S. (2004). *Principles of magnetohydrodynamics, with applications to laboratory and astrophysical plasmas*. Cambridge University Press (cited on p. 66).

- Griem H. R. (1997). *Principles of plasma spectroscopy*. Cambridge University Press (cited on pp. 27, 29, 97, 99).
- Griffiths D. J. (1999). *Introduction to electrodynamics*. 3rd edition. Prentice Hall (cited on p. 64).
- Gulec A., Bozduman F. and Hala A. M. (2015). Atmospheric pressure 2.45-ghz microwave helium plasma. *IEEE Transactions on Plasma Science* 43, 786–790. DOI: [10/f67rzc](https://doi.org/10/f67rzc) (cited on pp. 97, 99).
- Hammer M. R. (2008). A magnetically excited microwave plasma source for atomic emission spectroscopy with performance approaching that of the inductively coupled plasma. *Spectrochimica Acta Part B: Atomic Spectroscopy* 63, 456–464. DOI: [10/dx7qkc](https://doi.org/10/dx7qkc) (cited on p. 88).
- Hanson K. M. (1993). Special topics in test methodology: tomographic reconstruction of axially symmetric objects from a single dynamic radiograph. *Progress in Astronautics and Aeronautics* 155, 687–698 (cited on pp. 26, 29).
- Harilal S. S., Brumfield B. E. and Phillips M. C. (2015). Lifecycle of laser-produced air sparks. *Physics of Plasmas* 22, 063301. DOI: [10/f7jffs](https://doi.org/10/f7jffs) (cited on pp. 10, 14, 22, 23, 33, 45, 48).
- Haynes W. M. (2014). *CRC handbook of chemistry and physics*. 95th edition. CRC Press (cited on pp. 23, 32, 34, 45).
- Hutchinson I. H. (2002). *Principles of plasma diagnostics*. 2nd edition. Cambridge University Press (cited on pp. 21, 26, 27).
- IEA (2017). *World energy balances: data and statistics*. International Energy Agency, Paris (cited on p. 9).
- Irvine W. T. M. and Bouwmeester D. (2008). Linked and knotted beams of light. *Nature Physics* 4, 716–720. DOI: [10/d23t2b](https://doi.org/10/d23t2b) (cited on p. 10).
- Jackson J. D. (1998). *Classical electrodynamics*. 3rd edition. John Wiley & Sons (cited on p. 64).
- Jeans J. H. (1940). *An introduction to the kinetic theory of gases*. Cambridge University Press (cited on pp. 22, 32).
- Kabouzi Y., Calzada M. D., Moisan M., Tran K. C. and Trassy C. (2002). Radial contraction of microwave-sustained plasma columns at atmospheric pressure. *Journal of Applied Physics* 91, 1008. DOI: [10/ccpkj2](https://doi.org/10/ccpkj2) (cited on pp. 77, 81).
- Kamchatnov A. M. (1982). Topological solitons in magnetohydrodynamics. *Soviet Physics Journal of Experimental and Theoretical Physics* 55, 69–73 (cited on p. 10).
- Kedia H., Bialynicki-Birula I., Peralta-Salas D. and Irvine W. T. M. (2013). Tying knots in light fields. *Physical Review Letters* 111, 150404. DOI: [10/ggh6pb](https://doi.org/10/ggh6pb) (cited on p. 10).

- Kopp G. and Lean J. L. (2011). A new, lower value of total solar irradiance: evidence and climate significance. *Geophysical Research Letters* 38, L01706. DOI: [10/b9tkk2](https://doi.org/10.1029/2010GL014601) (cited on p. 9).
- Kramida A., Ralchenko Y., Reader J. and NIST ASD Team (2019). NIST atomic spectra database. *National Institute of Standards and Technology*. DOI: [10/dmk4](https://doi.org/10.18130/10.18130/10.18130) (cited on p. 21).
- Krehl P. (2011). Shock wave physics and detonation physics - a stimulus for the emergence of numerous new branches in science and engineering. *The European Physical Journal H* 36, 85–152. DOI: [10/bsdwtq](https://doi.org/10.1051/epjh/201136001) (cited on p. 57).
- Krehl P. and Van der Geest M. (1991). The discovery of the mach reflection effect and its demonstration in an auditorium. *Shock Waves* 1, 3–15. DOI: [10/bb23dx](https://doi.org/10.1007/BF01614800) (cited on p. 57).
- Kulsrud R. M. (2005). *Plasma physics for astrophysics*. Princeton University Press (cited on p. 66).
- Landau L. D. and Lifshitz E. M. (1987). *Fluid mechanics*. 2nd edition. Pergamon Press Oxford (cited on pp. 23, 34).
- Mach E. and Mach L. (1890). Ueber die interferenz der schallwellen von grosser excursion. *Annalen der Physik* 277, 140–143. DOI: [10/bvq5gs](https://doi.org/10.1007/BF02111400) (cited on p. 57).
- Marder M. P. (2010). *Condensed matter physics*. 2nd edition. John Wiley & Sons (cited on p. 66).
- Matusiewicz H. (1992). A novel microwave plasma cavity assembly for atomic emission spectrometry. *Spectrochimica Acta Part B: Atomic Spectroscopy* 47, 1221–1227. DOI: [10/dcmfcb](https://doi.org/10.1016/0883-2925(92)90000-0) (cited on p. 88).
- Miziolek A. W., Palleschi V. and Schechter I. (2006). *Laser induced breakdown spectroscopy*. Cambridge University Press (cited on pp. 23, 48).
- Moffatt H. K. (1969). The degree of knottedness of tangled vortex lines. *Journal of Fluid Mechanics* 35, 117–129. DOI: [10/fh75pf](https://doi.org/10.1017/jfm.1969.001) (cited on pp. 9, 13).
- Monfared S. K., Graham W. G., Morgan T. J. and Huwel L. (2011). Spectroscopic characterization of laser-produced atmospheric pressure helium microplasmas. *Plasma Sources Science and Technology* 20, 035001. DOI: [10/d8xg6g](https://doi.org/10.1088/0963-0252/20/3/035001) (cited on pp. 97, 99).
- Morsy M. H. and Chung S. H. (2002). Numerical simulation of front lobe formation in laser-induced spark ignition of ch<sub>4</sub>/air mixtures. *Proceedings of the Combustion Institute* 29, 1613–1619. DOI: [10/fmp2ht](https://doi.org/10.1016/S0085-4266(02)00000-0) (cited on pp. 22, 39, 45).
- Muller R. H. (1969). Definitions and conventions in ellipsometry. *Surface Science* 16, 14–33. DOI: [10/c3674v](https://doi.org/10.1016/0039-6028(69)90000-0) (cited on p. 71).

- Nassif D. and Hüwel L. (2000). Appearance of toroidal structure in dissipating laser-generated sparks. *Journal of Applied Physics* 87, 2127–2130. DOI: [10/dp3n7c](#) (cited on pp. 10, 14, 22, 23, 45).
- Nedanovska E., Nersisyan G., Morgan T. J., Huwel L., Murakami T., Lewis C. L. S., Riley D. and Graham W. G. (2015). Investigating the dynamics of laser induced sparks in atmospheric helium using rayleigh and thomson scattering. *Journal of Applied Physics* 117, 013302. DOI: [10/fj4v](#) (cited on pp. 45, 60, 82).
- Nishihara M., Freund J. B., Glumac N. G. and Elliott G. S. (2018). Influence of mode-beating pulse on laser-induced plasma. *Journal of Physics D: Applied Physics* 51, 135601. DOI: [10/dbmd](#) (cited on pp. 22, 46).
- Nyswander R. E. and Auger G. J. (1976). Nanosecond pulse generators for magnetron operation. *Naval Weapons Center China Lake CA*, ADA636968 (cited on p. 87).
- Ohno N., Razzak M. A., Ukai H., Takamura S. and Uesugi Y. (2006). Validity of electron temperature measurement by using boltzmann plot method in radio frequency inductive discharge in the atmospheric pressure range. *Plasma and Fusion Research* 1, 028. DOI: [10/dm7xwq](#) (cited on p. 99).
- Pokrzywka B., Mendys A., Dzierzega K., Grabiec M. and Pellerin S. (2012). Laser light scattering in a laser-induced argon plasma: investigations of the shock wave. *Spectrochimica Acta Part B: Atomic Spectroscopy* 74–75, 24–30. DOI: [10/fkxc](#) (cited on p. 60).
- Pretzler G. (1991). A new method for numerical abel-inversion. *Zeitschrift für Naturforschung A* 46, 639–641. DOI: [10/dmzs](#) (cited on p. 27).
- Pretzler G., Jäger H., Neger T., Philipp H. and Woisetschläger J. (1992). Comparison of different methods of abel inversion using computer simulated and experimental side-on data. *Zeitschrift für Naturforschung A* 47, 955–970. DOI: [10/gfz8fs](#) (cited on pp. 26, 27, 30).
- Quintero M. C., Rodero A., Garcia M. C. and Sola A. (1997). Determination of the excitation temperature in a nonthermodynamic-equilibrium high-pressure helium microwave plasma torch. *Applied Spectroscopy* 51, 778–784. DOI: [10/d3qzqb](#) (cited on p. 99).
- Raizer Y. P. (1991). *Gas discharge physics*. Springer-Verlag Berlin Heidelberg (cited on pp. 23, 42, 44, 48).
- Rañada A. F. (1989). A topological theory of the electromagnetic field. *Letters in Mathematical Physics* 18, 97–106. DOI: [10/b3c4jr](#) (cited on p. 10).

- Rayleigh L. (1917). On the pressure developed in a liquid during the collapse of a spherical cavity. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* 6, 94–98. DOI: [10/fwxkvm](https://doi.org/10/fwxkvm) (cited on p. 33).
- Richardson A. S. (2019). *NRL Plasma Formulary*. Naval Research Laboratory, Washington, DC (cited on pp. 77, 81, 82).
- Rybicki G. B. and Lightman A. P. (1979). *Radiative processes in astrophysics*. John Wiley & Sons (cited on p. 21).
- Schregel C. G., Carbone E. A. D., Luggenholscher D. and Czarnetzki U. (2016). Ignition and afterglow dynamics of a high pressure nanosecond pulsed helium micro-discharge: i. electron, rydberg molecules and he (23s) densities. *Plasma Sources Science and Technology* 25, 054003. DOI: [10/dmqc](https://doi.org/10/dmqc) (cited on pp. 21, 45).
- Sedov L. I. (1959). *Similarity and dimensional methods in mechanics*. Academic Press (cited on pp. 53, 54).
- Settles G. S. (2001). *Schlieren and shadowgraph techniques*. Springer-Verlag Berlin Heidelberg (cited on pp. 18, 51, 55).
- Seward C. (2014). Ball lightning events explained as self-stable spinning high-density plasma toroids or atmospheric spheromaks. *IEEE Access* 2, 153–159. DOI: [10/dkxt](https://doi.org/10/dkxt) (cited on pp. 10, 14).
- Shu F. H. (1992). *The physics of astrophysics: gas dynamics*. University Science Books (cited on p. 53).
- Sinibaldi G., Occhicone A., Alves Pereira F., Caprini D., Marino L., Michelotti F. and Casciola C. M. (2019). Laser induced cavitation: plasma generation and breakdown shockwave. *Physics of Fluids* 31, 103302. DOI: [10/dpvz](https://doi.org/10/dpvz) (cited on p. 33).
- Smerlak M. (2011). A blackbody is not a blackbox. *European Journal of Physics* 32, 1143. DOI: [10/bcznd2](https://doi.org/10/bcznd2) (cited on p. 21).
- Smiet C. B. (2017). *Knots in plasma*. Casimir PhD series Leiden Delft (cited on pp. 10, 59).
- Smiet C. B., Candelaresi S., Thompson A., Swearngin J., Dalhuisen J. W. and Bouwmeester D. (2015). Self-organizing knotted magnetic structures in plasma. *Physical Review Letters* 115, 095001. DOI: [10/dkts](https://doi.org/10/dkts) (cited on pp. 10, 11, 14).
- Smiet C. B., de Blank H. J., de Jong T. A., Kok D. N. L. and Bouwmeester D. (2019). Resistive evolution of toroidal field distributions and their relation to magnetic clouds. *Journal of Plasma Physics* 85, 905850107. DOI: [10/fbwt](https://doi.org/10/fbwt) (cited on pp. 10, 59).
- Smith L. M., Keefer D. R. and Sudharsanan S. I. (1988). Abel inversion using transform techniques. *Journal of Quantitative Spectroscopy and Radiative Transfer* 39, 367–373. DOI: [10/fhzgsq](https://doi.org/10/fhzgsq) (cited on pp. 26, 27).

- Spanel P., Hall E. F. H., Workman C. T. and Smith D. (2004). A directly coupled monolithic rectangular resonator forming a robust microwave plasma ion source for sift-ms. *Plasma Sources Science and Technology* 13, 282–284. DOI: [10/bsphgx](#) (cited on p. 88).
- Stepanyan S., Minesi N., Tibère-Inglesse A., Salmon A., Stancu G. D. and Laux C. O. (2019). Spatial evolution of the plasma kernel produced by nanosecond discharges in air. *Journal of Physics D: Applied Physics* 52, 295203. DOI: [10/dkxs](#) (cited on pp. 10, 14).
- Stevelfelt J. and Johansson J. (1971). Measurements of the spectral light emission from decaying high pressure helium plasmas. *AB Atomenergi, Nykoeping, Sweden, AE-418* (cited on p. 45).
- Tahir I., Dexter A. and Carter R. (2006). Frequency and phase modulation performance of an injection-locked cw magnetron. *IEEE Transactions on Electron Devices* 53, 1721–1729. DOI: [10/c52h72](#) (cited on p. 96).
- Taylor G. I. (1950a). The formation of a blast wave by a very intense explosion i. theoretical discussion. *The Proceedings of the Royal Society of London A* 201, 159–174. DOI: [10/fj2s7h](#) (cited on p. 53).
- Taylor G. I. (1950b). The formation of a blast wave by a very intense explosion. ii. the atomic explosion of 1945. *The Proceedings of the Royal Society of London A* 201, 175–186. DOI: [10/b7v6cp](#) (cited on p. 53).
- Taylor J. B. (1974). Relaxation of toroidal plasma and generation of reverse magnetic fields. *Physical Review Letters* 33, 1139–1141. DOI: [10/cnczw2](#) (cited on p. 10).
- Thompson A., Swearingin J. and Bouwmeester D. (2014). Linked and knotted gravitational radiation. *Journal of Physics A: Mathematical and Theoretical* 47, 355205. DOI: [10/fbfd](#) (cited on p. 10).
- Tonks L. and Langmuir I. (1929). Oscillations in ionized gases. *Physical Review* 33, 195. DOI: [10/bjfr8n](#) (cited on p. 61).
- Wagner W. G. and Lengyel B. A. (1963). Evolution of the giant pulse in a laser. *Journal of Applied Physics* 34, 2040. DOI: [10/brrss4](#) (cited on p. 48).
- Waldron R. A. (1960). Perturbation theory of resonant cavities. *Proceedings of the IEE - Part C: Monographs* 107, 272–274. DOI: [10/d6khgq](#) (cited on p. 96).
- Wesson J. (2004). *Tokamaks*. 3rd edition. Clarendon Press Oxford (cited on p. 61).
- Williams T. C. and Shaddix C. R. (2007). Simultaneous correction of flat field and nonlinearity response of intensified charge-coupled devices. *Review of Scientific Instruments* 78, 123702. DOI: [10/ft2r97](#) (cited on p. 17).



- Woltjer L. (1958). A theorem on force-free magnetic fields. *Proceedings of the National Academy of Sciences* 44, 489–491. DOI: [10/btg2w8](https://doi.org/10/btg2w8) (cited on pp. 9, 13).
- Ye Y., Spina D., Xing Y., Bogaerts W. and Dhaene T. (2018). Numerical modeling of a linear photonic system for accurate and efficient time-domain simulations. *Photonics Research* 6, 560–573. DOI: [10 / d3hz](https://doi.org/10/d3hz) (cited on p. 71).
- Zel'dovich Y. B. and Raizer Y. P. (1966). *Physics of shock waves and high-temperature hydrodynamic phenomena*. Academic Press (cited on pp. 53, 54).

